



DVT RISK REASSESSMENT

SR No	PARAMETERS	Date	4/16	22/16	23/16	24/16	25/16
		Time	9 AM	9 AM	5 PM	2 PM	9 AM
		Score	2	2	2	2	2
1	Active cancer (on-going treatment or diagnosed within 6 months or palliative care)	0/1	0	0	0	0	0
2	Bedridden recently >3 days or major surgery within four weeks.	0/1	1	1	1	1	1
3	Calf swelling >3 cm compared with asymptomatic side, measured at 10 cm below tibial tubercle (Assess for both legs)	0/1	0	0	0	0	0
4	Collateral (non varicose) superficial veins present (Assess for both legs)	0/1	0	0	0	0	0
5	Entire leg swelling (Assess for both legs)	0/1	0	0	0	0	0
6	Localized tenderness along the deep venous system (Assess for both legs)	0/1	0	0	0	0	0
7	Pitting edema, greater in the symptomatic leg (Assess for both legs)	0/1	0	0	0	0	0
8	Paralysis, paresis, or recent plaster immobilization of the lower extremity (Assess for both legs)	0/1	1	1	1	1	1
9	Previously documented DVT (Assess for both legs)	0/1	0	0	0	0	0
10	Alternative diagnosis to DVT as likely or more likely (Assess for both legs)/Co-morbidity like ESLD / Renal disease, Renal Failure, CCF Cellulitis (Commonly mistaken as DVT), Dependent (stasis) oedema, Lymphatic obstruction, Septic arthritis, cirrhosis, Nephrotic syndrome, Calf muscle tear or strain, Hematoma (collection of blood) in the muscle, Sprain or rupture of a leg tendon, Fracture	0/-2	0	0	0	0	0
FINAL SCORE			2	2	2	2	2
Interpretation of score			Med	Med	Med	Med	Med
DVT Prophylaxis activated			Yes	Yes	✓	—	✓
Initials & Emp. ID			R 2/CD	R 3/CD	Gh	JL	DR
DVT Prophylaxis: For Low risk patient (-2 to 0) Ambulation Moderate Risk : (1 to 2) Pharmacological prophylaxis / Mechanical Prophylaxis High risk patients : (3 to 7) Pharmacological prophylaxis along with mechanical Prophylaxis		Mechanical Prophylaxis - Intermittent Pneumatic compressions (IPC) Pharmacological Prophylaxis - Low-Molecular weight heparin Unfractionated Heparin Oral anticoagulant					