



Time	Patient Specific Nursing Needs	Measurable Goal	Nursing Interventions	M	E	N	Evaluation / Outcome	Time Date Name & ID
8AM 3PM 10L	<p>URINARY</p> <p>ELIMINATION</p> <ul style="list-style-type: none"> <input type="checkbox"/> Self voiding in Bathroom <input type="checkbox"/> Bedpan, Urinal <input type="checkbox"/> Condom Catheter <input checked="" type="checkbox"/> Foley's Catheter <p>BOWEL</p> <p>ELIMINATION</p> <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Bowel opened normally <input type="checkbox"/> Constipation <input type="checkbox"/> Diarrhoea <input type="checkbox"/> Colostomy <p>CHEST DRAINS</p> <p>Others _____</p>	<input type="checkbox"/> Patient will have normal elimination pattern	<ul style="list-style-type: none"> ● Assess for elimination pattern ● Maintain I/O chart ● Encourage fluid intake ● Encourage early ambulation ● Condom catheter is applied by measuring appropriate size (in case of condom catheter) ● Catheter care is given every shift with 0.9% NS ● Securing of condom/foleys is done by appropriate adhesive ● Urobag/urometer is kept below the hip level ● Administer laxatives / suppository enema as per orders ● Clean perineal area with cotton and water ● Change colostomy bag as prescribed. 				<input type="checkbox"/> Voided Freely <input type="checkbox"/> No constipation <input type="checkbox"/> Balanced I and O Record <input type="checkbox"/> _____ <input type="checkbox"/> _____	<i>Jan 2013</i> <i>3pm</i> <i>10L</i>
8AM 3PM 10L	<p>MOBILITY</p> <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Bedfast <input type="checkbox"/> Bedside mobility <input type="checkbox"/> Walks with assistance <input type="checkbox"/> Ambulatory <input type="checkbox"/> Others _____ 	<input type="checkbox"/> Patient will mobilized as advised and tolerated <input type="checkbox"/> Patients will be given passive exercises	<ul style="list-style-type: none"> ● Assess the level of activity that is tolerated ● Encourage passive exercises ● Apply splints/ support to prevent foot drop ● Encourage range of motion exercises ● Encourage regular ambulation ● Use Sequential Compression Devices to prevent DVT 				<input type="checkbox"/> Patient has ambulated without/with assistance. <input type="checkbox"/> Patient is given passive exercises	<i>Jan 2013</i> <i>3pm</i> <i>10L</i>