Time	Patient Specific Nursing Needs	Measurable Goal	Nursing Interventions	M	E	N	Evaluation / Outcome	Time Date Name & ID
8 2pm 10pm	**URINARY ELIMINATION** ☐ Self voiding in Bathroom ☐ Bedpan, Urinal ☐ Condom Catheter ☑ Foley's Catheter **BOWEL ELIMINATION** ☑ Bowel opened normally ☐ Constipation ☐ Diarrhoea ☐ Colostomy CHEST DRAINS Others_____	☐ Patient will have normal elimination pattern	● Assess for elimination pattern ● Maintain I/O chart ● Encourage fluid intake ● Encourage early ambulation ● Condom catheter is applied by measuring appropriate size (in case of condom catheter) ● Catheter care is given every shift with 0.9% NS ● Securing of condom/foleys is done by appropriate adhesive ● Urobag/urometer is kept below the hip level ● Administer laxatives / suppository enema as per orders ● Clean perineal area with cotton and water ● Change colostomy bag as prescribed.	☐ ☑ ☐ ☐ ☐ ☑ ☐ ☐ ☐ ☐ ☐	☑ ☑ ☐ ☐ ☑ ☑ ☐ ☑ ☐ ☑ ☐	☑ ☑ ☐ ☐ ☑ ☑ ☑ ☑ ☐ ☑ ☐	☐ Voided Freely ☐ No constipation ☑ Balanced I and O Record ☐ _____ ☐ _____	N3548 N3548 N3524 Jam
8 2pm 10pm	**MOBILITY** ☑ Bedfast ☐ Bedside mobility ☐ Walks with assistance ☐ Ambulatory ☐ Others _____	☑ Patient will mobilized as adviced and tolerated ☐ Patients will be given passive exercises	● Assess the level of activity that is tolerated ● Encourage passive exercises ● Apply splints/ support to prevent foot drop ● Encourage range of motion exercises ● Encourage regular ambulation ● Use Sequential Compression Devices to prevent DVT	☐ ☐ ☐ ☐ ☐ ☑	☐ ☑ ☑ ☐ ☐ ☑	☐ ☑ ☑ ☐ ☐ ☑	☐ Patient has ambulated without/with assistance. ☑ Patient is given passive exercises	N3548 N3524 Jam