

# A Study on The Effectiveness of Lock-down Measures to Control The Spread of COVID-19

Subhas Kumar Ghosh<sup>a,\*</sup>, Sachchit Ghosh<sup>b</sup>, Sai Shanmukha Narumanchi<sup>c</sup>

<sup>a</sup>*Commonwealth Bank of Australia, Sydney, New South Wales, 2000, Australia*

<sup>b</sup>*The University of Sydney, Camperdown, NSW 2006, Australia*

<sup>c</sup>*Department of Computer Science, Southern Illinois University, Carbondale, IL 62901, USA.*

---

## Abstract

The ongoing pandemic of coronavirus disease 2019-2020 (COVID-19), caused by Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). A pathogenic virus SARS-CoV-2 is able to spread asymptotically during its incubation stage through a vulnerable population. Given the state of health-care, policymakers were to contain the spread of infection, minimize stress on the health systems and ensure public safety. Taking measures to reduce stress on the health sector and diminishing the number of infected patients is important to end the pandemic. Most effective tool that was at disposal is to close non-essential business and issue stay home order. In this paper we consider techniques and measure the effectiveness of stringency measures adopted by governments across the world. Analyzing effectiveness of control measures like lock-down allows us to understand whether the decisions made were optimal and resulted in a reduction of burden on the healthcare system. In specific we consider using synthetic control to construct alternative scenarios and understand what would have been the effect on health if a less stringent measures were to adopt. We present analysis for State of New York, United States, Italy and Indian capital city Delhi and show how lock-down measures has helped and what counterfactual scenarios would have been in comparison to the current state of affairs. We show that in New York number of deaths could have been x times higher, and Italy the number of confirmed

---

\*Corresponding author

*Email addresses:* `subhas.ghosh@cba.com.au` (Subhas Kumar Ghosh),  
`sgho2841@uni.sydney.edu.au` (Sachchit Ghosh),  
`saishanmukha.narumanchi@cs.siu.edu` (Sai Shanmukha Narumanchi)

case could have been  $x$  and number of deaths could have been  $y$  by 26th of June, 2020.

*Keywords:* COVID-19, Lock down, Mathematical modeling, Epidemic, Economic Impact

---

## 1. Introduction

In December 2019, an outbreak occurred in Wuhan, China involving a zoonotic coronavirus, similar to SARS coronavirus and MERS coronavirus [1]. The virus has been named Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), and the disease caused by the virus has been named the coronavirus disease 2019 (COVID-19). Since then the ongoing pandemic has infected more than 9 million people and has caused more than 467 thousand deaths worldwide.

Since the initial outbreak, several different studies have tried to estimate the number of infections [2] that stem from a single infected patient in order to predict the potential for transmission of the COVID-19 virus. In most cases, it was seen that  $R_0 > 1$ , implying exponential growth through infection of a vulnerable population. Original estimates placed mortality rates for individuals at high risk at 4.46 % with those suffering from cardiovascular or kidney disease having even greater susceptibility [3].

The SARS-CoV-2 virus has no available treatment as the pathways for proliferation and pathogenesis are still unclear [4]. Current treatments are based on those effective on strains of the previous SARS coronavirus and MERS coronavirus. The SARS-CoV-2 virus is able to replicate rapidly during the asymptomatic phase and affect the lungs and respiratory tract, resulting in pneumonia, hypoxia, and acute respiratory distress [5]. Infected patients are directly dependent on external ventilation in most cases.

With the increasing pressure on the health systems due to reliance on intensive care units or non-invasive ventilation, health strategies like social distancing were implemented. The concern was to ensure the number of infected patients does not exceed the health system's ability to cope with it. It also focused on increasing the capacities of available health systems at the time.

Under the conditions at the time, with a highly pathogenic SARS-CoV-2 that is able to spread asymptotically during its incubation stage through a vulnerable population, policymakers were to contain the spread of the

infection, minimize stress on the health systems and ensure public safety. This was done by issuing orders for widespread lock-down and implementing social distancing measures. All non-essential businesses and services were shut down until further notice.

Taking measures to reduce stress on the health sector and diminishing the number of infected patients is important to end the pandemic, and understanding the effectiveness of a lock-down enables the distinction of good safety measures from bad ones. Analyzing effectiveness of control measures like lock-down allows us to understand whether the decisions made were optimal and resulted in a reduction of burden on the healthcare system, and broke chain of transmission, preventing its spread and reducing the reproductive rate of the virus.

Any optimal policy considers a trade off between the benefit associated with lock-down and cost of reduced aggregate output. Aggregate output decreases as a function of the stringency of the policy, commitment from the government to maintain the level of stringency and adherence of general population. Aggregate output decreases through lower supply of labor, lower consumption and hence through lower investment, which results from investors' expectation of a lower marginal product of capital. On the other hand benefit associated with lock-down can be seen through number of life potentially saved and in curbing the pandemic early so that economic activity can be restarted early.

Our objective in this work is to understand the benefits obtained from stringency measures adopted by governments across the world in terms of its health benefits. In the remaining of this section we describe our contribution and related works. Subsequently in Section 2 we provide a brief overview and mathematical underpinning of the tools that we use. In Section 3 we describe our data driven methodology and in Section 4 we present our results in three different geographic setup. Finally, in Section 5 we present some concluding remarks.

### *1.1. Our contribution*

In this work we consider stringency measures adopted by governments across the world and provide a counterfactual assessment of the benefit from those measures in terms of health benefits. In order to estimate the counterfactual metric (say number of deaths), we use a geographic location as a treatment unit (say Italy) and a set of other geographic location as donor group (say Brazil and United States). We take a data driven approach to

construct a synthetic control [11, 12, 13] using pre-innervation period data (say from early February, 2020 to March 9, 2020) of the donor units and their linear combination, such that the squared error between the estimated synthetic control and the treatment unit is minimized by the choice of weight parameters in this pre-innervation time period. Now synthetic control can be extrapolated to estimate the metric. We use Multi-dimensional Robust Synthetic Control (m-RSC) as a tool as described in [13].

However, there are few difficulties in applying the tool as it is. Firstly, different government adopted level of stringency measure and there were different level of compliance, and commitment. Secondly, there were no ‘pure’ donor group as stringency measures were nearly ubiquitous. So we have used various secondary sources of data to score the level of stringency measures , and level of compliance. This allowed us to determine donor groups relative to a choice of treatment unit.

Finally, we present our results in three different geographic setup - namely in the State of New York, Italy and Indian city of Delhi, and analyze them.

### *1.2. Related Works*

In [6], authors have shown that the reproductive rate of the SARS-CoV-2 had significantly decreased after government intervention. They show that the spread of disease is confined if measures were brought into effect early. In [7], authors use the differential timing of the introduction of stringency measures and changes in Google search for unemployment claims to establish a framework to estimate how each stringency measures contribute to unemployment. Authors show that early intervention efforts in the form of non-essential business closure have contributed to less than 8.5 percent of unemployment claims.

Another facet to measure the success of the lock-down is to observe its effects on the health systems. Late intervention in the case of Italy led to the flooding of hospitals and ICUs due to exponential spread. However, the national lock-down was effective in reducing the proliferation and decreased the stress on the national health system as observed by authors in [8]. In their paper [9], authors extend SIR model to include auxiliary state variables in the form of hospital capacity, contact with an infected person, etc. They use a system dynamics model of the outbreak to simulate various lock-down scenarios with recommendations for optimal strategy. In our work we consider possible outcome if such strategies were not adopted, and present counterfactual scenarios.

International travel has also been impacted as a result of efforts to reduce the spread of the coronavirus disease. On the basis of reported cases, models built by [10] show a significant decrease in the number of infections compared to predictions if no travel bans were adopted as an option. Their modeling results indicate that travel restriction must be combined with a transmission within the community to curb the spread.

## 2. Tools

As stated above, our objective is to study the effects of government response at an aggregate level in terms of life saved, and limiting the number of cases that requires hospitalization. Such interventions can effectively be studied at a comparative level. In other words, if we have data for the evolution of aggregate outcomes, e.g. number of confirmed cases and deaths, when policy is applied in a group under study versus when the same policy is not applied in a control group. However, government policies were applied at different level across a geographic region. We do not have a mechanism to conduct a randomized trial. Hence, we consider constructing synthetic control method[11, 12, 13]. In a synthetic control set up, where observational data is available for different groups, we can construct a synthetic or virtual control group by combining measurements from alternatives (or donors). In following we provide a brief overview of m-RSC [13].

Suppose that observations from  $N$  different geographically distinct groups or units are indexed by  $i \in [N]$  in  $T$  time periods (days) indexed by  $j \in [T]$ . Let  $k \in [K]$  be the metrics of interest (e.g. number of confirmed cases, number of deceased, number of tests conducted, etc.). By  $M_{ijk}$  we denote the ground-truth measurement of interest, and by  $X_{ijk}$ , an observation of this measurement with some noise. Let  $1 \leq T_0 \leq T$  be the time instance in which our group of interest experiences an intervention, namely a government response to control the spread (e.g. stay home order, school or business closure, or mass vaccination). Without loss of generality we consider unit  $i = 1$  (say, New York) and metric  $k = 1$  (say, number of deaths) as our unit and metric of interest respectively.

Our objective now is to estimate the trajectory of metric of interest  $k = 1$  for unit  $i = 1$  if no government response to control the spread had occurred. In order to do that we will use the trajectory associated with the donor units ( $2 \leq i \leq N$ ), and metrics  $k \in [K]$ . In following we make two assumptions: (1) for all  $2 \leq i \leq N$ ,  $k \in [K]$  and  $j \in [T]$ , we have  $X_{ijk} = M_{ijk} + \epsilon_{ijk}$




Figure 1: The Singular Value spectrum for all countries (of dimensions  $185 \times 150$ , showing top 20 singular values, in descending order.

where,  $\epsilon_{ijk}$  is the observational noise, and (2) Same model is obeyed by  $i = 1$  in pre-intervention period, i.e. for all  $j \in [T_0]$  and  $k \in [K]$  we have  $X_{1jk} = M_{1jk} + \epsilon_{1jk}$ . As described by authors in [13], in following we also assume that for unit  $i = 1$ , we only observe the measurement  $X_{1jk}$  for pre-intervention period, i.e. for all  $j \in [T_0]$  and  $k \in [K]$ . Our objective is to compute a counterfactual sequence of observation  $M_{1jk}$  for the time period  $j \in [T]$ , and  $k \in [K]$ , and in specific for  $T_0 \leq j \leq T$ , and  $k = 1$ , using synthetic version of unit  $i = 1$ .

Define  $\mathcal{M} = [M_{ijk}] \in \mathbb{R}^{N \times T \times K}$ .  $\mathcal{M}$  is assumed to have a few well behaved properties as required by the algorithm, namely, it must be approximately low-rank and boundedness of  $|M_{ijk}|$  (for details see [13]). To check whether our model assumption holds in practice, we consider  $N = 185$ ,  $T = 150$ ,  $K = 2$ , with 185 countries as units. We consider number of confirmed cases and number of deceased as two metrics over 150 days between January 22, 2020 and June 20, 2020. For assumption to hold, data matrix corresponding to number of confirmed cases and number of deceased and their combination should be approximated by a low-rank matrix.

As shown in Figure 1, the spectrum of the top 20 singular values (sorted

in descending order) for each matrix. The plots clearly support the implications that most of the spectrum is concentrated within the top 5 principal components. Same conclusion holds true when units are states of United States, and when we consider only countries in European Union.

Let  $\mathcal{Z} \in \mathbb{R}^{(N-1) \times T \times K}$  corresponding to donor units, and  $X_1 \in \mathbb{R}^{1 \times T_0 \times K}$  correspond to unit under intervention. We obtain  $\hat{\mathcal{M}}$  from  $\mathcal{Z}$  after applying a hard singular value thresholding. Subsequently, weights are learned using linear regression by computing

$$\hat{\beta} = \arg \min_{v \in \mathbb{R}^{(N-1)}} \left\| X_1 - v^T \hat{\mathcal{M}}_{T_0} \right\|_2^2$$

For every  $k \in [K]$ , the corresponding estimated counterfactual means for the treatment unit is then defined as

$$\hat{\mathcal{M}}_1^{(k)} = \hat{\beta}^T \hat{\mathcal{M}}^{(k)}$$

### 3. Methods

#### 3.1. Overview

As described in Section 1, we use m-RSC to construct a synthetic control for the treatment unit using data from multiple control units or donor group using pre-intervention period data. The synthetic control is then used for estimating the counterfactual in the post-intervention period. In our setup, intervention date is typically the date when a stay-home order or lock-down was declared for the treatment unit. However, government policy may have been applied over time with different levels of stringency measures.

To understand this we use stringency and policy indices data from Ox-CGRT [14], which records the strictness of policies that restrict people's behavior and includes 8 different measures - e.g. school and workplace closure, cancellation of public events, restrictions on gathering size etc. Figure 2, shows the plot of stringency index, with mobility data.

It can be observed that the level of lock-down varies over time and geographic region. We use this information in two different ways. First we choose the maximum level of index, first increased level of restriction index, and 14 days after the maximum level of index as various intervention dates and compare their effects. Second, we combine this information with mobility data to select the control groups for a treatment unit as discussed below.



Figure 2: Stringency index, and Mobility Trends in (a) Italy (b) Germany (c) Sweden (d) Spain (e) France.



In order to understand the effect of stringency on a treatment unit for a metric, we need to select a donor group where level of stringency was different or adherence to stringency was different. Since, there was a degree of stringency measures and adherence to such measures at different level - under any possible choice of donor group, we acknowledge, that we will be underestimating the counterfactual - i.e. what would have been without any stringency measures. To estimate the degree of adherence to lock-down measures, we use mobility data from Apple, Google and Facebook. Apple mobility data provides a relative volume of directions requests per region, sub-region or city compared to a baseline volume - i.e. percentage change over time from the baseline including weekly seasonality. Facebook data provides the relative percentage of population that is staying in the same place and also the percentage of population that moved from a region to another. In Facebook data, to quantify how much people move around measure is derived by counting the number of level-16 Bing tiles (which are approximately 600 meters by 600 meters in area at the equator) they are seen in within a day. Assuming  $U_{d,r}$  is the set of eligible users in region  $r$  on day  $d$ , and  $tiles(u)$  is the number of tiles visited by a given user  $u$  in  $U_{d,r}$ , total number of tiles visited for that region is given by  $totaltiles(U_{d,r}) = \sum_{u \in U_{d,r}} \min(tiles(u), 200)$ . Change in Movement measure is then the difference between a baseline and value on day  $d$  for average number of  $totaltiles(U_{d,r})$ . Similarly, Stay Put metric is calculated as the percentage of eligible people who are only observed in a single level-16 Bing tile during the course of a day on an average compared to a baseline. Finally, Google Community Mobility Report provides a percent change in visits to places like grocery stores and parks within a geographic area from baseline.

It can be seen from Figure 2, that the adherence and stringency level do not correspond. For example, in Sweden, with increasing level of government measures between March - April, there has not been any significant change in proportion of people staying put or moving between regions. Similarly, in other places, it can be observed that while government measures remain at the same level over April, number of people staying put at one place starts declining.

Selecting donor group: We combine metrics that allow the spread of the virus and similarly combine those that reduces the possibility by taking average of them and come with a single adherence score. For unit  $i \in [N]$ ,  $\forall j \neq i : j \in [N]$ ,  $j$  is a donor unit if adherence score, stringency Index of  $j$  is less than  $i$  and corresponding metric is higher (i.e. total number of deaths




Figure 3: Adherence score vs. Stringency index where size of each circle is determined by (a) total number of deaths till date (b) total number of confirmed cases till date.

till date, or total number of confirmed cases till date, respectively). Figure 3 shows this relation in graphical form for a selected set of countries. As per Figure 3, United States, Brazil and United Kingdom are in the donor group when we compute counterfactual estimate for Italy and metric under consideration is total number of confirmed cases till date.

**Statistical Performance Evaluation:** in Figure 4 we present the distribution of Mean and Median absolute percentage error statistic for the runs-forecasts from m-RSC algorithm with changing forecast horizon. We consider every Mondays between March 1, 2020 to June 21, 2020, both included to forecast the number of deaths at the end of the day on June 26, 2020 for all states in United States, and compare with actual data. For every state, donor group is selected using the method described above.

It can be seen from Figure 4, that both Mean and Median absolute percentage error statistic are larger, when larger forecast horizon has been con-




Figure 4: MAPE

sidered, and that is to be expected. Both MAPE and MdAPE converge to a less than 20 percent when horizon is about a week. MAPE being very large from MdAPE clearly indicates a right skew in the predicted values. This exactly corresponds to the places where exponential growth of the pandemic can be observed by March 22, 2020. The counterfactual prediction for June 26, 2020 on March 22, 2020 for these few places are several times higher as as expected, while, stricter stringency measures were being implemented in United States around those dates.

### 3.2. Data Source

We use following five sources of data as described in Table 1:

## 4. Results

In this section we present three examples of the application of m-RSC to derive the counterfactual estimates of possible number of deaths under the changed conditions like delaying or starting the stringency measures at earlier date. We consider three different units of treatments, namely: State of New York, Italy and Indian capital city Delhi. In some sense these place have also been termed as regional epicenters of the epidemic.

### 4.1. State of New York

### 4.2. Italy

### 4.3. Delhi, India

India had one of the most strict stay home order across the country in first phase of the lock-down between 25 March 2020 – 14 April 2020 (21 days), where entire 1.3 billion were put under restricted movement. Overall

Data source		Description
Daily update from JHU [15]		We use this data to derive metric for units: i.e. number of confirmed cases and number of deceased for each day and geographic locations.
Facebook Movement Range Maps [16]		The relative percentage of population that is staying put and also the percentage of population that moved from a region to another. We use this data in selection of donor units.
Apple Trends[17]	Mobility	Relative volume of directions requests per region, sub-region or city compared to a baseline volume, categorized by Driving, Walking or Public Transport. We use this data in selection of donor units.
OxCGRT [14]		Strictness of policies that restrict people’s behavior, 8 measures combined to provide a score between 0 and 100, where 100 being most stringent. We use this data in selection of donor units.
Google Mobility Report [18]	Community	Percent change in visits to places like grocery stores and parks within a geographic area. We use this data in selection of donor units.

Table 1: Data Source




Figure 5: New York State: (a) Actual number of deaths with linear trend line and points where trend changed (b) Actual number of deaths with counterfactual prediction estimated by m-RSC with actual lock-down date as intervention point (c) Actual number of confirmed cases with counterfactual prediction estimated by m-RSC.



the lock-down had multiple phases, second phase was from 15th of April 2020 to 3rd of May 2020, and third phase was 4th of May to 17th of May, 2020. We present the counterfactual for each of these dates. However, in this case we consider both daily number of confirmed cases as well as cumulative number of confirmed cases for phase three of the lock-down. We limit the donor group as all others states of India.

Figure 7 shows that counterfactual converges closely with the actual at third phase of the lock-down. It should be noted that there are a few discrepancies in reporting. First, there is a weekly seasonality - possibly due to lesser number of reporting over the weekends. Second a revised higher number of reporting on certain dates (high peak).

Linear trend-line fit shows two change points in the growth of number of cases - indicating that the exponential phase came at much later date and growth of the epidemic was under effective control in the earlier stages. Since, stay-home order was applicable across all states and adherence was almost uniform - trajectory of actual and counterfactual remains nearly same.

## 5. Discussion and Concluding Remarks

### References

- [1] Y. Liu, A. A. Gayle, A. Wilder-Smith, J. Rocklöv, The reproductive number of covid-19 is higher compared to sars coronavirus, *Journal of Travel Medicine* 27 (2). doi:10.1093/jtm/taaa021.  
URL <https://doi.org/10.1093/jtm/taaa021>
- [2] G. Viceconte, N. Petrosillo, Covid-19 r0: Magic number or conundrum?, *Infectious disease reports* 12 (1) (2020) 8516–8516.  
URL <https://pubmed.ncbi.nlm.nih.gov/32201554>
- [3] A. Banerjee, L. Pasea, S. Harris, A. Gonzalez-Izquierdo, A. Torralbo, L. Shallcross, M. Noursadeghi, D. Pillay, N. Sebire, C. Holmes, C. Pagel, W. K. Wong, C. Langenberg, B. Williams, S. Denaxas, H. Hemingway, Estimating excess 1-year mortality associated with the covid-19 pandemic according to underlying conditions and age: a population-based cohort study, *The Lancet* 395 (10238) (2020) 1715–1725.
- [4] A. Rismanbaf, Potential treatments for covid-19; a narrative literature review, *Archives of academic emergency medicine* 8 (1) (2020) e29–e29.




Figure 7: Delhi, India: (a) Actual number of confirmed cases with trend line and points where trend changed (b) Actual number of confirmed cases with counterfactual prediction estimated by m-RSC with First lock-down date as intervention point (c) On Second lock-down date (d) On third lock-down date



- [5] S. Price, S. Singh, S. Ledot, P. Bianchi, M. Hind, G. Tavazzi, P. Vranckx, Respiratory management in severe acute respiratory syndrome coronavirus 2 infection, *European Heart Journal: Acute Cardiovascular Care* 9 (3) (2020) 229–238. doi:10.1177/2048872620924613. URL <https://doi.org/10.1177/2048872620924613>
- [6] W. C. Koh, L. Naing, J. Wong, Estimating the impact of physical distancing measures in containing covid-19: an empirical analysis, *medRxiv* doi:10.1101/2020.06.11.20128074.
- [7] E. Kong, D. Prinz, The impact of shutdown policies on unemployment during a pandemic 24–72arXiv:<https://cepr.org/sites/default/files/news/CovidEconomics17.pdf>.
- [8] M. Supino, A. d’Onofrio, F. Luongo, G. Occhipinti, A. Dal Co, The effects of containment measures in the italian outbreak of covid-19, *medRxiv* doi:10.1101/2020.03.25.20042713.
- [9] D. Ibarra-Vega, Lockdown, one, two, none, or smart. modeling containing covid-19 infection. a conceptual model, *Science of The Total Environment* 730 (2020) 138917. doi:<https://doi.org/10.1016/j.scitotenv.2020.138917>.
- [10] M. Chinazzi, J. T. Davis, M. Ajelli, C. Gioannini, M. Litvinova, S. Merler, A. Pastore y Piontti, K. Mu, L. Rossi, K. Sun, C. Viboud, X. Xiong, H. Yu, M. E. Halloran, I. M. Longini, A. Vespignani, The effect of travel restrictions on the spread of the 2019 novel coronavirus (covid-19) outbreak, *Science* 368 (6489) (2020) 395–400. arXiv:<https://science.sciencemag.org/content/368/6489/395.full.pdf>, doi:10.1126/science.aba9757. URL <https://science.sciencemag.org/content/368/6489/395>
- [11] A. Abadie, A. Diamond, J. Hainmueller, Synthetic control methods for comparative case studies: Estimating the effect of california’s tobacco control program, *Journal of the American Statistical Association* 105 (490) (2010) 493–505. doi:10.1198/jasa.2009.ap08746.
- [12] M. Amjad, D. Shah, D. Shen, Robust synthetic control, *Journal of Machine Learning Research* 19 (22) (2018) 1–51.

- [13] M. Amjad, V. Misra, D. Shah, D. Shen, Mrsc: Multi-dimensional robust synthetic control, *Proc. ACM Meas. Anal. Comput. Syst.* 3 (2). doi:10.1145/3341617.3326152.  
URL <https://doi.org/10.1145/3341617.3326152>
- [14] T. Hale, S. Webster, A. Petherick, T. Phillips, B. Kira, Variation in government responses to covid-19.  
URL <https://www.bsg.ox.ac.uk/research/publications>
- [15] E. Dong, H. Du, L. Gardner, An interactive web-based dashboard to track COVID-19 in real time, *Lancet Infect Dis* 20 (5) (2020) 533–534.
- [16] K. Jin, L. McGorman, Data for good: New tools to help health researchers track and combat COVID-19 (4 2020).  
URL <https://about.fb.com/news/2020/04/data-for-good/>
- [17] Apple, Mobility trends reports.  
URL <https://www.apple.com/covid19/mobility/>
- [18] J. Fitzpatrick, K. DeSalvo, Helping public health officials combat covid-19.