

Supplementary Material

Ceremonial Entheogen Use in Puerto Rico: A Descriptive Survey of Participants and a Small Facilitator Subsample

All results below derive from the same analytic sample described in the main text ($N = 73$; participant branch $n = 67$ routed; facilitator branch $n = 8$, 7 dual-role). Structural-skip and “prefer not to answer” conventions follow the main text’s Methods and are applied uniformly. Counts are exact throughout, including cells below five. The bias-corrected ε^2 formula can return small negative values when an effect is indistinguishable from zero; these are displayed truncated to 0.00 and marked †, applied uniformly.

S1 Multiplicity correction: all four test families

The preparation-to-satisfaction comparison, designated in the analysis plan as the primary comparison before any test was run, is reported uncorrected in the main text and is excluded from every family below. The study was not preregistered. All other bivariate tests were assigned in advance to one of four thematically organized families and corrected jointly within family by the Benjamini–Hochberg false-discovery-rate procedure. Two variables, non-consensual contact and screening received, recur as predictor or outcome across three of the four families (see the safety-domain and legal-knowledge families below), so family-wise correction does not bound the error rate for conclusions about either variable across every test in which it appears; no q -value in any family is below 0.05, so this does not change any result. **No test in any family was significant at $q < 0.05$.**

Table S1: Benjamini–Hochberg correction, bivariate family. † marks an ε^2 whose bias-corrected value was negative and is displayed truncated to 0.00. ‡: a tie-aware permutation check (10,000 label permutations, seed 20260715) gave $p = 0.010$ for this test, the only one in this family reaching uncorrected significance; its “no”-background group contributes only 3 non-missing values.

Test	raw p	BH q	ε^2
Preparation (3-level) $\rightarrow$ satisfaction (KW)	0.120	0.300	0.04
Comfort/safety by prior knowledge of facilitator background (KW)	0.326	0.407	0.00
Comfort/safety by screening received (KW)	0.301	0.407	0.01
Trust in facilitator by prior knowledge of background (KW)	0.017‡	0.085	0.10
Trust in facilitator by screening received (KW)	0.679	0.679	0.00†

Table S2: Benjamini–Hochberg correction, substance-stratified family.

Test	raw p	BH q	effect size
Satisfaction by substance group (KW)	0.083	0.166	$\varepsilon^2 = 0.05$
Non-consensual contact by substance group (permutation χ^2)	0.486	0.486	$V = 0.17$

Table S3: Benjamini–Hochberg correction, safety-domain family. The non-consensual contact outcome is the lifetime item, while the three participant-side exposures crossed against it all ask about the most recent ceremony; 3 of the 5 respondents reporting the event ever placed it at an earlier ceremony, so for those three the exposure and the outcome describe different events. This referent mismatch applies to every row in the upper block and is the main reason these tests are hypothesis-generating only.

Test	raw p	BH q	effect size
Non-consensual contact $\times$ prior knowledge of facilitator background	0.047	0.188	$V = 0.38$
Non-consensual contact $\times$ screening received	0.094	0.188	$V = 0.32$
Witnessed crisis signs $\times$ distress-response protocol (facilitator branch)	0.143	0.191	OR undefined
Witnessed crisis signs $\times$ emergency plan (facilitator branch)	0.429	0.429	OR undefined

Table S4: Benjamini–Hochberg correction, legal-knowledge family. Predictors are three participant-side legal items; outcomes are disclosure to a health professional, screening received, and non-consensual contact. All tests Kruskal–Wallis. $\dagger$ marks a truncated ε^2 , as above.

Test	raw p	BH q	ε^2
General legal knowledge by disclosure to a health professional	0.258	0.581	0.01
Most-recent-ceremony legal knowledge by disclosure	0.164	0.495	0.03
Most-recent-ceremony legal worry by disclosure	0.031	0.283	0.09
General legal knowledge by screening received	0.412	0.741	0.00 †
Most-recent-ceremony legal knowledge by screening received	0.659	0.741	0.00 †
Most-recent-ceremony legal worry by screening received	0.914	0.914	0.00 †
General legal knowledge by non-consensual contact	0.617	0.741	0.00 †
Most-recent-ceremony legal knowledge by non-consensual contact	0.165	0.495	0.03
Most-recent-ceremony legal worry by non-consensual contact	0.611	0.741	0.00 †

The only legal-domain comparison reaching uncorrected significance was most-recent-ceremony legal worry by disclosure to a health professional ($H = 6.92$, $p = 0.031$, $\varepsilon^2 = 0.09$): median 1.5 (IQR 0–2) among the 8 reporting disclosure against median 0 (IQR 0–0) among the 47 reporting none. It did not survive correction ($q = 0.283$). Two further legal items were not testable: a general legal-worry item reached only six respondents because of a skip-logic error described in the main text’s Limitations, and a facilitator legal-knowledge item sits on the unlinked facilitator branch with no participant-side counterpart.

S2 Exploratory proportional-odds model

This model adjusts preparation for age and facilitator trust simultaneously. The full covariate set specified in the original protocol was not estimable: gender cells were too thin for a four-level categorical predictor, and screening overlapped substantively with the trust and background measures already in the model. A prespecified three-predictor fallback was used instead. The outcome was collapsed to three ordered levels for this model only, because the raw seven-level scale has cells as small as one to three respondents at the low end and risks quasi-complete separation at this sample size; every other table in the main text and this supplement retains the full 0–6 scale.

Table S5: Proportional-odds model for reported satisfaction. Complete-case $n = 62$ of the 63 participants with a non-missing outcome. Collapsed outcome: low/mixed (0–3) $n = 9$; mostly positive (4–5) $n = 19$; very positive (6) $n = 34$.

Predictor	OR	95% CI	p
Age (years)	0.99	0.94–1.03	0.527
Any preparation vs. none	5.21	0.84–32.41	0.077
Trust in facilitator (0–4)	2.20	1.06–4.53	0.033

Omnibus likelihood-ratio test against an intercept-and-cutpoint-only model: $\chi^2(3) = 8.68$, $p = 0.034$; McFadden pseudo- $R^2 = 0.07$. **Power caveat.** Sixty-two complete cases support five estimated parameters (three slopes and two cutpoints), roughly 12 cases per parameter, below conventional rules of thumb for stable ordinal-regression estimates. The proportional-odds assumption was not formally tested, the sample being too small to test it reliably, and is simply assumed. Every estimate in this table is exploratory and hypothesis-generating, and should not be presented without this caveat.

S3 Psychometric evidence for the instrument

The instrument was purpose-built, so this is its first reliability evidence. Two genuine same-respondent pairs are testable.

With only two items, α is a monotonic transform of the pairwise correlation and the item-rest correlations are not independent numbers. The corresponding Spearman correlations are $\rho = 0.39$ ($p = 0.002$, $n = 64$) for the trust-and-comfort pair and $\rho = 0.79$ ($p < 0.001$, $n = 55$) for the

Table S6: Reliability of the two testable same-respondent item pairs.

Scale	Cronbach's α	95% CI	n
Trust in facilitator + in-ceremony comfort/safety (2 items)	0.58	0.31–0.74	64
Legal knowledge, general + most-recent ceremony (2 items)	0.86	0.75–0.92	55

legal-knowledge pair; the latter two items required harmonization onto a shared 0–4 code because the most-recent-ceremony version omits the bottom category its general counterpart carries.

A third candidate pair was untestable: a perceived-life-impact item had zero variance, with all 67 respondents answering selecting the same option, so it cannot correlate with anything and α is undefined. This is reported as a data-quality finding, not a scale result.