9.10.4.3

Question: If two equal chords of a circle intersect within the circle, prove that the line joining the point of intersection to the centre makes equal angles with the chords.

Solution:

| Input Parameters | Description | Value |
|------------------|----------------------|-----------------|
| 0 | Center(at origin) | 0 |
| d | Length of the chords | 2 |
| r | Radius | 1 |
| θ_1 | $\angle ROP$ | $	heta_1^\circ$ |
| θ_2 | $\angle ROQ$ | -48° |
| θ_3 | $\angle ROR$ | 0° |
| θ_4 | $\angle ROS$ | -132° |

Table 1: Table of input parameters

| Output Parameters | Description | Value |
|-------------------|-------------|--|
| P | Point | $\begin{pmatrix} \cos \theta_1 \\ \sin \theta_1 \end{pmatrix}$ |
| Q | Point | $\begin{pmatrix} \cos \theta_2 \\ \sin \theta_2 \end{pmatrix}$ |
| R | Point | $\begin{pmatrix} \cos \theta_3 \\ \sin \theta_3 \end{pmatrix}$ |
| S | Point | $\begin{pmatrix} \cos \theta_4 \\ \sin \theta_4 \end{pmatrix}$ |

Table 2: Table of output parameters

$$\cos \angle RQS = \frac{(\mathbf{R} - \mathbf{Q})^{\top} (\mathbf{Q} - \mathbf{S})}{||\mathbf{R} - \mathbf{Q}||||\mathbf{Q} - \mathbf{S}||}$$
(1)

$$= \cos \frac{\theta_3 - \theta_4}{2} \tag{2}$$

$$\cos \angle PSQ = \frac{(\mathbf{P} - \mathbf{S})^{\top} (\mathbf{S} - \mathbf{Q})}{||\mathbf{P} - \mathbf{S}||||\mathbf{S} - \mathbf{Q}||}$$
(3)

$$=\cos\frac{\theta_1 - \theta_2}{2} \tag{4}$$

$$\cos \angle RQS = \cos \angle PSQ \tag{5}$$

$$or, \theta_1 = \theta_2 - \theta_3 + \theta_4 \tag{6}$$

$$= -180^{\circ} \tag{7}$$

The equation of PQ and RS is obtained by

$$\mathbf{n_1}^{\top}(\mathbf{x} - \mathbf{P}) = 0 \tag{8}$$

$$\mathbf{n_2}^{\top}(\mathbf{x} - \mathbf{R}) = 0 \tag{9}$$

$$or, \mathbf{n_1} = \begin{pmatrix} \sin \theta_2 - \sin \theta_1 \\ \cos \theta_1 - \cos \theta_2 \end{pmatrix}$$

$$or, \mathbf{n_2} = \begin{pmatrix} \sin \theta_4 - \sin \theta_3 \\ \cos \theta_3 - \cos \theta_4 \end{pmatrix}$$

$$(10)$$

$$or, \mathbf{n_2} = \begin{pmatrix} \sin \theta_4 - \sin \theta_3 \\ \cos \theta_3 - \cos \theta_4 \end{pmatrix} \tag{11}$$

(12)

The value of the point of the intersection is

$$\begin{pmatrix} \mathbf{n_1}^\top \\ \mathbf{n_2}^\top \end{pmatrix} \mathbf{X} = \begin{pmatrix} \mathbf{n_1}^\top \mathbf{P} \\ \mathbf{n_2}^\top \mathbf{R} \end{pmatrix}$$
 (13)

$$\mathbf{X} = \frac{1}{\begin{vmatrix} \cos \theta_3 - \cos \theta_4 & \cos \theta_2 - \cos \theta_1 \\ \sin \theta_3 - \sin \theta_4 & \sin \theta_2 - \sin \theta_1 \end{vmatrix}} \times \begin{pmatrix} \cos \theta_3 - \cos \theta_4 & \cos \theta_2 - \cos \theta_1 \\ \sin \theta_3 - \sin \theta_4 & \sin \theta_2 - \sin \theta_1 \end{pmatrix} \begin{pmatrix} \sin \theta_2 - \theta_1 \\ \sin \theta_4 - \theta_3 \end{pmatrix}$$
(14)

$$or, \mathbf{T} = \begin{pmatrix} 0 \\ -0.45 \end{pmatrix} \tag{15}$$

$$\cos \angle OTP = \frac{(\mathbf{T} - \mathbf{P})^{\top} (\mathbf{O} - \mathbf{T})}{||\mathbf{T} - \mathbf{P}||||\mathbf{T} - \mathbf{O}||}$$
(16)

$$or, \angle OTP = 66^{\circ} \tag{17}$$

Similarly,

$$\cos \angle OTR = \frac{(\mathbf{T} - \mathbf{R})^{\top} (\mathbf{T} - \mathbf{O})}{||\mathbf{T} - \mathbf{R}||||\mathbf{T} - \mathbf{O}||}$$
(18)

$$or, \angle OTR = 66^{\circ} \tag{19}$$

$$So, \angle OTP = \angle OTR (proved)$$
 (20)




Figure 1: