

# Assignment Part II

## Question 1

What is the optimal value of alpha for ridge and lasso regression? What will be the changes in the model if you choose double the value of alpha for both ridge and lasso? What will be the most important predictor variables after the change is implemented?

Ans: The optimal value of alpha for Ridge is 2 and for Lasso it is 0.001. With these alphas the R2 of the model was approximately 0.83.

After doubling the alpha values in the Ridge and Lasso, the prediction accuracy remains around 0.82 but there is a small change in the co-efficient values. The new model is created and demonstrated in the Jupiter notebook. Below are the changes in the co-efficient.

### Ridge Regression Model

Ridge Co-Efficient	
Total_sqr_footage	0.169122
GarageArea	0.101585
TotRmsAbvGrd	0.067348
OverallCond	0.047652
LotArea	0.043941
CentralAir_Y	0.032034
LotFrontage	0.031772
Total_porch_sf	0.031639
Neighborhood_StoneBr	0.029093
Alley_Pave	0.024270
OpenPorchSF	0.023148
MSSubClass_70	0.022995
RoofMatl_WdShngl	0.022586
Neighborhood_Veenker	0.022410
SaleType_Con	0.022293
HouseStyle_2.5Unf	0.021873
PavedDrive_P	0.020160
KitchenQual_Ex	0.019378
LandContour_HLS	0.018595
SaleType_Oth	0.018123

Ridge Doubled Alpha Co-Efficient	
Total_sqr_footage	0.149028
GarageArea	0.091803
TotRmsAbvGrd	0.068283
OverallCond	0.043303
LotArea	0.038824
Total_porch_sf	0.033870
CentralAir_Y	0.031832
LotFrontage	0.027526
Neighborhood_StoneBr	0.026581
OpenPorchSF	0.022713
MSSubClass_70	0.022189
Alley_Pave	0.021672
Neighborhood_Veenker	0.020098
BsmtQual_Ex	0.019949
KitchenQual_Ex	0.019787
HouseStyle_2.5Unf	0.018952
MasVnrType_Stone	0.018388
PavedDrive_P	0.017973
RoofMatl_WdShngl	0.017856
PavedDrive_Y	0.016840

## Lasso Regression Model

Lasso Co-Efficient		Lasso Doubled Alpha Co-Efficient	
Total_sqr_footage	0.202244	Total_sqr_footage	0.204642
GarageArea	0.110863	GarageArea	0.103822
TotRmsAbvGrd	0.063161	TotRmsAbvGrd	0.064902
OverallCond	0.046686	OverallCond	0.042168
LotArea	0.044597	CentralAir_Y	0.033113
CentralAir_Y	0.033294	Total_porch_sf	0.030659
Total_porch_sf	0.028923	LotArea	0.025909
Neighborhood_StoneBr	0.023370	BsmtQual_Ex	0.018128
Alley_Pave	0.020848	Neighborhood_StoneBr	0.017152
OpenPorchSF	0.020776	Alley_Pave	0.016628
MSSubClass_70	0.018898	OpenPorchSF	0.016490
LandContour_HLS	0.017279	KitchenQual_Ex	0.016359
KitchenQual_Ex	0.016795	LandContour_HLS	0.014793
BsmtQual_Ex	0.016710	MSSubClass_70	0.014495
Condition1_Norm	0.015551	MasVnrType_Stone	0.013292
Neighborhood_Veenker	0.014707	Condition1_Norm	0.012674
MasVnrType_Stone	0.014389	BsmtCond_TA	0.011677
PavedDrive_P	0.013578	SaleCondition_Partial	0.011236
LotFrontage	0.013377	LotConfig_CulDSac	0.008776
PavedDrive_Y	0.012363	PavedDrive_Y	0.008685

Overall, since the alpha values are small, **we do not see a huge change in the model after doubling the alpha.**

## Question 2

You have determined the optimal value of lambda for ridge and lasso regression during the assignment. Now, which one will you choose to apply and why?

- The optimum lambda value in case of Ridge and Lasso is as follows: -
  - Ridge – 2
  - Lasso – 0.0001
- The Mean Squared Error in case of Ridge and Lasso are:
  - Ridge - 0.0018396090787924262
  - Lasso - 0.0018634152629407766
- The Mean Squared Error of both the models are almost same.
- Since Lasso helps in feature reduction (as the coefficient value of some of the features become zero), **Lasso has a better edge over Ridge and should be used as the final model.**

### Question 3

After building the model, you realised that the five most important predictor variables in the lasso model are not available in the incoming data. You will now have to create another model excluding the five most important predictor variables. Which are the five most important predictor variables now?

Ans: The five most important predictor variables in the current lasso model is:-

1. Total\_sqr\_footage
2. GarageArea
3. TotRmsAbvGrd
4. OverallCond
5. LotArea

We build a Lasso model in the Jupiter notebook after removing these attributes from the dataset.

The R2 of the new model without the top 5 predictors drops to .73

The Mean Squared Error increases to 0.0028575670906482538 .

**The new Top 5 predictors are: -**

Lasso Co-Efficient	
LotFrontage	0.146535
Total_porch_sf	0.072445
HouseStyle_2.5Unf	0.062900
HouseStyle_2.5Fin	0.050487
Neighborhood_Veenker	0.042532

### Question 4

How can you make sure that a model is robust and generalisable? What are the implications of the same for the accuracy of the model and why?

Ans: As Per, Occam's Razor— given two models that show similar 'performance' in the finite training or test data, we should pick the one that makes fewer on the test data due to following reasons:-

- Simpler models are usually more 'generic' and are more widely applicable
- Simpler models require fewer training samples for effective training than the more complex ones and hence are easier to train.
- Simpler models are more robust.
  - Complex models tend to change wildly with changes in the training data set

- Simple models have low variance, high bias and complex models have low bias, high variance
- Simpler models make more errors in the training set. Complex models lead to overfitting — they work very well for the training samples, fail miserably when applied to other test samples.

Therefore, to make the model more robust and generalizable, make the model simple but not simpler which will not be of any use.

Regularization can be used to make the model simpler. Regularization helps to strike the delicate balance between keeping the model simple and not making it too naïve to be of any use. For regression, regularization involves adding a regularization term to the cost that adds up the absolute values or the squares of the parameters of the model.

Also, making a model simple may lead to Bias-Variance Trade-off:

- A complex model will need to change for every little change in the dataset and hence is very unstable and extremely sensitive to any changes in the training data.
- A simpler model that abstracts out some pattern followed by the data points given is unlikely to change wildly even if more points are added or removed.

**Bias** quantifies how accurate is the model likely to be on test data. A complex model can do an accurate job prediction provided there is enough training data. Models that are too naïve, for e.g., a model that gives same answer to all test inputs and makes no discrimination whatsoever has a very large bias as its expected error across all test inputs are very high.

**Variance** refers to the degree of changes in the model itself with respect to changes in the training data.

Therefore, accuracy of the model can be maintained by keeping the balance between Bias and Variance as it minimizes the total error as shown in the below graph.

