

Assignment 19

Mtech in AI Department
Priya Bhatia
AI20MTECH14015

Abstract

This document illustrates the concept of orthonormal basis.

Download the latex-tikz codes from

https://github.com/priya6971/matrix_theory_EE5609/tree/master/Assignment19

1 PROBLEM

Let $\{u_1, u_2, \dots, u_n\}$ be an orthonormal basis of C^n as column vectors. Let $M = \{u_1, u_2, \dots, u_k\}$ and $N = \{u_{k+1}, u_{k+2}, \dots, u_n\}$ and P be the diagonal $k \times k$ matrix with diagonal entries $\alpha_1, \alpha_2, \dots, \alpha_k \in R$. Then which of the following is true?

1. $\text{Rank}(\mathbf{M}\mathbf{P}\mathbf{M}^*) = k$ whenever $\alpha_i \neq \alpha_j$, $1 \leq i, j \leq k$
2. $\text{Trace}(\mathbf{M}\mathbf{P}\mathbf{M}^*) = \sum_{i=1}^k \alpha_i$
3. $\text{Rank}(\mathbf{M}^*\mathbf{N}) = \min(k, n - k)$
4. $\text{Rank}(\mathbf{M}\mathbf{M}^* + \mathbf{N}\mathbf{N}^*) < n$

2 DEFINITIONS

Orthonormal Basis	$B = \{u_1, u_2, \dots, u_n\}$ is the Orthonormal basis for C^n if it generates every vector C^n and the inner product $\langle u_i, u_j \rangle = 0$ if $i \neq j$. That is the vectors are mutually perpendicular and $\langle u_i, u_j \rangle = 1$ otherwise.
Trace	Trace of a square matrix A , denoted by $\text{tr}(\mathbf{A})$ is defined to be the sum of elements on the main diagonal (from the upper left to lower right) of A Some useful properties of Trace : $\text{tr}(\mathbf{AB}) = \text{tr}(\mathbf{BA})$, where A is the $m \times n$ matrix and B is the $n \times m$ matrix
Basis Theorem	A nonempty subset of nonzero vectors in R^n is called an orthogonal set if every pair of distinct vectors in the set is orthogonal. Any Orthogonal sets of vectors are automatically linearly independent and if A matrix columns are linearly independent, then it is invertible.

TABLE 1: Definitions

3 SOLUTION

$\text{Rank}(\mathbf{MPM}^*) = k$	<p>M and M^* vectors are linearly independent and thus it is invertible (Since the elementary matrices are invertible, such multiplication does not change the rank of a matrix) $\implies \text{Rank}(\mathbf{MPM}^*) = \text{Rank}(\mathbf{P})$ Now \mathbf{P} be the diagonal $k \times k$ matrix with diagonal entries $\alpha_1, \alpha_2, \dots, \alpha_k \in R$. $\text{Rank}(\mathbf{P})$ is not always k. It can be less than k if any of the entries in $\alpha_1, \alpha_2, \dots, \alpha_k$ is 0. Thus, $\text{Rank}(\mathbf{MPM}^*) \neq k$ Thus, the given statement is false</p>
$\text{Trace}(\mathbf{MPM}^*) = \sum_{i=1}^k \alpha_i$	<p>$\text{Trace}(\mathbf{MPM}^*) = \text{Trace}(\mathbf{M}^*\mathbf{MP})$ (According to properties of trace mentioned in Definitions) $\mathbf{M} = (u_1 \ u_2 \ u_3 \ \dots \ u_k)$ $\mathbf{M}^* = \begin{pmatrix} u_1^T \\ u_2^T \\ u_3^T \\ \vdots \\ u_k^T \end{pmatrix}$ $\mathbf{M}^*\mathbf{M} = \begin{pmatrix} u_1^T u_1 & 0 & 0 & \dots & 0 \\ 0 & u_2^T u_2 & 0 & \dots & 0 \\ 0 & 0 & u_3^T u_3 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & u_k^T u_k \end{pmatrix}$ (Refer to Properties mentioned in Orthonormal Basis in Definition section that is $\langle u_i, u_j \rangle = 0$ if $i \neq j$) $\mathbf{M}^*\mathbf{M} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix}$ (Refer to Properties mentioned in Orthonormal Basis in Definition section that is $\langle u_i, u_j \rangle = 1$ if $i = j$) $\mathbf{M}^*\mathbf{M} = \mathbf{I}^k$ $\mathbf{M}^*\mathbf{MP} = \mathbf{I}^k\mathbf{P} = \mathbf{P}$ $\text{Trace}(\mathbf{M}^*\mathbf{MP}) = \text{Trace}(\mathbf{I}^k\mathbf{P}) = \text{Trace}(\mathbf{P}) = \sum_{i=1}^k \alpha_i$ (Refer Definition section of Trace, it is sum of elements on the main diagonal) So, the given statement is true</p>
$\text{Rank}(\mathbf{M}^*\mathbf{N}) = \min(k, n - k)$	$M = \{u_1, u_2, \dots, u_k\}$ and $N = \{u_{k+1}, u_{k+2}, \dots, u_n\}$

	$\mathbf{M}^* \mathbf{N} = \begin{pmatrix} u_{11}^T & u_{21}^T & \cdot & \cdot & u_{n1}^T \\ u_{12}^T & u_{22}^T & \cdot & \cdot & u_{n2}^T \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ u_{1K}^T & u_{2K}^T & \cdot & \cdot & u_{nK}^T \end{pmatrix} \begin{pmatrix} u_{(k+1)1} & u_{(k+1)2} & \cdot & \cdot & u_{(k+1)n} \\ u_{(k+2)1} & u_{(k+2)2} & \cdot & \cdot & u_{(k+2)n} \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ u_{K1} & u_{K2} & \cdot & \cdot & u_{Kn} \end{pmatrix}$
$\text{Rank}(\mathbf{MM}^* + \mathbf{NN}^*) < n$	<p>M and M^* vectors are linearly independent and thus it is invertible (Since the elementary matrices are invertible, such multiplication does not change the rank of a matrix) $\implies \text{Rank}(\mathbf{MPM}^*) = \text{Rank}(\mathbf{P})$ Now \mathbf{P} be the diagonal $k \times k$ matrix with diagonal entries $\alpha_1, \alpha_2, \dots, \alpha_k \in R$. $\text{Rank}(\mathbf{P})$ is not always k. It can be less than k if any of the entries in $\alpha_1, \alpha_2, \dots, \alpha_k$ is 0. Thus, $\text{Rank}(\mathbf{MPM}^*) \neq k$ Thus, the given statement is false</p>

TABLE 2: Finding of True and False Statements

4 CONCLUSION

$\text{Rank}(\mathbf{MPM}^*) = k$	False
$\text{Trace}(\mathbf{MPM}^*) = \sum_{i=1}^k \alpha_i$	True
$\text{Rank}(\mathbf{M}^* \mathbf{N}) = \min(k, n - k)$	False
$\text{Rank}(\mathbf{MM}^* + \mathbf{NN}^*) < n$	False

TABLE 3: Conclusion of above Solutions