

Comprehensive Guide to Managing Two-Spotted Spider Mites on Tomato Plants

1. Overview

The cultivation of tomato plants is one of the most rewarding and globally significant agricultural endeavors. However, it is frequently challenged by a microscopic, highly destructive pest known as the two-spotted spider mite, scientifically classified as *Tetranychus urticae*. This microscopic organism is an incredibly pervasive pest, boasting a cosmopolitan distribution that affects outdoor garden beds, massive commercial agricultural fields, and highly controlled indoor greenhouse environments alike. Despite the common term "mite," it is essential to understand that the spider mite is not a true insect. Instead, it belongs to the arachnid family, specifically the subclass Acari, making it a close relative of spiders, ticks, scorpions, and daddy-longlegs. This biological distinction is crucial because the physiology of arachnids dictates how they reproduce, how they damage plant cellular structures, and, most importantly, why traditional insect-killing chemicals often fail to control them. While insects typically have three distinct body parts and six legs, adult arachnids like the spider mite possess two main body parts and eight legs.

The two-spotted spider mite is widely considered one of the most economically damaging pests worldwide, capable of devastating a tomato crop in a matter of weeks if left unchecked. These mites are incredibly small, averaging only 1/64 to 1/50 of an inch in length (approximately 0.5 millimeters), which is roughly the size of a period at the end of a printed sentence. Due to their minuscule size and their tendency to hide on the undersides of leaves, a severe infestation can quietly build up over a long period before a grower or farmer even realizes there is a problem in the crop canopy.

In terms of physical appearance, adult two-spotted spider mites range in color from pale yellow, translucent cream, or light green to dark brown. Occasionally, they can appear bright orange or red, particularly when environmental conditions become unfavorable for their growth or during the colder overwintering months. As their common name suggests, they have two distinct, dark-pigmented spots or blotches on their backs, located on either side of the abdomen near the head. These spots are not actual pigmentation on the skin; rather, they are the accumulation of ingested plant food particles and waste that are visible through the mite's highly translucent body wall.

To effectively manage these resilient pests, one must thoroughly understand their rapid and complex life cycle, which is heavily influenced by ambient temperature. The two-spotted spider mite develops through five distinct life stages: the egg, the larva, two nymphal stages (the protonymph and the deutonymph), and finally, the mature adult.

- **The Egg Stage:** The life cycle begins when an adult female lays tiny, spherical, pearl-like eggs. These eggs are initially clear or pale green and become opaque white as they age. Just before hatching, the red eyespots of the developing larvae become visible through the eggshell. Females typically deposit these eggs singly on the undersides of the tomato leaves, often securely attached within the protective, fine silk webbing spun by the adult

colony.

- **The Larval Stage:** Within three to nineteen days, depending almost entirely on the surrounding temperature, the egg hatches into a tiny larva. Interestingly, at this initial stage of life, the larva is nearly translucent and possesses only six legs, making it look somewhat different from the adult. Despite its size, the larva immediately begins feeding aggressively on the plant's leaf tissues.
- **The Nymph Stages:** After a period of feeding, the larva enters a brief resting phase where it stops moving entirely in preparation for molting. It then sheds its skin to become an eight-legged protonymph. It feeds, rests, and molts a second time to become an eight-legged deutonymph. During these highly active nymphal stages, the mites consume vast amounts of plant sap to fuel their growth.
- **The Adult Stage:** After a final resting phase and molt, the mite emerges as a fully mature, reproductive adult. The male mites are generally smaller, more highly active, and feature a narrower, more pointed abdomen compared to the females. Female mites, which are slightly larger and rounder, have a remarkable and terrifying reproductive capacity. They utilize a form of asexual reproduction known as parthenogenesis, specifically a type called arrhenotoky. This means that if a female lays unfertilized eggs, they will still hatch successfully, but they will exclusively develop into male mites. If she mates and lays fertilized eggs, they will develop into female mites.

The speed at which this entire life cycle is completed is dictated by the weather. In cooler climates averaging around 60 degrees Fahrenheit, it takes approximately 28 days for an egg to develop into a mature adult. However, in ideal, warm conditions (averaging 82 to 85 degrees Fahrenheit), an egg can mature into a reproducing adult in as little as eight to fourteen days. Because a single female can lay approximately 20 eggs per day, resulting in a total of over 100 to 120 eggs in her short lifespan, populations can explode at an exponential rate. Under perfect conditions, a spider mite population can multiply its numbers seventy-fold in a matter of weeks, allowing them to rapidly overwhelm a healthy tomato crop. This intense biological drive and rapid generational turnover are what make the two-spotted spider mite such a formidable opponent in the agricultural world.

2. Symptoms

Detecting a two-spotted spider mite infestation in its earliest stages is the single most critical step in saving a tomato crop from total devastation. Because the mites are microscopic, growers and farmers rarely see the pests themselves during the initial phases of an invasion. Instead, the first indicators of trouble are the visible physiological symptoms that manifest on the tomato plant's foliage. Understanding exactly how spider mites feed helps explain the progressive symptoms they cause to the plant's cellular structure.

Mites do not chew on leaves like caterpillars or beetles; instead, they are equipped with specialized, needle-like mouthparts. They use these microscopic mouthparts to pierce individual plant parenchyma cells. Once a cell is successfully punctured, the mite injects saliva and sucks out the cell's internal contents, including the vital green pigment known as chlorophyll. Every single minute, a feeding spider mite can destroy one to two dozen individual plant cells.

When examining a perfectly healthy tomato plant, the leaves should be a deep, solid, vibrant green, indicating that the cells are full of chlorophyll and actively photosynthesizing. However, when mites begin feeding in large numbers, they leave behind millions of empty, dead plant cells.

Here are the visible signs of a disease state or mite infestation, detailed in simple, recognizable symptoms as they progress from mild to severe:

- **Stippling (Tiny Pale Specks):** The very first visible symptom of spider mite damage is a phenomenon known as stippling. As individual leaf cells are drained of their green contents and die, they turn white, yellow, or silvery-transparent. To the naked eye, this looks like hundreds or thousands of tiny, pale yellow or white dots scattered across the normally smooth, green surface of the leaf. These specks usually appear first around the midrib and the larger structural veins of the leaf before spreading outward toward the leaf edges. At this stage, the damage is often mistaken for a mild nutrient deficiency or viral speckling.
- **Yellowing, Bronzing, and Chlorosis:** As the mites continue to multiply and feed relentlessly, the thousands of tiny white dots begin to merge and coalesce together. The empty cells give large areas of the leaf a sickly, whitish or silvery-transparent appearance. The overall aesthetic of the leaf will shift from a healthy green to a pale green, and eventually to a widespread, sickly yellow or reddish-brown bronze color. This widespread loss of chlorophyll drastically reduces the leaf's ability to undergo photosynthesis, robbing the tomato plant of its energy source.
- **Crusty or Gritty Leaf Texture:** If you physically touch and gently rub the underside of a heavily infested tomato leaf, it will no longer feel smooth, supple, or slightly fuzzy. Instead, it will feel excessively dry, crusty, or gritty to the touch. This rough, dusty texture is caused by a microscopic accumulation of biological debris. It is a combination of the dried-out plant tissue, hundreds of tiny, clear spider mite eggshells, the pale, discarded cast skins from molting nymphs, and the dark waste products (feces) left behind by the feeding colonies.
- **Fine Silk Webbing:** This is one of the most definitive and easily recognizable signs of a severe spider mite infestation. As the mite colonies grow in population density, the adults spin ultra-fine silk threads, much like a tiny spider. Initially, these webs may only be visible on the undersides of the leaves or in the tight, protective corners where the leaf stems meet the main branches of the tomato plant. However, in severe cases of neglect, the webbing can become incredibly dense, eventually covering the entire plant, engulfing whole leaves, flower clusters, and the delicate growing tips. This webbing serves multiple vital purposes for the mites: it creates a microclimate that controls humidity, it physically protects the delicate eggs from harsh weather and chemical sprays, it shields the colony from larger predatory insects, and it provides a network of silky "highways" for the mites to travel rapidly across the plant.
- **Desiccation and Premature Leaf Drop:** Tomato plants rely entirely on their leaves for the energy required for rapid growth and fruit production. When leaves are heavily damaged by the constant piercing and draining of cells by mites, the foliage quickly loses its ability to retain moisture. The leaves become completely desiccated, drying out, turning brown, and dying. The plant, in an effort to save itself, will eventually shed these dead leaves. In hot and dry weather conditions, a severely infested tomato plant can lose a massive portion of its foliage in just a few short weeks, leaving the stems bare.
- **Stunted Growth and Poor Fruit Quality:** The ultimate consequence of a spider mite infestation is the catastrophic impact on the harvest. If a tomato plant is heavily attacked while it is still young, its growth will slow dramatically, and it may remain permanently undersized and weak. If the plant has managed to bear fruit, the consequences are equally dire. Heavy mite feeding directly on the developing fruit can cause strange, yellowish speckling across the skin of the tomato. Furthermore, because the plant has lost

the protective canopy of leaves it needs to provide shade, the resulting tomatoes are often heavily sunburned (sunscaled). The defoliated plants lack the energy to size up the fruit, leading to yields of incredibly small, poor-quality, and tasteless tomatoes, rendering the entire crop unmarketable for commercial farmers and deeply disappointing for home gardeners.

To definitively confirm that these symptoms are caused by spider mites—and not by a fungal disease, a bacterial infection, or a soil nutrient deficiency—gardeners can use a simple, traditional scouting technique known as the paper test. Take a clean sheet of white printer paper, a piece of white cardboard, or a white cloth tray, and hold it directly underneath a symptomatic, stippled tomato leaf. Tap or shake the leaf stem sharply a few times. If two-spotted spider mites are present on the plant, the physical shock will dislodge them, and they will fall onto the bright white surface. Because they are so small, they will initially look like tiny specks of dark dirt, debris, or moving dust. If these dark or reddish specks begin to slowly crawl across the paper, an active spider mite infestation is absolutely confirmed. For a more detailed inspection, a handheld magnifying glass or a jeweler's loupe should be used to look directly at the undersides of the discolored leaves to spot the slow-moving live mites, the translucent spherical eggs, and the characteristic silk webbing.

3. Causes & Spread

Understanding exactly how two-spotted spider mites arrive on tomato plants and the specific environmental conditions that cause their microscopic populations to explode into crop-destroying swarms is the key to controlling them. Spider mites are opportunistic pests; they are often naturally present in the background of many outdoor gardens, weed borders, and agricultural fields in small, virtually harmless numbers. However, specific environmental triggers and human errors can cause their populations to grow completely out of control.

Environmental and Climatic Triggers:

- **Prolonged Hot and Dry Weather:** The single most significant and universal cause of a severe spider mite outbreak is a period of prolonged hot, arid weather. Two-spotted spider mites thrive and multiply best when ambient temperatures climb above 85 degrees Fahrenheit (29 degrees Celsius) and when the relative air humidity drops significantly below 90 percent. Under these hot, dry conditions, the mites' internal life cycle accelerates rapidly, reducing the time it takes to go from an egg to a reproducing adult down to just a few days. In addition to speeding up their breeding capacity, dry weather prevents the growth of natural, insect-killing fungal diseases that normally thrive in wet weather and help keep mite populations in check naturally.
- **Drought Stress in the Host Plant:** When a tomato plant does not receive enough consistent irrigation, it enters a state of severe drought stress. Dehydrated plants experience profound changes in their internal chemical composition. As the plant loses water, its sap becomes highly concentrated, becoming sweeter and much richer in certain essential amino acids and complex sugars. This chemical shift actually makes the tomato plant's sap far more nutritious, palatable, and attractive to spider mites. A well-watered, fully hydrated tomato plant with high internal cell pressure (turgor) is much more capable of fending off a minor mite attack than a thirsty, struggling plant whose concentrated sap acts as a superfood for the pests.
- **Dusty Conditions:** Spider mites absolutely love dusty environments. Dust settling heavily on tomato leaves from nearby dirt roads, dry pathways, or windy fields provides two

distinct survival advantages for the mites. First, the layer of dust creates a protective micro-environment on the leaf surface that helps shield the mites' soft bodies and delicate eggs from drying out completely. Second, and much more importantly, abrasive dust is highly detrimental to the soft bodies and delicate wings of the natural predators that actively hunt spider mites, such as beneficial predatory mites and tiny parasitic wasps. A dusty tomato plant effectively becomes a safe, predator-free haven for spider mites.

Human and Agricultural Causes:

- **The Overuse of Broad-Spectrum Pesticides:** Paradoxically, the act of spraying harsh chemical insecticides is one of the most common causes of a severe spider mite outbreak in both home gardens and commercial farms. Broad-spectrum chemical insecticides—particularly older classes of chemicals like organophosphates, carbamates, or synthetic pyrethroids (such as malathion, bifenthrin, or cypermethrin)—are designed to kill a massive variety of insects. When a grower sprays these harsh chemicals to control an unrelated pest like aphids, whiteflies, or hornworms, the toxic spray also completely wipes out the beneficial predatory insects that naturally eat spider mites (such as ladybugs, lacewings, bigeyed bugs, and predatory mites). Because spider mites are arachnids and possess a different biology than true insects, they are often naturally immune or highly resistant to these insect-killing chemicals. With all of their natural predators suddenly dead and the mites completely unharmed by the spray, the spider mite population experiences a massive, unchecked explosion, a phenomenon commonly referred to in agriculture as a "mite flare" or "hormoligosis". In some cases, chemicals like pyrethroids actually stimulate the female spider mites to lay eggs faster than they normally would.
- **Weed Hosts and Overwintering Behavior:** Two-spotted spider mites do not simply vanish when the growing season ends. As temperatures begin to drop in the late autumn, the adult female mites seek out safe, sheltered areas to survive the cold. Their bodies change color to a reddish-orange hue, and they enter a state of biological dormancy (diapause), overwintering deep in the soil, hidden in fallen leaf litter, or buried within the ground cover and perennial weeds at the edges of fields and gardens. When spring arrives and temperatures warm up, these overwintering females wake up, become active, and begin feeding on early spring weeds. If the garden borders are full of broadleaf weeds, these wild plants act as a bridge, allowing the mites to feed and build up enormous population numbers before finally migrating onto the newly planted, tender tomato crops.
- **Physical and Mechanical Spread:** Because they are so tiny, lightweight, and easily dislodged, spider mites are incredibly easy to spread from plant to plant. In outdoor environments, they can easily be blown from an infested plant to a healthy one by a strong gust of wind, a dispersal method known as "ballooning," where they use a strand of silk to catch the breeze. They also frequently hitch rides on the clothing, gloves, and agricultural tools of gardeners and farmers who brush past an infested plant and then move down the row to work on a healthy one. Furthermore, purchasing new, already-infested tomato transplants or ornamental propagation stock from a commercial nursery is a very common way these pests are accidentally introduced into a previously clean greenhouse or garden space.

4. Treatment / Remedies

Treating a two-spotted spider mite infestation requires immense persistence, strict attention to detail, and a thorough understanding of the pest's biology. Because their life cycle is so incredibly fast, a single, one-time treatment is almost never enough to eradicate the problem; eggs that survive the first chemical or natural treatment will simply hatch a few days later, starting the destructive cycle all over again. A multi-pronged Integrated Pest Management (IPM) approach utilizing natural home remedies, biological predators, and, if absolutely necessary, targeted chemical remedies is the most effective strategy for long-term control.

Natural and Home Remedies

For home gardeners, organic farmers, and commercial growers seeking environmentally friendly and sustainable solutions, natural remedies can be highly effective if they are applied thoroughly and consistently. The goal of natural remedies is not just to kill the visible mites, but to physically break down their protective webbing, ruin their eggs, and create a hostile microclimate on the plant leaf, all while preserving the health of the tomato plant and the surrounding ecosystem.

- **The High-Pressure Water Spray (Physical Removal):** One of the absolute simplest, cheapest, and most overlooked treatments is the use of plain water. Because spider mites are tiny, soft-bodied, and delicate, a strong, high-pressure jet of water from a standard garden hose can physically blast them off the tomato plant, causing them to fall to the wet soil where they will likely die.
 - *How to apply:* Use a hose wand or an adjustable nozzle set to a strong, steady spray. It is absolutely essential to aim the water stream directly upward at the undersides of the leaves where the mites actually live and breed. The forceful water physically dislodges the adults, washes away the protective silk webbing, and crushes or washes away the delicate eggs.
 - *Frequency:* This physical washing should be done daily, or at the very least twice a week, during periods when spider mites are highly active. This method also has the highly beneficial side effect of washing abrasive dust off the leaves, making the environment much less hospitable for future mite generations.
- **Pruning and Strict Sanitation:** For highly localized or light, early-stage infestations, physical removal of the pest's habitat is highly effective. Inspect the tomato plant closely and use clean, sharp pruning shears to cut off the most heavily infested, severely yellowed, or web-covered leaves and stems.
 - *Disposal Warning:* Never throw these infested leaves on the ground near the plants, and absolutely never put them into a garden compost pile. Composting does not always reach temperatures high enough to kill spider mites, and you may inadvertently allow the mites to survive and return to the garden when you spread the compost later. Instead, place the infested plant material directly into a tightly sealed plastic bag and dispose of it in the municipal household garbage.
- **Homemade Insecticidal Soaps and Oil Emulsions:** Standard, mild liquid soaps can be used to create an incredibly inexpensive and safe organic spray. Soap works purely on contact by chemically breaking down the waxy, protective outer coating (the cuticle) of the spider mite's body. Once this protective moisture barrier is dissolved, the mite rapidly dehydrates, shrivels up, and dies.
 - *The Recipe:* A highly effective homemade concentrate can be made by mixing a 1:2 ratio of mild soap and cooking oil. For example, vigorously shake together 500 milliliters of a light plant oil (like sunflower oil) and 250 milliliters of a pure, mild liquid soap (such as liquid Castile soap or a very mild, non-degreasing dish liquid)

until the mixture turns white and creamy. Avoid heavy-duty dish detergents containing strong degreasers or bleach, as these will strip the natural protective oils from the tomato leaves and cause severe chemical burns.

- *Application:* Take just one tablespoon of this creamy concentrate and dilute it thoroughly into one liter of clean water. Pour the mixture into a garden sprayer and thoroughly drench the plant until it drips, paying special attention to completely coat the undersides of the leaves. Because soap only kills on direct, wet contact, it must physically hit the mites to work. Apply the spray in the early morning or late evening when temperatures are cooler. Spraying soap or oil in the extreme heat of the midday sun can cause the wet leaves to rapidly scorch and burn. Repeat this process every 5 to 7 days to disrupt the hatching cycle and kill newly hatched larvae.
- **Neem Oil Botanical Extract:** Neem oil is a powerful, globally recognized organic pesticide pressed from the seeds of the tropical neem tree. The primary active chemical compound found naturally in neem oil is known as azadirachtin. Neem oil works very differently than simple soap. It acts as a suffocant when wet, but more importantly, when spider mites ingest the azadirachtin left on the leaf surface, it severely disrupts their internal hormonal systems. It stops them from feeding normally, drastically reduces the female's ability to lay viable eggs, and prevents the young, six-legged larvae from successfully molting into eight-legged adults, breaking the generational cycle.
 - *Application:* Pure, cold-pressed neem oil should be mixed with water and a few drops of mild Castile soap, which acts as an emulsifier to help the oil and water mix together evenly. Spray it generously over the entire plant. Like soap, neem oil should never be applied when the weather is extremely hot (over 90 degrees Fahrenheit) or when the tomato plant is suffering from drought stress, as the heavy oil can physically clog and suffocate the plant's breathing pores (stomata), causing leaf burn and dropping. Neem oil provides a much longer-lasting residual protection compared to plain soap sprays.
- **Advanced Botanical Extracts and Essential Oils:** Many plants in nature produce their own potent defensive chemicals to repel or kill pests. Scientific research has shown that extracts and essential oils derived from plants like rosemary, peppermint, clove, garlic, and cayenne pepper are highly effective natural acaricides (mite-killers).
 - *Aloe Vera Distillate:* Research has demonstrated that a potent extract can be made using Aloe Vera. By mixing 100 grams of dried aloe leaf powder with 500 milliliters of a solvent like acetone or water, evaporating it, and dissolving the remaining distillate into clean water, researchers achieved a 100% mortality rate against spider mites in laboratory settings.
 - *Cilantro Seed Extract:* A much simpler home preparation involves cilantro. Crushing 200 grams of fresh cilantro seeds and boiling them in one liter of water for ten minutes creates a highly potent, all-natural spray. After straining out the crushed seeds, this infused water can be sprayed directly and safely onto the undersides of tomato leaves to kill mites and prevent new colonies from taking hold.
 - *Bugleweed (Ajuga) Extract:* Crude extracts made by simply blending a handful of fresh bugleweed (*Ajuga remota*) leaves with water and straining the liquid have been scientifically shown to cause massive mortality rates in spider mites. Studies indicate an 84% mortality rate within just 24 hours, with the fresh plant juice also proving highly toxic to unhatched eggs and developing nymphs, while the scent repels new adults from landing on the plant.

- **Biological Control (Predatory Insects and Mites):** In a balanced natural ecosystem, two-spotted spider mites are constantly hunted by a fierce army of natural predators. By encouraging these natural enemies to stay in the garden, or by actively purchasing and releasing them onto the tomato crop, growers can establish a self-sustaining, chemical-free defense system.
 - *Predatory Mites:* The absolute most famous and effective biological control agent is *Phytoseiulus persimilis*, a tiny, fast-moving predatory mite that almost exclusively hunts, kills, and devours two-spotted spider mites and their eggs. These predators are highly active, possess enormous appetites, and can quickly clean up an infestation. They can be purchased commercially from biological control companies and sprinkled directly onto the leaves of the tomato plants. Other predatory mites, such as the western predatory mite, are also highly effective.
 - *Other Insect Predators:* A wide variety of insects actively hunt spider mites, including ladybugs (especially the specialized "spider mite destroyer" lady beetle), green lacewings, brown lacewings, minute pirate bugs, rove beetles, and bigeyed bugs. Certain species of thrips, like the Western flower thrips (*Frankliniella occidentalis*), are also known to eat spider mite eggs, though they can sometimes damage the plant if the mite population disappears. To keep these incredibly beneficial insects alive and working, it is absolutely paramount to completely avoid using harsh chemical insecticide sprays.
- **Biological Control (Fungal Pathogens):** The realm of biological control also extends to the use of highly specialized, insect-killing fungi. There are organic, commercially available biopesticide products (often sold as wettable powders) that contain the dormant spores of naturally occurring fungal pathogens, most notably *Beauveria bassiana* and *Metarhizium anisopliae*. When mixed with water and sprayed onto the tomato plant canopy, these microscopic fungal spores physically attach to the bodies of the spider mites. The spores then germinate, penetrate the mite's hard outer shell, and grow rapidly inside the pest's body, effectively killing the mite from the inside out within a few days. Because these products rely on infection rather than chemical toxicity, they are completely safe for humans, pets, beneficial pollinating insects, and the tomato plants themselves, making them perfect for organic fruit production.

Chemical Remedies

For large-scale commercial agricultural farmers facing a catastrophic infestation that threatens massive economic loss, or for home growers where all natural remedies have completely failed, chemical treatments may become necessary. However, chemical control of the two-spotted spider mite is notoriously difficult, complex, and fraught with risk.

Because their life cycle is so incredibly short, and because they produce so many offspring, random genetic mutations occur very frequently within the population. Furthermore, spider mites reproduce via arrhenotokous parthenogenesis, which rapidly passes genetic traits down the line. If a farmer applies a chemical pesticide that kills 99% of the mites, the tiny 1% that survive likely possess a natural genetic mutation that makes them immune to that specific chemical. Within just a few weeks, those survivors will breed a massive new population that is now entirely resistant to the spray. As a result, the two-spotted spider mite is officially recognized by scientists and agricultural bodies as one of the "most resistant species" in the world, having documented resistance to almost every major class of pesticide ever invented.

Therefore, if chemicals must be used, it is absolutely essential to practice strict Resistance

Management. This means farmers must actively rotate chemical products that belong to different Insecticide Resistance Action Committee (IRAC) groups. Different groups have completely different "Modes of Action" (MOA)—meaning they kill the mite using different biological mechanisms, such as attacking the nervous system versus attacking the energy production system.

- **Abamectin (Avermectins - IRAC Group 6):** This is one of the most powerful, highly relied upon, and widely used commercial acaricides in global agriculture. Derived originally from a naturally occurring soil bacterium, Abamectin acts as a potent nerve poison, paralyzing the mite and stopping it from feeding immediately.
 - *Usage Instructions:* Abamectin is typically sold as a highly concentrated emulsifiable concentrate (EC) and must be diluted properly for safety and efficacy. The general application rate for foliar sprays is precisely 10 to 20 milliliters of the chemical mixed into 100 liters of water. For larger commercial farm applications, this equates to roughly 300 to 450 milliliters of product per hectare. In some specific agricultural guidelines, it is mixed at a rate of 35 ml along with 250 ml of a light horticultural mineral oil per 100 liters of water to increase its ability to penetrate the leaf.
 - *Application Rules and Limits:* Thorough, high-volume coverage of the dense plant canopy is strictly required. A non-ionic or silicone-based surfactant is often legally required to be added to the spray tank to help the liquid stick, spread evenly, and penetrate the waxy surface of the tomato leaves. Due to the severe risk of the mites developing resistance, growers are strictly advised by product labels not to apply Abamectin more than twice sequentially per growing season, and never to exceed a total maximum application of 3.0 Liters per hectare in a single year.
- **Other Commercial Miticides and Chemical Classes:** To successfully rotate chemicals and prevent resistance, growers utilize a variety of other targeted compounds.
 - *Spiromesifen (IRAC Group 23):* This chemical (often sold under the brand name Oberon) works by inhibiting the mite's ability to produce vital lipids (fats), which kills the eggs and prevents the young nymphs from molting properly.
 - *METI Acaricides (IRAC Group 21A):* Chemicals like Pyridaben disrupt the mite's cellular mitochondria, stopping them from producing energy.
 - *Other Active Ingredients:* Compounds like diafenthiuron, fenazaquin, chlorfenapyr, and bifenazate are also highly effective at knocking down severe, chemically resistant mite populations when used correctly according to local agricultural extensions. These specific chemicals are categorized strictly as miticides (killing mites) rather than broad-spectrum insecticides, making them slightly safer for the beneficial predator insects.
- **Understanding Chemical Resistance Mechanisms:** When spider mites become immune to a chemical, it occurs at a microscopic level. For example, resistance to Pyrethroids (IRAC Group 3A) or Organophosphates (IRAC Group 1B) often happens through a "target site point mutation," where the physical shape of the mite's nerve receptors changes so the poison can no longer bind to it like a key in a lock. Other forms of resistance involve the mite's body overproducing powerful detoxifying enzymes, such as P450 monooxygenases, esterases, and Glutathione S-Transferases (GST). These enzymes act like a highly efficient waste disposal system, chemically breaking down and neutralizing the toxic pesticide before it can reach the mite's nervous system.
- **Elemental Sulfur Dust:** A very traditional, ancient, yet highly effective preventative chemical remedy is the application of elemental sulfur dust or wettable sulfur sprays.

Applying a light, even coating of sulfur dust (or spraying wettable sulfur at a rate of 2 grams per liter) to the tomato plants two to three weeks after planting can act as an excellent, low-cost preventative measure. Sulfur has the incredible dual benefit of creating an acidic environment on the leaf that prevents spider mites from establishing colonies, while simultaneously preventing the germination of deadly fungal disease spores like blight. However, extreme care must be taken: applying sulfur when temperatures exceed 90 degrees Fahrenheit will severely scorch and burn the tomato foliage.

- **Chemicals to Strictly Avoid:** Unless absolutely necessary to treat a different, severe pest emergency, tomato growers should completely avoid using old, broad-spectrum insecticides like malathion, carbamates, or any pyrethroid-based sprays (such as bifenthrin, cyfluthrin, or zeta-cypermethrin). As heavily emphasized previously, these chemicals not only fail to kill resistant mites, but they also actively stimulate the surviving spider mites to increase their reproduction rates while simultaneously killing off 100% of the natural predators that keep the mite populations from destroying the crop.

5. Preventive Measures

Because completely eliminating spider mites once they have established massive colonies and spun their protective silk webs is so incredibly labor-intensive and costly, the old farming adage "an ounce of prevention is worth a pound of cure" applies perfectly to tomato cultivation.

Establishing a growing environment that favors the robust health of the tomato plant while actively discouraging the establishment of mites is the absolute best strategy for success.

- **Consistent, Deep Watering Practices:** Because spider mites actively seek out and thrive on drought-stressed plants with highly concentrated, sugary sap, ensuring that tomato plants receive adequate, consistent hydration is the most fundamental preventative step a grower can take. Watering should be done deeply, directly at the root zone, rather than shallowly at the surface. Deep watering encourages the roots to grow far down into the soil profile so that the plant can access consistent moisture even during the absolute hottest parts of the summer day. A fully hydrated plant maintains healthy cell pressure, keeping its sap dilute, and produces natural defensive chemicals that deter pests.
- **Environmental Dust Management:** Because abrasive dust provides a safe, dry haven for mites and kills their delicate natural predators, actively reducing dust in the garden or farm environment is essential. Applying a thick, even layer of organic mulch (such as clean straw, wood chips, or dried leaves) over the bare soil entirely around the base of the tomato plants prevents dry dirt from splashing or blowing up onto the lower leaves. In highly arid, dry regions, farmers can periodically wash down the surrounding dirt pathways and the foliage of non-flowering border trees with water to help keep ambient dust levels to an absolute minimum.
- **Strict Sanitation and Weed Control:** Maintaining a meticulously clean garden space eliminates the hiding spots where overwintering, dormant mites wait for spring. Growers should clear away all dead plant debris, old roots, and fallen leaves at the very end of every growing season. A deep summer or fall plowing of the soil can also help bury and destroy overwintering pests. Throughout the spring and hot summer, the area immediately surrounding the tomato plants must be kept entirely weed-free. Broadleaf weeds act as alternative food sources, breeding grounds, and convenient stepping-stones for mites migrating toward the main, valuable tomato crop.

- **Intercropping and Companion Planting Strategies:** Planting certain strong-smelling or repellent crops directly alongside or alternating with tomatoes can significantly reduce the severity of mite populations. Extensive agricultural research, such as field trials conducted in Eastern Ethiopia, demonstrates that intercropping tomatoes with specific vegetables creates an environment that naturally lowers mite infestations compared to planting massive, unbroken fields of solitary tomatoes. In these studies, planting tomatoes alongside onions resulted in significantly lower spider mite populations. Furthermore, the application of natural tobacco leaf extract alongside the tomatoes provided massive reductions in pest density, performing almost as well as harsh commercial chemical sprays. The strong, pungent odors and root exudates of companion plants like onions, garlic, and brassicas (like cabbage) can mask the chemical scent of the tomato plants, making it much harder for migrating pests to locate their preferred host.
- **Choosing and Breeding Genetically Resistant Tomato Varieties:** Not all tomato plants are equally susceptible to spider mite attacks. Modern agricultural science, combined with traditional plant breeding, has identified specific genetic varieties that naturally fight back against pests.
 - *The Deep Science of Genetic Resistance:* Some wild, uncultivated relatives of the modern tomato, most notably a species from South America called *Solanum habrochaites* (specifically accession LA2329), are highly, naturally resistant to a wide variety of arthropod pests. This remarkable resistance comes from the presence of extremely high densities of microscopic, hair-like structures covering their leaves and stems, known scientifically as trichomes. In the *Solanum* plant genus, there are eight distinct types of trichomes. Four of these are "glandular" (Types I, IV, VI, and VII), meaning they secrete chemicals, while the other four are "nonglandular" (Types II, III, V, and VIII), which just act as physical spikes. The glandular trichomes (specifically Type IV and Type VI) act as a dense physical barrier, like a microscopic forest, making it incredibly difficult for the tiny mites to physically walk across the leaf surface. More importantly, these specific glandular trichomes store and secrete sticky, highly toxic secondary metabolites—complex chemical compounds such as the sesquiterpene hydrocarbons known as zingiberene, 7-epizingiberene, and zingiberene alcohol. When a spider mite encounters the sticky zingiberene coating, it acts as a toxic trap. It severely deters the mite from attempting to feed, halts its desire to lay eggs, and, in the case of pure zingiberene alcohol, is directly, lethally toxic to the mite itself. Plant geneticists and breeders are continually working to cross these highly valuable, toxic wild traits into everyday, delicious cultivated tomatoes (*Solanum lycopersicum*) without ruining the flavor of the fruit.
 - *Commercially Available Tolerant Varieties:* While true, complete immunity in a great-tasting garden tomato is currently rare, choosing highly vigorous, robust commercial varieties heavily mitigates damage. Varieties known for producing massive, deeply penetrating root systems and incredibly vigorous foliage—such as the Beefmaster, Big Beef, and Lemon Boy hybrids—are widely known to survive and continue producing fruit even under the severe stress of a heavy spider mite attack. These robust plants do not possess toxic trichomes; they simply grow so fast and pull so much water that they outlast and outgrow the physical damage much longer than more delicate, sensitive heirloom varieties. Furthermore, specific agricultural strains developed by global research institutes, such as the "Arka Abhed" or "Arka Ananya" varieties developed by the Indian Council of Agricultural

Research (ICAR), possess stacked, multiple genetic resistances to bacterial wilts, leaf curl viruses, and blights, ensuring the plant remains as healthy and robust as possible to survive secondary mite attacks in the field.

6. Suggestions & Best Practices

To successfully integrate all these biological, natural, and preventative strategies into a highly successful growing season, gardeners and commercial farmers alike should adopt a rigorous routine of daily best practices focused on early vigilance and highly timely interventions.

- **Establish a Rigorous Scouting Routine:** You cannot fight an invisible enemy you do not actively look for. During the hot, dry months of peak summer, do not wait for the upper canopy leaves to turn yellow or for massive silk webs to appear before taking action. Walk the garden or the crop rows and carefully inspect the tomato plants every three to five days without fail. Look very closely at the lowest leaves near the soil first, as spider mites almost always begin their initial infestation near the dry ground and slowly work their way up the plant canopy as their numbers swell. Perform the white paper shake test regularly across different sections of the field. Catching a localized infestation when it is limited to just a few lower leaves allows you to easily prune away the problem with zero chemical intervention before it spreads to the rest of the crop.
- **Time Your Spray Applications Perfectly:** Whether you are applying simple water blasts, organic soapy emulsions, heavy botanical neem oil, or complex chemical sprays, the time of day you apply them is critical to the survival of the plant. Never apply heavy oils, concentrated soaps, or chemicals during the middle of the day when the ambient air is hot and the sun is intensely bright. The liquid droplets can act like tiny magnifying glasses, focusing the sun's rays and causing severe, irreversible phototoxicity (sunburn) to the tomato plant's delicate leaves. Instead, always apply treatments early in the morning while temperatures are cool and the plant's cellular structure is fully rehydrated from the night, or late in the evening as the sun begins to go down and the temperatures drop.
- **Master Proper Application Technology:** A very common, frustrating mistake made by novice gardeners and rushed farmers is spraying pesticide only onto the tops of the plant leaves, simply walking down the row and waving a wand. Spider mites almost exclusively live, feed, hide, and lay their eggs on the heavily protected *underside* of the tomato leaves. Ensure that any manual spray wand, backpack sprayer, or tractor-mounted hose nozzle is specifically angled upward so that the liquid physically hits and completely coats the bottom surface of the foliage. For tomatoes that are heavily staked or grown on tall vertical trellises, high-volume spraying equipment that pushes a massive amount of water into the deep canopy is absolutely required to achieve the thorough coverage necessary for eradication. For smaller, non-trellised bush tomatoes, specialized droppers can be used to direct the spray deep into the plant while keeping it out of the surrounding dirt rows.
- **Mandatory Post-Treatment Monitoring:** Never assume a single, initial treatment permanently solved the pest problem. Spider mite eggs are highly durable and naturally resistant to many botanical and chemical sprays. Always return to check the treated plants seven to ten days after the very first application. Perform the white paper test again. If tiny, crawling mites are still visible, a second application is immediately required to kill the newly hatched generation of larvae before they are biologically old enough to lay a new generation of eggs.

- **Carefully Support Post-Infestation Plant Recovery:** If a tomato plant has suffered a moderate, damaging mite infestation but the pests have finally been successfully eradicated, the traumatized plant will need careful, balanced help to recover its strength. Do not make the mistake of over-fertilizing the soil with heavy, synthetic nitrogen immediately after an attack. Massive doses of nitrogen push the plant to rapidly create incredibly weak, highly sappy, dark green new growth that is irresistible to any surviving mites or new pests moving into the area. Instead, ensure the soil moisture remains highly consistent, provide very gentle, balanced organic compost to slowly rebuild soil health, and manually prune away the most severely damaged, completely dead leaves so the plant can focus its precious energy reserves on growing new, healthy foliage and ripening the existing fruit.

7. References / Sources

The information synthesized within this comprehensive guide is derived from the extensive research, field trials, and published pest management literature of the following highly reputable agricultural institutions, universities, and scientific journals:

- **University of California Agriculture and Natural Resources (UC IPM):** Guidelines on the life cycle, symptoms, and integrated pest management of the twospotted spider mite (*Tetranychus urticae*).
- **North Carolina State University (NC State Extension):** Research on biological control strategies, specifically the utilization of the predatory mite *Phytoseiulus persimilis* in tomato crops.
- **Indian Council of Agricultural Research (ICAR) & Tamil Nadu Agricultural University (TNAU):** Field guides on eco-friendly management, chemical efficacy (diafenthiuron, neem oil), and the development of multiple disease-resistant tomato varieties like Arka Abhed and Arka Ananya.
- **University of Kentucky & Colorado State University:** Entomological studies detailing the specific overwintering behaviors, identification markers, and physiological damage caused by severe mite infestations in vegetable crops.
- **The World Vegetable Center & Academic Journals (e.g., HortScience):** In-depth genetic research regarding the role of glandular trichomes (Types I-VIII) and toxic secondary metabolites (zingiberene) in wild tomato species (*Solanum habrochaites*) for natural pest resistance.
- **National Center for Appropriate Technology (NCAT - ATTRA):** Formulations and efficacy studies on botanical extracts, including cilantro, bugleweed, and aloe vera distillates for organic mite control.
- **Insecticide Resistance Action Committee (IRAC):** Global database profiling the extensive chemical resistance mechanisms (P450 monooxygenase, point mutations) of *Tetranychus urticae* across various pesticide classes (Groups 1A, 3A, 6, 10A, 20B, 21A).
- **Haramaya University (Eastern Ethiopia Field Trials):** Agricultural research on the significant reduction of spider mite populations utilizing intercropping techniques (tomato with onion) and natural tobacco leaf extracts.
- **Various State Agricultural Extensions (Utah State, Univ. of Minnesota, Texas A&M):** Practical recommendations for scouting techniques, physical water removal, insecticidal soap recipes, and strict chemical rotation guidelines.

8. Key Takeaways

- **Understand the Biology of the Pest:** The two-spotted spider mite is an eight-legged arachnid, not a true insect. This biological distinction explains why traditional insect bug sprays often fail to kill them. They reproduce incredibly fast in hot, dry weather, completing a full life cycle from egg to adult in as little as eight days while laying over a hundred eggs.
- **Spot the Cellular Symptoms Early:** Mites do not chew holes; they pierce plant cells and suck out the green chlorophyll. Look closely for stippling (hundreds of tiny pale or white specks) on the leaves, which rapidly progresses to widespread yellowing, bronzing, and leaf death. Check for a rough, gritty texture under the leaves and fine silk webbing spun tightly around the stems.
- **Perform the Simple Paper Test:** You can easily confirm an invisible infestation by shaking a suspect, yellowing leaf over a piece of stark white paper. If tiny, dirt-like specks fall off the leaf and immediately begin to crawl across the page, you have an active spider mite colony.
- **Control the Environment, Not Just the Bug:** Mites exclusively target and thrive on drought-stressed, dusty plants. Prevent massive outbreaks entirely by keeping your tomato plants deeply and consistently watered, using heavy organic mulch to keep dry dirt down, and completely removing surrounding broadleaf weeds where the mites spend the winter.
- **Strictly Avoid Broad-Spectrum Pesticides:** Harsh chemicals (like malathion or pyrethroids) that kill a wide range of bugs will instantly destroy the highly beneficial predator insects (like ladybugs, lacewings, and predatory mites) that naturally eat spider mites. Without predators, the surviving mite population will explode out of control.
- **Start with Gentle, Physical Remedies:** For early infestations, use physical removal by spraying the undersides of the tomato leaves with a highly pressurized jet of water daily. Prune away and immediately throw out heavily infested foliage in sealed garbage bags—never compost them.
- **Utilize Powerful Organic Sprays:** Apply homemade insecticidal emulsions (a mixture of sunflower oil, liquid Castile soap, and water) or natural, cold-pressed neem oil directly to the undersides of the leaves. These natural remedies suffocate the mites and disrupt their hormones without leaving toxic, long-lasting chemical residues on your food.
- **Employ Natural Biological Predators:** Consider purchasing and releasing beneficial predators like *Phytoseiulus persimilis* mites, or utilizing highly specific organic fungal sprays like *Beauveria bassiana*, to naturally hunt, infect, and eliminate the pests entirely without chemicals.
- **Rotate Chemical Controls if Necessary:** If you must utilize commercial chemical miticides (such as Abamectin) to save a massive crop, follow all dilution and dosage instructions perfectly. You must actively rotate between different chemical classes (IRAC groups) to prevent the mites from mutating and building permanent genetic resistance to the poison.
- **Choose Vigorous and Genetically Resistant Varieties:** Plant commercial tomato varieties known for massive root systems and highly vigorous growth (like Beefmaster), which can physically survive minor pest damage better than delicate heirloom plants. Look for advanced varieties bred with natural pest-repellent leaf hairs (trichomes) or

multiple disease resistances.

Works cited

1. Twospotted Spider Mite [fact sheet] - UNH Extension, <https://extension.unh.edu/resource/twospotted-spider-mite-fact-sheet>
2. Tetranychus urticae - Wikipedia, https://en.wikipedia.org/wiki/Tetranychus_urticae
3. How to Get Rid of Spider Mites Effectively (Complete Guide) - Planet Natural, <https://www.planetnatural.com/pest-problem-solver/houseplant-pests/spider-mite-control/>
4. Twospotted Spider Mite Archives | Insecticide Resistance Action Committee - IRAC, <https://irac-online.org/pests/tetranychus-urticae/>
5. Twospotted Spider Mite, Tetranychus urticae - Wisconsin Horticulture, <https://hort.extension.wisc.edu/articles/twospotted-spider-mite-tetranychus-urticae/>
6. Twospotted Spider Mites | Integrated Crop Management, <https://crops.extension.iastate.edu/encyclopedia/twospotted-spider-mites>
7. Spider mites and other pest mites - Biocontrol, Damage and Life Cycle - Koppert Canada, <https://www.koppert.ca/plant-pests/spider-mites-and-other-mites/>
8. Twospotted Spider Mites / Floriculture and Ornamental Nurseries ..., <https://ipm.ucanr.edu/agriculture/floriculture-and-ornamental-nurseries/twospotted-spider-mite/>
9. Spider Mites - Utah State University Extension, https://extension.usu.edu/pests/ipm/notes_ag/veg-spider-mites
10. How to Control Two-Spotted Mites in Tomato Plants Organically, <https://octaorganics.com/blogs/news/how-to-control-two-spotted-mites-in-tomato-plants-organically>
11. Red spider mite: Tetranychus spp - TNAU Agritech Portal :: Crop Protection, https://agritech.tnau.ac.in/crop_protection/tomato/tomato_7.html
12. Two-Spotted Spider Mite - UK Entomology - University of Kentucky, <https://entomology.mgcafe.uky.edu/ef310>
13. Tomato – Spider Mite Damage - Arkansas Cooperative Extension Service, <https://www.uaex.uada.edu/yard-garden/plant-health-clinic/disease-notes/posts/tomato-spider-mite-damage.aspx>
14. Twospotted spider mites in home gardens - University of Minnesota Extension, <https://extension.umn.edu/yard-and-garden-insects/spider-mites>
15. Spider Mites—Botanical Control Formulations - ATTRA – Sustainable Agriculture, <https://attra.ncat.org/publication/spider-mites-botanical-control-formulations/>
16. Spider Mites / Home and Landscape / UC Statewide IPM Program (UC IPM), <https://ipm.ucanr.edu/home-and-landscape/spider-mites/>
17. How to Get Rid of Spider Mites on Plants | The Old Farmer's Almanac, <https://www.almanac.com/pest/spider-mites>
18. Tomato Russet Mites, Spider Mites, Tomato Spotted Wilt Virus, and Melon Aphids, <https://pestadvisories.usu.edu/2021/07/21/tomato-russet-mites-spider-mites-tomato-spotted-wilt-virus-and-melon-aphids/>
19. Problem: Spider Mites on Tomato Brand names appearing in this publication are for product identification purposes only. No endor - Kansas State University, <https://hnr.k-state.edu/extension/horticulture-resource-center/common-pest-problems/document/s/Tomato%20Spider%20Mites.pdf>
20. How to Get Rid of Spider Mites - Garden Design, <https://www.gardendesign.com/how-to/spider-mites.html>
21. The solution for spider mite infestations that I never see mentioned here.... - Reddit, https://www.reddit.com/r/houseplants/comments/t7njzi/the_solution_for_spider_mite_infestations_that_i/
22. Managing Mites On Tomato Crops - Growing Produce, <https://www.growingproduce.com/crop-protection/insect-control/managing-mites-on-tomato-crops/>
23. Tomato-Spider mite | Pacific Northwest Pest Management Handbooks, <https://pnwhandbooks.org/insect/vegetable/vegetable-pests/hosts-pests/tomato-spider-mite>
- 24.

How To Mix Neem Oil For Plants: 3 DIY Neem Oil Spray Recipes - pinch of seeds, <https://pinchofseeds.com/neem-oil-insecticide-fungicide-soil-drench-recipes/> 25. How to Manage Spider Mites in Tomatoes with Predatory Mites - YouTube, <https://www.youtube.com/watch?v=TNMoLhT2A14> 26. Tomato Cultivars Resistant or Susceptible to Spider Mites Differ in Their Biosynthesis and Metabolic Profile of the Monoterpenoid Pathway - PMC, <https://pmc.ncbi.nlm.nih.gov/articles/PMC7952643/> 27. Biological Control of Spider Mites in Tomatoes - Entomology | NC State Extension, <https://entomology.ces.ncsu.edu/biological-control-of-spider-mites-in-tomatoes/> 28. Ecofriendly Management of Two Spotted Spider Mites on Tomato (*Solanum lycopersicum* (Mill.) in Eastern Ethiopia | American Journal of Agricultural and Biological Sciences, <https://thescipub.com/abstract/ajabssp.2024.8.15> 29. Grower Fact Sheet: Twospotted Spider Mite Management Guide for Horticulture Crops - AgriLife Extension Entomology, <https://extensionentomology.tamu.edu/wp-content/uploads/2025/03/twospotted-spider-mite-management-for-horticulture-crops-1.pdf> 30. Sustainable Management of *Tetranychus urticae* and *Trialeurodes* ..., <https://pmc.ncbi.nlm.nih.gov/articles/PMC11431958/> 31. News - How to Use Abamectin - Herbicide, Insecticide, Fungicide - POMAIS, <https://www.bigpesticides.com/news/how-to-use-abamectin/> 32. ADAMA ABAMECTIN, https://www.adama.com/south-africa/sites/adama_south_africa/files/product-documents/2023-07/GHS_LABEL_ADAMA_ABAMECTIN_V2.0.pdf 33. genfarm abamectin 18ec, https://www.genfarm.com.au/sites/genfarm/files/2021-07/documents/LABEL_GNF%20ABAMECTIN%2018EC%20INSECTICIDE-MITICIDE.pdf 34. Abamectin 0.15 EC Insecticide - Graco Fertilizer Company, <https://gracofertilizer.com/wp-content/uploads/2023/08/ABAMECTIN.pdf> 35. US EPA, Pesticide Product Label, NUP-13026 ABAMECTIN 0.15 MITICIDE/INSECTICIDE, 03/15/2022, https://www3.epa.gov/pesticides/chem_search/ppls/000228-00734-20220315.pdf 36. Pesticidal management of two spotted spider mite (*Tetranychus urticae*) infesting tomato under protected cultivation - Indian Agricultural Research Journals, <https://epubs.icar.org.in/index.php/IJPP/article/view/115042> 37. Magnitude of Resistance and Metabolism of Acaricides in Two-spotted Spider Mite, *Tetranychus urticae* Koch on Vegetables in Southern Districts of Tamil Nadu - ARCC Journals, <https://arccjournals.com/journal/indian-journal-of-agricultural-research/A-5932> 38. Preventing spider mites on tomato plants - Grow Tomatoes Easily, <https://growtomatoeseasily.com/preventing-spider-mites-on-tomato-plants/> 39. ToMATo, <https://ppqs.gov.in/sites/default/files/tomato.pdf> 40. Breeding for Tomato Resistance to Spider Mite *Tetranychus urticae* Koch (Acari: Tetranychidae) - UKnowledge - University of Kentucky, https://uknowledge.uky.edu/pss_etds/114/ 41. Sources of Resistance for Two-spotted Spider Mite (*Tetranychus urticae*) in Scarlet (*Solanum aethiopicum* L.) and Gboma (*S. macrocarpon* L.) Eggplant Germplasms in - ASHS Journals, <https://journals.ashs.org/view/journals/hortsci/54/2/article-p240.xml> 42. Tomato most resistant to Spider Mites?, https://www.reddit.com/r/tomatoes/comments/13weemo/tomato_most_resistant_to_spider_mites/ 43. Field Day on Multiple Disease-Resistant Tomato Cultivation | ICAR, <https://www.icar.org.in/en/field-day-multiple-disease-resistant-tomato-cultivation>