

Tomato Target Spot: A Comprehensive Guide to Identification, Management, and Prevention

1. Overview

Tomatoes represent one of the most frequently cultivated, economically valuable, and widely consumed vegetables in the world. They are a staple of global agriculture, prized equally by massive commercial farming operations and backyard home gardeners. However, the successful cultivation of tomatoes is continuously threatened by a wide variety of environmental stressors, insect pests, and microscopic pathogens. Among the most challenging, aggressive, and destructive of these threats is a fungal disease known to plant pathologists and farmers as Target Spot.

Target Spot is a severe foliar (leaf) and fruit disease caused by a highly adaptable fungal pathogen known scientifically as *Corynespora cassiicola*. Historically, this disease was primarily restricted to the tropical and subtropical regions of the world, where constant heat and high humidity provided the perfect breeding ground for fungal spores. For example, in the state of Florida, where the fresh-market tomato industry generates nearly half a billion dollars annually, Target Spot is considered one of the primary limiting factors for crop yields. Similarly, in tropical island nations like Mauritius, where tomatoes are the third most consumed vegetable, sudden outbreaks of Target Spot in super-humid zones have been known to infect up to 80% of entire greenhouse crops in a single season.

However, in recent years, the geographical footprint of Target Spot has expanded significantly. It is no longer just a tropical problem. The disease has increasingly found its way into temperate summer gardens and controlled agricultural environments, such as high-tunnel greenhouses, across the globe. For instance, the disease was recently identified causing significant damage in high-tunnel tomato operations as far north as Indiana. This expansion highlights the urgent need for growers in all climates to understand and recognize this pathogen.

The disease derives its common name from the highly distinct visual pattern it leaves on the foliage and the fruit of the infected tomato plant. As the microscopic fungal infection progresses and consumes the plant's tissue, it creates large, circular, sunken craters and dark brown spots. These spots feature distinct concentric rings, closely resembling a bullseye or a traditional archery target.

If left unmanaged, Target Spot can completely devastate a tomato crop within a matter of weeks. The disease follows a specific upward trajectory. It begins by attacking the lowest leaves closest to the damp soil, eventually causing severe defoliation—a destructive process where the plant becomes so damaged that it sheds its leaves prematurely. Leaves are the solar panels of the plant; without them, the tomato plant loses its ability to generate energy through photosynthesis. Furthermore, without a thick canopy of leaves to shade the developing fruit, the tomatoes become highly susceptible to environmental damage like sunscald (severe sunburn), and the overall harvest is drastically reduced or entirely lost.

Understanding Target Spot requires a careful examination of the precise environmental conditions that allow the fungus to wake up, reproduce, and thrive. The pathogen is highly

dependent on extended periods of continuous moisture and warm temperatures. Consequently, the disease often emerges as a severe, fast-moving problem during wet summer months, in regions characterized by high natural humidity, or in home gardens where poor, overhead watering practices inadvertently create a damp, stagnant microclimate around the plant's leaves.

Because the visual symptoms of Target Spot can very easily be confused with other extremely common tomato ailments—such as Early Blight, Septoria Leaf Spot, or Bacterial Spot—accurate identification is the most critical first step in saving a crop. The approach to managing this disease must be multifaceted. Because the fungus is highly adaptable, there is currently no single "cure-all" spray for Target Spot, nor are there fully immune commercial tomato varieties widely available on the market just yet, though scientists are getting closer.

Instead, successful management relies entirely on an integrated pest management (IPM) strategy. This involves a proactive combination of early symptom recognition, precise environmental control, careful garden sanitation, optimal plant nutrition, and the strategic, timely application of natural home remedies or chemical treatments to protect healthy plant tissue from initial infection. This comprehensive report provides a detailed, easy-to-understand examination of Target Spot, designed specifically for those without a deep background in plant pathology. It explores the biological behavior of the fungus, the precise visual symptoms to look for, and exhaustive, research-backed methodologies for treating active infections and implementing long-term preventative measures to ensure a healthy, highly productive tomato harvest.

2. Symptoms

The successful management of Target Spot relies entirely on early and accurate visual identification. The fungus is incredibly aggressive and is capable of infecting all above-ground components of the tomato plant, including the broad leaves, the main structural stems, the petioles (the small, delicate stalks that attach the leaves to the main stems), and the fleshy fruit itself.

While a plant can be infected at any point in its life cycle, tomatoes are particularly vulnerable during their early, fragile seedling stage, and later, during the highly energy-intensive period when they are actively producing blossoms and ripening fruit.

The disease typically begins deep within the interior of the plant canopy, starting on the oldest, lowest leaves that hang closest to the damp garden soil. Because these initial symptoms are hidden beneath thick layers of green foliage, early detection can be incredibly difficult for the casual observer. Gardeners and farmers must make it a habit to actively part the branches and deeply inspect the interior and lower sections of the plant to catch the disease in its infancy before it explodes outward.

Foliar (Leaf) Symptoms

The destruction of the tomato plant's leaves is the most common and visibly debilitating aspect of a Target Spot infection. The symptom progression on the foliage occurs in distinct, observable stages that escalate as the fungus consumes more tissue:

- **Initial Stage:** The very first signs of an infection are tiny, pinpoint-sized spots appearing on the upper surface of the lower leaves. These spots often appear "water-soaked," meaning they look like tiny, translucent drops of dark liquid trapped just beneath the waxy surface of the leaf. At this very early stage, the plant is already under attack, but the

damage is microscopic.

- **Intermediate Stage:** As the fungus releases enzymes to consume the leaf tissue, the water-soaked points expand into small, dead (necrotic) lesions. At this stage, the spots typically develop a light brown or tan center surrounded by a much darker, harsh brown or black margin. Because these spots are relatively small and dark, they are frequently misidentified by home gardeners as Bacterial Spot, another common tomato disease caused by the *Xanthomonas* bacteria.
- **Advanced Stage:** If the environment remains favorable for the fungus—meaning it stays warm and wet—the lesions will continue to enlarge rapidly. They become distinctively circular, taking on a gray to pale brown color in the center. It is during this advanced stage that the disease's namesake characteristic appears: dark, concentric circles form within the lesion, giving it a distinct "target" or bullseye appearance. Depending on the specific variety of tomato being grown, the plant's immune system may react by producing a bright yellow "halo" of dying, chlorotic tissue immediately surrounding the brown fungal spots.
- **Progression and Defoliation:** As the disease advances unchecked, individual target spots will grow until they touch one another, merging (coalescing) into massive areas of blighted, dead, crispy tissue. Once a leaf is sufficiently damaged by these merged spots, the entire leaf will turn completely yellow, wither, and fall off the plant. Because the disease starts at the bottom and relies on splashing rain to move upward, it creates a visual phenomenon known to farmers as "melting out." The interior and lower canopy of the tomato plant becomes completely bare and stripped of foliage, leaving only a few healthy green leaves struggling at the very top of the plant.

Stem and Petiole Symptoms

While less visually prominent than the large bullseyes on the broad leaves, Target Spot actively degrades the structural framework of the plant, which can lead to catastrophic branch failure.

- The infection on stems and petioles begins exactly like the leaf symptoms: as tiny, brown, pinpoint lesions.
- However, unlike the perfectly round spots on the flat leaves, stem lesions are forced to grow along the vertical axis of the plant. As they develop, they become elongated, oval, or oblong in shape.
- In severe cases, these oblong lesions will expand horizontally until they completely encircle, or "girdle," the stem or petiole. When a stem is girdled, the internal vascular system of the plant is destroyed. The flow of water and nutrients is entirely cut off, causing all the leaves, blossoms, and fruit attached above that specific point to collapse, wilt, and die instantly.

Fruit Symptoms

The most economically devastating aspect of Target Spot is its direct impact on the tomato fruit. While some diseases only hurt the leaves, Target Spot actively rots the fruit, rendering it completely inedible, unsellable, and prone to rapid, total post-harvest decay.

- **Early Fruit Symptoms:** The infection first appears on the smooth skin of the developing tomato as small, slightly sunken brown flecks. To the untrained eye, these tiny pits can closely resemble minor physical or abiotic injury, such as the scars left by blowing sand or small hail (a phenomenon often referred to as sandblasting).
- **Advanced Fruit Symptoms:** As the tomato matures and swells with water, the fungus

continues to aggressively digest the fruit tissue beneath the skin. The small flecks rapidly expand into large, perfectly circular, deeply sunken craters. These craters typically have pale-brown centers and clearly visible concentric rings, perfectly mirroring the bullseye pattern seen on the leaves.

- **Cracking and Rot:** As the fungal crater grows larger, the dead tissue in the center loses its natural elasticity. As the healthy parts of the tomato continue to grow and expand, the dead, rigid tissue in the target spots will tear and crack open. These deep, open cracks bypass the tomato's protective skin, providing a wide-open entry point for secondary bacteria, fruit flies, and other opportunistic fungi. This leads to severe, rapid, and total post-harvest fruit rot. In advanced, highly humid stages, dark-gray to black fuzzy fungal growth (the actual spores of the pathogen) may become visible rising out of the center of the sunken craters.

Distinguishing Target Spot from Similar Diseases

One of the greatest challenges in successfully managing tomato diseases in a home garden or farm is correctly identifying the pathogen. Misidentification leads to the wrong treatments being applied, wasting time and money while the plant continues to die. Target Spot (*Corynespora cassiicola*) is exceptionally easy to confuse with Early Blight (*Alternaria solani*) and Septoria Leaf Spot (*Septoria lycopersici*). All three of these fungal pathogens thrive in similar wet, humid environments, and all three predictably start their attack on the lower leaves of the plant. However, careful observation reveals distinct differences.

- **Differentiating from Early Blight:** Early Blight is the disease most commonly confused with Target Spot, primarily because both pathogens produce brown leaf spots that feature concentric, target-like rings. However, Early Blight lesions are often irregular, asymmetrical, or jagged in shape, whereas Target Spot lesions are typically much more perfectly circular. Furthermore, Early Blight is usually accompanied by a widespread, general yellowing (chlorosis) of the entire leaf structure, whereas the yellowing in Target Spot is generally confined to a distinct, tight halo immediately surrounding the fungal lesion. Finally, while Early Blight can cause leathery, dark spots on the fruit (usually concentrated near the top stem area), Target Spot causes distinctly pitted, deeply sunken craters anywhere on the fruit's surface.
- **Differentiating from Septoria Leaf Spot:** Septoria Leaf Spot is generally easier to distinguish once you know what to look for. Septoria creates much smaller leaf spots—usually only 1/16 to 1/8 of an inch in diameter—compared to the larger half-inch spots of Target Spot. Septoria spots have distinct light tan or gray centers with very dark purple or brown margins, but crucially, they do *not* have concentric rings inside them. If you inspect the center of a Septoria spot closely with a 10X magnifying glass or hand lens, you will see tiny, raised black specks. These are the fungal fruiting bodies, known as pycnidia, which are entirely absent in Target Spot lesions. Additionally, Septoria very rarely infects the tomato fruit itself, restricting its damage almost entirely to the foliage and stems.
- **Differentiating from Late Blight:** Late Blight (*Phytophthora infestans*) is the infamous water mold pathogen responsible for the catastrophic Irish Potato Famine in the 1840s. Despite the confusingly similar name to Early Blight, it behaves entirely differently than Target Spot. Late Blight does not start as small circles with target rings. Instead, it creates massive, irregularly shaped, pale green to brown splotches that look like the plant has suffered sudden frost damage. Furthermore, Late Blight produces a distinct white, fluffy,

fuzzy mold on the underside of the leaves during humid weather, a feature entirely absent in Target Spot. Late Blight is highly aggressive and can strike anywhere on the plant at any time of the season, whereas Target Spot consistently begins at the bottom and works its way up.

3. Causes & Spread

To effectively combat and prevent Target Spot, it is absolutely necessary to understand the biology of the organism causing it, how it manages to survive the freezing winters from year to year, and the specific environmental triggers that cause it to wake up, multiply, and attack the tomato plant.

The Pathogen: *Corynespora cassiicola*

Target Spot is caused by the ascomycete fungus *Corynespora cassiicola*. This microscopic organism has a long and complex history in agricultural science. It was first documented and described by researchers Berkeley and Curtis in 1868, originally classified under the name *Helminthosporium cassiicola* due to its physical characteristics under a microscope. It wasn't until decades later, in 1950, that it was formally reclassified to its current scientific nomenclature by C.T. Wei. While it shares some visual similarities with the *Helminthosporium* family, advanced modern genetic and phylogenetic studies have shown that it actually belongs to the class Dothidiomyceta, sharing a closer evolutionary relationship with the *Pyrenophora* family of fungi. *Corynespora cassiicola* is renowned and feared among plant pathologists for its incredible genetic diversity, immense adaptability, and aggressive nature. Unlike some highly specialized pathogens that have evolved to only infect a single specific type of plant (like certain rusts or mildews), this fungus possesses an enormous, varied host range. It is documented to be capable of infecting over 500 different plant species across the globe. Beyond tomatoes, it is a severe agricultural pest that routinely decimates crops of soybeans, cucumbers, cotton, and papayas. On papayas, for example, it causes small angular spots where the center of the leaf completely falls out, producing a dramatic "shot-hole" effect.

It is also highly capable of infecting a massive variety of common ornamental plants and wild garden weeds. However, recent genetic research indicates that the fungus exhibits a high degree of what is known as "host specialization." This means that over time, different populations of the fungus adapt to their preferred food source. A strain of the fungus that has spent generations adapting to attack soybeans is most dangerous to other soybeans, while strains that have adapted to tomatoes are exceptionally aggressive and fast-moving when they find another tomato plant, though they can cross over if necessary.

Pathogen Survival Mechanisms

The incredible resilience of the Target Spot fungus lies in its ability to operate in three entirely distinct biological roles depending on the environment and the availability of resources. This versatility makes it incredibly difficult to eradicate completely from a farm or garden.

- **As a Necrotroph:** When actively infecting a living tomato plant, the fungus acts as a necrotroph. This means it actively kills the living plant cells using secreted chemical enzymes and toxins, and then feeds on the nutrients released by the dead tissue. This active, localized killing of plant cells is exactly what causes the visible brown spots, yellow

halos, and deep craters on the plant.

- **As a Saprophyte:** When the active summer growing season ends, the temperatures drop, and the tomato plant ultimately dies, the fungus does not perish with it. Instead, it undergoes a biological transition, shifting into a saprophyte. A saprophyte is an organism that acts like nature's recycling crew, surviving by digesting dead, decaying organic matter. In this state, the pathogen can safely overwinter within the dead tomato leaves, discarded stems, and fallen rotting fruit left behind in the garden soil. It is remarkably durable in this dormant state, capable of surviving deep within the soil and crop debris for up to two full years, patiently waiting for a new, susceptible tomato plant to be planted nearby.
- **As an Epiphyte:** Perhaps its stealthiest and most insidious survival mechanism is its ability to act as an epiphyte. An epiphyte is an organism that lives on the physical surface of another plant without actually infecting it, feeding from it, or harming it in any way. *Corynespora cassiicola* can live completely harmlessly on the surface of common garden weeds, such as nightshade or horsenettle. This allows the local, unseen weed population surrounding a garden to act as a hidden reservoir—a constant source of infectious inoculum—for the fungus. When the weather conditions are right, the fungus will easily jump from the unharmed weed directly onto a highly susceptible tomato plant.

Environmental Conditions for Infection

Like all fungi, *Corynespora cassiicola* requires a highly specific set of environmental conditions to wake up from its dormancy, produce infectious spores, and successfully invade a host plant. Outbreaks of this disease are almost entirely dictated by two factors: moisture and temperature.

- **Temperature Requirements:** The fungus is active and capable of growing across a moderately broad temperature spectrum, generally surviving anywhere between 60°F and 90°F (15°C to 32°C). Laboratory studies testing various strains of the fungus have shown that if temperatures drop below 60°F or spike drastically above 90°F, the biological growth of the fungus slows down considerably, though the organism does not die. However, the fungus is at its absolute most aggressive, and the disease develops most rapidly, in the optimal "Goldilocks" temperature zone of 68°F to 82°F (20°C to 28°C). This explains why the disease often explodes during warm, mid-summer days.
- **Moisture and Leaf Wetness:** While temperature dictates the overall speed of the disease, moisture is the absolute trigger required for infection. Target Spot requires exceptionally high ambient humidity levels (often upwards of 70%) and, most importantly, continuous, unbroken periods of "leaf wetness". For the microscopic fungal spores to germinate, grow a physical appendage called a germ tube, and penetrate the tough, waxy outer skin of the tomato leaf, the leaf surface must remain physically covered in a film of water for an extended, uninterrupted period ranging from 16 to 44 hours. If the leaf dries out after only 10 hours, the spore will fail to penetrate and the infection cycle will be broken. This sustained moisture allows the spore to anchor itself, dissolve the plant's defenses, and burrow its way into the vulnerable interior tissue.

How the Disease Spreads

Once a fungal spore successfully penetrates and infects a leaf, it immediately begins to feed and grow. Within just a few days, the fungal colony will mature and produce a massive new generation of reproductive spores (called conidia) right in the center of the brown target spot.

Under a light microscope, these spores appear as long, thin, pale brown, cylindrical shapes that are straight or slightly curved, featuring multiple internal segments (pseudosepta). They are entirely invisible to the naked eye.

The primary mechanisms for spreading these newly minted spores to other plants are wind and water.

- **Wind:** Airborne conidia are the primary means of spread across large distances. The dry spores are incredibly lightweight. They are easily picked up by passing breezes and can be carried significant distances across a massive commercial farm or a neighborhood community garden, eventually landing on neighboring, perfectly healthy plants. * **Rain and Irrigation Splash:** Water is the primary driver of the disease moving vertically up the plant. When heavy raindrops or water droplets from an overhead sprinkler hit an infected leaf, the physical impact of the water dislodges the microscopic spores and splashes them forcefully upward and outward onto higher foliage. Because the disease overwinters in the soil and starts on the lowest leaves, this mechanical rain splash is the primary mechanism that drives the infection higher and higher up the plant canopy with every passing summer storm. This splashing effect is also what moves spores from the infected soil debris directly onto the lowest hanging leaves, starting the initial infection cycle in the spring.

Furthermore, any physical damage to the plant drastically increases its vulnerability to these windblown and splashed spores. Tomato fruit that has been wounded by blowing sand, heavy hail, insect bites, or aggressive handling by gardeners has broken its natural, protective skin. These open wounds bypass the need for the fungus to dissolve the outer layer, allowing the fungal spores direct, effortless entry into the fleshy interior, practically guaranteeing an immediate and severe infection.

4. Treatment / Remedies

Managing an active outbreak of Target Spot requires swift, decisive, and consistent action. It is vital to understand that once the fungus has successfully penetrated the plant tissue and a brown, necrotic lesion has formed, that specific piece of plant tissue is entirely dead. It cannot be healed, reversed, or turned green again by any treatment. The overarching goal of all treatments—whether they are natural home remedies or synthetic chemical sprays—is to stop the existing fungal spores from germinating and spreading to the healthy, unaffected parts of the plant, thereby slowing the progression of the disease enough to allow the plant to produce a secure harvest.

Because the fungus relies so heavily on long periods of wetness to travel and infect, treatment efficacy is heavily dependent on the weather. Protective sprays must be reapplied periodically throughout the growing season, especially immediately following heavy rain events that can wash the protective chemical coating off the leaves, leaving the plant vulnerable once again.

Natural / Home Remedies

For home gardeners, organic farmers, or those seeking to minimize their reliance on synthetic chemical applications, several highly effective natural remedies exist. These organic solutions rely on biological science: altering the pH of the leaf surface to make it inhospitable, physically disrupting the cellular walls of the fungal spores, or introducing highly competitive, beneficial predatory microbes that protect the plant.

- **Hydrogen Peroxide Sprays:** Hydrogen peroxide is an inexpensive, accessible, and highly effective natural remedy for stopping active fungal spread. Plants actually naturally produce small amounts of hydrogen peroxide internally as a defense mechanism to fight off bacterial and fungal infections on their own; introducing a diluted spray gives the plant a massive defensive boost. When a standard 3% hydrogen peroxide solution is sprayed onto an infected tomato plant, it acts as a powerful oxidizing agent. The extra oxygen molecules react violently with the cellular walls of the fungus and bacteria on the surface of the leaf, essentially burning and destroying them on contact. The ultraviolet rays of the sun then quickly activate and break down the hydrogen peroxide into completely harmless water and oxygen, leaving absolutely no toxic residue behind on the food. A highly recommended home recipe involves mixing 8 ounces of standard 3% hydrogen peroxide into one gallon of water. This mixture must be applied thoroughly to the top and bottom of the leaves, and the application should be repeated after any rain events, and generally every two weeks as a maintenance routine.
- **Baking Soda Solutions:** Baking soda (sodium bicarbonate) is a traditional, time-tested preventative treatment that works by dramatically altering the pH of the leaf surface. Fungal spores prefer a slightly acidic environment to germinate successfully. By coating the leaf in an alkaline baking soda solution, the gardener creates a harsh, inhospitable environment where the spores simply cannot survive, take root, or grow a germ tube. A widely used, effective home recipe involves mixing 3 tablespoons of standard baking soda with 1 gallon of water. However, because water simply rolls off the waxy surface of a tomato leaf, sticking agents must be used. Gardeners should gently mix in 1 tablespoon of vegetable oil (acting as a natural binder to stick the powder to the leaf) and 2 drops of mild liquid dish soap (acting as a surfactant to break the surface tension, allowing the liquid to spread evenly across the entire leaf). This solution must be sprayed until the leaves are dripping wet, meticulously targeting both the upper and lower surfaces.
- **Organic Copper Fungicides:** Copper has been utilized in agriculture as an antimicrobial agent for centuries and remains one of the most potent, reliable organic treatments available today. Commercial products utilizing active ingredients like copper octanoate, basic copper sulfate, or copper sulfide are heavily regulated but approved for organic use. They work by releasing active copper ions that penetrate the fungal spores and disrupt their internal enzymes and proteins, effectively killing the organism. Copper is considered a broad-spectrum protectant, meaning it will also guard the plant against destructive bacterial diseases like Bacterial Spot and Bacterial Speck. This is highly beneficial, as distinguishing between fungal and bacterial spots with the naked eye in the field can be incredibly difficult. While highly effective, copper must be used carefully and strictly according to the manufacturer's label instructions, as heavy, continuous overuse can lead to copper building up to toxic levels in the garden soil over many years. Formulations like Liquid Copper Fungicide by Southern Ag are highly concentrated and cost-effective for small-scale growers.
- **OMRI-Listed Biofungicides:** The Organic Materials Review Institute (OMRI) is a highly respected organization that reviews and certifies products for use in certified organic farming. They have approved several highly advanced, naturally derived products for disease management that rely on cutting-edge biological science.
 - **Polyoxin D Zinc Salts (Oso):** This is a novel, naturally derived biofungicide. Field trials conducted by university extensions have shown that weekly sprays of Oso at 6.5 fluid ounces per acre, beginning at the very first sign of symptom onset, provide a highly significant reduction in the severity of leaf spots.

- **Botanical Extracts (EcoSwing & Sporatec):** These products utilize natural plant oils that suppress fungal growth. EcoSwing utilizes an 82% extract of the *Swinglea glutinosa* plant, while Sporatec relies on a potent blend of rosemary, clove, and thyme essential oils.
- **Beneficial Microbes:** Products containing living bacteria or fungi—such as *Bacillus subtilis* (sold under brands like Serenade, Cease, and Taegro), *Bacillus mycoides* (Lifegard), or *Trichoderma asperellum*—introduce massive numbers of harmless, beneficial microbes to the tomato plant. These biological products work through competitive exclusion. The beneficial microbes rapidly colonize the leaf surface, consuming all available physical space and trace nutrients. When the Target Spot spores eventually land on the leaf, they are entirely out-competed and starve to death before they can cause an infection. Furthermore, products like Lifegard activate "induced systemic resistance" (ISR), essentially acting like a vaccine that tricks the tomato plant's immune system into turning on its natural defenses before the actual pathogen attacks.
- **Biosurfactants (Zonix):** Products containing rhamnolipid biosurfactants, extracted from the bacterium *Pseudomonas aeruginosa*, physically disrupt and burst the cellular walls of the fungal spores upon contact.

Chemical Remedies

In large-scale commercial agricultural settings, or in severe home garden outbreaks where natural remedies fail to halt the explosive progression of the disease, synthetic chemical fungicides remain the primary, most reliable method of control.

- **Protectant Fungicides:** The most common and reliable chemical treatments utilize powerful protectant active ingredients, most notably chlorothalonil (commonly sold to home gardeners under the brand name Daconil) and mancozeb. These chemicals are "non-systemic." This means they do not absorb into the plant's vascular system or travel through the sap. Instead, they act like a coat of paint, forming a protective, highly toxic chemical shield on the exterior surface of the leaf. When a windblown fungal spore lands on a leaf coated in chlorothalonil, the chemical immediately prevents the spore from germinating, stopping the infection before it starts. Because these are purely contact barriers, the application technique is everything. Achieving 100% full coverage of all plant surfaces—including the often-ignored underside of the leaves and deep within the dense inner canopy—is absolutely mandatory. If a spore lands on an untreated patch of leaf, the plant will become infected. Note that mancozeb has strict harvest intervals and generally cannot be applied within 5 days of harvesting fruit to ensure consumer safety.
- **Application Routines:** Chemical fungicides must be applied preventatively to be effective. The protective layer must be present on the plant *before* the spore lands. Applications generally begin early in the season, just before the weather becomes highly favorable for the disease (humid and warm), or immediately upon spotting the very first tiny, pinpoint lesion. The sprays must be reapplied at regular, strict intervals, typically every 7 to 14 days. This consistent routine ensures that all newly grown, fresh foliage is instantly protected, and it replaces any chemicals that have naturally degraded in the intense summer sun or washed away in heavy rainstorms.
- **The Crisis of Fungicide Resistance:** A major, growing crisis in the global management of Target Spot is the pathogen's incredibly rapid development of chemical resistance. Just as human bacteria can become immune to overused antibiotics, *Corynespora cassiicola*

has a highly diverse and adaptable genetic makeup. This allows it to rapidly mutate and survive continuous exposure to certain synthetic chemicals. In recent years, widespread, documented resistance has been discovered against a specific, highly advanced class of chemicals known as QoI fungicides (also known as strobilurins, which fall under the Fungicide Resistance Action Committee classification FRAC Group 11). Because the fungus in many agricultural regions is now completely immune to these once-powerful chemicals, agricultural university extensions strongly advise against relying on Group 11 fungicides for Target Spot control. Growers are encouraged to utilize older, multi-site protectants like chlorothalonil, which are harder for the fungus to mutate against, or use carefully planned chemical rotation programs utilizing newer active ingredients like azoxystrobin, pyraclostrobin, or boscalid to ensure ongoing efficacy and prevent further resistance.

5. Preventive Measures

Because treating an active fungal infection is exceptionally difficult, labor-intensive, and expensive, the absolute cornerstone of all Target Spot management is prevention. By deliberately altering the environment, physically disrupting the pathogen's life cycle, and selecting tough plant genetics, growers can drastically reduce the likelihood of a severe outbreak ever occurring.

Crop Rotation and Sanitation

The easiest and most organic way to prevent a severe disease outbreak is to ensure the pathogen is simply not present in the soil when the spring planting season begins. Because *Corynespora cassiicola* is a robust saprophyte that can survive by eating dead plant debris for up to two full years, planting tomatoes in the exact same spot year after year is a guaranteed recipe for agricultural disaster.

- **Rotation Strategies:** Growers must practice strict, multi-year crop rotation. Tomatoes should never be planted in the same plot of soil for at least two, and ideally three, consecutive years. Furthermore, because the Target Spot fungus possesses such a wide host range, tomatoes should not be rotated with other susceptible plants. Gardeners must avoid planting tomatoes where other Solanaceous crops (the nightshade family, which includes peppers, eggplants, or potatoes) or cucurbits (like cucumbers, melons, or squash) were recently grown, as they all share the same diseases. Instead, rotate the tomato plot with completely immune crops like sweet corn, beans, peas, or brassicas (cabbage, broccoli) to naturally starve the fungus completely out of the soil profile over time.
- **End-of-Season Sanitation:** At the end of the autumn harvest, the garden must be meticulously cleaned. All dead tomato vines, fallen leaves, and rotting, unharvested fruit must be completely removed from the growing area. This material should be burned, thrown away in municipal trash, or tilled deeply into the earth to accelerate rapid decomposition. It should not be put in a cold home compost pile, as the temperatures may not get hot enough to kill the spores. Leaving a dead, infected tomato plant standing in the garden through the winter guarantees a massive release of fresh fungal spores the following spring.
- **Weed Management:** Because the fungus can live silently as an epiphyte on the surface

of common weeds without killing them, maintaining a completely weed-free perimeter around the tomato patch eliminates this hidden reservoir of disease.

Genetic Resistance and Variety Selection

Planting disease-resistant varieties is universally considered by horticulturists to be the most efficient, cheapest, and eco-friendly method of disease management. However, for decades, the tomato industry has faced a major hurdle: there have been zero commercial tomato varieties with known, documented genetic resistance specifically tailored to Target Spot.

- **The Breeding Breakthrough:** In recent years, researchers at agricultural institutions, most notably at the University of Florida's Gulf Coast Research and Education Center, have made massive, exciting breakthroughs. By screening ancient, wild ancestors of the modern domesticated tomato—specifically wild accessions like *Solanum galapagense* (originating from the Galapagos Islands), *Solanum cheesmaniae*, and *Solanum pimpinellifolium*—scientists have discovered hidden genetic traits that provide natural immunity to Target Spot. These genetic markers are known as quantitative trait loci (QTLs). When these tiny, wild plants were tested in rigorous field trials, they successfully fought off the Target Spot fungus entirely on their own, without the need for any chemical fungicide sprays. Plant breeders are currently in the painstaking, multi-year process of "introgression." This involves cross-breeding these highly resistant wild plants with modern, large, tasty commercial tomato hybrids over many generations. The goal is to transfer the resistance genes while keeping the desirable traits of a large, slicing tomato.
- **Current Variety Selection:** While specific *Corynespora cassiicola* resistant varieties are still undergoing final development and commercialization, growers in the meantime should proactively select robust hybrid varieties that carry broad-spectrum disease resistance. Varieties that have proven to be exceptionally tough against other fungal wilts (like Fusarium or Verticillium wilt) and blights tend to possess overall greater plant vigor, allowing them to survive environmental stress and fungal attacks longer.
 - **Modern Hybrids:** For extreme resilience, modern hybrids are highly recommended. 'Chef's Choice Red' is an award-winning hybrid noted for highly disease-resistant vines. 'Celebrity' is widely considered a benchmark for success in difficult conditions due to its semi-determinate, highly resistant nature. Other highly resistant hybrids include Granadero, Juliet, and Early Girl.
 - **Resistant Heirlooms:** Heirloom varieties are prized worldwide for their superior, complex flavor profiles, but they are notoriously finicky and highly susceptible to fungal diseases. However, some heirlooms possess excellent natural resistance to a variety of blights and wilts, making them safer choices. Varieties like Cherokee Purple, Roma, Ace 55, Brandywine, Green Zebra, Marglobe, and Mortgage Lifter all carry documented resistance to various common wilts, root-knot nematodes, and blights, providing them a fighting chance in a difficult summer.

Spacing and Airflow Management

Because Target Spot spores strictly require 16 to 44 hours of continuous leaf wetness to germinate successfully, manipulating the physical space around the plant to ensure rapid drying is a highly powerful, entirely free preventative tool.

- **Plant Spacing:** Tomatoes must be planted with adequate physical space between them, generally 24 to 36 inches apart depending on the variety. Crowding plants tightly together

to maximize space is a false economy; it traps humid air, prevents drying sunlight from reaching the lower leaves, and stops prevailing winds from penetrating the canopy.

- **Staking and Trellising:** Tomato plants should absolutely never be allowed to sprawl wildly on the bare earth. Sprawling places the most vulnerable leaves directly in physical contact with the damp soil and the millions of overwintering fungal spores hidden within it. Plants must be lifted upward and supported using wooden stakes, heavy-duty wire cages, or vertical string trellises (often used in high tunnels). Growing vertically pulls the leaves high into the path of drying winds and maximizes intense sunlight exposure, which works together to evaporate morning dew rapidly. Choosing open, elevated planting sites on hills or exposed areas where prevailing winds can sweep through the foliage constantly is almost as effective as chemical fungicide programs in certain weather conditions.

6. Suggestions & Best Practices

Beyond direct disease treatment and genetic selection, the everyday cultural habits and routines of the farmer or gardener play a massive, outsized role in dictating the ultimate severity of a Target Spot outbreak. How a plant is watered, how it is fed, and how it is pruned directly influences its microclimate and vulnerability.

Irrigation Practices

Watering techniques are the single most controllable factor in Target Spot management. Since mechanical rain splash and extended leaf wetness are the primary drivers of the disease, introducing water to the garden incorrectly will literally manufacture an outbreak out of thin air.

- **Avoid Overhead Watering:** Traditional lawn sprinklers or hose nozzle attachments that spray water indiscriminately over the top of the plant canopy should be strictly and permanently avoided in the tomato patch. Overhead watering artificially mimics a rainstorm. It splashes invisible spores from the dirt directly onto the lower leaves, spreads existing spores across the canopy, and systematically washes off expensive, carefully applied protective fungicide coatings.
- **Implement Drip Irrigation:** The absolute gold standard for tomato cultivation, utilized by all major commercial operations, is drip irrigation or the use of porous soaker hoses. These efficient systems deliver water slowly and directly to the soil exactly at the base of the plant. This method deeply soaks the root zone while leaving the entire above-ground foliage completely bone dry, entirely robbing the fungus of the moisture it needs to germinate. For advanced subsurface irrigation, maintaining a constant water table between 18 and 24 inches below the top of the bed is recommended.
- **Timing of Watering:** If drip irrigation is unavailable and overhead watering with a hose or watering can is absolutely unavoidable, it must be done exclusively in the early morning, just as the sun is rising. This precise timing allows the intense heat of the morning sun and the daytime breezes to quickly evaporate the excess moisture off the leaves long before the 16-hour fungal germination window can be met. Watering in the late afternoon or evening is highly destructive. The leaves will remain wet throughout the cool, dark, stagnant night, providing the exact prolonged wetness the fungus requires to launch a successful attack.

Fertilization and Canopy Management

The nutritional health of the tomato plant dictates how it grows, which in turn heavily influences its microclimate. Proper fertilization is a delicate balancing act that requires adherence to the "4Rs" of nutrient management: right rate, right timing, right placement, and right source.

- **The Danger of Excess Nitrogen:** While nitrogen is an essential building block for leafy green plant growth, over-fertilizing a tomato plant with excessive nitrogen creates a highly dangerous scenario for disease management. High nitrogen forces the plant into overdrive, producing a massive, lush, incredibly dense canopy of vegetative leaves. A canopy that is too thick acts like a dense jungle umbrella; it blocks out sunlight and wind, preventing the interior and lower leaves from ever drying out. Field research indicates that Target Spot is significantly more severe and harder to control on plants with these overly lush canopies, simply because the dense leaves physically trap humidity and dry far too slowly.
- **Fertigation and Balance:** To prevent explosive, overly dense vegetative growth, commercial growers utilize a technique known as "fertigation"—the process of injecting precise, highly diluted doses of liquid fertilizer directly into the drip irrigation water lines. This allows the grower to feed the plant small, easily digestible amounts on a daily or weekly basis, rather than dumping large amounts of granular fertilizer into the soil all at once, which causes growth spikes. A balanced approach utilizing proper ratios is required. For soils testing low in nutrients, application rates up to 150 lbs of P₂O₅ (Phosphorus) and 200 lbs of K₂O (Potassium) per acre are appropriate, alongside carefully metered Nitrogen, ensuring a strong structural plant with heavy fruit set without generating a dangerously overgrown, disease-prone canopy.

Pruning and Monitoring

Active, physical intervention throughout the growing season helps drastically mitigate disease pressure before it gets out of control.

- **Bottom Pruning:** As the indeterminate tomato plant grows vertically up its stake or trellis, the lowest branches closest to the soil should be purposefully and cleanly pruned off. Removing the bottom 10 to 12 inches of foliage creates a physical air gap between the damp soil and the plant canopy. This gap dramatically improves air circulation at ground level, allowing the soil to dry faster, and completely removes the leaves that are most likely to be hit by splashing rain containing soil-borne fungal spores.
- **Stem Pruning:** Depending on the variety, tomato plants should be pruned to maintain a single, double, or triple main stem. Removing excess "suckers" (the small shoots that grow in the "V" joint between the main stem and a branch) prevents the plant from becoming an unmanageable, dense bush, further improving critical airflow.
- **Scouting and Leaf Removal:** Gardeners must actively monitor (scout) their plants at least once a week. This involves physically parting the branches to inspect the hidden interior and the lowest, oldest leaves. If a single leaf is found exhibiting the characteristic water-soaked spots, tiny pinpoint lesions, or advanced bullseye patterns of Target Spot, it must be immediately pinched or snipped off and removed from the garden entirely. Removing the very first infected leaves physically extracts the millions of developing, microscopic spores from the environment before the wind or the next rainstorm can spread them higher up the plant, effectively short-circuiting the disease cycle.

7. References / Sources

The information synthesized within this report is derived from extensive field research, agricultural extension publications, and plant pathology data provided by leading academic and agricultural institutions. Core data regarding the biological lifecycle of *Corynespora cassiicola*, optimal environmental conditions for infection, and visual symptom progression were sourced from the University of Florida Institute of Food and Agricultural Sciences (UF/IFAS), including research by G. Vallad, L.J. Smith, and L. Datnoff, as well as extensive agronomic spotlights published by Bayer Crop Science. Information detailing the emergence of Target Spot in new geographical locations, such as Indiana and Mauritius, was provided by the Purdue University Vegetable Crops Hotline and the journal *Plant Disease* (APS Journals).

Strategies for organic and chemical disease management, including the efficacy of OMRI-listed biofungicides and the growing concern of QoI (strobilurin) fungicide resistance, were heavily informed by the Cornell Cooperative Extension (including field trials at the Hudson Valley Farm Hub), the University of Connecticut IPM program (J. Boucher), and the University of Kentucky College of Agriculture, Food and Environment. Insights into genetic resistance breeding programs, specifically the introgression of wild tomato traits from *Solanum galapagense* and *Solanum pimpinellifolium*, were drawn from doctoral research conducted at the UF/IFAS Gulf Coast Research and Education Center (E. Sierra) and the *Journal of the American Society for Horticultural Science*. Finally, best practices regarding fertigation, irrigation management, and cultural prevention techniques were corroborated by data from the University of California Agriculture and Natural Resources (UC ANR), Mississippi State University Extension, and the USDA National Institute of Food and Agriculture.

8. Key Takeaways

- **Accurate Identification is Crucial:** Target Spot (*Corynespora cassiicola*) is distinctively characterized by brown, circular lesions featuring concentric rings (resembling an archery bullseye) on the foliage, and large, deeply pitted, cracking craters on the fruit. It consistently begins its attack on the oldest, lowest leaves and progresses vertically upward.
- **Distinguish from Similar Pathogens:** Do not confuse it with Early Blight or Septoria. Target Spot lesions are highly circular without widespread general leaf yellowing (unlike Early Blight). Unlike Septoria (which has tiny spots with no rings), Target Spot features large concentric rings and actively rots the tomato fruit itself.
- **Moisture is the Absolute Trigger:** The fungus strictly requires 16 to 44 hours of continuous, unbroken leaf wetness and warm temperatures (optimally 68°F to 82°F) to successfully infect the plant. Keeping the foliage dry is the single most critical preventative measure.
- **Water at the Base:** Never use overhead sprinklers, which splash spores and keep leaves wet. Utilize drip irrigation or soaker hoses to apply water directly to the soil. If overhead watering is absolutely necessary, do it exclusively in the early morning to allow rapid daytime evaporation.
- **Sanitation Breaks the Cycle:** The pathogen survives as a saprophyte for up to two years on dead plant material in the soil. Strict, multi-year crop rotation (avoiding other nightshades) and the complete removal and destruction of all dead vines and debris at

the end of the season are mandatory.

- **Natural Interventions Work:** For organic management, early and consistent intervention with 3% hydrogen peroxide sprays, baking soda solutions, organic copper, or OMRI-listed biofungicides (like *Bacillus* species and Polyoxin D) can effectively halt the spread of the fungus by outcompeting it or destroying the cell walls.
- **Chemical Care Requires Complete Coverage:** If utilizing chemical protectants like chlorothalonil, applications must be made preventatively and reapplied every 7 to 14 days. Avoid Group 11 (QoI) strobilurin fungicides, as the pathogen has developed widespread, documented chemical resistance to them globally.
- **Manage the Canopy Actively:** Avoid excessive nitrogen fertilization. Over-feeding creates a massive, overly lush canopy that blocks drying wind and sunlight, physically trapping humidity and encouraging rapid fungal growth. Prune the lowest leaves to create an air gap and improve airflow near the damp soil.
- **The Future is Genetic Resistance:** While no fully immune commercial varieties currently exist on the market, ongoing breeding programs are successfully integrating natural Target Spot immunity discovered in ancient, wild tomato ancestors into future commercial hybrid varieties. In the meantime, select vigorous hybrids or heirlooms with broad-spectrum disease resistance.

Works cited

1. A Review of *Corynespora cassiicola* and Its Increasing Relevance to Tomato in Florida, <https://apsjournals.apsnet.org/doi/10.1094/PHP-05-18-0023-RV>
2. Target Spot of Tomato - Vegetables by Bayer, <https://www.vegetables.bayer.com/ca/en-ca/resources/agronomic-spotlights/target-spot-of-tomato.html>
3. An overview of target spot of tomato caused by *Corynespora cassiicola*, https://repository.lsu.edu/plantcrop_pubs/105/
4. First Report of Target Spot on Tomato Caused by *Corynespora cassiicola* in Mauritius, <https://apsjournals.apsnet.org/doi/10.1094/PDIS-05-20-1119-PDN>
5. New Disease Report – Target Spot of Tomato | Purdue University Vegetable Crops Hotline, <https://vegcropshotline.org/article/new-disease-report-target-spot-of-tomato/>
6. Common tomato diseases in the garden and how to manage them | Good Growing, <https://extension.illinois.edu/blogs/good-growing/2025-07-11-common-tomato-diseases-garden-and-how-manage-them>
7. Foliar Diseases of Tomato: Early Blight - Septoria Leaf Spot - Plant Pathology, <https://plantpathology.mgcafe.uky.edu/sites/plantpathology.ca.uky.edu/files/PPFS-VG-25.pdf>
8. Bacterial Spot of Pepper and Tomato | NC State Extension Publications, <https://content.ces.ncsu.edu/bacterial-spot-of-pepper-and-tomato>
9. Early blight in tomato and potato | UMN Extension, <https://extension.umn.edu/disease-management/early-blight-tomato-and-potato>
10. Tomato - Early Blight and Septoria Leaf Spot - Kansas State University, <https://hnr.k-state.edu/extension/horticulture-resource-center/common-pest-problems/documents/Tomato%20-%20Early%20Blight%20and%20Septoria%20Leaf%20Spot.pdf>
11. Tomato Plant Diseases and How to Stop Them - USDA, <https://www.usda.gov/about-usda/news/blog/tomato-plant-diseases-and-how-stop-them>
12. Bullseye! Scientists Center in on Tomato Plants With Resistance to Target Spot, <https://www.growingproduce.com/crop-protection/disease-control/bullseye-scientists-center-in-on-tomato-plants-with-resistance-to-target-spot/>
13. Take the Right Aim to Tame Target Spot of

Tomato - Growing Produce,
<https://www.growingproduce.com/vegetables/take-the-right-aim-to-tame-target-spot-of-tomato/>
 14. What You Need to Know About Tomato IPM | Integrated Pest Management | College of Agriculture, Health and Natural Resources | University of Connecticut,
<https://ipm.cahnر.uconn.edu/what-you-need-to-know-about-tomato-ipm/> 15. Tomatoes - Nutrient Management Program - University of Florida, Institute of Food and Agricultural Sciences - UF/IFAS, <https://bnp.ifas.ufl.edu/crop-recommendations/tomatoes/> 16. Tomato Leaf Spot Diseases - University of Vermont,
<https://www.uvm.edu/extension/news/tomato-leaf-spot-diseases> 17. Fact sheet - Tomato target spot (163) - Lucid Apps,
https://apps.lucidcentral.org/ppp/text/web_full/entities/tomato_target_spot_163.htm 18. PP351/PP351: Target Spot of Tomato in Florida - Ask IFAS,
<https://ask.ifas.ufl.edu/publication/PP351> 19. Managing Tomato Diseases, Disorders, and Pests | Yard and Garden - Iowa State University,
<https://yardandgarden.extension.iastate.edu/how-to/managing-tomato-diseases-disorders-and-pests> 20. How to Identify, Control and Prevent Blight on Your Tomatoes - GardenTech,
<https://www.gardentech.com/blog/pest-id-and-prevention/fight-blight-on-your-tomatoes> 21. Identification of Resistance to Target Spot of Tomato Caused by *Corynespora cassicola* in Wild Tomato Accessions in - ASHS Journals,
<https://journals.ashs.org/view/journals/jashs/149/2/article-p99.xml> 22. Characteristics of *Corynespora cassicola*, the causal agent of tobacco *Corynespora* leaf spot, revealed by genomic and metabolic phenomic analysis - PMC,
<https://pmc.ncbi.nlm.nih.gov/articles/PMC11306238/> 23. How to Prevent Tomato Plant Fungal Diseases (Leaf Examples & 4 Sprays) - YouTube,
<https://www.youtube.com/watch?v=oEmY2aHUuoA> 24. What is the most effective homemade fungicide for tomatoes? - Reddit,
https://www.reddit.com/r/tomatoes/comments/14shrcu/what_is_the_most_effective_homemade_fungicide_for/ 25. Tomato Blight Treatment & Identification - Roots and Refuge - Jessica Sowards, <https://rootsandrefuge.com/tomato-blight-treatment/> 26. Hydrogen Peroxide & Baking Soda Home Remedies for Garden Tomato Plant Diseases: Mix, Routine, Theory - YouTube,
<https://www.youtube.com/watch?v=OANcSiKZzyQ> 27. Tomato Blight (Identification, Prevention and Treatment) - Mlgardener,
<https://miggardener.com/blogs/blog/the-difference-between-early-and-late-blight> 28. Efficacy of Selected Fungicides Listed for Organic Production for Tomato Disease Management | eOrganic,
<https://eorganic.org/node/35737> 29. How to Manage Tomato Leaf Spot Diseases - LSU AgCenter, <https://www.lsuagcenter.com/profiles/mhferguson/articles/page1590779196725> 30. Managing Field Tomato Diseases with OMRI-Listed Fungicides 2024 Annual Report, <https://hvfarmhub.org/wp-content/uploads/2025/08/2024-Field-Tomato-OSO-Report-.pdf> 31. PPFS-VG-43-organic tomato spray guide (2-21-26) - Plant Pathology,
<https://plantpathology.mgcafe.uky.edu/sites/plantpathology.ca.uky.edu/files/PPFS-VG-43.pdf> 32. Biopesticides for Managing Diseases of Tomatoes Organically - Cornell Vegetables,
<https://www.vegetables.cornell.edu/pest-management/disease-factsheets/biopesticides/biopesticides-for-managing-diseases-of-tomatoes-organically/> 33. Selected Organic Fungicides & Bactericides Labeled for Vegetable Bedding plants/transplants - Integrated Pest Management - University of Connecticut,
<https://ipm.cahnр.uconn.edu/wp-content/uploads/sites/3216/2022/12/2020organicfungicidevegetransplantsfinal.pdf> 34. Need for disease resistance breeding against *Corynespora cassicola* in crops - Frontiers,

<https://www.frontiersin.org/journals/agronomy/articles/10.3389/fagro.2023.1275906/full> 35.
Tomatoes 'Я' Us 2024 Tomato Planting List Updated 01/16/24,
<https://extension.arizona.edu/sites/extension.arizona.edu/files/programs/Tomato%20List.pdf> 36.
TOMATO VARIETIES 2025,
<https://ucanr.edu/sites/default/files/2025-03/tomato-varieties-2025.pdf> 37. 5 Tomatoes You Have
to Grow in 2025 - Rural Sprout, <https://www.ruralsprout.com/best-tomato-varieties-2025/> 38.
Disease Resistant Heirloom Tomato Varieties,
<https://www.marysheirloomseeds.com/blogs/news/disease-resistant-heirloom-tomato-varieties>
39. Fertilization / Tomato / Agriculture - UC IPM,
<https://ipm.ucanr.edu/agriculture/tomato/fertilization/> 40. tomato-agronomics-workbook.pdf -
Netafim USA,
<https://www.netafimusa.com/globalassets/tomato-agronomics-workbook.pdf?v=4aeae4> 41.
Fertigation: The Basics of Injecting Fertilizer for Field-Grown Tomatoes | Mississippi State
University Extension Service,
<https://extension.msstate.edu/publications/fertigation-the-basics-injecting-fertilizer-for-field-grow>
n-tomatoes 42. CA Fertilization Guidelines - Processing Tomatoes - CDFA - FREP,
<https://www.cdfa.ca.gov/is/ffldrs/frep/FertilizationGuidelines/Tomato.html>