Question	A fibrous root system adapts better in comparison to tap root system for
Type	multiple choice
Option	Food storage
Option	Solid anchorage
Option	Water and mineral absorption
Option	Organic food translocation
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the incorrectly matched
Type	multiple choice
Option	Prop root – Banayan
Option	Stilt root – Sugarcane
Option	Pneumatophore – Rhizophora
Option	Tap root modification – Asparagus
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Arrange the root regions in their correct manner 1. Region of meristematic activity 2. Region of maturation 3. Region of elongation 4. Root cap
Type	multiple choice
Option	1, 2, 3, 4
Option	4, 1, 3, 2
Option	3, 1, 2, 4
Option	3, 4, 1, 2
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	What is incorrect about pneumatophore?
Type	multiple choice
Option	Grows in swampy areas
Option	Grows vertically upward the ground
Option	Helps in respiration
Option	Present in garden plants
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Which part of root is responsible for rapid elongation and enlargement of root
Type	multiple choice
Option	Region of meristematic activity
Option	Zone of elongation
Option	Region of maturation
Option	Region of root hairs
Answer	2
Solution	
Positive Marks	4

Negative Marks	1
Question	Mark the correct one
Type	multiple choice
Option	Sweet potato - adventitious root modification for storage food
Option	Turnip - adventitious root modification for respiration
Option	Potato - tap root modification for respiration
Option	Banyan tree - pneumatophore modification for storage food
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	Consider the following statements (A) Root hair develop from zone of elongation in primary root (B) Largest zone in primary root is zone of maturation (C) Root cap protect zone of elongation How many are correct –
Type	multiple choice
Option	Only A
Option	Only B
Option	Both A and B
Option	All three
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Consider the following statement – (I) Leaf develop from apical bud in acropetal manner (II) Leaf sheath is modification of petiole How many statements are correct –
Type	multiple choice
Option	only I
Option	only II
Option	both I and II
Option	both wrong
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	Which of the following is incorrect about leaf?
Type	multiple choice
Option	It is arranged in acropetal
Option	It develops from a node
Option	It is generally flattened in shape
Option	It is arranged in basipetal order
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	In Australian acacia, modification of …… develop into spines and …… is photosynthetic.
Type	multiple choice
Option	Lamina, petiole
Option	Leaf base, petiole

Option	Petiole, leaf base
Option	Leaf base, lamina
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	Stolon and runners resemble each other in
Type	multiple choice
Option	Possessing node and internodes
Option	Helping to find new niches
Option	Possessing tuft of roots and rosette of leaves at nodes
Option	More than one option is correct
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	In banana, pineapple and Chrysanthemum, the lateral branches originate from the basal and underground portion of main stem and then come obliquely upward giving rise to leaf shoots
Type	multiple choice
Option	Runner
Option	Corm
Option	Bulb
Option	Sucker
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Select the incorrect match
Type	multiple choice
Option	Thorn - Citrus, Bougainvillea
Option	Flattened stem – Opuntia
Option	Underground stem – Turmeric
Option	Stem tendril – Colocasia
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	What is true about flowering plant stem
Type	multiple choice
Option	Stems modify as they perform different functions
Option	Stem modification can be only into tendril, thorns, and spines.
Option	Stem has points called node where leaves are borne
Option	Both (a) and (c)
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Consider the following statement: I. Stem bears buds either terminal or axillary II. Mature stem is green in colour III. Arrangement of leaves on stem or branch is called phyllotaxy.
Type	multiple choice

Option	Only I
Option	Only III
Option	Only I and II
Option	Only I and III
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the incorrectly matched -
Type	multiple choice
Option	Pea – racemose
Option	Mustard – cymose
Option	Cassia – zygomorphic
Option	China rose – Hypogynous
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Difference between solitary and racemose inflorescence is
Type	multiple choice
Option	Solitary has only one flower at apex but racemose has group of flowers
Option	Solitary inflorescence can be axillary or apical type but racemose is only axillary type
Option	Solitary is only apical type but racemose is also apical type
Option	Both (a) and (b)
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the incorrect for basipetal arrangement of inflorescence.
Type	multiple choice
Option	Floral axis terminates at apex into flower
Option	Floral axis has definite growth
Option	It is in the form of flower cluster.
Option	It has old flower at the base and new flower at the apex.
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	What type of meristem turn into floral meristem?
Type	multiple choice
Option	Lateral meristem
Option	Apical meristem
Option	Intercalary meristem
Option	Both (a) and (b)
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Which of the following has a racemose type of inflorescence?
Type	multiple choice
Option	Sunhemp
Option	Petunia

Option	Colchicum
Option	Aloe
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the incorrectly matched -
Type	multiple choice
Option	Petunia – Gamopetalous
Option	Chilli – Racemose
Option	Tobacco – Gamopetalous
Option	Tomato – Epipetalous
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Statement 1: Inflorescence is arrangement of flowers on floral axis. Statement 2: Inflorescence has modified shoot apical meristem transformation into floral meristem.
Type	multiple choice
Option	Only statement 1 is correct.
Option	Only statement 2 is correct.
Option	Both statements are correct
Option	Both statements incorrect
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Math the following. **Column I (Parts of flower)** / **Column II (Description)** A. Calyx — 1. Female reproductive part B. Corolla — 2. Male reproductive part C. Androeciu m — 3. Outermost whorl of lower D. Gynoecium — 4. Composed of petals
Type	multiple choice
Option	A-4, B-1, C-3, D-2
Option	A-3, B-2, C-1, D-4
Option	A-3, B-4, C-2, D-1
Option	A-2, B-1, C-3, D-4
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Based on the position of calyx, corolla and androecium with respect to the ovary on thalamus, the flower is of how many types?

Type	multiple choice
Option	One
Option	Two
Option	Three
Option	Four
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Hypogynous flower is present in
Type	multiple choice
Option	Pea and Sunflower
Option	Rose and plum
Option	Peach and cucumber
Option	Mustard and chilli
Answer	41
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the correctly matched -
Type	multiple choice
Option	Gulmohar – Actinomorphic
Option	Bean – Zygomorphic
Option	Castor – Non-endospermous
Option	Rose – Syncarpous
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the incorrectly matched -
Type	multiple choice
Option	Syncarpous – Cabbage
Option	Apocarpous – Rose
Option	Many ovule – Groundnut
Option	Inferior ovary – Coconut
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Few examples are given- Petunia, Makoi, Cabbage, Mustard, Lily, Colchicum, Sesbania and Sweet pea- How many show flowers with radial symmetry-
Type	multiple choice
Option	Four
Option	Six
Option	Five
Option	Eight
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	China rose and brinjal have similarity in-
Type	multiple choice

Option	Aestivation
Option	Placentation
Option	Monadelphous
Option	Monocot
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Match the following columns.

	Column I (Placentation)		Column II (Examples)
A.	Basal	1.	Mustard
B.	Axile	2.	China rose
C.	Parietal	3.	*Dianthus*
D.	Free-central	4.	Sunflower

Type	multiple choice
Option	A-2, B-3, C-4, D-1
Option	A-1, B-2, C-3, D-4
Option	A-4, B-2, C-1, D-3
Option	A-3, B-4, C-1, D-2
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	A fruit
Type	multiple choice
Option	Is a mature ovary
Option	Develops after fertilization
Option	Sometimes develops without Fertilisation
Option	All of them
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Fruit is not fleshy in-
Type	multiple choice
Option	Apple
Option	Mango
Option	Coconut
Option	Pea
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Endosperm, a product of double fertilization in angiosperms is absent in the seeds of
Type	multiple choice
Option	Coconut
Option	Orchids

Option	Maize
Option	Castor
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Thick and fleshy pericarp is differentiated into
Type	multiple choice
Option	Epicarp
Option	Mesocarp
Option	Endocarp
Option	All of them
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	What is incorrect about the aleurone layer?
Type	multiple choice
Option	It is a proteinaceous layer.
Option	Found in dicotyledons
Option	Outer covering of endosperm
Option	None of them
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Statement 1: Hilum is a point on the seed coat where developing seed were attached to fruit. Statement 2: Micropyle is small pore below the hilum.
Type	multiple choice
Option	Only statement 1 is correct.
Option	Only statement 2 is correct.
Option	Both statements are correct.
Option	Both statements are incorrect.
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	Drupe fruit is present in…… and have edible part is……
Type	multiple choice
Option	Coconut, endocarp
Option	Mango, epicarp
Option	Coconut, endosperm
Option	Mango, Mesocarp and epicarp
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Epicarp is a
Type	multiple choice
Option	Outer part of mango fruit
Option	Thin layered covering
Option	Inside seed part

Option	Both (a) and (b)
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	A dicot seed is identified by
Type	multiple choice
Option	Two layers called testa and tegmen
Option	Presence of hilum
Option	Presence of two cotyledons and one embryonal axis
Option	All (a), (b) and (c)
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the incorrect one
Type	multiple choice
Option	Seeds outer covering – Testa
Option	Pea seed - endosperm seed
Option	Endosperm - food storing tissue
Option	Bean seed - non endosperm seed
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Scutellum is part of
Type	multiple choice
Option	Monocotyledon seed
Option	Angiosperm seed
Option	Monocot embryo cotyledon
Option	All of them
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Monocots have radicals which give rise to roots in monocots. It is also covered by
Type	multiple choice
Option	Coleoptile
Option	Coleorrhiza
Option	Scutellum
Option	Endosperm
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Statement 1: Fruit is a feature of flowering plants where seeds are formed from ovules. Statement 2: When a fruit formed after without fertilization it is called normal fruit.
Type	multiple choice
Option	Both statements are correct
Option	Only Statement 1 is correct

Option	Only Statement 2 is correct
Option	Both statements are incorrect
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Identify A, B, C in given diagram 
Type	multiple choice
Option	A-cotyledon, B-radicle
Option	A-plumule, C-cotyledon
Option	A-cotyledon, B-plumule
Option	B-radicle, C-plumule
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Match the following

A	Fruit outer covering	I	Stony endocarp
B	Mango fruit	II	Fibrous mesocarp
C	Coconut fruit	III	Form embryo
D	Embryonal axis	IV	Pericarp

Type	multiple choice
Option	A-II, B-III, C-IV, D-I
Option	A-III, B-I, C-IV, D-II
Option	A-IV, B-I, C-II, D-III
Option	A-II, B-IV, C-III, D-I
Answer	3
Solution	
Positive Marks	4
Negative Marks	1