Question	The reason that plants retain the capacity for unlimited growth throughout their life is that -
Type	multiple choice
Option	All Cells can divide in plants
Option	Due to the presence of meristems at certain locations in their body
Option	No differentiation occurs
Option	Dead cells are absent in Plants
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the incorrectly matched pair
Type	multiple choice
Option	Cork cambium - Product of dedifferentiation
Option	Intercalary meristem – Primary meristem
Option	Plant growth – Open
Option	Plant Growth – Close
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	In given equation $W_1 = W_0 e^{rt}, r$ is -
Type	multiple choice
Option	Growth index
Option	Geometric growth rate
Option	Growth term
Option	Growth efficiency
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	All are the characteristics of the cells in meristematic region, except
Type	multiple choice
Option	Rich in protoplasm
Option	Possess large conspicuous nuclei
Option	Their cell walls are primary in nature
Option	Thin and cellulosic with less plasmodesmata connections
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Intercellular intrinsic factor which controls growth and development is:
Type	multiple choice
Option	Gene and Plant growth regulators
Option	Gene
Option	Plant growth regulators
Option	Temperature
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	S-I: Primary growth is the increase in girth of axis due to activity of Apical meristem

	S-II: In arithmetic growth, following mitotic cell division, only one daughter cell continues to divide while the other differentiates and matures.
Type	multiple choice
Option	Only S-I is correct
Option	Only S-II is correct
Option	Both S-I and S-II are correct
Option	Both S-I and S-II are incorrect
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the incorrectly matched pair-
Type	multiple choice
Option	$N^6 -$ furfurylami no purine – Adenine derivative
Option	Abscisic acid – Carotenoids derivative
Option	Gibberellic acid – Terpenes
Option	Kinetin – Carotenoids derivative
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Developmental heterophylly is seen in:
Type	multiple choice
Option	Coriander and Larkspur
Option	Cotton and buttercup
Option	Larkspur and Buttercup
Option	Coriander and buttercup
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	S-I: Auxin was isolated by F.W. Went from tips of coleoptiles of oat seedlings S-II: Development is considered as the sum of growth and differentiation
Type	multiple choice
Option	Only S-I is correct
Option	Only S-II is correct
Option	Both S-I and S-II are correct
Option	Both S-I and S-II are incorrect
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Few parameters are given- Increase in fresh weight, dry weight, length, area, increase in protoplasm of cell How many of the above are difficult to measure directly?
Type	multiple choice
Option	One
Option	Two
Option	Three
Option	Four
Answer	1
Solution	
Positive Marks	4

Negative Marks	1
Question	Cork cambium in dicot stem is a product of dedifferentiation of -
Type	multiple choice
Option	Epidermis
Option	Hypodermis
Option	Cortex
Option	Endodermis
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the incorreotly matched pair (a) Dormin - Abscisic acid (b) Ethyle ne - Discovered by H.H Cousins (c) Kinetin - Term given by Francis Darwin (d) Phototro pism - Coleoptiles of Canary grass.
Type	multiple choice
Option	Dormin — Discovered by H.H Cousins
Option	Ethyle ne — Discovered by H.H Cousins
Option	Kinetin — Term given by Francis Darwin
Option	Phototropism — Coleoptiles of Canary grass
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Few examples are given- Secondary cortex, vascular cambium, cork layer, cork cambium, secondary xylem, secondary phloem How many are products of redifferentiation?
Type	multiple choice
Option	Four
Option	Two
Option	Three
Option	Five
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	W_1 represents final size in Geometric Growth rate equation $W_1 = W_0 e^{rt}$, which means:
Type	multiple choice
Option	Height
Option	Number
Option	Weight
Option	All of these
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Which phytohormone discovered in relation (PGR) was to Bakana disease?

Type	multiple choice
Option	IAA
Option	IBA
Option	GA
Option	ABA
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Which of the following application of auxin widely used for plant propagation?
Type	multiple choice
Option	Herbicidal action
Option	Apical dominance
Option	Initiate rooting in stem cutting
Option	None of these
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Auxin promote flowering in -
Type	multiple choice
Option	Pineapples
Option	Apple
Option	Cucumber
Option	Orange
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	S-I: All GAs are acidic S-II: Gibberellins delay senescence
Type	multiple choice
Option	Only S-I is correct
Option	Only S-II is correct
Option	Both S-I and S-II are correct
Option	Both S-I and S-II are incorrect
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Spraying …….. crop with gibberellins increases the length of the stem
Type	multiple choice
Option	Tomato
Option	Beet
Option	Wheat
Option	Sugarcane
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the incorrectly matching pair:
Type	multiple choice
Option	Auxin —— Xylem differentiation

Option	Auxin —— Used in brewing industry
Option	Auxin —— Induce parthenocarpy
Option	GA_3 —— First gibberellin to be discovered
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Gibberellic acid can be obtained from
Type	multiple choice
Option	Fungi
Option	Bacteria
Option	Higher plants
Option	Both (a) and (c)
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Auxin was first isolated from
Type	multiple choice
Option	Human urine
Option	Corn germ oil
Option	Tobacco stems
Option	Fungus
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	In a plant, decapitation will:
Type	multiple choice
Option	Increase apical dominance
Option	Increase auxin level
Option	Promote lateral branch
Option	Both (b) and (c)
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Select the incorrect statement -
Type	multiple choice
Option	NAA and 2,4-D are synthetic auxins
Option	Auxin helps in cell division
Option	Auxin promote parthenocarpy in tomatoes
Option	NAA is used to overcome apical dominance
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Spraying …..A….. conifers with Gas ….B….. the maturity period
Type	multiple choice
Option	A- mature, B-slows
Option	A-juvenile, B-hastens
Option	A-mature, B-hastens
Option	A-juvenile, B- slows

Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	S-I: There are more than 100 gibberellins reported from widely different organisms such as fungi and higher plants S-II: 2, 4-D, widely used to kill monocotyledonous weeds
Type	multiple choice
Option	Only S-I is correct
Option	Only S-II is correct
Option	Both S-I and S-II are correct
Option	Both S-I and S-II are incorrect
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	Hormone which delay senescence and thus the fruits can be left on the tree longer so as to extend the market period is
Type	multiple choice
Option	GA
Option	Auxin
Option	Cytokinin
Option	Zeatin
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	Hormone which is used to speed up the malting process in brewing industry is
Type	multiple choice
Option	Cytokinin
Option	IAA
Option	GA_3
Option	Kinetin
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	The ability of Gibberellin to increase length of grape stalks is by
Type	multiple choice
Option	Decreasing apical dominance
Option	Increasing length of axis
Option	Increasing apically
Option	Both (b) and (c)
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Genetic dwarfism can be overcome by hormone -
Type	multiple choice
Option	Zeatin
Option	IBA
Option	Kinetin
Option	GA

Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Hormone which promotes adventitious shoot formation is-
Type	multiple choice
Option	Ethylene
Option	ABA
Option	Auxin
Option	Cytokinin
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	The application of ethephon is-
Type	multiple choice
Option	Cell division
Option	Develops lateral branch
Option	Flowering in cucumber
Option	Act as weedicide
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Hormone purified from autoclaved herring sperm DNA is -
Type	multiple choice
Option	NAA
Option	2, 4-D
Option	Kinetin
Option	Zeatin
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Mark the correct matching pair w.r.t hormones and their applications
Type	multiple choice
Option	Ethylene — Promote nutrient mobilization
Option	Abscisic acid —– Role in seed development
Option	Cytokinin —– Promotes rapid internode elongation in deep water rice plants
Option	Ethylene —– Produce new leaves
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Richmond lang effect is related to which hormone-
Type	multiple choice
Option	Cytokinin
Option	Auxin
Option	Ethylene
Option	ABA
Answer	1
Solution	
Positive Marks	4

Negative Marks	1
Question	SI: Natural cytokinins are synthesized in regions where slow cell division occurs S-II: Kinetin does not occur naturally in plants
Type	multiple choice
Option	Only S-I is correct
Option	Only S-II is correct
Option	Both S-I and S-II are correct
Option	Both S-I and S-II are incorrect
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Hormone which is inhibitor of plant growth inhibitor and of metabolism is-
Type	multiple choice
Option	Cytokinin
Option	Auxin
Option	Ethylene
Option	ABA
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	S-I: ABA inhibits seed germination S-II: Ethylene is used to initiate flowering and for synchronizing fruitset in pineapples
Type	multiple choice
Option	Only S-I is correct
Option	Only S-II is correct
Option	Both S-I and S-II are correct
Option	Both S-I and S-II are incorrect
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Hormone which helps the plants to increase their absorption surface is-
Type	multiple choice
Option	IBA
Option	Zeatin
Option	Ethylene
Option	Cytokinin
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Hormone in relation with triple response bioassay is-
Type	multiple choice
Option	Ethylene
Option	Auxin
Option	Cytokinin
Option	ABA
Answer	1
Solution	
Positive Marks	4

Negative Marks	1
Question	Mark the incorrect matching pair-
Type	multiple choice
Option	Ethephon —— Hastens fruit ripening in tomatoes
Option	Cytokinin —— Overcome the apical dominance
Option	ABA —— Acts as an antagonist to Gas
Option	Cytokinin —— Initiate germination in peanut
Answer	4
Solution	
Positive Marks	4
Negative Marks	1
Question	Which of the following is a role of Ethylene?
Type	multiple choice
Option	Respiratory climactic
Option	Weedicide
Option	Apical dominance
Option	Closing of stomata
Answer	1
Solution	
Positive Marks	4
Negative Marks	1
Question	The application of Cytokinin which is useful in tissue culture is-
Type	multiple choice
Option	Nutrient mobilization
Option	Lateral branch development
Option	Development of adventitious shoot
Option	Flowering
Answer	3
Solution	
Positive Marks	4
Negative Marks	1
Question	Search for natural substances with cytokinin-like activities led to the isolation of …….. from cornkernels and coconut milk.
Type	multiple choice
Option	Kinetin
Option	Zeatin
Option	Modified cytokinin
Option	All of these
Answer	2
Solution	
Positive Marks	4
Negative Marks	1
Question	Amino acid which can be precursor of Ethylene is -
Type	multiple choice
Option	Threonine
Option	Alanine
Option	Tryptophan
Option	Methionine
Answer	4
Solution	
Positive Marks	4
Negative Marks	1