

2024 PRIOR AUTHORIZATION CRITERIA

TABLE OF CONTENTS

Abiraterone Acetate	190
Actimmune.....	10
Adempas	168
Akeega.....	190
Alecensa	190
Alosetron Hydrochloride.....	11
Alunbrig.....	190
Alyq	178
Ambrisentan.....	170
Apokyn	22
Apomorphine Hydrochloride	22
Arcalyst.....	23
Aripiprazole.....	20
Aripiprazole Odt.....	20
Armodafinil	24
Asenapine Maleate Sl.....	20
Atovaquone.....	27
Augtyro.....	190
Austedo	28
Austedo Xr.....	28
Austedo Xr Titration Kit.....	28
Avita	211
Ayvakit.....	190
Balversa	190
Benlysta IV	29
Benlysta SC.....	31
Benzotropine Mesylate	108
Besremi	190
Betaseron	132
Bexarotene Cap.....	190
Bexarotene Gel	38
Bosentan	172
Bosulif	190
Braftovi.....	190
Brukinsa	190
Budesonide	62
Budesonide Er	63
Bydureon Bcise	64
Byetta	65
Cabometyx	190
Calquence.....	190

Caprelsa.....	190
Carglumic Acid	66
Chenodal	67
Chlorpromazine Hydrochloride.....	20
Cinacalcet Hydrochloride	68
Cinryze.....	101
Clemastine Fumarate	108
Clobazam.....	33
Clorazepate Dipotassium	34
Clozapine.....	20
Clozapine Odt.....	20
Cobenfy	69
Cometriq	190
Copaxone	135
Copiktra.....	190
Corlanor	73
Cosentyx.....	40
Cosentyx Sensoready Pen	40
Cosentyx Uno Inj	40
Cotellic	190
Cresemba	74
Cyltezo.....	42
Cyltezo Kit	42
Cyltezo Pen.....	42
Cyltezo Pen Kit	42
Cyltezo Pen-Cd/Uc/Hs Starter Pack	42
Cyltezo Pen-Ps Starter Pack	42
Cyproheptadine Hydrochloride	108
Cystadrops	75
Cystagon.....	77
Cystaran	76
Dalfampridine Er	78
Danazol	13
Dasatinib	190
Daurismo.....	190
Deferasirox (Exjade).....	113
Deferasirox (Jadenu)	114
Depo-Testosterone	14
Dexmethylphenidate Hydrochloride.....	95
Diazepam	35
Diazepam Intensol	35
Dicyclomine Hydrochloride.....	108
Dihydroergotamine Mesylate Spray	129
Dimethyl Fumarate	133
Dimethyl Fumarate Starterpack.....	133
Diphenoxylate Hydrochloride/Atropine Sulfate	108
Droxidopa.....	79
Dupixent.....	80

Eligard	120
Emgality.....	83
Emsam.....	84
Enbrel	44
Enbrel Mini.....	44
Enbrel Sureclick.....	44
Endari	86
Epclusa	87
Epidiolex.....	88
Erivedge	190
Erleada	190
Erlotinib Hydrochloride	190
Everolimus.....	190
Exkivity	190
Fanapt	20
Fanapt Titration Pack	20
Farydak.....	190
Fentanyl	147
Fentanyl Citrate Oral Transmucosal.....	93
Fingolimod	134
Fintepla	94
Fluphenazine Decanoate	20
Fluphenazine Hydrochloride.....	20
Forteo.....	206
Fotivda	190
Fruzaqla.....	190
Gammaflex	60
Gamunex-C.....	96
Gattex.....	98
Gavreto	190
Gefitinib	190
Gilotrif	190
Granix.....	71
Hadlima	46
Hadlima Pushtouch Pen	46
Haegarda	103
Haloperidol.....	20
Haloperidol Decanoate	20
Haloperidol Lactate.....	20
Harvoni.....	107
Humira	48
Humira Pediatric Crohns Disease Starter Pack	48
Humira Pen	48
Humira Pen-Cd/Uc/Hs Starter.....	48
Humira Pen-Pediatric Uc Starter Pack	48
Humira Pen-Ps/Uv Starter.....	48
Hydroxyzine Hydrochloride.....	108
Ibrance	190

Icatibant Acetate.....	105
Iclusig	190
Idhifa	190
Imatinib Mesylate	190
Imbruvica	190
Imiquimod	109
Inbrija	110
Ingrezza	111
Inlyta	190
Inqovi	190
Inrebic	190
Ivabradine	73
Ivermectin Cream.....	115
Ivermectin Tablet	116
Iwilfin.....	190
Jakafi	190
Javygtor	189
Jaypirca	191
Kalydeco	117
Kerendia	118
Kesimpta	136
Kisqali	191
Kisqali Femara 200 Dose	191
Kisqali Femara 400 Dose	191
Kisqali Femara 600 Dose	191
Korlym	119
Koselugo.....	191
Krazati	191
Lapatinib Ditosylate	191
Lazcluze	191
Lenalidomide.....	191
Lenvima 10 Mg Daily Dose	191
Lenvima 12Mg Daily Dose.....	191
Lenvima 14 Mg Daily Dose	191
Lenvima 18 Mg Daily Dose	191
Lenvima 20 Mg Daily Dose	191
Lenvima 24 Mg Daily Dose	191
Lenvima 4 Mg Daily Dose	191
Lenvima 8 Mg Daily Dose	191
Leukine	72
Leuprolide Acetate	120
L-glutamine	86
Lidocaine Patch	121
Lidocaine Solution.....	122
Lidocaine/Prilocaine.....	123
Lidocan	121
Lidocan III	121
Linezolid	124

Lonsurf	191
Lorazepam.....	36
Lorazepam Intensol.....	36
Lorbrena	191
Loxapine	20
Lumakras	191
Lupron Depot (1-Month).....	120
Lupron Depot (3-Month).....	120
Lupron Depot (4-Month).....	120
Lupron Depot (6-Month).....	120
Lupron Depot-Ped (1-Month)	120
Lupron Depot-Ped (3-Month)	120
Lupron Depot-Ped (6-Month)	120
Lurasidone.....	20
Lybalvi	20
Lynparza	191
Lytgobi.....	191
Margenza	112
Matulane.....	191
Mayzent	137
Mayzent Starter Pack	137
Mekinist	191
Mektovi	191
Memantine Hcl Titration Pak	127
Memantine Hydrochloride.....	127
Memantine Hydrochloride Er	126
Methylphenidate Hydrochloride (Ritalin)	186
Methylphenidate Hydrochloride Er Tablet	128
Mifepristone	119
Miglustat	203
Modafinil	130
Molindone Hydrochloride.....	20
Morphine Sulfate Er	149
Mounjaro	131
Myalept	139
Natpara	140
Nelarabine.....	112
Nerlynx	191
Ninlaro.....	191
Noxafil Pak	161
Nubeqa.....	191
Nuedexta	141
Nuplazid	142
Nurtec	143
Ocaliva.....	144
Octreotide Acetate.....	199
Odomzo.....	191
Ofev.....	146

Ogsiveo.....	191
Ojemda.....	191
Ojjaara.....	191
Olanzapine	20
Olanzapine Odt	20
Omnitrope.....	99
Onureg	191
Opsumit.....	174
Orgovyx	191
Orkambi	153
Orserdu	191
Otezla	154
Ozempic	156
Paliperidone Er.....	20
Palynziq	157
Panretin.....	158
Pazopanib.....	191
Pegasys.....	159
Pemazyre.....	191
Perphenazine	20
Pimecrolimus	25
Pimozide.....	20
Piqray 200Mg Daily Dose	191
Piqray 250Mg Daily Dose	191
Piqray 300Mg Daily Dose	191
Pirfenidone.....	160
Pomalyst.....	191
Posaconazole Dr.....	161
Posaconazole Susp	161
Procrit.....	89
Prolastin-C.....	12
Prolia	163
Promacta	166
Promethazine Hcl Plain	108
Promethazine Hydrochloride	108
Pyrimethamine.....	182
Qinlock	191
Quetiapine Fumarate	20
Quetiapine Fumarate Er.....	20
Quinine Sulfate	183
Repatha	184
Repatha Pushtronex System	184
Repatha Sureclick.....	184
Retacrit.....	91
Retevmo	191
Revlimid	192
Rexulti	20
Rezlidhia.....	192

Rinvoq	50
Rinvoq Lq.....	52
Risperidone	20
Risperidone Odt	20
Roflumilast	187
Rozlytrek	192
Rubraca	192
Rybelsus	188
Rydapt	192
Sajazir	105
Sapropterin Dihydrochloride	189
Scemblix	192
Scopolamine.....	108
Secuado.....	20
Signifor	195
Sildenafil.....	176
Sivextro	196
Skyrizi	53
Skyrizi Pen	53
Sodium Oxybate.....	198
Sodium Phenylbutyrate	218
Somavert.....	201
Sorafenib.....	192
Sovaldi.....	202
Sprycel.....	192
Stelara	55
Stivarga	192
Sunitinib Malate.....	192
Sympazan	37
Synribo	112
Tabrecta	192
Tacrolimus.....	26
Tadalafil 20Mg Tablet	178
Tafinlar	192
Tagrisso	192
Talzenna	192
Tasigna	192
Tasimelteon.....	205
Tazarotene	210
Tazorac.....	210
Tazverik	192
Tepmetko	192
Teriparatide	206
Testosterone	18
Testosterone Cypionate.....	14
Testosterone Enanthate	16
Testosterone Pump.....	18
Tetrabenazine	208

Thalomid	192
Thioridazine Hcl.....	20
Thiothixene	20
Tibsovo	192
Tobramycin Neb.....	209
Torpenz	192
Tracleer	172
Tramadol Hcl Er	151
Trelstar Mixject	212
Tretinoin 10Mg Cap	192
Tretinoin Cream, Gel.....	211
Tridacaine.....	121
Trientine Hydrochloride.....	213
Trifluoperazine Hcl	20
Trihexyphenidyl Hydrochloride.....	108
Trikafta	214
Trulicity	215
Truqap	192
Tukysa	192
Turalio	192
Tymlos.....	216
Vanflyta	192
Venclexta.....	192
Venclexta Starting Pack.....	192
Ventavis.....	180
Versacloz	20
Verzenio	192
Victoza.....	219
Vitrakvi	192
Vizimpro	192
Vonjo	192
Voranigo.....	192
Voriconazole	220
Vosevi.....	222
Votrient	192
Vumerity	138
Vyndaqel	223
Welireg.....	192
Xalkori	192
Xeljanz Solution.....	57
Xeljanz Tablet.....	58
Xeljanz Xr.....	59
Xermelo.....	224
Xgeva.....	225
Xifaxan.....	227
Xolair	228
Xospata	192
Xpovio	192

Xpovio 100 Mg Once Weekly	192
Xpovio 40 Mg Once Weekly	192
Xpovio 40 Mg Twice Weekly	192
Xpovio 60 Mg Once Weekly	192
Xpovio 60 Mg Twice Weekly	192
Xpovio 80 Mg Once Weekly	192
Xpovio 80 Mg Twice Weekly	192
Xtandi	193
Yargesa	203
Zejula.....	193
Zelboraf	193
Zepatier	231
Ziprasidone Mesylate.....	20
Zokinvy	232
Zolinza	193
Zydelig.....	193
Zykadia	193
Zyprexa Relprevv.....	20

Prior Authorization Group Description:

Actimmune PA

Drug Name(s)

Actimmune

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Alosetron PA

Drug Name(s)

Alosetron Hydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of irritable bowel syndrome with severe diarrhea (IBS-D) AND
2. Patient's sex is female AND
3. Patient exhibits at least ONE of the following:
 - a. Frequent and severe abdominal pain/discomfort OR
 - b. Frequent bowel urgency or fecal incontinence OR
 - c. Disability or restriction of daily activities due to IBS AND
4. Prescriber has ruled out anatomic or biochemical abnormalities of the gastrointestinal tract

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Alpha-1-Proteinase Inhibitor PA - Prolastin-C

Drug Name(s)

Prolastin-C

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of alpha-1 antitrypsin deficiency (AATD) with clinically evident emphysema AND
2. Patient has a pre-treatment serum alpha-1 antitrypsin (AAT) level less than 11 micromol/L (80 mg/dL by immunodiffusion or 57 mg/dL using nephelometry) AND
3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of alpha-1 antitrypsin deficiency (AATD) with clinically evident emphysema AND
3. Patient has had clinical benefit with the requested agent AND
4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Anabolic Steroid PA – Danazol

Drug Name(s)

Danazol

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has ONE of the following diagnoses:

A. Patient has an FDA labeled indication for the requested agent OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR

B. Prescriber has provided information in support of therapy with more than one agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Androgen Injectable PA - testosterone cypionate

Drug Name(s)

Depo-Testosterone

Testosterone Cypionate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:

A. Patient's sex is male with AIDS/HIV-associated wasting syndrome AND BOTH of the following:

i. ONE of the following:

a. Unexplained involuntary weight loss (greater than 10% baseline body weight within 12 months, or 7.5% within 6 months) OR

b. Body mass index less than 20 kg/m² OR

c. At least 5% total body cell mass (BCM) loss within 6 months OR

d. BCM less than 35% of total body weight and BMI less than 27 kg/m² AND

ii. All other causes of weight loss have been ruled out OR

B. Patient's sex is female with metastatic/inoperable breast cancer OR

C. Patient's sex is male with primary or secondary (hypogonadotropic) hypogonadism OR

D. Patient's sex is male and is an adolescent with delayed puberty AND

2. If the patient's sex is a male, ONE of the following:

A. Patient is NOT currently receiving testosterone replacement therapy AND has ONE of the following pretreatment levels:

i. Total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR

ii. Free serum testosterone level that is below the testing laboratory's lower limit of the normal range OR

B. Patient is currently receiving testosterone replacement therapy AND has ONE of the following current levels:

i. Total serum testosterone level that is within OR below the testing laboratory's lower limit of the normal range OR is less than 300 ng/dL OR

ii. Free serum testosterone level is within OR below the testing laboratory's normal range AND

3. ONE of the following:

A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR

B. Prescriber has provided information in support of therapy with more than one agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be 6 months for delayed puberty, 12 months for all other indications

Other Criteria:

Prior Authorization Group Description:

Androgen Injectable PA - testosterone enanthate

Drug Name(s)

Testosterone Enanthate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:

A. Patient's sex is male with AIDS/HIV-associated wasting syndrome AND BOTH of the following:

i. ONE of the following:

- a. Unexplained involuntary weight loss (greater than 10% baseline body weight within 12 months, or 7.5% within 6 months) OR
- b. Body mass index less than 20 kg/m² OR
- c. At least 5% total body cell mass (BCM) loss within 6 months OR
- d. BCM less than 35% of total body weight and BMI less than 27 kg/m² AND

ii. All other causes of weight loss have been ruled out OR

B. Patient's sex is female with metastatic/inoperable breast cancer OR

C. Patient's sex is male with primary or secondary (hypogonadotropic) hypogonadism OR

D. Patient's sex is male and is an adolescent with delayed puberty AND

2. If the patient's sex is a male, ONE of the following:

A. Patient is NOT currently receiving testosterone replacement therapy AND has ONE of the following pretreatment levels:

- i. Total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR
- ii. Free serum testosterone level that is below the testing laboratory's lower limit of the normal range OR

B. Patient is currently receiving testosterone replacement therapy AND has ONE of the following current levels:

- i. Total serum testosterone level that is within OR below the testing laboratory's lower limit of the normal range OR is less than 300 ng/dL OR
- ii. Free serum testosterone level is within OR below the testing laboratory's normal range AND

3. ONE of the following:

A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR

B. Prescriber has provided information in support of therapy with more than one agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be 6 months for delayed puberty, 12 months for all other indications

Other Criteria:

Prior Authorization Group Description:

Androgen Topical PA

Drug Name(s)

Testosterone

Testosterone Pump

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:

A. Patient has AIDS/HIV-associated wasting syndrome AND BOTH of the following:

i. ONE of the following:

a. Unexplained involuntary weight loss (greater than 10% baseline body weight within 12 months, or 7.5% within 6 months) OR

b. Body mass index less than 20 kg/m² OR

c. At least 5% total body cell mass (BCM) loss within 6 months OR

d. In men: BCM less than 35% of total body weight and BMI less than 27 kg/m² OR

e. In women: BCM less than 23% of total body weight and BMI less than 27 kg/m² AND

ii. All other causes of weight loss have been ruled out OR

B. Patient's sex is male with primary or secondary (hypogonadotropic) hypogonadism AND

2. If the patient's sex is male, ONE of the following:

A. Patient is NOT currently receiving testosterone replacement therapy AND has ONE of the following pretreatment levels:

i. Total serum testosterone level that is below the testing laboratory's lower limit of the normal range or is less than 300 ng/dL OR

ii. Free serum testosterone level that is below the testing laboratory's lower limit of the normal range OR

B. Patient is currently receiving testosterone replacement therapy AND has ONE of the following current levels:

i. Total serum testosterone level that is within OR below the testing laboratory's lower limit of the normal range OR is less than 300 ng/dL OR

ii. Free serum testosterone level is within OR below the testing laboratory's normal range AND

3. ONE of the following:

A. Patient will NOT be using the requested agent in combination with another androgen or anabolic steroid OR

B. Prescriber has submitted information in support of therapy with more than one agent

Age Restriction:**Prescriber Restrictions:**

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Antipsychotics PA

Drug Name(s)

Aripiprazole
Aripiprazole Odt
Asenapine Maleate Sl
Chlorpromazine Hydrochloride
Clozapine
Clozapine Odt
Fanapt
Fanapt Titration Pack
Fluphenazine Decanoate
Fluphenazine Hydrochloride
Haloperidol
Haloperidol Decanoate
Haloperidol Lactate
Loxapine
Lurasidone
Lybalvi
Molindone Hydrochloride
Olanzapine
Olanzapine Odt
Paliperidone Er
Perphenazine
Pimozide
Quetiapine Fumarate
Quetiapine Fumarate Er
Rexulti
Risperidone
Risperidone Odt
Secuado
Thioridazine Hcl
Thiothixene
Trifluoperazine Hcl
Versacloz
Ziprasidone Mesylate
Zyprexa Relprevv

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

PA does NOT apply to patients less than 65 years of age.

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent OR
 - c. ONE of the following:
 - i. Patient has a diagnosis other than dementia-related psychosis or dementia related behavioral symptoms OR
 - ii. Patient has dementia-related psychosis or dementia related behavioral symptoms AND BOTH of the following:
 1. Dementia-related psychosis is determined to be severe or the associated behavior puts the patient or others in danger AND
 2. Prescriber has documented that s/he has discussed the risk of increased mortality with the patient and/or the patient's surrogate decision maker

Approval authorizations will apply to the requested medication only.

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Apomorphine Inj PA

Drug Name(s)

Apokyn

Apomorphine Hydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ALL of the following:

1. The requested agent will be used to treat acute, intermittent hypomobility, “off” episodes (“end of dose wearing off” and unpredictable “on/off” episodes) associated with advanced Parkinson’s disease AND
2. The requested agent will be used in combination with agents used for therapy in Parkinson’s disease (e.g., levodopa, dopamine agonist, monoamine oxidase B inhibitor) AND
3. Patient will NOT be using the requested agent in combination with a 5-HT3 antagonist (e.g., ondansetron, granisetron, dolasetron, palonosetron, alosetron)

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient’s diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient’s diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Arcalyst PA

Drug Name(s)

Arcalyst

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require BOTH of the following:

1. ONE of the following:

A. Patient has been diagnosed with Cryopyrin-Associated Periodic Syndromes (CAPS) including Familial Cold Auto-inflammatory Syndrome (FCAS) or Muckle-Wells Syndrome (MWS) OR

B. BOTH of the following:

i. Patient has a diagnosis of deficiency of interleukin-1 receptor antagonist AND

ii. The requested agent is being used for maintenance of remission OR

C. BOTH of the following:

i. Patient has a diagnosis of recurrent pericarditis AND

ii. The requested agent is being used to reduce the risk of recurrence AND

2. Patient will NOT be using the requested agent in combination with another biologic agent

Age Restriction:

For diagnosis of CAPS including FCAS or MWS, patient is 12 years of age or over

For diagnosis of recurrent pericarditis and reduction in risk of recurrence, patient is 12 years of age or over

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Armodafinil PA

Drug Name(s)

Armodafinil

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. Patient will NOT be using the requested agent in combination with another target agent (i.e., modafinil)

Age Restriction:

Patient is 17 years of age or over

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Atopic Dermatitis PA – Pimecrolimus

Drug Name(s)

Pimecrolimus

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ONE of the following:

1. Patient has a diagnosis of atopic dermatitis or vulvar lichen sclerosus AND ONE of the following:
 - A. Patient has tried and had an inadequate response to a topical corticosteroid or topical corticosteroid combination preparation (e.g., hydrocortisone, triamcinolone) OR
 - B. Patient has an intolerance or hypersensitivity to a topical corticosteroid or topical corticosteroid combination preparation OR
 - C. Patient has an FDA labeled contraindication to a topical corticosteroid or topical corticosteroid combination preparation OR
2. Patient has a diagnosis of facial seborrheic dermatitis associated with HIV infection AND BOTH of the following:
 - A. Patient is currently on an antiretroviral treatment regimen AND
 - B. ONE of the following:
 - i. Patient has tried and had an inadequate response to a topical corticosteroid or topical antifungal treatment (e.g., hydrocortisone, triamcinolone, ketoconazole) OR
 - ii. Patient has an intolerance or hypersensitivity to a topical corticosteroid or topical antifungal treatment OR
 - iii. Patient has an FDA labeled contraindication to a topical corticosteroid or topical antifungal treatment OR
3. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Atopic Dermatitis PA – Tacrolimus

Drug Name(s)

Tacrolimus

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ONE of the following:

1. Patient has a diagnosis of atopic dermatitis AND ONE of the following:
 - A. Patient has tried and had an inadequate response to a topical corticosteroid or topical corticosteroid combination preparation (e.g., hydrocortisone, triamcinolone) OR
 - B. Patient has an intolerance or hypersensitivity to a topical corticosteroid or topical corticosteroid combination preparation OR
 - C. Patient has an FDA labeled contraindication to a topical corticosteroid or topical corticosteroid combination preparation OR
2. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Atovaquone PA

Drug Name(s)

Atovaquone

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

1. Patient has a diagnosis of mild-to-moderate *Pneumocystis jirovecii* pneumonia OR
2. Patient is using the requested agent for prevention of *Pneumocystis jirovecii* pneumonia AND

ii. ONE of the following:

1. Patient has an intolerance or hypersensitivity to trimethoprim/sulfamethoxazole (TMP/SMX) OR
2. Patient has an FDA labeled contraindication to trimethoprim/sulfamethoxazole (TMP/SMX) OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Austedo PA

Drug Name(s)

Austedo

Austedo Xr

Austedo Xr Titration Kit

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. ONE of the following:

A. Patient has a diagnosis of chorea associated with Huntington's disease AND BOTH of the following:

i. ONE of the following:

1. Patient does NOT have a current diagnosis of depression OR
2. Patient has a current diagnosis of depression and is being treated for depression AND

ii. ONE of the following:

1. Patient does NOT have a diagnosis of passive suicidal ideation OR
2. Patient has a diagnosis of passive suicidal ideation and must NOT be actively suicidal OR

B. Patient has a diagnosis of tardive dyskinesia AND ONE of the following:

i. Prescriber has reduced the dose of or discontinued any medications known to cause tardive dyskinesia (i.e., dopamine receptor blocking agents) OR

ii. Prescriber has provided clinical rationale indicating that a reduced dose or discontinuation of any medications known to cause tardive dyskinesia is not appropriate AND

2. Patient will NOT be using the requested agent in combination with a monoamine oxidase inhibitor (MAOI) AND

3. Patient will NOT be using the requested agent in combination with reserpine

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Benlysta IV PA

Drug Name(s)

Benlysta IV

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. ONE of the following:
 - a. Patient has a diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:
 - i. Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR
 - b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:
 - i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND
2. Patient will NOT be using the requested agent in combination with another biologic agent AND
3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. ONE of the following:
 - a. Patient has diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:
 - i. Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR
 - b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:
 - i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND
3. Patient has had clinical benefit with the requested agent AND
4. Patient will NOT be using the requested agent in combination with another biologic agent AND
5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

For diagnosis of active systemic lupus erythematosus (SLE) disease, patient is 5 years of age or over. For diagnosis of active lupus nephritis (LN), patient is 5 years of age or over.

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Benlysta SC PA

Drug Name(s)

Benlysta SC

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. ONE of the following:
 - a. Patient has a diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:
 - i. Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR
 - b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:
 - i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND
2. Patient will NOT be using the requested agent in combination with another biologic agent AND
3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. ONE of the following:
 - a. Patient has diagnosis of active systemic lupus erythematosus (SLE) disease AND the following:
 - i. Patient will continue standard SLE therapy [corticosteroids (e.g., methylprednisolone, prednisone), hydroxychloroquine, immunosuppressives (e.g., azathioprine, methotrexate, oral cyclophosphamide)] in combination with the requested agent OR
 - b. Patient has a diagnosis of active lupus nephritis (LN) AND the following:
 - i. Patient will continue standard LN therapy [corticosteroids (e.g., methylprednisolone, prednisone), immunosuppressives (e.g., azathioprine, mycophenolate)] in combination with the requested agent AND
3. Patient has had clinical benefit with the requested agent AND
4. Patient will NOT be using the requested agent in combination with another biologic agent AND
5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

For diagnosis of active systemic lupus erythematosus (SLE) disease, patient is 5 years of age or over. For diagnosis of active lupus nephritis (LN), patient is 18 years of age or over.

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Benzodiazepines PA – Clobazam

Drug Name(s)

Clobazam

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently being treated with the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Seizure disorder OR

b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Benzodiazepines PA – Clorazepate

Drug Name(s)

Clorazepate Dipotassium

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- b. Prescriber states the patient is currently being treated with the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

- a. Seizure disorder OR
- b. Anxiety disorder AND ONE of the following:
 - 1) Patient has tried and has an inadequate response to a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin norepinephrine reuptake inhibitor (SNRI) OR
 - 2) Patient has an intolerance or hypersensitivity to a formulary SSRI or SNRI OR
 - 3) Patient has an FDA labeled contraindication to a formulary SSRI or SNRI OR
- c. Alcohol withdrawal OR
- d. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Benzodiazepines PA – Diazepam

Drug Name(s)

Diazepam

Diazepam Intensol

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- b. Prescriber states the patient is currently being treated with the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

- a. Seizure disorder OR
- b. Anxiety disorder AND ONE of the following:
 - 1) Patient has tried and had an inadequate response to a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin norepinephrine reuptake inhibitor (SNRI) OR
 - 2) Patient has an intolerance or hypersensitivity to a formulary SSRI or SNRI OR
 - 3) Patient has an FDA labeled contraindication to a formulary SSRI or SNRI OR
- c. Skeletal muscle spasms OR
- d. Alcohol withdrawal OR
- e. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Benzodiazepines PA – Lorazepam

Drug Name(s)

Lorazepam

Lorazepam Intensol

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- b. Prescriber states the patient is currently being treated with the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Anxiety disorder AND ONE of the following:

- 1) Patient has tried and had an inadequate response to a formulary selective serotonin reuptake inhibitor (SSRI) or serotonin norepinephrine reuptake inhibitor (SNRI) OR
- 2) Patient has an intolerance or hypersensitivity to a formulary SSRI or SNRI OR
- 3) Patient has an FDA labeled contraindication to a formulary SSRI or SNRI OR

b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Benzodiazepines PA – Sympazan

Drug Name(s)

Sympazan

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR

b. Prescriber states the patient is currently being treated with the requested agent AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. BOTH of the following:

i. Patient has ONE of the following diagnoses:

a. Seizure disorder OR

b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Bexarotene Gel PA

Drug Name(s)

Bexarotene Gel

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

- A. Patient has a diagnosis of stage IA or IB cutaneous T-cell lymphoma (CTCL) with cutaneous lesions OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. ALL of the following:
 - i. ONE of the following:
 - 1. BOTH of the following:
 - a. Patient has a diagnosis of stage IA or IB cutaneous T-cell lymphoma (CTCL) with cutaneous lesions AND
 - b. ONE of the following:
 - i. Patient has refractory or persistent disease despite a previous treatment trial with a skin-directed therapy (e.g., topical corticosteroid, topical imiquimod) OR
 - ii. Patient has an intolerance or hypersensitivity to a previous treatment trial with a skin-directed therapy (e.g., topical corticosteroid, topical imiquimod) OR
 - iii. Patient has an FDA labeled contraindication to a previous treatment trial with a skin-directed therapy (e.g., topical corticosteroid, topical imiquimod) OR
 - 2. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., dermatologist, oncologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND
 - iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. ONE of the following:
 - A. Patient has a diagnosis of stage IA or IB cutaneous T-cell lymphoma (CTCL) with cutaneous lesions OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
3. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Patient has had clinical benefit with the requested agent AND
 - ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., dermatologist, oncologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND
 - iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Prior Authorization Group Description:

Biologic Immunomodulators PA – Cosentyx

Drug Name(s)

Cosentyx

Cosentyx Sensoready Pen

Cosentyx Uno Inj

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
 - E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis or plaque psoriasis

NO prerequisites are required for diagnoses of ankylosing spondylitis, enthesitis related arthritis, hidradenitis suppurativa, or non-radiographic axial spondyloarthritis

Formulary conventional agents for psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Prior Authorization Group Description:

Biologic Immunomodulators PA – Cyltezo

Drug Name(s)

Cyltezo

Cyltezo Kit

Cyltezo Pen

Cyltezo Pen Kit

Cyltezo Pen-Cd/Uc/Hs Starter Pack

Cyltezo Pen-Ps Starter Pack

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
 - E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:**Coverage Duration:**

Approval will be 12 weeks for initial use for ulcerative colitis, 12 months for all others

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile idiopathic arthritis, Crohn's disease, or moderate ulcerative colitis

NO prerequisites are required for diagnoses of ankylosing spondylitis, hidradenitis suppurativa, severe ulcerative colitis, or uveitis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, or psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, or mercaptopurine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, or mercaptopurine

Prior Authorization Group Description:

Biologic Immunomodulators PA – Enbrel

Drug Name(s)

Enbrel

Enbrel Mini

Enbrel Sureclick

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
 - E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile psoriatic arthritis, or juvenile idiopathic arthritis

NO prerequisites are required for a diagnoses of ankylosing spondylitis or severe juvenile psoriatic arthritis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, juvenile psoriatic arthritis, or psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Prior Authorization Group Description:

Biologic Immunomodulators PA – Hadlima

Drug Name(s)

Hadlima

Hadlima Pushtouch Pen

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
 - E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be 12 weeks for initial use for ulcerative colitis, 12 months for all others

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile idiopathic arthritis, Crohn's disease, or moderate ulcerative colitis

NO prerequisites are required for diagnoses of ankylosing spondylitis, hidradenitis suppurativa, severe ulcerative colitis, or uveitis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, or psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, or mercaptopurine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, or mercaptopurine

Prior Authorization Group Description:

Biologic Immunomodulators PA – Humira

Drug Name(s)

Humira

Humira Pediatric Crohns Disease Starter Pack

Humira Pen

Humira Pen-Cd/Uc/Hs Starter

Humira Pen-Pediatric Uc Starter Pack

Humira Pen-Ps/Uv Starter

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
 - E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:**Coverage Duration:**

Approval will be 12 weeks for initial use for ulcerative colitis, 12 months for all others

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, rheumatoid arthritis, juvenile idiopathic arthritis, Crohn's disease, or moderate ulcerative colitis

NO prerequisites are required for diagnoses of ankylosing spondylitis, hidradenitis suppurativa, severe ulcerative colitis, or uveitis

Formulary conventional agents for rheumatoid arthritis, juvenile idiopathic arthritis, or psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, or mercaptopurine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, or mercaptopurine

Prior Authorization Group Description:

Biologic Immunomodulators PA – Rinvoq

Drug Name(s)

Rinvoq

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. ONE of the following:
 - i. BOTH of the following:
 - a. Patient has an FDA labeled indication other than moderate to severe atopic dermatitis for the requested agent AND
 - b. ONE of the following:
 1. Patient's medication history indicates use of preferred TNF agent(s) OR
 2. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
 3. Patient has an FDA labeled contraindication to preferred TNF agent(s) OR
 4. The request is for an FDA labeled indication that is not covered by preferred TNF agent(s) OR
 - ii. Patient has a diagnosis of moderate to severe atopic dermatitis AND ONE of the following:
 1. Patient's medication history indicates use of TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) for the requested indication OR
 2. Patient has an intolerance or hypersensitivity to TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) for the requested indication OR
 3. Patient has an FDA labeled contraindication to TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) for the requested indication AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Use of ONE preferred TNF (Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnoses of ankylosing spondylitis, rheumatoid arthritis, adult psoriatic arthritis, or juvenile idiopathic arthritis

Use of ONE preferred TNF (Cyltezo, Hadlima, or Humira) is required for diagnoses of ulcerative colitis or Crohn's disease

Use of TWO conventional prerequisite agents (i.e., ONE formulary topical corticosteroid AND ONE formulary topical calcineurin inhibitor) are required for diagnosis of moderate to severe atopic dermatitis

NO preferred TNF agents are required for diagnosis of pediatric psoriatic arthritis

NO preferred TNF agents are required for diagnosis of non-radiographic Axial Spondyloarthritis

Prior Authorization Group Description:

Biologic Immunomodulators PA – Rinvoq Solution

Drug Name(s)

Rinvoq Lq

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. ONE of the following:
 - i. Patient's medication history indicates use of preferred TNF agent(s) OR
 - ii. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
 - iii. Patient has an FDA labeled contraindication to preferred TNF agent(s) OR
 - iv. The request is for an FDA labeled indication that is not covered by preferred TNF agent(s) AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Use of ONE preferred TNF (Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnoses adult psoriatic arthritis or juvenile idiopathic arthritis

NO preferred TNF agents are required for diagnosis of pediatric psoriatic arthritis

Prior Authorization Group Description:

Biologic Immunomodulators PA – Skyrizi

Drug Name(s)

Skyrizi

Skyrizi Pen

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
 - E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of Crohn's disease, plaque psoriasis, psoriatic arthritis, or moderate ulcerative colitis

NO prerequisites are required for diagnosis of severe ulcerative colitis

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, or mercaptopurine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for psoriatic arthritis include leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, or mercaptopurine

Prior Authorization Group Description:

Biologic Immunomodulators PA – Stelara

Drug Name(s)

Stelara

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. Patient's medication history indicates use of another biologic immunomodulator agent for the same FDA labeled indication OR
 - D. Patient's diagnosis does NOT require a conventional prerequisite agent OR
 - E. Patient's medication history indicates use of ONE formulary conventional prerequisite agent for the requested indication OR
 - F. Patient has an intolerance or hypersensitivity to at least ONE formulary conventional prerequisite agent for the requested indication OR
 - G. Patient has an FDA labeled contraindication to at least ONE formulary conventional prerequisite agent for the requested indication AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Use of ONE conventional prerequisite agent is required for diagnoses of psoriatic arthritis, plaque psoriasis, moderate ulcerative colitis, or Crohn's disease

NO prerequisites are required for diagnosis of severe ulcerative colitis

Formulary conventional agents for psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

Formulary conventional agents (topical or systemic) for plaque psoriasis include acitretin, calcipotriene, methotrexate, tazarotene, or topical corticosteroids

Formulary conventional agents for Crohn's disease include methotrexate, sulfasalazine, corticosteroids, azathioprine, mercaptopurine

Formulary conventional agents for moderate ulcerative colitis include 5-aminosalicylates, corticosteroids, azathioprine, mercaptopurine

Prior Authorization Group Description:

Biologic Immunomodulators PA - Xeljanz Solution

Drug Name(s)

Xeljanz Solution

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. ONE of the following:
 - i. Patient's medication history indicates use of preferred TNF agent(s) OR
 - ii. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
 - iii. Patient has an FDA labeled contraindication to preferred TNF agent(s) AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Use of ONE preferred TNF (Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnosis of juvenile idiopathic arthritis

Prior Authorization Group Description:

Biologic Immunomodulators PA - Xeljanz Tablet

Drug Name(s)

Xeljanz Tablet

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. ONE of the following:
 - i. Patient's medication history indicates use of preferred TNF agent(s) OR
 - ii. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
 - iii. Patient has an FDA labeled contraindication to preferred TNF agent(s) AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Use of ONE preferred TNF (Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnoses of psoriatic arthritis, rheumatoid arthritis, juvenile idiopathic arthritis, or ankylosing spondylitis

Use of ONE preferred TNF (Cyltezo, Hadlima, or Humira) is required for diagnosis of ulcerative colitis

Prior Authorization Group Description:

Biologic Immunomodulators PA - Xeljanz XR

Drug Name(s)

Xeljanz Xr

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. ONE of the following:
 - i. Patient's medication history indicates use of preferred TNF agent(s) OR
 - ii. Patient has an intolerance or hypersensitivity to preferred TNF agent(s) OR
 - iii. Patient has an FDA labeled contraindication to preferred TNF agent(s) AND
3. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical improvement (slowing of disease progression or decrease in symptom severity and/or frequency) AND
4. Patient will NOT be using the requested agent in combination with another biologic immunomodulator

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Use of ONE preferred TNF (Cyltezo, Enbrel, Hadlima, or Humira) is required for diagnoses of psoriatic arthritis, rheumatoid arthritis, or ankylosing spondylitis

Use of ONE preferred TNF (Cyltezo, Hadlima, or Humira) is required for diagnosis of ulcerative colitis

Prior Authorization Group Description:

Bivigam/Flebogamma/Gammaplex/Octagam/Privigen PA

Drug Name(s)

Gammaplex

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ONE of the following:

1. Patient has ONE of the following diagnoses:

- A. Primary immunodeficiency [e.g., congenital agammaglobulinemia, common variable immunodeficiency (CVID), severe combined immunodeficiency, Wiskott-Aldrich Syndrome, X-linked agammaglobulinemia (XLA), humoral immunodeficiency, IgG subclass deficiency with or without IgA deficiency] OR
- B. B-cell chronic lymphocytic leukemia OR multiple myeloma AND ONE of the following:
 - i. Patient has a history of infections OR
 - ii. Patient has evidence of specific antibody deficiency OR
 - iii. Patient has hypogammaglobulinemia OR
- C. Idiopathic thrombocytopenia purpura AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone), or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- D. Dermatomyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- E. Polymyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- F. Severe rheumatoid arthritis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., tumor necrosis factor antagonists (e.g., Humira), DMARDS (e.g., methotrexate)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

Criteria continues: see Other Criteria

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 6 months for indications in Other Criteria, 12 months for all others

Other Criteria:

G. Myasthenia gravis (MG) AND ONE of the following:

- i. Patient is in acute myasthenic crisis OR
- ii. Patient has severe refractory MG (e.g., major functional disability/weakness) AND ONE of the following:
 - a) Patient has failed ONE immunomodulator therapy (i.e., corticosteroid, pyridostigmine, or azathioprine) OR
 - b) Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE immunomodulator therapy OR

H. Multiple sclerosis (MS) AND BOTH of the following:

- i. Patient has a diagnosis of relapsing remitting MS (RRMS) AND
- ii. Patient has had an insufficient response, documented failure, or FDA labeled contraindication to TWO MS agents (e.g., Betaseron, Copaxone, dimethyl fumarate, fingolimod, Mayzent, Vumerity) OR

I. Acquired von Willebrand hemophilia AND ONE of the following:

- i. Patient has failed ONE conventional therapy (e.g., desmopressin solution, von Willebrand factor replacement therapy, corticosteroids, cyclophosphamide, FEIBA, or recombinant factor VIIa) OR
- ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

J. Refractory pemphigus vulgaris AND ONE of the following:

- i. Patient has failed ONE conventional immunosuppressive therapy (e.g., azathioprine, cyclophosphamide, mycophenolate, corticosteroids) OR
- ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional immunosuppressive therapy OR

2. ONE of the following:

- A. Patient has another FDA labeled indication for the requested agent OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Indications with 6 months approval duration: Acquired von Willebrand hemophilia, Guillain-Barre Syndrome, Lambert-Eaton myasthenia syndrome, Kawasaki disease, CMV induced pneumonitis in solid organ transplant, Toxic shock syndrome due to invasive group A streptococcus, Toxic epidermal necrolysis and Stevens-Johnson syndrome

Drug is also subject to Part B versus Part D review.

Prior Authorization Group Description:

Budesonide Oral ER PA – Entocort

Drug Name(s)

Budesonide

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Budesonide Oral ER PA – Uceris

Drug Name(s)

Budesonide Er

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Bydureon PA

Drug Name(s)

Bydureon Bcise

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Byetta PA

Drug Name(s)

Byetta

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Carglumic PA

Drug Name(s)

Carglumic Acid

Indications:

All FDA-Approved Indications, Some Medically-Accepted Indications.

Off-Label Uses:

Acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA)

Exclusion Criteria:**Required Medical Information:**

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of ONE of the following:
 - a. Acute hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS) OR
 - b. Chronic hyperammonemia due to the deficiency of the hepatic enzyme N-acetylglutamate synthase (NAGS) OR
 - c. Acute hyperammonemia due to propionic acidemia (PA) or methylmalonic acidemia (MMA)AND
2. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., geneticist, nephrologist, metabolic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Chenodal PA

Drug Name(s)

Chenodal

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of radiolucent stones in a well-opacifying gallbladder AND
2. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Cinacalcet PA

Drug Name(s)

Cinacalcet Hydrochloride

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has ONE of the following:

- A. A diagnosis of hypercalcemia due to parathyroid carcinoma OR
- B. A diagnosis of primary hyperparathyroidism (HPT) AND BOTH of the following:
 - i. Patient has a pretreatment serum calcium level that is above the testing laboratory's upper limit of normal AND
 - ii. Patient is unable to undergo parathyroidectomy OR
- C. Another indication that is FDA approved or supported in CMS approved compendia for the requested agent not otherwise excluded from Part D [i.e., secondary hyperparathyroidism due to end-stage renal disease (ESRD) on dialysis]

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Cobenfy PA

Drug Name(s)

Cobenfy

Indications:

All FDA-Approved Indications

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Prescriber has assessed the patient's liver enzymes and bilirubin prior to starting therapy with the requested agent AND
 - ii. Prescriber has assessed the patient's heart rate prior to starting therapy with the requested agent AND
 - iii. ONE of the following:
 - a. Patient has tried and had an inadequate response to TWO antipsychotic agents (e.g., aripiprazole, olanzapine, quetiapine, risperidone) for the requested indication OR
 - b. Patient has an intolerance or hypersensitivity to TWO antipsychotic agents (e.g., aripiprazole, olanzapine, quetiapine, risperidone) OR
 - c. Patient has an FDA labeled contraindication to TWO antipsychotic agents (e.g., aripiprazole, olanzapine, quetiapine, risperidone) AND
 - iv. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:

- i. Prescriber has assessed the patient's liver enzymes and bilirubin as clinically indicated during treatment with the requested agent AND
- ii. Prescriber has assessed the patient's heart rate as clinically indicated during treatment with the requested agent AND
- iii. Patient does NOT have any FDA labeled contraindications to the requested agent AND
- iv. Patient has had clinical benefit with the requested agent

Prior Authorization Group Description:

Colony Stimulating Factors PA – Granix

Drug Name(s)

Granix

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Other Criteria:

Prior Authorization Group Description:

Colony Stimulating Factors PA – Leukine

Drug Name(s)

Leukine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, hematologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 6 months

Other Criteria:

Prior Authorization Group Description:

Corlanor PA

Drug Name(s)

Corlanor

Ivabradine

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has stable, symptomatic chronic heart failure (e.g., NYHA Class II, III, IV: ACCF/AHA Class C, D)
AND

2. ONE of following:

a. ALL of the following:

- i. The requested agent is for a pediatric patient, 6 months of age or over AND
- ii. Patient has heart failure due to dilated cardiomyopathy (DCM) AND
- iii. Patient is in sinus rhythm with an elevated heart rate OR

b. ALL of the following:

- i. The requested agent is for an adult patient AND
- ii. Patient has a baseline OR current left ventricular ejection fraction of 35% or less AND
- iii. Patient is in sinus rhythm with a resting heart rate of 70 beats or greater per minute prior to initiating therapy with the requested agent AND
- iv. ONE of the following:

- 1. Patient is on a maximally tolerated dose of beta blocker (e.g., bisoprolol, carvedilol, metoprolol) OR
- 2. Patient has an intolerance, FDA labeled contraindications, or hypersensitivity to a beta blocker

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Cresemba PA

Drug Name(s)

Cresemba

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of invasive aspergillosis OR
- B. Patient has a diagnosis of invasive mucormycosis OR
- C. Patient has another indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

- A. Patient has a diagnosis of invasive aspergillosis and patient has continued indicators of active disease (e.g., continued radiologic findings, direct microscopy findings, histopathology findings, positive cultures, positive serum galactomannan assay) OR
- B. Patient has a diagnosis of invasive mucormycosis and patient has continued indicators of active disease (e.g., continued radiologic findings, direct microscopy findings, histopathology findings, positive cultures, positive serum galactomannan assay) OR
- C. BOTH of the following:
 - i. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient has had clinical benefit with the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 6 months

Other Criteria:

Prior Authorization Group Description:

Cystadrops PA

Drug Name(s)

Cystadrops

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., ophthalmologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Cystaran PA

Drug Name(s)

Cystaran

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., ophthalmologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Cystinosis Agents PA – Cystagon

Drug Name(s)

Cystagon

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of nephropathic cystinosis AND
2. Prescriber has performed a baseline white blood cell (WBC) cystine level test AND
3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of nephropathic cystinosis AND
3. Patient has had clinical benefit with the requested agent (e.g., decrease in WBC cystine levels from baseline) AND
4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Dalfampridine PA

Drug Name(s)

Dalfampridine Er

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has a diagnosis of multiple sclerosis (MS) AND
2. ONE of the following:
 - A. The requested agent will be used in combination with a disease modifying agent [e.g., Aubagio, Avonex, Bafiertam, Betaseron, dimethyl fumarate (e.g., Tecfidera), Extavia, fingolimod, glatiramer (e.g., Copaxone, Glatopa), Kesimpta, Mavenclad, Mayzent, Plegridy, Ponvory, Rebif, Vumerity, Zeposia] OR
 - B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to a disease modifying agent OR
 - C. Prescriber has provided information indicating that a disease modifying agent is not clinically appropriate for the patient

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of multiple sclerosis (MS) AND
3. ONE of the following:
 - A. The requested agent will be used in combination with a disease modifying agent [e.g., Aubagio, Avonex, Bafiertam, Betaseron, dimethyl fumarate (e.g., Tecfidera), Extavia, fingolimod, glatiramer (e.g., Copaxone, Glatopa), Kesimpta, Mavenclad, Mayzent, Plegridy, Ponvory, Rebif, Tecfidera, Vumerity, Zeposia] OR
 - B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to a disease modifying agent OR
 - C. Prescriber has provided information indicating that a disease modifying agent is not clinically appropriate for the patient AND
4. Patient has had improvements or stabilization from baseline in timed walking speed (timed 25-foot walk)

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Initial approval will be for 3 months, renewal approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Droxidopa PA

Drug Name(s)

Droxidopa

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of neurogenic orthostatic hypotension (nOH) AND
2. Prescriber has performed baseline blood pressure readings while the patient is sitting or supine (lying face up), AND also within three minutes of standing from a supine position AND
3. Patient has a decrease of at least 20 mmHg in systolic blood pressure or 10 mmHg diastolic blood pressure within three minutes after standing AND
4. Patient has persistent and consistent symptoms of neurogenic orthostatic hypotension (nOH) caused by ONE of the following:
 - A. Primary autonomic failure [Parkinson's disease (PD), multiple system atrophy, or pure autonomic failure] OR
 - B. Dopamine beta-hydroxylase deficiency OR
 - C. Non-diabetic autonomic neuropathy AND
5. Prescriber has assessed the severity of the patient's baseline symptoms of dizziness, lightheadedness, feeling faint, or feeling like the patient may black out AND
6. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of neurogenic orthostatic hypotension (nOH) AND
3. Patient has had improvements or stabilization with the requested agent as indicated by improvement in severity from baseline symptoms of ONE of the following:
 - A. Dizziness
 - B. Lightheadedness
 - C. Feeling faint
 - D. Feeling like the patient may black out AND
4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cardiologist, neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be 1 month for initial, 3 months for renewal

Other Criteria:

Prior Authorization Group Description:

Dupixent PA

Drug Name(s)

Dupixent

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

A. Patient has a diagnosis of moderate-to-severe atopic dermatitis AND ALL of the following:

i. ONE of the following:

a. Patient has tried and failed a topical steroid (e.g., triamcinolone) OR

b. Patient has an intolerance, hypersensitivity, or an FDA labeled contraindication to a topical steroid AND

ii. For patients 2 years of age or over, ONE of the following:

a. Patient has tried and failed a topical calcineurin inhibitor (e.g., pimecrolimus, tacrolimus) OR

b. Patient has an intolerance, hypersensitivity, or an FDA labeled contraindication to a topical calcineurin inhibitor AND

iii. Patient will NOT be using the requested agent in combination with another biologic agent or a JAK inhibitor for the requested indication (e.g., Adbry, Cibinqo, Opzelura, Rinvoq) OR

B. Patient has a diagnosis of moderate-to-severe asthma with an eosinophilic phenotype or oral corticosteroid dependent asthma AND BOTH of the following:

i. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LRTA, LAMA, theophylline) in combination with the requested agent AND

ii. Patient will NOT be using the requested agent in combination with Xolair or with an injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication OR

C. Patient has a diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) AND the following:

i. BOTH of the following:

a. ONE of the following:

1. Patient has tried and had an inadequate response to an oral systemic corticosteroid AND an intranasal corticosteroid (e.g., fluticasone) OR

2. Patient has an intolerance, hypersensitivity, or an FDA labeled contraindication to an oral systemic corticosteroid AND an intranasal corticosteroid AND

Initial criteria continues: see Other Criteria

Age Restriction:

For diagnosis of moderate-to-severe atopic dermatitis, patient (pt) is 6 months of age or over. For diagnosis of moderate-to-severe asthma with an eosinophilic phenotype or oral corticosteroid dependent asthma, pt is 6 years of age or over. For diagnosis of CRSwNP, pt is 12 years of age or over. For diagnosis of EoE, pt is 1 year of age or over. For diagnosis of PN, pt is 18 years of age or over. For diagnosis of COPD with an eosinophilic phenotype, pt is 18 years of age or over.

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., allergist, dermatologist, immunologist, gastroenterologist, otolaryngologist, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- b. Patient will continue standard maintenance therapy (e.g., intranasal corticosteroid) in combination with the requested agent OR
 - D. Patient has a diagnosis of eosinophilic esophagitis (EoE) confirmed by esophageal biopsy OR
 - E. Patient has a diagnosis of prurigo nodularis (PN) OR
 - F. Patient has a diagnosis of chronic obstructive pulmonary disease (COPD) with an eosinophilic phenotype AND BOTH of the following:
 - i. Patient is currently being treated with AND will continue COPD control therapy (e.g., ICS, LABA, LAMA) in combination with the requested agent AND
 - ii. Patient will NOT be using the requested agent in combination with Xolair or with an injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication AND
2. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. ONE of the following:
 - A. Patient has a diagnosis of moderate-to-severe atopic dermatitis AND the following:
 - i. Patient will NOT be using the requested agent in combination with another biologic agent or a JAK inhibitor for the requested indication (e.g., Adbry, Cibinqo, Opzelura, Rinvoq) OR
 - B. Patient has a diagnosis of moderate-to-severe asthma with an eosinophilic phenotype or oral corticosteroid dependent asthma AND BOTH of the following:
 - i. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LTRA, LAMA, theophylline) in combination with the requested agent AND
 - ii. Patient will NOT be using the requested agent in combination with Xolair or with an injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication OR
 - C. Patient has a diagnosis of chronic rhinosinusitis with nasal polyposis (CRSwNP) AND the following:
 - i. Patient will continue standard maintenance therapy (e.g., intranasal corticosteroid) in combination with the requested agent OR

- D. Patient has a diagnosis of eosinophilic esophagitis (EoE) OR
- E. Patient has a diagnosis of prurigo nodularis (PN) OR
- F. Patient has a diagnosis of chronic obstructive pulmonary disease (COPD) with an eosinophilic phenotype AND BOTH of the following:
 - i. Patient is currently being treated with AND will continue COPD control therapy (e.g., ICS, LABA, LAMA) in combination with the requested agent AND
 - ii. Patient will NOT be using the requested agent in combination with Xolair or with an injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Prior Authorization Group Description:

Emgality PA

Drug Name(s)

Emgality

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require the following:

1. ONE of the following:
 - A. Patient has a diagnosis of migraine AND ALL of the following:
 - i. The requested agent is being used for migraine prophylaxis AND
 - ii. Patient has 4 migraine headaches or more per month AND
 - iii. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis OR
 - B. Patient has a diagnosis of episodic cluster headache AND BOTH of the following:
 - i. Patient has had at least 5 cluster headache attacks AND
 - ii. Patient has had at least two cluster periods lasting 7 days to one year and separated by pain-free remission periods of 3 months or more

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. ONE of the following:
 - A. ALL of the following:
 - i. Patient has a diagnosis of migraine AND
 - ii. The requested agent is being used for migraine prophylaxis AND
 - iii. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis OR
 - B. Patient has a diagnosis of episodic cluster headache AND
3. Patient has had clinical benefit with the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Emsam PA

Drug Name(s)

Emsam

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. ONE of the following:
 - A. Patient has a diagnosis of major depressive disorder (MDD) OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. BOTH of the following:
 - i. ONE of the following:
 - a. BOTH of the following:
 - i. Patient has a diagnosis of major depressive disorder (MDD) AND
 - ii. ONE of the following:
 1. Patient has tried and had an inadequate response to at least two different oral antidepressants (e.g., SSRIs, SNRIs, mirtazapine, bupropion) OR
 2. Patient has an intolerance or hypersensitivity to at least two different oral antidepressants (e.g., SSRIs, SNRIs, mirtazapine, bupropion) OR
 3. Patient has an FDA labeled contraindication to at least two different oral antidepressants (e.g., SSRIs, SNRIs, mirtazapine, bupropion) OR
 - b. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. ONE of the following:

- A. Patient has a diagnosis of major depressive disorder (MDD) OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
3. ONE of the following:
- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days
 - OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. BOTH of the following:
 - i. Patient has had clinical benefit with the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Prior Authorization Group Description:

Endari PA

Drug Name(s)

Endari

L-glutamine

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of sickle cell disease AND
2. Patient is using the requested agent to reduce the acute complications of sickle cell disease AND
3. ONE of the following:
 - A. Patient has tried and had an inadequate response to maximally tolerated dose of hydroxyurea OR
 - B. Patient has an intolerance or hypersensitivity to hydroxyurea OR
 - C. Patient has an FDA labeled contraindication to hydroxyurea AND
4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Epclusa PA

Drug Name(s)

Epclusa

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. ONE of the following:

- A. Patient has a diagnosis of hepatitis C confirmed by serological markers OR
- B. Patient is a hepatitis C virus (HCV) - uninfected solid organ transplant recipient AND BOTH of the following:
 - i. Patient received an HCV - viremic donor organ AND
 - ii. The requested agent is being used for prophylaxis AND

2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND

3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND

4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group Description:

Epidiolex PA

Drug Name(s)

Epidiolex

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of seizures associated with ONE of the following:

- A. Lennox-Gastaut syndrome OR
- B. Dravet syndrome OR
- C. Tuberous sclerosis complex AND

2. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Erythropoietin Stimulating Agents PA - Epogen/Procrit

Drug Name(s)

Procrit

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. The requested agent is being prescribed for ONE of the following:

A. To reduce the possibility of allogeneic blood transfusion in a surgery patient AND the patient's hemoglobin level is greater than 10 g/dL but 13 g/dL or less OR

B. Anemia due to myelosuppressive chemotherapy for a non-myeloid malignancy AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR less than 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. Patient is being concurrently treated with chemotherapy with or without radiation (treatment period extends to 8 weeks post chemotherapy) AND

iii. The intent of chemotherapy is non-curative OR

C. Anemia associated with chronic kidney disease in a patient NOT on dialysis AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR 11 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. The rate of hemoglobin decline indicates the likelihood of requiring a RBC transfusion AND

iii. The intent of therapy is to reduce the risk of alloimmunization and/or other RBC transfusion related risks OR

D. Anemia due to myelodysplastic syndrome AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) OR

E. Anemia resulting from zidovudine treatment of HIV infection AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks)

OR

Initial criteria continues: see Other Criteria

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

1 month for surgery (reduce transfusion possibility), 6 months for chemo, 12 months for other

Other Criteria:

- F. Another indication that is supported in CMS approved compendia for the requested agent AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND
- 2. Patient's transferrin saturation and serum ferritin have been evaluated

Drug is also subject to Part B versus Part D review.

Prior Authorization Group Description:

Erythropoietin Stimulating Agents PA – Retacrit

Drug Name(s)

Retacrit

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. The requested agent is being prescribed for ONE of the following:

A. To reduce the possibility of allogeneic blood transfusion in a surgery patient AND the patient's hemoglobin level is greater than 10 g/dL but 13 g/dL or less OR

B. Anemia due to myelosuppressive chemotherapy for a non-myeloid malignancy AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR less than 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

ii. Patient is being concurrently treated with chemotherapy with or without radiation (treatment period extends to 8 weeks post chemotherapy) AND

iii. The intent of chemotherapy is non-curative OR

C. Anemia associated with chronic kidney disease in a patient NOT on dialysis AND ALL of the following:

i. Patient's hemoglobin level is less than 10 g/dL for patients initiating ESA therapy OR 11 g/dL or less for patients stabilized on therapy (measured within the previous 4 weeks)

AND

ii. The rate of hemoglobin decline indicates the likelihood of requiring a RBC transfusion AND

iii. The intent of therapy is to reduce the risk of alloimmunization and/or other RBC transfusion related risks OR

D. Anemia resulting from zidovudine treatment of HIV infection AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks)

OR

E. Another indication that is supported in CMS approved compendia for the requested agent AND the patient's hemoglobin level is less than 12 g/dL for patients initiating ESA therapy OR less than or equal to 12 g/dL for patients stabilized on therapy (measured within the previous 4 weeks) AND

2. Patient's transferrin saturation and serum ferritin have been evaluated

Drug is also subject to Part B versus Part D review.

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

1 month for surgery (reduce transfusion possibility), 6 months for chemo, 12 months for other

Other Criteria:

Prior Authorization Group Description:

Fentanyl Oral PA - Fentanyl lozenge

Drug Name(s)

Fentanyl Citrate Oral Transmucosal

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require BOTH of the following:

1. ONE of the following:

a. Patient has a documented diagnosis (i.e., medical records) of chronic cancer pain due to an active malignancy AND the following:

i. There is evidence of a claim that the patient is currently being treated with a long-acting opioid with the requested agent within the past 90 days OR

b. Patient has a diagnosis that is supported in CMS approved compendia for the requested agent AND

2. Patient will NOT be using the requested agent in combination with any other oral or nasal fentanyl agent

Age Restriction:

Patient is 16 years of age or over

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Fintepla PA

Drug Name(s)

Fintepla

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of seizures associated with Dravet syndrome (DS) or Lennox-Gastaut syndrome (LGS) AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. An echocardiogram assessment will be obtained before and during treatment with the requested agent, to evaluate for valvular heart disease and pulmonary arterial hypertension AND
 - ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND
 - iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Focalin PA

Drug Name(s)

Dexmethylphenidate Hydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Gammagard/Gammaked/Gamunex-C PA

Drug Name(s)

Gamunex-C

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ONE of the following:

1. Patient has ONE of the following diagnoses:

- A. Primary immunodeficiency [e.g., congenital agammaglobulinemia, common variable immunodeficiency (CVID), severe combined immunodeficiency, Wiskott-Aldrich Syndrome, X-linked agammaglobulinemia (XLA), humoral immunodeficiency, IgG subclass deficiency with or without IgA deficiency] OR
- B. B-cell chronic lymphocytic leukemia OR multiple myeloma AND ONE of the following:
 - i. Patient has a history of infections OR
 - ii. Patient has evidence of specific antibody deficiency OR
 - iii. Patient has hypogammaglobulinemia OR
- C. Idiopathic thrombocytopenia purpura AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone), or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- D. Dermatomyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- E. Polymyositis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., corticosteroids (e.g., methylprednisolone) or immunosuppressants (e.g., azathioprine)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR
- F. Severe rheumatoid arthritis AND ONE of the following:
 - i. Patient has failed ONE conventional therapy [e.g., tumor necrosis factor antagonists (e.g., Humira), DMARDS (e.g., methotrexate)] OR
 - ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

Criteria continues: see Other Criteria

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 6 months for indications in Other Criteria, 12 months for all others

Other Criteria:

G. Myasthenia gravis (MG) AND ONE of the following:

- i. Patient is in acute myasthenic crisis OR
- ii. Patient has severe refractory MG (e.g., major functional disability/weakness) AND ONE of the following:
 - a) Patient has failed ONE immunomodulator therapy (i.e., corticosteroid, pyridostigmine, or azathioprine) OR
 - b) Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE immunomodulator therapy OR

H. Multiple sclerosis (MS) AND BOTH of the following:

- i. Patient has a diagnosis of relapsing remitting MS (RRMS) AND
- ii. Patient has had an insufficient response, documented failure, or FDA labeled contraindication to TWO MS agents (e.g., Betaseron, Copaxone, dimethyl fumarate, fingolimod, Mayzent, Vumerity) OR

I. Acquired von Willebrand hemophilia AND ONE of the following:

- i. Patient has failed ONE conventional therapy (e.g., desmopressin solution, von Willebrand factor replacement therapy, corticosteroids, cyclophosphamide, FEIBA, or recombinant factor VIIa) OR
- ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional therapy OR

J. Refractory pemphigus vulgaris AND ONE of the following:

- i. Patient has failed ONE conventional immunosuppressive therapy (e.g., azathioprine, cyclophosphamide, mycophenolate, corticosteroids) OR
- ii. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ONE conventional immunosuppressive therapy OR

2. ONE of the following:

- A. Patient has another FDA labeled indication for the requested agent OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Indications with 6 months approval duration: Acquired von Willebrand hemophilia, Guillain-Barre Syndrome, Lambert-Eaton myasthenia syndrome, Kawasaki disease, CMV induced pneumonitis in solid organ transplant, Toxic shock syndrome due to invasive group A streptococcus, Toxic epidermal necrolysis and Stevens-Johnson syndrome

Drug is also subject to Part B versus Part D review.

Prior Authorization Group Description:

Gattex PA

Drug Name(s)

Gattex

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of short bowel syndrome (SBS) AND
2. Patient is dependent on parenteral nutrition OR intravenous (PN/IV) fluids AND
3. ONE of the following:
 - A. Patient is aged 1 year to 17 years AND BOTH of the following:
 - i. A fecal occult blood test has been performed within 6 months prior to initiating treatment with the requested agent AND
 - ii. ONE of the following:
 - a. There was no unexplained blood in the stool OR
 - b. There was unexplained blood in the stool AND a colonoscopy or a sigmoidoscopy was performed OR
 - B. Patient is 18 years of age or over AND BOTH of the following:
 - i. Patient has had a colonoscopy within 6 months prior to initiating treatment with the requested agent AND
 - ii. If polyps were present at this colonoscopy, the polyps were removed AND
4. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of short bowel syndrome (SBS) AND
3. Patient has had a reduction from baseline in parenteral nutrition OR intravenous (PN/IV) fluids AND
4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal

Other Criteria:

Prior Authorization Group Description:

Growth Hormone PA – Omnitrope

Drug Name(s)

Omnitrope

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

For Children – Criteria for initial approval require the following:

1. ONE of the following:

- a. Patient has a diagnosis of Turner Syndrome OR
- b. Patient has a diagnosis of Prader-Willi Syndrome OR
- c. Patient has a diagnosis of panhypopituitarism AND BOTH of the following:
 - i. Deficiencies in 3 or more pituitary axes AND
 - ii. Measured serum IGF-1 (insulin-like growth factor-1) levels are below the age and sex-appropriate reference range when off GH therapy OR
- d. Patient has a diagnosis of growth hormone deficiency (GHD) or short stature AND BOTH of the following:
 - i. Patient has ONE of the following:
 - a) Height more than 2 standard deviations (SD) below the mean for age and sex OR
 - b) Height more than 1.5 SD below the midparental height OR
 - c) A decrease in height SD of more than 0.5 over one year in children at least 2 years of age OR
 - d) Height velocity more than 2 SD below the mean over one year or more than 1.5 SD sustained over two years AND
 - ii. Failure of at least 2 growth hormone (GH) stimulation tests (e.g., peak GH value of less than 10 mcg/L after stimulation, or otherwise considered abnormal as determined by testing lab) OR
- e. Patient has a diagnosis of small for gestational age (SGA) AND ALL of the following:
 - i. Patient is at least 2 years of age AND
 - ii. Documented birth weight and/or length that is 2 or more SD below the mean for gestational age AND
 - iii. At 24 months of age, the patient fails to manifest catch-up growth evidenced by a height that remains 2 or more SD below the mean for age and sex

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

For Children – Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the preferred agent through the plan's Prior Authorization criteria AND
2. Patient has been diagnosed with ONE of the following:
 - a. Growth Hormone Deficiency, Short Stature OR
 - b. Panhypopituitarism OR
 - c. Prader-Willi Syndrome OR
 - d. Small for Gestational Age (SGA) OR
 - e. Turner Syndrome AND
3. ALL of the following:
 - a. Patient does NOT have closed epiphyses AND
 - b. Patient is being monitored for adverse effects of therapy with the requested agent AND
 - c. Patient's height has increased or height velocity has improved since initiation or last approval of the requested agent

For Adults – Criteria for initial approval require the following:

1. Patient has been diagnosed with ONE of the following:
 - a. Childhood growth hormone deficiency (GHD) with genetic or organic origin AND ONE of the following:
 - i. Low IGF-1 (insulin-like growth factor-1) level without GH replacement therapy OR
 - ii. Failure of at least one growth hormone (GH) stimulation test as an adult (e.g., peak GH value of 5 mcg/L or lower after stimulation, or otherwise considered abnormal as determined by testing lab) OR
 - b. Acquired adult GHD secondary to structural lesions or trauma AND ONE of the following:
 - i. Patient has a diagnosis of panhypopituitarism AND BOTH of the following:
 - a) Deficiencies in 3 or more pituitary axes AND
 - b) Low IGF-1 level without GH replacement therapy OR
 - ii. Patient has failed at least one growth hormone (GH) stimulation test as an adult OR
 - c. Idiopathic GHD (adult or childhood onset) AND the patient has failed at least two growth hormone (GH) stimulation tests as an adult

For Adults – Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the preferred agent through the plan's Prior Authorization criteria AND
2. Patient has been diagnosed with ONE of the following:
 - a. Childhood growth hormone deficiency (GHD) with genetic or organic origin OR
 - b. Acquired adult GHD secondary to structural lesions or trauma OR
 - c. Idiopathic GHD (adult or childhood onset) AND
3. Patient is being monitored for adverse effects of therapy with the requested agent AND
4. Patient's IGF-1 level has been evaluated to confirm the appropriateness of the current dose AND
5. Patient has had clinical benefit with the requested agent (i.e., body composition, hip-to-waist ratio, cardiovascular health, bone mineral density, serum cholesterol, physical strength, or quality of life)

Prior Authorization Group Description:

HAE PA – Cinryze

Drug Name(s)

Cinryze

Indications:

All FDA-Approved Indications, Some Medically-Accepted Indications.

Off-Label Uses:

Acute HAE attacks

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hereditary angioedema (HAE) which has been confirmed via two confirmatory tests of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:
 - a. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type I)]: decreased quantities of C4 and C1-INH (antigenic and functional level) OR
 - b. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type II)]: decreased quantities of C4 and C1-INH function (C1-INH protein level may be normal) OR
 - c. Hereditary angioedema (HAE) with normal C1INH [HAE-nI-C1INH (Type III)]: Normal levels of C4 and C1-INH [antigenic and functional level (at baseline and during an attack)] AND ONE of the following:
 - i. BOTH of the following:
 1. Family history of angioedema AND
 2. ALL other causes of angioedema have been ruled out OR
 - ii. Patient demonstrates a Factor XII mutation, angiopoietin-1 (ANGPT1) mutation, plasminogen (PLG) mutation, kininogen1 mutation, heparan sulfate 3-O-sulfotransferase 6 gene mutation, or myoferlin gene mutation that is associated with the disease AND
2. Medications known to cause angioedema (e.g., ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND
3. ONE of the following:
 - a. The requested agent will be used to treat acute HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks OR
 - b. The requested agent will be used for prophylaxis against HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria
AND
2. Patient has a diagnosis of hereditary angioedema (HAE) AND ONE of the following:
 - a. The requested agent will be used to treat acute HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks OR
 - b. The requested agent will be used for prophylaxis against HAE attacks AND the patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks AND
3. Patient has had a decrease in the frequency or severity of acute attacks or has had stabilization of disease from use of the requested agent

Prior Authorization Group Description:

HAE PA – Haegarda

Drug Name(s)

Haegarda

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hereditary angioedema (HAE) which has been confirmed via two confirmatory tests of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:
 - a. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type I)]: decreased quantities of C4 and C1-INH (antigenic and functional level) OR
 - b. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type II)]: decreased quantities of C4 and C1-INH function (C1-INH protein level may be normal) OR
 - c. Hereditary angioedema (HAE) with normal C1INH [HAE-nI-C1INH (Type III)]: Normal levels of C4 and C1-INH [antigenic and functional level (at baseline and during an attack)] AND ONE of the following:
 - i. BOTH of the following:
 1. Family history of angioedema AND
 2. ALL other causes of angioedema have been ruled out OR
 - ii. Patient demonstrates a Factor XII mutation, angiopoietin-1 (ANGPT1) mutation, plasminogen (PLG) mutation, kininogen1 mutation, heparan sulfate 3-O-sulfotransferase 6 gene mutation, or myoferlin gene mutation that is associated with the disease AND
2. Medications known to cause angioedema (e.g., ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND
3. The requested agent will be used for prophylaxis against HAE attacks AND
4. Patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of hereditary angioedema (HAE) AND
3. The requested agent is being used for prophylaxis against HAE attacks AND
4. Patient has had a decrease in the frequency or severity of acute attacks or has had stabilization of disease from use of the requested agent AND

5. Patient will NOT be using the requested agent in combination with another HAE agent indicated for prophylaxis against HAE attacks

Prior Authorization Group Description:

HAE PA – Icatibant

Drug Name(s)

Icatibant Acetate

Sajazir

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hereditary angioedema (HAE) which has been confirmed via two confirmatory tests of C1-INH antigenic level, C1-INH functional level, and C4 level as follows:
 - a. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type I)]: decreased quantities of C4 and C1-INH (antigenic and functional level) OR
 - b. Hereditary angioedema (HAE) due to C1INH deficiency [HAE-C1INH (Type II)]: decreased quantities of C4 and C1-INH function (C1-INH protein level may be normal) OR
 - c. Hereditary angioedema (HAE) with normal C1INH [HAE-nI-C1INH (Type III)]: Normal levels of C4 and C1-INH [antigenic and functional level (at baseline and during an attack)] AND ONE of the following:
 - i. BOTH of the following:
 1. Family history of angioedema AND
 2. ALL other causes of angioedema have been ruled out OR
 - ii. Patient demonstrates a Factor XII mutation, angiopoietin-1 (ANGPT1) mutation, plasminogen (PLG) mutation, kininogen1 mutation, heparan sulfate 3-O-sulfotransferase 6 gene mutation, or myoferlin gene mutation that is associated with the disease AND
2. Medications known to cause angioedema (e.g., ACE-Inhibitors, estrogens, angiotensin II receptor blockers) have been evaluated and discontinued when appropriate AND
3. The requested agent will be used to treat acute HAE attacks AND
4. Patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of hereditary angioedema (HAE) AND
3. The requested agent will be used to treat acute HAE attacks AND

4. Patient will NOT be using the requested agent in combination with another HAE agent indicated for treatment of acute HAE attacks AND
5. Patient has had a decrease in the frequency or severity of acute attacks or stabilization of disease from use of the requested agent

Prior Authorization Group Description:

Harvoni PA

Drug Name(s)

Harvoni

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA dosing for the requested indication

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group Description:

High Risk Medication PA - All Starts

Drug Name(s)

Benztropine Mesylate

Clemastine Fumarate

Cyproheptadine Hydrochloride

Dicyclomine Hydrochloride

Diphenoxylate Hydrochloride/Atropine Sulfate

Hydroxyzine Hydrochloride

Promethazine Hcl Plain

Promethazine Hydrochloride

Scopolamine

Trihexyphenidyl Hydrochloride

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

PA does NOT apply to patients less than 65 years of age.

Criteria for approval require ALL of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested high-risk medication AND
2. Prescriber has indicated that the benefits of the requested high-risk medication outweigh the risks for the patient AND
3. Prescriber has indicated that the risks and potential side effects of the requested high-risk medication have been discussed with the patient

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Imiquimod PA

Drug Name(s)

Imiquimod

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. Patient has ONE of the following diagnoses:

- A. Actinic keratosis OR
- B. Superficial basal cell carcinoma OR
- C. External genital and/or perianal warts/condyloma acuminata OR
- D. Squamous cell carcinoma OR
- E. Basal cell carcinoma OR
- F. Another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

4 months for Actinic keratosis, other diagnoses - see Other Criteria

Other Criteria:

2 months for Superficial basal cell carcinoma, Squamous cell carcinoma, or Basal cell carcinoma

4 months for External genital and/or perianal warts/condyloma acuminata

12 months for All other diagnoses

Prior Authorization Group Description:

Inbrija PA

Drug Name(s)

Inbrija

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ALL of the following:

1. The requested agent will be used for intermittent treatment of OFF episodes in patients with Parkinson's disease AND
2. Patient is receiving concurrent therapy with carbidopa/levodopa AND
3. Patient will NOT be using a nonselective monoamine oxidase (MAO) inhibitor (e.g., phenelzine, tranylcypromine) in combination with, or within 2 weeks of, the requested agent

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Ingrezza PA

Drug Name(s)

Ingrezza

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. ONE of the following:

A. Patient has a diagnosis of chorea associated with Huntington's disease AND BOTH of the following:

i. ONE of the following:

1. Patient does NOT have a current diagnosis of depression OR
2. Patient has a current diagnosis of depression and is being treated for depression AND

ii. ONE of the following:

1. Patient does NOT have a diagnosis of passive suicidal ideation and/or behavior OR
2. Patient has a diagnosis of passive suicidal ideation and/or behavior and must NOT be actively suicidal OR

B. Patient has a diagnosis of tardive dyskinesia AND ONE of the following:

i. Prescriber has reduced the dose of or discontinued any medications known to cause tardive dyskinesia (i.e., dopamine receptor blocking agents) OR

ii. Prescriber has provided clinical rationale indicating that a reduced dose or discontinuation of any medications known to cause tardive dyskinesia is not appropriate

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Injectable Oncology PA

Drug Name(s)

Margenza

Nelarabine

Synribo

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. ONE of the following:
 - a. The requested agent is FDA labeled or supported by CMS approved compendia as first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or CMS approved compendia supported therapy that are indicated as first-line therapy for the requested indication OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
 - iii. Patient does NOT have any FDA labeled contraindications to the requested agent AND
 - iv. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines

May also be subject to Part B versus Part D review.

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Iron Chelating Agents PA – Exjade

Drug Name(s)

Deferasirox (Exjade)

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome AND ONE of the following:

- i. A liver iron (Fe) concentration (LIC) of at least 5 mg Fe per gram of dry weight OR
- ii. A serum ferritin greater than 300 mcg/L OR
- iii. MRI confirmation of iron deposition OR

B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND

2. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome OR

B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent for the requested indication

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Iron Chelating Agents PA – Jadenu

Drug Name(s)

Deferasirox (Jadenu)

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome AND ONE of the following:

- i. A liver iron (Fe) concentration (LIC) of at least 5 mg Fe per gram of dry weight OR
- ii. A serum ferritin greater than 300 mcg/L OR
- iii. MRI confirmation of iron deposition OR

B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND

2. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of chronic iron overload due to a non-transfusion dependent thalassemia syndrome OR

B. Patient has a diagnosis of chronic iron overload due to blood transfusions AND

3. Patient has had clinical benefit with the requested agent AND

4. Patient will NOT be using the requested agent in combination with another iron chelating agent (e.g., deferiprone) for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent for the requested indication

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Ivermectin Cream PA

Drug Name(s)

Ivermectin Cream

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Ivermectin Tablet PA

Drug Name(s)

Ivermectin Tablet

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 4 months

Other Criteria:

Prior Authorization Group Description:

Kalydeco PA

Drug Name(s)

Kalydeco

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cystic fibrosis AND
2. ONE of the following:
 - A. Patient has ONE of the CFTR gene mutations or a mutation in the CFTR gene that is responsive based on in vitro data, as indicated in the FDA label, confirmed by genetic testing OR
 - B. Patient has another CFTR gene mutation(s) that is responsive to the requested agent, as indicated in the FDA label, confirmed by genetic testing AND
3. Patient is NOT homozygous for the F508del mutation AND
4. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of cystic fibrosis AND
3. Patient has had improvement or stabilization with the requested agent [e.g., improvement in FEV1 from baseline, increase in weight/BMI, improvement from baseline Cystic Fibrosis Questionnaire-Revised (CFQ-R) Respiratory Domain score, improvements in respiratory symptoms (cough, sputum production, and difficulty breathing), and/or reduced number of pulmonary exacerbations] AND
4. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Kerendia PA

Drug Name(s)

Kerendia

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Korlym PA

Drug Name(s)

Korlym

Mifepristone

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of Cushing's syndrome AND
2. ONE of the following:
 - A. Patient has type 2 diabetes mellitus OR
 - B. Patient has glucose intolerance as defined by a 2-hour glucose tolerance test plasma glucose value of 140-199 mg/dL AND
3. ONE of the following:
 - A. Patient has failed surgical resection OR
 - B. Patient is NOT a candidate for surgical resection

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of Cushing's syndrome AND
3. Patient has had clinical benefit with the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Leuprolide PA

Drug Name(s)

Eligard

Leuprolide Acetate

Lupron Depot (1-Month)

Lupron Depot (3-Month)

Lupron Depot (4-Month)

Lupron Depot (6-Month)

Lupron Depot-Ped (1-Month)

Lupron Depot-Ped (3-Month)

Lupron Depot-Ped (6-Month)

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ALL of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. BOTH of the following:
 - i. Patient is NOT currently being treated with the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent AND
3. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Lidocaine Topical PA - Lidocaine Patch

Drug Name(s)

Lidocan

Lidocan III

Lidocaine Patch

Tridacaine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require BOTH of the following:

1. Patient has ONE of the following diagnoses:

- A. Pain associated with postherpetic neuralgia (PHN) OR
 - B. Pain associated with diabetic neuropathy OR
 - C. Neuropathic pain associated with cancer, or cancer treatment OR
 - D. Another diagnosis that is supported in CMS approved compendia for the requested agent
- AND

2. ONE of the following:

- A. Patient has tried and had an inadequate response to a conventional therapy [e.g., gabapentin, pregabalin, oral prescription NSAID (non-steroidal anti-inflammatory drug)] for the requested indication OR
- B. Patient has an intolerance or hypersensitivity to a conventional therapy OR
- C. Patient has an FDA labeled contraindication to a conventional therapy

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Lidocaine Topical PA - Lidocaine Solution

Drug Name(s)

Lidocaine Solution

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. The requested agent will be used for ONE of the following:
 - A. Topical anesthesia of accessible mucous membranes of the oral and nasal cavities OR
 - B. Topical anesthesia of accessible mucous membranes of proximal portions of the digestive tract OR
 - C. Another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Lidocaine Topical PA - Lidocaine/prilocaine Cream

Drug Name(s)

Lidocaine/Prilocaine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. The requested agent will be used for ONE of the following:
 - A. Local analgesia on normal intact skin OR
 - B. Topical anesthetic for dermal procedures OR
 - C. Adjunctive anesthesia prior to local anesthetic infiltration in adult male genital skin OR
 - D. Anesthesia for minor procedures on female external genitalia OR
 - E. Another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Linezolid PA

Drug Name(s)

Linezolid

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND ONE of the following:

- a. The requested agent is prescribed by an infectious disease specialist or the prescriber has consulted with an infectious disease specialist on treatment of this patient OR
- b. Patient has a documented infection due to vancomycin-resistant *Enterococcus faecium* OR
- c. Patient has a diagnosis of pneumonia caused by *Staphylococcus aureus* or *Streptococcus pneumoniae* AND ONE of the following:
 - i. Patient has a documented infection that is resistant to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole, OR that is resistant to vancomycin OR
 - ii. Patient has an intolerance or hypersensitivity to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR
 - iii. Patient has an FDA labeled contraindication to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR
 - iv. Patient has an intolerance or hypersensitivity to vancomycin OR
 - v. Patient has an FDA labeled contraindication to vancomycin OR
- d. Patient has a documented skin and skin structure infection, including diabetic foot infections, caused by *Staphylococcus aureus*, *Streptococcus pyogenes*, or *Streptococcus agalactiae* AND ONE of the following:
 - i. Patient has a documented infection that is resistant to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole, OR that is resistant to vancomycin at the site of infection OR
 - ii. Patient has an intolerance or hypersensitivity to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR
 - iii. Patient has an FDA labeled contraindication to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR

Criteria continues: see Other Criteria

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 3 months

Other Criteria:

- iv. Patient has an intolerance or hypersensitivity to vancomycin OR

- v. Patient has an FDA labeled contraindication to vancomycin AND
- 2. Patient will NOT be using the requested agent in combination with Sivextro (tedizolid) for the same infection AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Prior Authorization Group Description:

Memantine ER PA

Drug Name(s)

Memantine Hydrochloride Er

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

PA does NOT apply to patients greater than or equal to 30 years of age

Criteria for approval require the following:

1. Patient is younger than 30 years of age AND ONE of the following:
 - A. Patient has a diagnosis of moderate to severe dementia of the Alzheimer's type OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Memantine PA

Drug Name(s)

Memantine Hcl Titration Pak

Memantine Hydrochloride

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

PA does NOT apply to patients greater than or equal to 30 years of age

Criteria for approval require the following:

1. Patient is younger than 30 years of age AND ONE of the following:
 - A. Patient has a diagnosis of moderate to severe dementia of the Alzheimer's type OR
 - B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Methylphenidate ER Tablet PA

Drug Name(s)

Methylphenidate Hydrochloride Er Tablet

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Migranal PA

Drug Name(s)

Dihydroergotamine Mesylate Spray

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. The requested agent will be used for the treatment of acute migraine with or without aura AND
2. ONE of the following:
 - A. Patient has tried and had an inadequate response to TWO triptan agents with differing active ingredients (e.g., sumatriptan, rizatriptan) OR
 - B. Patient has an intolerance or hypersensitivity to TWO triptan agents with differing active ingredients OR
 - C. Patient has an FDA labeled contraindication to TWO triptan agents with differing active ingredients AND
3. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, acute CGRP)

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. The requested agent will be used for the treatment of acute migraine with or without aura AND
3. Patient has had clinical benefit with the requested agent AND
4. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, acute CGRP)

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Modafinil PA

Drug Name(s)

Modafinil

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. Patient will NOT be using the requested agent in combination with another target agent (i.e., armodafinil)

Age Restriction:

Patient is 17 years of age or over

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Mounjaro PA

Drug Name(s)

Mounjaro

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

MS PA – Betaseron

Drug Name(s)

Betaseron

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical benefit with the requested agent AND
4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

MS PA - Dimethyl Fumarate

Drug Name(s)

Dimethyl Fumarate

Dimethyl Fumarate Starterpack

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical benefit with the requested agent AND
4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

MS PA – Fingolimod

Drug Name(s)

Fingolimod

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication AND
3. Prescriber has performed an electrocardiogram within 6 months prior to initiating treatment

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical benefit with the requested agent AND
4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

MS PA – Glatiramer

Drug Name(s)

Copaxone

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical benefit with the requested agent AND
4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

MS PA – Kesimpta

Drug Name(s)

Kesimpta

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical benefit with the requested agent AND
4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

MS PA – Mayzent

Drug Name(s)

Mayzent

Mayzent Starter Pack

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical benefit with the requested agent AND
4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

MS PA – Vumerity

Drug Name(s)

Vumerity

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical benefit with the requested agent AND
4. Patient will NOT be using the requested agent in combination with another disease modifying agent (DMA) or a biologic immunomodulator for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Myalept PA

Drug Name(s)

Myalept

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has leptin deficiency associated with a diagnosis of either congenital generalized lipodystrophy (CGL) or acquired generalized lipodystrophy (AGL) AND
2. Prescriber has provided the patient's baseline levels for HbA1C, triglycerides, and fasting insulin, measured prior to beginning therapy with the requested agent AND
3. Patient also has at least ONE of the complications related to lipodystrophy: diabetes mellitus, hypertriglyceridemia (200 mg/dL or higher), and/or high fasting insulin (30µU/mL or higher) AND
4. Patient has tried and had an inadequate response to maximum tolerable dosing of a conventional agent for the additional diagnosis AND
5. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has leptin deficiency associated with a diagnosis of either congenital generalized lipodystrophy (CGL) or acquired generalized lipodystrophy (AGL) AND
3. Patient has had improvement or stabilization with the requested agent as indicated by change from baseline level of at least ONE of the following:
 - A. HbA1C
 - B. Triglycerides
 - C. Fasting insulin AND
4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Conventional agent examples include:

Hypertriglyceridemia: statins, fenofibrates, Omega-3-Acid Ethyl Esters (generic Lovaza)

Diabetes/high fasting insulin: insulin, sulfonylurea/sulfonylurea combination, metformin/metformin combination

Prior Authorization Group Description:

Natpara PA

Drug Name(s)

Natpara

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

Increased baseline risk for osteosarcoma (e.g., Paget's disease of bone, unexplained elevations of alkaline phosphatase, hereditary disorders predisposing to osteosarcoma, history of external beam or implant radiation therapy involving the skeleton, pediatric and young adult patients with open epiphyses)

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of hypocalcemia associated with hypoparathyroidism AND
2. Patient does NOT have a baseline vitamin D level below the testing laboratory's lower limit of normal AND
3. Patient's baseline serum calcium level (albumin-corrected) is above 7.5 mg/dL AND
4. Patient will NOT be using the requested agent in combination with alendronate for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of hypocalcemia associated with hypoparathyroidism AND
3. Patient has had clinical benefit with the requested agent AND
4. Patient will NOT be using the requested agent in combination with alendronate for the requested indication

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, nephrologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Nuedexta PA

Drug Name(s)

Nuedexta

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. ONE of the following:

A. Patient has a diagnosis of pseudobulbar affect OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. Patient will NOT be using the requested agent in combination with a monoamine oxidase inhibitor (MAOI) [e.g., Marplan (isocarboxazid), Nardil (phenelzine), and Parnate (tranylcypromine)]

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Nuplazid PA

Drug Name(s)

Nuplazid

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Nurtec PA

Drug Name(s)

Nurtec

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require BOTH of the following:

1. Patient has a diagnosis of migraine AND
2. ONE of the following:
 - A. The requested agent is being used for the treatment of acute migraine with or without aura AND BOTH of the following:
 - i. ONE of the following:
 - a. Patient has tried and had an inadequate response to a triptan (e.g., sumatriptan, rizatriptan) agent OR
 - b. Patient has an intolerance, or hypersensitivity to a triptan OR
 - c. Patient has an FDA labeled contraindication to a triptan AND
 - ii. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, ergotamine, acute CGRP) OR
 - B. The requested agent is being used for migraine prophylaxis AND BOTH of the following:
 - i. Patient has 4 migraine headaches or more per month AND
 - ii. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 Months

Other Criteria:

Criteria for renewal require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of migraine AND
3. ONE of the following:
 - A. The requested agent is being used for the treatment of acute migraine with or without aura AND BOTH of the following:
 - i. Patient has had clinical benefit with the requested agent AND
 - ii. Patient will NOT be using the requested agent in combination with another acute migraine agent (e.g., triptan, 5HT-1F, ergotamine, acute CGRP) OR
 - B. The requested agent is being used for migraine prophylaxis AND BOTH of the following:
 - i. Patient has had clinical benefit with the requested agent AND
 - ii. Patient will NOT be using the requested agent in combination with another calcitonin gene-related peptide (CGRP) agent for migraine prophylaxis

Prior Authorization Group Description:

Ocaliva PA

Drug Name(s)

Ocaliva

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of primary biliary cholangitis (PBC) confirmed by at least TWO of the following:
 - A. There is biochemical evidence of cholestasis with an alkaline phosphatase (ALP) elevation
 - B. Presence of antimitochondrial antibody (AMA): a titer greater than 1:80 OR a level that is above the testing laboratory's upper limit of the normal range
 - C. If the AMA is negative or present only in low titer (less than or equal to 1:80), presence of other PBC-specific autoantibodies, including sp100 or gp210
 - D. Histologic evidence of nonsuppurative destruction cholangitis and destruction of interlobular bile ducts AND
2. ONE of the following:
 - A. Patient does NOT have cirrhosis OR
 - B. Patient has compensated cirrhosis with NO evidence of portal hypertension AND
3. Prescriber has measured the patient's alkaline phosphatase (ALP) level AND total bilirubin level AND
4. ONE of the following:
 - A. BOTH of the following:
 - i. Patient has tried and had an inadequate response to ursodiol AND
 - ii. The requested agent will be used in combination with ursodiol OR
 - B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ursodiol

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of primary biliary cholangitis (PBC) AND
3. ONE of the following:
 - A. Patient does NOT have cirrhosis OR
 - B. Patient has compensated cirrhosis with NO evidence of portal hypertension AND
4. ONE of the following:
 - A. The requested agent will be used in combination with ursodiol OR
 - B. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to ursodiol AND

5. Patient has had improvements or stabilization with the requested agent as indicated by BOTH of the following:

- A. Decrease in alkaline phosphatase (ALP) level from baseline AND
- B. Total bilirubin is less than or equal to the upper limit of normal (ULN)

Prior Authorization Group Description:

Ofev PA

Drug Name(s)

Ofev

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require the following:

1. ONE of the following:
 - A. BOTH of the following:
 - i. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND
 - ii. Patient has no known explanation for interstitial lung disease (ILD) or pulmonary fibrosis (e.g., radiation, drugs, metal dusts, sarcoidosis, or any connective tissue disease known to cause ILD) OR
 - B. BOTH of the following:
 - i. Patient has a diagnosis of systemic sclerosis-associated interstitial lung disease (SSc-ILD) AND
 - ii. Patient's diagnosis has been confirmed on high-resolution computed tomography (HRCT) or chest radiography scans OR
 - C. BOTH of the following:
 - i. Patient has a diagnosis of chronic fibrosing interstitial lung disease (ILD) with a progressive phenotype AND
 - ii. Patient's diagnosis has been confirmed on high-resolution computed tomography (HRCT)

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of ONE of the following:
 - A. Idiopathic pulmonary fibrosis (IPF) OR
 - B. Systemic sclerosis-associated interstitial lung disease (SSc-ILD) OR
 - C. Chronic fibrosing interstitial lung disease (ILD) with a progressive phenotype AND
3. Patient has had clinical benefit with the requested agent

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., pathologist, pulmonologist, radiologist, rheumatologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Opioids ER PA - Fentanyl Patch

Drug Name(s)

Fentanyl

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient has a diagnosis of pain due to sickle cell disease OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - iii. ALL of the following:
 - a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - d. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR
 - 3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND
 - e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND
 - f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Opioids ER PA – Morphine

Drug Name(s)

Morphine Sulfate Er

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient has a diagnosis of pain due to sickle cell disease OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - iii. ALL of the following:
 - a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - d. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR
 - 3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND
 - e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND
 - f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Opioids ER PA – Tramadol

Drug Name(s)

Tramadol Hcl Er

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of chronic cancer pain due to an active malignancy OR
- B. Patient has a diagnosis of pain due to sickle cell disease OR
- C. Patient is undergoing treatment of chronic non-cancer pain AND ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - iii. ALL of the following:
 - a. Prescriber has provided documentation of a formal, consultative evaluation including BOTH of the following:
 - 1. Diagnosis AND
 - 2. A complete medical history which includes previous and current pharmacological and non-pharmacological therapy AND
 - b. The requested agent is NOT prescribed as an as-needed (prn) analgesic AND
 - c. Prescriber has confirmed that a patient-specific pain management plan is on file for the patient AND
 - d. ONE of the following:
 - 1. Patient's medication history includes use of an immediate-acting opioid OR
 - 2. Patient has an intolerance or hypersensitivity to an immediate-acting opioid OR
 - 3. Patient has an FDA labeled contraindication to an immediate-acting opioid AND
 - e. Prescriber has reviewed the patient's records in the state's prescription drug monitoring program (PDMP) AND has determined that the opioid dosages and combinations of opioids and other controlled substances within the patient's records do NOT indicate the patient is at high risk for overdose AND
 - f. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Orkambi PA

Drug Name(s)

Orkambi

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cystic fibrosis AND
2. ONE of the following:
 - A. Patient has the presence of the F508del mutation on both alleles (homozygous) of the CFTR gene confirmed by genetic testing OR
 - B. Patient has another CFTR gene mutation(s) that is responsive to the requested agent, as indicated in the FDA label, confirmed by genetic testing AND
3. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of cystic fibrosis AND
3. Patient has had improvement or stabilization with the requested agent [e.g., improvement in FEV1 from baseline, increase in weight/BMI, improvement from baseline Cystic Fibrosis Questionnaire-Revised (CFQ-R) Respiratory Domain score, improvements in respiratory symptoms (cough, sputum production, and difficulty breathing), and/or reduced number of pulmonary exacerbations] AND
4. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Otezla PA

Drug Name(s)

Otezla

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ONE of the following:

1. BOTH of the following:
 - A. Patient has ONE of the following diagnoses:
 - i. Plaque psoriasis OR
 - ii. Active psoriatic arthritis AND
 - B. ONE of the following:
 - i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - ii. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - iii. Patient's medication history indicates use of a biologic immunomodulator agent for the same FDA labeled indication OR
 - iv. Patient has tried and had an inadequate response to at least ONE conventional prerequisite agent for the requested indication OR
 - v. Patient has an intolerance or hypersensitivity to at least ONE conventional prerequisite agent for the requested indication OR
 - vi. Patient has an FDA labeled contraindication to at least ONE conventional prerequisite agent for the requested indication OR
2. Patient has a diagnosis of oral ulcers associated with Behcet's disease (BD)

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of plaque psoriasis, active psoriatic arthritis, or oral ulcers associated with Behcet's disease (BD) AND
3. Patient has had clinical benefit with the requested agent (slowing of disease progression or decrease in symptom severity and/or frequency)

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Formulary conventional agent required for diagnoses of plaque psoriasis or active psoriatic arthritis

Formulary conventional agents for plaque psoriasis include cyclosporine, methotrexate, tazarotene, topical calcitriol, or topical corticosteroids

Formulary conventional agents for active psoriatic arthritis include cyclosporine, leflunomide, methotrexate, or sulfasalazine

NO prerequisites are required for a diagnosis of oral ulcers associated with Behcet's disease (BD)

Prior Authorization Group Description:

Ozempic PA

Drug Name(s)

Ozempic

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 5. BOTH of the following:
 - a. Patient has a diagnosis of established cardiovascular disease [e.g., myocardial infarction, stroke, any revascularization procedure, transient ischemic attack, unstable angina, amputation, symptomatic or asymptomatic coronary artery disease] AND
 - b. The requested agent will be used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Palynziq PA

Drug Name(s)

Palynziq

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of phenylketonuria (PKU) AND
2. Patient has a baseline blood Phe level greater than 600 micromol/L (10 mg/dL) AND
3. Patient will NOT be using the requested agent in combination with sapropterin for the requested indication AND
4. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of phenylketonuria (PKU) AND
3. ONE of the following:
 - a. Patient's blood Phe levels are being maintained within the acceptable range OR
 - b. Patient has had a decrease in blood Phe level from baseline AND
4. Patient will NOT be using the requested agent in combination with sapropterin for the requested indication AND
5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., metabolic or genetic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Initial approval will be for 9 months, renewal approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Panretin PA

Drug Name(s)

Panretin

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require BOTH of the following:

1. ONE of the following:

- A. Patient has a diagnosis of cutaneous lesions associated with AIDS-related Kaposi's sarcoma (KS) OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
- B. Prescriber states the patient is currently being treated with the requested agent OR
- C. ALL of the following:
 - i. ONE of the following:
 - 1. BOTH of the following:
 - a. Patient has a diagnosis of cutaneous lesions associated with AIDS-related Kaposi's sarcoma (KS) AND
 - b. Patient does NOT require systemic anti-Kaposi's sarcoma therapy OR
 - 2. Patient has an indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Prescriber is a specialist in the area of the patient's diagnosis (e.g., oncologist, dermatologist, infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis AND
 - iii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Pegylated Interferon PA

Drug Name(s)

Pegasys

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

A. Patient has a diagnosis of chronic hepatitis B AND BOTH of the following:

- i. The chronic hepatitis B infection has been confirmed by serological markers AND
- ii. Patient has NOT been administered the requested agent for more than 48 weeks for the treatment of chronic hepatitis B OR

B. BOTH of the following:

- i. Patient has a diagnosis of chronic hepatitis C confirmed by serological markers AND
- ii. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype OR

C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

12 months for all other diagnoses. For hep B, hep C see Other Criteria

Other Criteria:

No prior peginterferon alfa use, approve 48 weeks for hepatitis B infection. Prior peginterferon alfa use, approve remainder of 48 weeks of total therapy for hepatitis B infection

Duration of therapy for hepatitis C: Based on FDA approved labeling or AASLD/IDSA guideline supported

Prior Authorization Group Description:

Pirfenidone PA

Drug Name(s)

Pirfenidone

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require BOTH of the following:

1. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND
2. Patient has no known explanation for interstitial lung disease (ILD) or pulmonary fibrosis (e.g., radiation, drugs, metal dusts, sarcoidosis, or any connective tissue disease known to cause ILD)

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of idiopathic pulmonary fibrosis (IPF) AND
3. Patient has had clinical benefit with the requested agent

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., pathologist, pulmonologist, radiologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Posaconazole PA

Drug Name(s)

Noxafil Pak

Posaconazole Dr

Posaconazole Susp

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. Patient has a diagnosis of oropharyngeal candidiasis AND ONE of the following:

- i. Patient has tried and had an inadequate response to fluconazole or an alternative antifungal agent OR
- ii. Patient has an intolerance or hypersensitivity to fluconazole or an alternative antifungal agent OR
- iii. Patient has an FDA labeled contraindication to fluconazole or an alternative antifungal agent OR

B. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or Candida AND patient is severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR

C. Patient has a diagnosis of invasive Aspergillus AND ONE of the following:

- i. Patient has tried and had an inadequate response to an alternative antifungal agent OR
- ii. Patient has an intolerance or hypersensitivity to an alternative antifungal agent OR
- iii. Patient has an FDA labeled contraindication to an alternative antifungal agent OR

D. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

One month for oropharyngeal candidiasis, 6 months for all other indications

Other Criteria:

Criteria for renewal approval require BOTH of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:

- A. The requested agent is being prescribed for prophylaxis of invasive *Aspergillus* or *Candida* and patient continues to be severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR
- B. Patient has a diagnosis of invasive *Aspergillus* AND patient has continued indicators of active disease (e.g., continued radiologic findings, positive cultures, positive serum galactomannan assay for *Aspergillus*) OR
- C. BOTH of the following:
 - i. Patient has a diagnosis of oropharyngeal candidiasis AND
 - ii. Patient has had clinical benefit with the requested agent OR
- D. BOTH of the following:
 - i. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient has had clinical benefit with the requested agent

Prior Authorization Group Description:

Prolia PA

Drug Name(s)

Prolia

Indications:

All FDA-Approved Indications, Some Medically-Accepted Indications.

Off-Label Uses:

Osteopenia (osteoporosis prophylaxis)

Exclusion Criteria:

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of:

1. ONE of:

A. Patient's (pt) sex is male or the pt is postmenopausal with a diagnosis of osteoporosis AND BOTH of:

i. Pt's diagnosis was confirmed by ONE of:

1. A fragility fracture in the hip or spine OR
2. A T-score of -2.5 or lower OR
3. A T-score of -1.0 to -2.5 AND ONE of:
 - a. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR
 - b. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR
 - c. A FRAX 10-year probability of hip fracture of 3% or greater AND

ii. ONE of:

1. Pt is at a very high fracture risk as defined by ONE of:
 - a. Pt had a recent fracture (within the past 12 months) OR
 - b. Pt had fractures while on FDA approved osteoporosis therapy OR
 - c. Pt has had multiple fractures OR
 - d. Pt had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR
 - e. Pt has a very low T-score (less than -3.0) OR
 - f. Pt is at high risk for falls or has a history of injurious falls OR
 - g. Pt has a very high fracture probability by FRAX (e.g., major osteoporosis fracture greater than 30%, hip fracture greater than 4.5%) or by other validated fracture risk algorithm OR
2. ONE of:
 - a. Pt's medication history includes use of a bisphosphonate OR
 - b. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR

B. Pt is requesting the agent for osteopenia (osteoporosis prophylaxis) AND ALL of:

i. ONE of:

1. Pt's sex is male and the pt is 50 years of age or over OR
2. Pt is postmenopausal AND

- ii. Pt has a T-score between -1.0 to -2.50 AND
- iii. ONE of:
 - a. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR
 - b. 10-year probability of a hip fracture 3% and greater per FRAX OR
 - c. 10-year probability of a major OP-related fracture 20% and greater per FRAX AND
- iv. ONE of:
 - a. Pt's medication history includes use of a bisphosphonate OR

Criteria continues: See Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- b. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR
- C. Pt's sex is a female with a diagnosis of breast cancer who is receiving aromatase inhibitor therapy AND ONE of:
 - i. Pt's medication history includes use of a bisphosphonate OR
 - ii. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR
- D. Pt's sex is male with a diagnosis of prostate cancer receiving androgen deprivation therapy (ADT) AND ONE of:
 - i. Pt's medication history includes use of a bisphosphonate OR
 - ii. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate OR
- E. Pt has a diagnosis of glucocorticoid-induced osteoporosis AND ALL of:
 - i. Pt is either initiating or continuing systemic glucocorticoids in a daily dose equivalent to 7.5 mg or greater of prednisone AND
 - ii. Pt is expected to remain on glucocorticoids for at least 6 months AND
 - iii. Pt's diagnosis was confirmed by ONE of:
 - 1. A fragility fracture in the hip or spine OR
 - 2. A T-score of -2.5 or lower OR
 - 3. A T-score of -1.0 to -2.5 AND ONE of the following:
 - a. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR
 - b. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR
 - c. A FRAX 10-year probability of hip fracture of 3% or greater AND
 - iv. ONE of:
 - 1. Pt is at a very high fracture risk as defined by ONE of the following:
 - a. Pt had a recent fracture (within the past 12 months) OR
 - b. Pt had fractures while on FDA approved osteoporosis therapy OR

- c. Pt has had multiple fractures OR
- d. Pt had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR
- e. Pt has a very low T-score (less than -3.0) OR
- f. Pt is at high risk for falls or has a history of injurious falls OR
- g. Pt has a very high fracture probability by FRAX (e.g., major osteoporosis fracture greater than 30%, hip fracture greater than 4.5%) or by other validated fracture risk algorithm OR

2. ONE of:

- a. Pt's medication history includes use of a bisphosphonate OR
- b. Pt has an intolerance, FDA labeled contraindication, or hypersensitivity to a bisphosphonate AND

2. ONE of:

- A. Pt has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR
- B. Pt has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR
- C. Prescriber has indicated that the pt is not at risk for hypocalcemia (not including risk associated with the requested agent) AND

3. Pt will NOT be using the requested agent in combination with a bisphosphonate, another form of denosumab (e.g., Xgeva), romosozumab-aqqg, or parathyroid hormone analog (e.g., abaloparatide, teriparatide) for the requested indication AND

4. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Prior Authorization Group Description:

Promacta PA

Drug Name(s)

Promacta

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ONE of the following:

1. Patient (pt) has a diagnosis of persistent or chronic immune (idiopathic) thrombocytopenia (ITP) AND ONE of the following:
 - A. Pt has tried and had an insufficient response to a corticosteroid or immunoglobulin (IVIg or anti-D) OR
 - B. Pt has an intolerance or hypersensitivity to a corticosteroid or immunoglobulin (IVIg or anti-D) OR
 - C. Pt has an FDA labeled contraindication to a corticosteroid or immunoglobulin (IVIg or anti-D) OR
 - D. Pt has had an insufficient response to a splenectomy OR
2. Pt has a diagnosis of hepatitis C associated thrombocytopenia AND ONE of the following:
 - A. Pt's platelet count is less than $75 \times 10^9/L$ AND the intent is to increase platelet counts sufficiently to initiate pegylated interferon therapy OR
 - B. Pt is on concurrent therapy with a pegylated interferon and ribavirin AND is at risk for discontinuing hepatitis C therapy due to thrombocytopenia OR
3. Pt has a diagnosis of severe aplastic anemia (SAA) AND ALL of the following:
 - A. Pt has at least 2 of the following blood criteria:
 - i. Neutrophils less than $0.5 \times 10^9/L$ OR
 - ii. Platelets less than $30 \times 10^9/L$ OR
 - iii. Reticulocyte count less than $60 \times 10^9/L$ AND
 - B. Pt has at least 1 of the following marrow criteria:
 - i. Severe hypocellularity is less than 25% OR
 - ii. Moderate hypocellularity is 25-50% with hematopoietic cells representing less than 30% of residual cells AND
 - C. ONE of the following:
 - i. Pt has tried and had an insufficient response to BOTH antithymocyte globulin (ATG) AND cyclosporine therapy OR
 - ii. BOTH of the following:
 1. Pt will be using the requested agent as first-line treatment (i.e., has not been treated with ATG and/or cyclosporine) AND
 2. Pt will use the requested agent in combination with standard immunosuppressive therapy (i.e., ATG AND cyclosporine) OR
4. Pt has another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Initial: 6 months for ITP. Renewal: 12 months for ITP. Other indications, see Other Criteria.

Other Criteria:

Criteria for renewal approval require BOTH of the following:

1. Pt has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Pt has a diagnosis of persistent or chronic immune (idiopathic) thrombocytopenia (ITP) AND ONE of the following:

i. Pt's platelet count is $50 \times 10^9/L$ or greater OR

ii. Pt's platelet count has increased sufficiently to avoid clinically significant bleeding OR

B. Pt has a diagnosis of hepatitis C associated thrombocytopenia AND BOTH of the following:

i. ONE of the following:

1. Pt will be initiating hepatitis C therapy with pegylated interferon and ribavirin OR

2. Pt will be maintaining hepatitis C therapy with pegylated interferon and ribavirin at the same time as the requested agent AND

ii. ONE of the following:

1. Pt's platelet count is $90 \times 10^9/L$ or greater OR

2. Pt's platelet count has increased sufficiently to initiate or maintain pegylated interferon based therapy for the treatment of hepatitis C OR

C. Pt has a diagnosis of severe aplastic anemia (SAA) AND the pt has had clinical benefit with the requested agent OR

D. Pt has another indication that is supported in CMS approved compendia and the pt has had clinical benefit with the requested agent

Initial: 48 weeks for hepatitis C associated thrombocytopenia, 6 months for first-line therapy in severe aplastic anemia, 16 weeks for SAA, 12 months for All other indications

Renewal: 48 weeks for hepatitis C associated thrombocytopenia, 12 months for SAA, 12 months for All other indications

Prior Authorization Group Description:

Pulmonary Hypertension PA – Adempas

Drug Name(s)

Adempas

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
- b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of chronic thromboembolic pulmonary hypertension (CTEPH), WHO Group 4, as determined by a ventilation-perfusion scan and a confirmatory selective pulmonary angiography AND ALL of the following:

i. ONE of the following:

- a. Patient is NOT a candidate for surgery OR
- b. Patient has had pulmonary endarterectomy AND has persistent or recurrent disease AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units OR

C. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

Initial criteria continues: see Other Criteria

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

- v. ONE of the following:
 - a. The requested agent will be utilized as monotherapy OR
 - b. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), AND BOTH of the following:
 - 1. Patient has unacceptable or deteriorating clinical status despite established pharmacotherapy AND
 - 2. The requested agent is in a different therapeutic class OR
 - c. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:
 - 1. ONE of the following:
 - i. A prostanoid has been started as one of the agents in the triple therapy OR
 - ii. Patient has an intolerance or hypersensitivity to a prostanoid OR
 - iii. Patient has an FDA labeled contraindication to a prostanoid AND
 - 2. Patient has unacceptable or deteriorating clinical status despite established pharmacotherapy AND
 - 3. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent

Prior Authorization Group Description:

Pulmonary Hypertension PA – Ambrisentan

Drug Name(s)

Ambrisentan

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
- b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

v. ONE of the following:

a. The requested agent will be utilized as monotherapy OR

b. The requested agent will be used in combination with a phosphodiesterase 5 (PDE5) inhibitor for dual therapy ONLY OR

c. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), [except for dual therapy requests for a phosphodiesterase 5 (PDE 5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:

1. ONE of the following:

i. A prostanoid has been started as one of the agents in the triple therapy OR

ii. Patient has an intolerance or hypersensitivity to a prostanoid OR

iii. Patient has an FDA labeled contraindication to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent

Prior Authorization Group Description:

Pulmonary Hypertension PA – Bosentan

Drug Name(s)

Bosentan

Tracleer

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

Elevated liver enzymes accompanied by signs or symptoms of liver dysfunction/injury or a bilirubin level of 2 times the ULN (upper limit of normal) or greater AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1, as determined by right heart catheterization, AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

v. ONE of the following:

a. The requested agent will be utilized as monotherapy OR

b. The requested agent will be used in combination with a phosphodiesterase 5 (PDE5) inhibitor for dual therapy ONLY OR

c. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), [except for dual therapy requests for a phosphodiesterase 5 inhibitor (PDE5) plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:

1. ONE of the following:

i. A prostanoid has been started as one of the agents in the triple therapy OR

ii. Patient has an intolerance or hypersensitivity to a prostanoid OR

iii. Patient has an FDA labeled contraindication to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic class OR

C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. Patient has had clinical benefit with the requested agent

Prior Authorization Group Description:

Pulmonary Hypertension PA – Opsumit

Drug Name(s)

Opsumit

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
- b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

v. ONE of the following:

a. The requested agent will be utilized as monotherapy OR

b. The requested agent will be used in combination with a phosphodiesterase 5 (PDE5) inhibitor for dual therapy ONLY OR

c. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), [except for dual therapy requests for a phosphodiesterase 5 (PDE 5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:

1. ONE of the following:

i. A prostanoid has been started as one of the agents in the triple therapy OR

ii. Patient has an intolerance or hypersensitivity to a prostanoid OR

iii. Patient has an FDA labeled contraindication to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic class

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication for the requested agent AND

3. Patient has had clinical benefit with the requested agent

Prior Authorization Group Description:

Pulmonary Hypertension PA – Sildenafil

Drug Name(s)

Sildenafil

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

Concurrently taking another phosphodiesterase 5 (PDE 5) inhibitor with the requested agent AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:
 - A. BOTH of the following:
 - i. ONE of the following:
 - a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND
 - ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR
 - B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:
 - i. Patient's World Health Organization (WHO) functional class is II or greater AND
 - ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND
 - iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND
 - iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND
 - v. ONE of the following:
 - a. The requested agent will be utilized as monotherapy OR
 - b. The requested agent will be used in combination with an endothelin receptor antagonist (ERA) for dual therapy ONLY OR
 - c. The requested agent will be utilized for add-on therapy to existing monotherapy, [except for dual requests for a phosphodiesterase 5 (PDE5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:
 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria

Age Restriction:**Prescriber Restrictions:**

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy) AND ALL of the following:
 - 1. ONE of the following:
 - i. A prostanoid has been started as one of the agents in the triple therapy OR
 - ii. Patient has an intolerance or hypersensitivity to a prostanoid OR
 - iii. Patient has an FDA labeled contraindication to a prostanoid AND
 - 2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 3. All three agents in the triple therapy are from a different therapeutic class OR
- C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent

Prior Authorization Group Description:

Pulmonary Hypertension PA – Tadalafil

Drug Name(s)

Alyq

Tadalafil 20Mg Tablet

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

Concurrently taking another phosphodiesterase 5 (PDE 5) inhibitor with the requested agent AND FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR

b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

i. Patient's World Health Organization (WHO) functional class is II or greater AND

ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND

iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND

iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

v. ONE of the following:

a. The requested agent will be utilized as monotherapy OR

b. The requested agent will be used in combination with an endothelin receptor antagonist (ERA) for dual therapy ONLY OR

c. The requested agent will be utilized for add-on therapy to existing monotherapy, [except for dual requests for a phosphodiesterase 5 (PDE5) inhibitor plus an endothelin receptor antagonist (ERA)], AND BOTH of the following:

1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

2. The requested agent is in a different therapeutic class OR

Initial criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

d. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy) AND ALL of the following:

1. ONE of the following:

i. A prostanoid has been started as one of the agents in the triple therapy OR

ii. Patient has an intolerance or hypersensitivity to a prostanoid OR

iii. Patient has an FDA labeled contraindication to a prostanoid AND

2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND

3. All three agents in the triple therapy are from a different therapeutic class OR

C. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND

3. Patient has had clinical benefit with the requested agent

Prior Authorization Group Description:

Pulmonary Hypertension PA – Ventavis

Drug Name(s)

Ventavis

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

i. ONE of the following:

- a. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
- b. Prescriber states the patient is currently being treated with the requested agent within the past 90 days AND

ii. Patient has an FDA labeled indication for the requested agent OR

B. Patient has a diagnosis of pulmonary arterial hypertension (PAH), WHO Group 1 as determined by right heart catheterization AND ALL of the following:

- i. Patient's World Health Organization (WHO) functional class is II or greater AND
- ii. Patient has a mean pulmonary arterial pressure greater than 20 mmHg AND
- iii. Patient has a pulmonary capillary wedge pressure less than or equal to 15 mmHg AND
- iv. Patient has a pulmonary vascular resistance greater than or equal to 3 Wood units AND

v. ONE of the following:

- a. The requested agent will be utilized as monotherapy OR
- b. The requested agent will be utilized for add-on therapy to existing monotherapy (dual therapy), AND BOTH of the following:
 - 1. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 2. The requested agent is in a different therapeutic class OR
- c. The requested agent will be utilized for add-on therapy to existing dual therapy (triple therapy), AND ALL of the following:
 - 1. Patient is WHO functional class III or IV AND
 - 2. Patient has unacceptable or deteriorating clinical status despite established PAH pharmacotherapy AND
 - 3. All three agents in the triple therapy are from a different therapeutic class

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has an FDA labeled indication for the requested agent AND
3. Patient has had clinical benefit with the requested agent

Drug is also subject to Part B versus Part D review.

Prior Authorization Group Description:

Pyrimethamine PA

Drug Name(s)

Pyrimethamine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 6 months

Other Criteria:

Prior Authorization Group Description:

Quinine PA

Drug Name(s)

Quinine Sulfate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has ONE of the following diagnoses:
 - a. Uncomplicated malaria OR
 - b. Babesiosis OR
 - c. An indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

7 days for malaria, 10 days for babesiosis, 12 months for all other diagnoses

Other Criteria:

Prior Authorization Group Description:

Repatha PA

Drug Name(s)

Repatha

Repatha Pushtronex System

Repatha Sureclick

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has ONE of the following:

- A. A diagnosis of heterozygous familial hypercholesterolemia (HeFH) AND ONE of the following:
 - i. Genetic confirmation of one mutant allele at the LDLR, Apo-B, PCSK9, or 1/LDLRAP1 gene OR
 - ii. History of LDL-C greater than 190 mg/dL (greater than 4.9 mmol/L) (pretreatment) OR
 - iii. Patient has clinical manifestations of HeFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthoma, or xanthelasma) OR
 - iv. Patient has “definite” or “possible” familial hypercholesterolemia as defined by the Simon Broome criteria OR
 - v. Patient has a Dutch Lipid Clinic Network criteria score of greater than 5 OR
 - vi. Patient has a treated low-density lipoprotein cholesterol (LDL-C) level 100 mg/dL or greater after treatment with antihyperlipidemic agents but prior to PCSK9 inhibitor therapy OR
- B. A diagnosis of homozygous familial hypercholesterolemia (HoFH) AND ONE of the following:
 - i. Genetic confirmation of two mutant alleles at the LDLR, Apo-B, PCSK9, or LDLRAP1 gene OR
 - ii. History of untreated LDL-C greater than 500 mg/dL (greater than 13 mmol/L) or treated LDL-C 300 mg/dL or greater (7.76 mmol/L or greater) OR
 - iii. Patient has clinical manifestations of HoFH (e.g., cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas, or xanthelasma) OR
- C. A diagnosis of established cardiovascular disease [angina pectoris, coronary heart disease, myocardial infarction, transient ischemic attacks, cerebrovascular disease (CeVD) or peripheral vascular disease (PVD) or after coronary revascularization or carotid endarterectomy] AND the requested agent will be used to reduce the risk of myocardial infarction, stroke, and coronary revascularization OR
- D. A diagnosis of primary hyperlipidemia (not associated with HeFH, HoFH, or established cardiovascular disease) OR

Initial criteria continues: see Other Criteria

Age Restriction:**Prescriber Restrictions:**

The agent was prescribed by, or in consultation with, a cardiologist, an endocrinologist, and/or a physician who focuses in the treatment of cardiovascular (CV) risk management and/or lipid disorders

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- E. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
- 3. ONE of the following:
 - A. Patient has tried and had an inadequate response to a high-intensity statin (i.e., rosuvastatin 20-40 mg or atorvastatin 40-80 mg) OR
 - B. Patient has an intolerance to TWO different statins OR
 - C. Patient has an FDA labeled contraindication to a statin AND
- 4. Patient will NOT be using the requested agent in combination with another PCSK9 agent

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
- 3. Patient has had clinical benefit with the requested agent AND
- 4. Patient will NOT be using the requested agent in combination with another PCSK9 agent

Prior Authorization Group Description:

Ritalin PA

Drug Name(s)

Methylphenidate Hydrochloride (Ritalin)

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has an FDA labeled indication for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Roflumilast PA

Drug Name(s)

Roflumilast

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication for the requested agent AND
2. ONE of the following:
 - A. Patient has tried and had an inadequate response to an agent from two of the following categories:
 - i. long-acting beta-2 agonist (LABA) [e.g., salmeterol]
 - ii. long-acting muscarinic antagonist (LAMA) [e.g., umeclidinium]
 - iii. inhaled corticosteroid (ICS) [e.g., fluticasone] OR
 - B. Patient has an intolerance or hypersensitivity to an agent from two of the following categories:
 - i. long-acting beta-2 agonist (LABA) [e.g., salmeterol]
 - ii. long-acting muscarinic antagonist (LAMA) [e.g., umeclidinium]
 - iii. inhaled corticosteroid (ICS) [e.g., fluticasone] OR
 - C. Patient has an FDA labeled contraindication to an agent from two of the following categories:
 - i. long-acting beta-2 agonist (LABA) [e.g., salmeterol]
 - ii. long-acting muscarinic antagonist (LAMA) [e.g., umeclidinium]
 - iii. inhaled corticosteroid (ICS) [e.g., fluticasone]

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Rybelsus PA

Drug Name(s)

Rybelsus

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Sapropterin PA

Drug Name(s)

Javygtor

Sapropterin Dihydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of phenylketonuria (PKU) AND
2. Prescriber has submitted a baseline blood Phe level measured prior to initiation of therapy with the requested agent, which is above the recommended levels indicated for the patient's age range or condition AND
3. Patient will NOT be using the requested agent in combination with Palynziq (pegvaliase-pqpz) for the requested indication AND
4. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of phenylketonuria (PKU) AND
3. ONE of the following:
 - a. Patient's blood Phe levels are being maintained within the acceptable range OR
 - b. Patient has had a decrease in blood Phe level from baseline AND
4. Patient will NOT be using the requested agent in combination with Palynziq (pegvaliase-pqpz) for the requested indication AND
5. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., metabolic or genetic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Initial: 2 months if dose is 5 to less than 20 mg/kg/day, 1 month if 20 mg/kg/day Renewal: 12 months

Other Criteria:

Prior Authorization Group Description:

Self - Administered Oncology PA

Drug Name(s)

Abiraterone Acetate
Akeega
Alecensa
Alunbrig
Augtyro
Ayvakit
Balversa
Besremi
Bexarotene Cap
Bosulif
Braftovi
Brukinsa
Cabometyx
Calquence
Caprelsa
Cometriq
Copiktra
Cotellic
Dasatinib
Daurismo
Erivedge
Erleada
Erlotinib Hydrochloride
Everolimus
Exkivity
Farydak
Fotivda
Fruzaqla
Gavreto
Gefitinib
Gilotrif
Ibrance
Iclusig
Idhifa
Imatinib Mesylate
Imbruvica
Inlyta
Inqovi
Inrebic
Iwilfin
Jakafi

Jaypirca
Kisqali
Kisqali Femara 200 Dose
Kisqali Femara 400 Dose
Kisqali Femara 600 Dose
Koselugo
Krazati
Lapatinib Ditosylate
Lazcluze
Lenalidomide
Lenvima 10 Mg Daily Dose
Lenvima 12Mg Daily Dose
Lenvima 14 Mg Daily Dose
Lenvima 18 Mg Daily Dose
Lenvima 20 Mg Daily Dose
Lenvima 24 Mg Daily Dose
Lenvima 4 Mg Daily Dose
Lenvima 8 Mg Daily Dose
Lonsurf
Lorbrena
Lumakras
Lynparza
Lytgobi
Matulane
Mekinist
Mektovi
Nerlynx
Ninlaro
Nubeqa
Odomzo
Ogsiveo
Ojemda
Ojjaara
Onureg
Orgovyx
Orserdu
Pazopanib
Pemazyre
Piqray 200Mg Daily Dose
Piqray 250Mg Daily Dose
Piqray 300Mg Daily Dose
Pomalyst
Qinlock
Retevmo

Revlimid
Rezlidhia
Rozlytrek
Rubraca
Rydapt
Scemblix
Sorafenib
Sprycel
Stivarga
Sunitinib Malate
Tabrecta
Tafinlar
Tagrisso
Talzenna
Tasigna
Tazverik
Tepmetko
Thalomid
Tibsovo
Torpenz
Tretinoin 10Mg Cap
Truqap
Tukysa
Turalio
Vanflyta
Venclexta
Venclexta Starting Pack
Verzenio
Vitrakvi
Vizimpro
Vonjo
Voranigo
Votrient
Welireg
Xalkori
Xospata
Xpovio
Xpovio 100 Mg Once Weekly
Xpovio 40 Mg Once Weekly
Xpovio 40 Mg Twice Weekly
Xpovio 60 Mg Once Weekly
Xpovio 60 Mg Twice Weekly
Xpovio 80 Mg Once Weekly
Xpovio 80 Mg Twice Weekly

Xtandi
Zejula
Zelboraf
Zolinza
Zydelig
Zykadia

Indications:

All Medically-Accepted Indications.

Off-Label Uses:

Exclusion Criteria:

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. ALL of the following:
 - i. Genetic testing has been completed, if required, for therapy with the requested agent and results indicate the requested agent is appropriate AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent AND
 - iii. ONE of the following:
 - a. The requested agent is FDA labeled or supported by compendia as a first-line therapy for the requested indication OR
 - b. Patient has tried appropriate FDA labeled or compendia supported therapy that are indicated as first-line therapy for the requested indication OR
 - c. Patient has an intolerance or hypersensitivity to the first-line therapy for the requested indication OR
 - d. Patient has an FDA labeled contraindication to the first-line therapy for the requested indication AND
 - iv. Patient does NOT have any FDA labeled limitations of use that is not otherwise supported in NCCN guidelines AND

Criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

Approval will be for 12 months

Other Criteria:

- v. ONE of the following:
 - a. The requested agent is not Bosulif OR
 - b. The requested agent is Bosulif AND ONE of the following:

1. Patient's medication history indicates use of imatinib OR Sprycel for the requested indication (if applicable) OR
 2. Patient has an intolerance or hypersensitivity to imatinib OR Sprycel OR
 3. Patient has an FDA labeled contraindication to imatinib OR Sprycel OR
 4. CMS approved compendia does not support the use of imatinib OR Sprycel for the requested indication OR
 5. Prescriber has provided information in support of use of Bosulif over imatinib OR Sprycel for the requested indication AND
- vi. ONE of the following:
- a. The requested agent is not Calquence OR
 - b. The requested agent is Calquence AND ONE of the following:
 1. Patient's medication history indicates use of Brukinsa OR Imbruvica for the requested indication (if applicable) OR
 2. Patient has an intolerance or hypersensitivity to Brukinsa OR Imbruvica OR
 3. Patient has an FDA labeled contraindication to Brukinsa OR Imbruvica OR
 4. CMS approved compendia do not support the use of Brukinsa OR Imbruvica for the requested indication OR
 5. Prescriber has provided information in support of use of Calquence over Brukinsa OR Imbruvica for the requested indication

Prior Authorization Group Description:

Signifor PA

Drug Name(s)

Signifor

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

Severe hepatic impairment (i.e., Child Pugh C)

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. Patient has a diagnosis of Cushing's disease (CD) AND ONE of the following:

i. Patient had an inadequate response to pituitary surgical resection OR

ii. Patient is NOT a candidate for pituitary surgical resection OR

B. Patient has an indication that is supported in CMS approved compendia for the requested agent

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of Cushing's disease (CD) AND BOTH of the following:

i. Patient has a urinary free cortisol level less than or equal to the upper limit of normal AND

ii. Patient has had improvement in at least ONE of the following clinical signs and symptoms:

1. Fasting plasma glucose OR

2. Hemoglobin A1c OR

3. Hypertension OR

4. Weight OR

B. BOTH of the following:

i. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

ii. Patient has had clinical benefit with the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Initial approval: 6 months for CD, 12 months for all other diagnoses, Renewal approval: 12 months

Other Criteria:

Prior Authorization Group Description:

Sivextro PA

Drug Name(s)

Sivextro

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ALL of the following:

1. Patient has ONE of the following:

A. BOTH of the following:

- i. A documented acute bacterial skin and skin structure infection (ABSSSI) defined as a bacterial infection of the skin with a lesion size area of at least 75 cm² (lesion size measured by the area of redness, edema, or induration) AND
- ii. The infection is due to *Staphylococcus aureus*, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus anginosus*, *Streptococcus intermedius*, *Streptococcus constellatus*, or *Enterococcus faecalis* OR

B. Another indication that is supported in CMS approved compendia for the requested agent
AND

2. ONE of the following:

A. The requested agent is prescribed by an infectious disease specialist or the prescriber has consulted with an infectious disease specialist on treatment of this patient OR

B. The requested agent is NOT prescribed by an infectious disease specialist or the prescriber has NOT consulted with an infectious disease specialist on treatment of this patient AND ONE of the following:

- i. There is documentation of resistance to TWO of the following: beta-lactams, macrolides, clindamycin, tetracycline, or co-trimoxazole at the site of infection OR
- ii. Patient has an intolerance or hypersensitivity to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR
- iii. Patient has an FDA labeled contraindication to TWO of the following: beta-lactams, macrolides, clindamycin, tetracyclines, or co-trimoxazole OR
- iv. There is documentation of resistance to vancomycin at the site of infection OR
- v. Patient has an intolerance or hypersensitivity to vancomycin OR
- vi. Patient has an FDA labeled contraindication to vancomycin AND

3. Patient will NOT be using the requested agent in combination with linezolid for the same infection
AND

4. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:**Coverage Duration:**

Approval will be 6 days for ABSSSI or 30 days for all other indications

Other Criteria:

Prior Authorization Group Description:

Sodium Oxybate PA

Drug Name(s)

Sodium Oxybate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of narcolepsy with cataplexy OR
- B. Patient has a diagnosis of narcolepsy with excessive daytime sleepiness AND BOTH of the following:

- i. ONE of the following:

- a. Patient is under 18 years of age OR

- b. ONE of the following:

- 1. Patient has tried and had an inadequate response to modafinil or armodafinil OR
 - 2. Patient has an intolerance or hypersensitivity to modafinil or armodafinil OR
 - 3. Patient has an FDA labeled contraindication to modafinil or armodafinil AND

- ii. ONE of the following:

- a. Patient has tried and had an inadequate response to ONE standard stimulant agent (e.g., methylphenidate) OR
 - b. Patient has an intolerance or hypersensitivity to ONE standard stimulant agent (e.g., methylphenidate) OR
 - c. Patient has an FDA labeled contraindication to ONE standard stimulant agent (e.g., methylphenidate) OR

- C. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:

Patient is 7 years of age or over

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Somatostatin Analogs PA – Octreotide

Drug Name(s)

Octreotide Acetate

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require BOTH of the following:

1. ONE of the following:

A. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND ONE of the following:

- i. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
- ii. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR

B. ONE of the following:

i. Patient has a diagnosis of acromegaly AND ONE of the following:

- a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR
- b. The requested agent is for adjunctive therapy with pituitary radiation therapy OR
- c. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by growth hormone levels or serum IGF-1 levels that are above the reference range OR

ii. Patient has severe diarrhea and/or flushing episodes associated with metastatic carcinoid tumors OR

iii. Patient has profuse watery diarrhea associated with Vasoactive Intestinal Peptide (VIP) secreting tumors OR

iv. Patient has a diagnosis of dumping syndrome AND ONE of the following:

- a. Patient has tried and had an inadequate response to acarbose OR
- b. Patient has an intolerance or hypersensitivity to acarbose OR
- c. Patient has an FDA labeled contraindication to acarbose OR

v. Patient has another indication that is supported in CMS approved compendia for the requested agent AND

2. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be 6 months for initial, 12 months for renewal

Other Criteria:

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. ONE of the following:

A. Patient has a diagnosis of acromegaly OR

B. Patient has severe diarrhea and/or flushing episodes associated with metastatic carcinoid tumors OR

C. Patient has profuse watery diarrhea associated with Vasoactive Intestinal Peptide (VIP) secreting tumors OR

D. Patient has a diagnosis of dumping syndrome OR

E. Patient has another indication that is supported in CMS approved compendia for the requested agent

AND

3. Patient has had clinical benefit with the requested agent AND

4. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Prior Authorization Group Description:

Somatostatin Analogs PA – Somavert

Drug Name(s)

Somavert

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require BOTH of the following:

1. Patient has a diagnosis of acromegaly AND ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent AND provided clinical justification to support that the patient is at risk if therapy is changed OR
 - C. BOTH of the following:
 - i. ONE of the following:
 - a. Patient is not a candidate for surgical resection or pituitary radiation therapy OR
 - b. The requested agent is for adjunctive therapy with pituitary radiation therapy OR
 - c. Patient had an inadequate response to surgery or pituitary radiation therapy as indicated by serum IGF-1 levels that are above the reference range AND
 - ii. ONE of the following:
 - a. Patient has tried and had an inadequate response to octreotide or Somatuline Depot (lanreotide) OR
 - b. Patient has an intolerance or hypersensitivity to octreotide or Somatuline Depot (lanreotide) OR
 - c. Patient has an FDA labeled contraindication to octreotide or Somatuline Depot (lanreotide) AND
2. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of acromegaly AND
3. Patient has had clinical benefit with the requested agent AND
4. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be 6 months for initial, 12 months for renewal

Other Criteria:

Prior Authorization Group Description:

Sovaldi PA

Drug Name(s)

Sovaldi

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication AND
5. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. Patient has an FDA labeled contraindication or hypersensitivity to TWO preferred agents: Epclusa and Harvoni for supported genotypes OR
 - D. Prescriber has provided information based on FDA approved labeling or AASLD/IDSA guidelines supporting the use of the non-preferred agent for the patient's diagnosis and genotype over TWO preferred agents: Epclusa and Harvoni for supported genotypes

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group Description:

Substrate Reduction Therapy PA – Miglustat

Drug Name(s)

Miglustat

Yargesa

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of Gaucher disease type 1 (GD1) confirmed by ONE of the following:
 - A. A baseline glucocerebrosidase enzyme activity of less than or equal to 15% of mean normal in peripheral blood leukocytes, fibroblasts, or other nucleated cells OR
 - B. Confirmation of genetic mutation of GBA gene with two disease-causing alleles AND
2. Prescriber has drawn baseline measurements of hemoglobin level, platelet count, liver volume, and spleen volume AND
3. Prior to any treatment for the intended diagnosis, the patient has had at least ONE of the following clinical presentations:
 - A. Anemia [defined as mean hemoglobin (Hb) level below the testing laboratory's lower limit of the normal range based on age and gender] OR
 - B. Thrombocytopenia (defined as platelet count of less than 100,000 per microliter) OR
 - C. Hepatomegaly OR
 - D. Splenomegaly OR
 - E. Growth failure (i.e., growth velocity is below the standard mean for age) OR
 - F. Evidence of bone disease with other causes ruled out

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of Gaucher disease type 1 (GD1) AND
3. Patient has had improvements or stabilization with the requested agent as indicated by ONE of the following:
 - A. Spleen volume OR
 - B. Hemoglobin level OR
 - C. Liver volume OR
 - D. Platelet count OR
 - E. Growth OR
 - F. Bone pain or crisis

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., endocrinologist, geneticist, hematologist, specialist in metabolic diseases) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Tasimelteon Capsule PA

Drug Name(s)

Tasimelteon

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require the following:

1. ONE of the following:

A. BOTH of the following:

- i. Patient has a diagnosis of Non-24-hour sleep-wake disorder AND
- ii. Patient is totally blind (i.e., no light perception) OR

B. BOTH of the following:

- i. Patient has a diagnosis of Smith-Magenis Syndrome (SMS) confirmed by the presence of ONE of the following genetic mutations:
 - A. A heterozygous deletion of 17p11.2 OR
 - B. A heterozygous pathogenic variant involving RAI1 AND
- ii. The requested agent is being used to treat nighttime sleep disturbances associated with SMS

Age Restriction:

For diagnosis of Non-24-hour sleep-wake disorder, patient is 18 years of age or over. For diagnosis of Smith-Magenis Syndrome (SMS), patient is 16 years of age or over.

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., neurologist, sleep specialist, psychiatrist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Teriparatide PA

Drug Name(s)

Forteo

Teriparatide

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ALL of the following:

1. Patient has ONE of the following diagnoses:
 - A. Postmenopausal with osteoporosis OR
 - B. Patient's sex is male with primary or hypogonadal osteoporosis OR
 - C. Osteoporosis with sustained systemic glucocorticoid therapy AND
2. Patient's diagnosis was confirmed by ONE of the following:
 - A. A fragility fracture in the hip or spine OR
 - B. A T-score of -2.5 or lower OR
 - C. A T-score of -1.0 to -2.5 AND ONE of the following:
 - i. A fragility fracture of the proximal humerus, pelvis, or distal forearm OR
 - ii. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR
 - iii. A FRAX 10-year probability of hip fracture of 3% or greater AND
3. ONE of the following:
 - A. Patient is at a very high fracture risk as defined by ONE of the following:
 - i. Patient had a recent fracture (within the past 12 months) OR
 - ii. Patient had fractures while on FDA approved osteoporosis therapy OR
 - iii. Patient has had multiple fractures OR
 - iv. Patient had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR
 - v. Patient has a very low T-score (less than -3.0) OR
 - vi. Patient is at high risk for falls or has a history of injurious falls OR
 - vii. Patient has a very high fracture probability by FRAX (e.g., major osteoporosis fracture greater than 30%, hip fracture greater than 4.5%) or by other validated fracture risk algorithm OR
 - B. ONE of the following:
 - i. Patient has tried and had an inadequate response to a bisphosphonate OR
 - ii. Patient has an intolerance or hypersensitivity to a bisphosphonate OR
 - iii. Patient has an FDA labeled contraindication to a bisphosphonate AND
4. Patient will NOT be using the requested agent in combination with a bisphosphonate, denosumab (e.g., Prolia, Xgeva), romosozumab-aqqg, or another parathyroid hormone analog (e.g., abaloparatide) for the requested indication AND

Criteria continues: see Other Criteria

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

No prior teriparatide and/or Tymlos use approve 2 years, Prior use - see Other Criteria

Other Criteria:

5. The requested dose is within FDA labeled dosing for the requested indication AND

6. ONE of the following:

A. Patient has never received treatment with teriparatide or Tymlos (abaloparatide) OR

B. Patient has been previously treated with teriparatide or Tymlos (abaloparatide) AND ONE of the following:

i. The total cumulative duration of treatment with teriparatide and Tymlos (abaloparatide) has NOT exceeded 2 years OR

ii. Patient has received 2 years or more of treatment with teriparatide, and remains at or has returned to having a high risk for fracture

Prior teriparatide and/or Tymlos use approve remainder of 2 years of total cumulative therapy. Approve 1 year if patient has received 2 years or more teriparatide

Prior Authorization Group Description:

Tetrabenazine PA

Drug Name(s)

Tetrabenazine

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. ONE of the following:

- A. Patient has a diagnosis of chorea associated with Huntington's disease OR
- B. Patient has an indication that is supported in CMS approved compendia for the requested agent AND

2. ONE of the following:

- A. Patient does NOT have a current diagnosis of depression OR
- B. Patient has a current diagnosis of depression and is being treated for depression AND

3. ONE of the following:

- A. Patient does NOT have a diagnosis of suicidal ideation and/or behavior OR
- B. Patient has a diagnosis of suicidal ideation and/or behavior and must NOT be actively suicidal AND

4. Patient will NOT be using the requested agent in combination with a monoamine oxidase inhibitor (MAOI) AND

5. Patient will NOT be using the requested agent in combination with reserpine

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Tobramycin neb PA

Drug Name(s)

Tobramycin Neb

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of cystic fibrosis AND
2. Documentation has been provided that indicates the patient has a *Pseudomonas aeruginosa* respiratory infection AND
3. ONE of the following:
 - a. Patient is NOT currently (within the past 60 days) being treated with another inhaled antibiotic (e.g., inhaled aztreonam, inhaled tobramycin) OR
 - b. Patient is currently (within the past 60 days) being treated with another inhaled antibiotic (e.g., inhaled aztreonam, inhaled tobramycin) AND ONE of the following:
 - i. Prescriber has confirmed that the other inhaled antibiotic will be discontinued, and that therapy will be continued only with the requested agent OR
 - ii. Prescriber has provided information in support of another inhaled antibiotic therapy used concurrently with or alternating with (i.e., continuous alternating therapy) the requested agent

Drug is also subject to Part B versus Part D review.

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Topical Retinoids PA – Tazarotene

Drug Name(s)

Tazarotene

Tazorac

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

Requested agent will be used for cosmetic purposes

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

- a. Patient has an FDA labeled indication for the requested agent OR
- b. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Topical Retinoids PA – Tretinoin

Drug Name(s)

Avita

Tretinoin Cream, Gel

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

Requested agent will be used for cosmetic purposes

Required Medical Information:

Criteria for approval require the following:

1. ONE of the following:

- a. Patient has an FDA labeled indication for the requested agent OR
- b. Patient has an indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Trelstar PA

Drug Name(s)

Trelstar Mixject

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ALL of the following:

1. Patient has an FDA labeled indication or an indication that is supported in CMS approved compendia for the requested agent AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 180 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent OR
 - C. BOTH of the following:
 - i. Patient is NOT currently being treated with the requested agent AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent AND
3. The requested dose is within FDA labeled dosing or supported in CMS approved compendia dosing for the requested indication

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Trientine PA

Drug Name(s)

Trientine Hydrochloride

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require BOTH of the following:

1. Patient has a diagnosis of Wilson's disease confirmed by ONE of the following:
 - A. Confirmation of genetic mutation of the ATP7B gene OR
 - B. Patient has TWO or more of the following:
 - i. Presence of hepatic abnormality (e.g., acute liver failure, cirrhosis, fatty liver)
 - ii. Presence of Kayser-Fleischer rings
 - iii. Serum ceruloplasmin level less than 20 mg/dL
 - iv. Basal urinary copper excretion greater than 40 mcg/24 hours or the testing laboratory's upper limit of normal
 - v. Hepatic parenchymal copper content greater than 40 mcg/g dry weight
 - vi. Presence of neurological symptoms (e.g., dystonia, hypertonia, rigidity with tremors, dysarthria, muscle spasms, dysphasia, polyneuropathy, dysautonomia) AND
2. ONE of the following:
 - A. Patient has tried and had an inadequate response to penicillamine OR
 - B. Patient has an intolerance or hypersensitivity to penicillamine OR
 - C. Patient has an FDA labeled contraindication to penicillamine

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of Wilson's disease AND
3. Patient has had clinical benefit with the requested agent as evidenced by ONE of the following:
 - A. Improvement and/or stabilization in hepatic abnormality OR
 - B. Reduction in Kayser-Fleischer rings OR
 - C. Improvement and/or stabilization in neurological symptoms (e.g., dystonia, hypertonia, rigidity with tremors, dysarthria, muscle spasms, dysphasia, polyneuropathy, dysautonomia) OR
 - D. Basal urinary copper excretion greater than 200 mcg/24 hours

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist, neurologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Trikafta PA

Drug Name(s)

Trikafta

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cystic fibrosis AND
2. ONE of the following:
 - A. Patient has the presence of the F508del mutation in at least ONE allele (heterozygous OR homozygous) of the CFTR gene confirmed by genetic testing OR
 - B. Patient has ONE of the CFTR gene mutations or a mutation in the CFTR gene that is responsive based on in vitro data, as indicated in the FDA label, confirmed by genetic testing OR
 - C. Patient has another CFTR gene mutation(s) that is responsive to the requested agent, as indicated in the FDA label, confirmed by genetic testing AND
3. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of cystic fibrosis AND
3. Patient has had improvement or stabilization with the requested agent [e.g., improvement in FEV1 from baseline, increase in weight/BMI, improvement from baseline Cystic Fibrosis Questionnaire-Revised (CFQ-R) Respiratory Domain score, improvements in respiratory symptoms (cough, sputum production, and difficulty breathing), and/or reduced number of pulmonary exacerbations] AND
4. Patient will NOT be using the requested agent in combination with another CFTR modulator agent for the requested indication

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cystic fibrosis, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Trulicity PA

Drug Name(s)

Trulicity

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 5. BOTH of the following:
 - a. Patient has a diagnosis of established cardiovascular disease [e.g., myocardial infarction, stroke, any revascularization procedure, transient ischemic attack, unstable angina, amputation, symptomatic or asymptomatic coronary artery disease] AND
 - b. The requested agent will be used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Tymlos PA

Drug Name(s)

Tymlos

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for approval require ALL of the following:

1. Patient (pt) has ONE of the following:
 - A. Postmenopausal with osteoporosis OR
 - B. Pt's sex is male with osteoporosis AND
2. BOTH of the following:
 - A. Pt's diagnosis was confirmed by ONE of the following:
 - i. A fragility fracture in the hip or spine OR
 - ii. A T-score of -2.5 or lower OR
 - iii. A T-score of -1.0 to -2.5 AND ONE of the following:
 - a. A fragility fracture of proximal humerus, pelvis, or distal forearm OR
 - b. A FRAX 10-year probability for major osteoporotic fracture of 20% or greater OR
 - c. A FRAX 10-year probability of hip fracture of 3% or greater AND
 - B. ONE of the following:
 - i. Pt is at a very high fracture risk as defined by ONE of the following:
 - a. Pt had a recent fracture (within the past 12 months) OR
 - b. Pt had fractures while on FDA approved osteoporosis therapy OR
 - c. Pt has had multiple fractures OR
 - d. Pt had fractures while on drugs causing skeletal harm (e.g., long-term glucocorticoids) OR
 - e. Pt has a very low T-score (less than -3.0) OR
 - f. Pt is at high risk for falls or has a history of injurious falls OR
 - g. Pt has a very high fracture probability by FRAX (e.g., major osteoporosis fracture greater than 30%, hip fracture greater than 4.5%) or by other validated fracture risk algorithm OR
 - ii. ONE of the following:
 - a. Pt has tried and had an inadequate response to a bisphosphonate OR
 - b. Pt has an intolerance or hypersensitivity to a bisphosphonate OR
 - c. Pt has an FDA labeled contraindication to a bisphosphonate AND
3. Pt will NOT be using the requested agent in combination with a bisphosphonate, denosumab (e.g., Prolia, Xgeva), romosozumab-aqqg, or another parathyroid hormone analog (e.g., teriparatide) for the requested indication AND
4. The requested dose is within FDA labeled dosing for the requested indication AND
5. The total cumulative duration of treatment with teriparatide and Tymlos (abaloparatide) has not exceeded 2 years

Age Restriction:

Prescriber Restrictions:

Coverage Duration:

No prior Tymlos and/or teriparatide use approve 2 years, Prior use - see Other Criteria

Other Criteria:

Prior Tymlos and/or teriparatide use approve remainder of 2 years of total cumulative therapy

Prior Authorization Group Description:

Urea Cycle Disorders PA - Sodium Phenylbutyrate

Drug Name(s)

Sodium Phenylbutyrate

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of ONE of the following:
 - a. Urea cycle disorder with neonatal-onset involving deficiencies of carbamylphosphate synthetase, ornithine transcarbamylase, or argininosuccinic acid synthetase OR
 - b. Urea cycle disorder with late-onset and history of hyperammonemic encephalopathy involving deficiencies of carbamylphosphate synthetase, ornithine transcarbamylase, or argininosuccinic acid synthetase AND
2. The requested dose is within FDA labeled dosing for the requested indication

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., geneticist, metabolic disorders) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Victoza PA

Drug Name(s)

Victoza

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

Requested agent will be used for weight loss alone

Required Medical Information:

Criteria for approval require BOTH of the following:

1. Patient has a diagnosis of type 2 diabetes mellitus AND
2. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. BOTH of the following:
 - i. ONE of the following:
 1. Patient's medication history includes use of an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) within the past 90 days OR
 2. Patient had an ineffective treatment response to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 3. Patient has an intolerance or hypersensitivity to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 4. Patient has an FDA labeled contraindication to an oral diabetes medication (e.g., metformin, an agent containing metformin, glipizide) OR
 5. BOTH of the following:
 - a. Patient has a diagnosis of established cardiovascular disease [e.g., myocardial infarction, stroke, any revascularization procedure, transient ischemic attack, unstable angina, amputation, symptomatic or asymptomatic coronary artery disease] AND
 - b. The requested agent will be used to reduce the risk of major adverse cardiovascular events (cardiovascular death, non-fatal myocardial infarction, or non-fatal stroke) AND
 - ii. Patient does NOT have any FDA labeled contraindications to the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Voriconazole PA

Drug Name(s)

Voriconazole

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

- A. Patient has a diagnosis of invasive *Aspergillus* OR
- B. Patient has a serious infection caused by *Scedosporium apiospermum* or *Fusarium* species OR
- C. Patient has a diagnosis of esophageal candidiasis or candidemia in nonneutropenic patient

AND ONE of the following:

- i. Patient has tried and had an inadequate response to fluconazole or an alternative antifungal agent OR
 - ii. Patient has an intolerance or hypersensitivity to fluconazole or an alternative antifungal agent OR
 - iii. Patient has an FDA labeled contraindication to fluconazole or an alternative antifungal agent OR
- D. Patient has a diagnosis of blastomycosis AND ONE of the following:
- i. Patient has tried and had an inadequate response to itraconazole OR
 - ii. Patient has an intolerance or hypersensitivity to itraconazole OR
 - iii. Patient has an FDA labeled contraindication to itraconazole OR
- E. The requested agent is being prescribed for prophylaxis of invasive *Aspergillus* or *Candida* AND patient is severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR
- F. Patient has another indication that is supported in CMS approved compendia for the requested agent

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

One month for esophageal candidiasis, 6 months for all other indications

Other Criteria:

Criteria for renewal approval require BOTH of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. ONE of the following:

- A. Patient has a diagnosis of invasive Aspergillus, a serious infection caused by *Scedosporium apiospermum* or *Fusarium* species, esophageal candidiasis, candidemia in nonneutropenic patient, or blastomycosis and patient has continued indicators of active disease (e.g., continued radiologic findings, positive cultures, positive serum galactomannan assay for *Aspergillus*) OR
- B. The requested agent is being prescribed for prophylaxis of invasive Aspergillus or *Candida* and patient continues to be severely immunocompromised, such as a hematopoietic stem cell transplant [HSCT] recipient, or hematologic malignancy with prolonged neutropenia from chemotherapy, or is a high-risk solid organ (lung, heart-lung, liver, pancreas, small bowel) transplant patient, or long term use of high dose corticosteroids (greater than 1 mg/kg/day of prednisone or equivalent) OR
- C. BOTH of the following:
- i. Patient has another indication that is supported in CMS approved compendia for the requested agent AND
 - ii. Patient has had clinical benefit with the requested agent

Prior Authorization Group Description:

Vosevi PA

Drug Name(s)

Vosevi

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication AND
5. If genotype 1, the patient's subtype has been identified and provided

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group Description:

Vyndaqel PA

Drug Name(s)

Vyndaqel

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND
2. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND
3. Patient will NOT be using the requested agent in combination with another tafamidis agent for the requested indication

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of cardiomyopathy of wild type or hereditary transthyretin-mediated amyloidosis (ATTR-CM) AND
3. The requested agent will be used to reduce cardiovascular mortality and cardiovascular-related hospitalization AND
4. Patient has had clinical benefit with the requested agent AND
5. Patient will NOT be using the requested agent in combination with another tafamidis agent for the requested indication

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cardiologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Xermelo PA

Drug Name(s)

Xermelo

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. Patient has a diagnosis of carcinoid syndrome diarrhea AND
2. Patient has tried and had an inadequate response to treatment with a somatostatin analog (e.g., Sandostatin [octreotide], Sandostatin LAR [octreotide], Somatuline Depot [lanreotide]) AND
3. The requested agent will be used in combination with a somatostatin analog (e.g., Sandostatin [octreotide], Sandostatin LAR [octreotide], Somatuline Depot [lanreotide])

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
2. Patient has a diagnosis of carcinoid syndrome diarrhea AND
3. Patient has had clinical benefit with the requested agent (e.g., reduction in the average number of daily bowel movements) AND
4. The requested agent will be used in combination with a somatostatin analog (e.g., Sandostatin [octreotide], Sandostatin LAR [octreotide], Somatuline Depot [lanreotide])

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Xgeva PA

Drug Name(s)

Xgeva

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. ONE of the following:

A. Patient has a diagnosis of multiple myeloma AND BOTH of the following:

i. The requested agent will be used for the prevention of skeletal-related events AND

ii. ONE of the following:

1. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR

2. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR

3. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR

B. Patient has a diagnosis of prostate cancer AND ALL of the following:

i. The requested agent will be used for the prevention of skeletal-related events AND

ii. Patient has bone metastases AND

iii. ONE of the following:

1. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR

2. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR

3. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR

Criteria continues: see Other Criteria

Age Restriction:**Prescriber Restrictions:****Coverage Duration:**

Approval will be for 12 months

Other Criteria:

C. Patient has a solid tumor cancer diagnosis (e.g., thyroid, non-small cell lung, kidney cancer, or breast cancer) AND ALL of the following:

i. The requested agent will be used for the prevention of skeletal-related events AND

ii. Patient has bone metastases AND

- iii. ONE of the following:
 - 1. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR
 - 2. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR
 - 3. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR
- D. Patient has a diagnosis of giant cell tumor of bone AND ONE of the following:
 - i. Patient has a pretreatment or current calcium level that is NOT below the limits of the testing laboratory's normal range OR
 - ii. Patient has a pretreatment or current calcium level that is below the limits of the testing laboratory's normal range AND it will be corrected prior to use of the requested agent OR
 - iii. Prescriber has indicated that the patient is not at risk for hypocalcemia (not including risk associated with the requested agent) OR
- E. Patient has a diagnosis of hypercalcemia of malignancy AND
- 2. Patient will NOT be using the requested agent in combination with Prolia (denosumab) AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Prior Authorization Group Description:

Xifaxan PA

Drug Name(s)

Xifaxan

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require the following:

1. Patient has ONE of the following:
 - a. A diagnosis of irritable bowel syndrome with diarrhea (IBS-D) OR
 - b. A diagnosis of hepatic encephalopathy [reduction in risk of overt hepatic encephalopathy (HE) recurrence] OR
 - c. BOTH of the following:
 - i. A diagnosis of traveler's diarrhea (TD) AND
 - ii. The traveler's diarrhea is caused by noninvasive strains of Escherichia coli

Age Restriction:

For diagnosis of traveler's diarrhea (TD), patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:**Coverage Duration:**

Approval will be for 12 months

Other Criteria:

Prior Authorization Group Description:

Xolair PA

Drug Name(s)

Xolair

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:****Required Medical Information:**

Criteria for initial approval require ALL of the following:

1. ONE of the following:

A. Patient has a diagnosis of moderate to severe persistent asthma AND ALL of the following:

i. ONE of the following:

a. Patient is 6 to less than 12 years of age AND BOTH of the following:

I. Patient's pretreatment IgE level is 30 IU/mL to 1300 IU/mL AND

II. Patient's weight is 20 kg to 150 kg OR

b. Patient is 12 years of age or over AND BOTH of the following:

I. Patient's pretreatment IgE level is 30 IU/mL to 700 IU/mL AND

II. Patient's weight is 30 kg to 150 kg AND

iii. Allergic asthma has been confirmed by a positive skin test or in vitro reactivity test (RAST) to a perennial aeroallergen AND

iv. ONE of the following:

a. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LRTA, LAMA, theophylline) in combination with the requested agent OR

b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an asthma control therapy OR

B. Patient has a diagnosis of chronic idiopathic urticaria AND BOTH of the following:

i. Patient has had over 6 weeks of hives and itching AND

ii. ONE of the following:

a. Patient has tried and had an inadequate response to maximum tolerable H1 antihistamine therapy OR

b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to H1 antihistamine therapy OR

C. Patient has a diagnosis of nasal polyps AND BOTH of the following:

i. ONE of the following:

a. Patient has tried and had an inadequate response to an intranasal corticosteroid OR

b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an intranasal corticosteroid AND

ii. ONE of the following:

a. The requested agent will be used in combination with an intranasal corticosteroid OR

- b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an intranasal corticosteroid OR

Initial criteria continues: see Other Criteria

Age Restriction:

For diagnosis of moderate to severe persistent asthma, patient is 6 years of age or over. For diagnosis of chronic idiopathic urticaria, patient is 12 years of age or over. For diagnosis of nasal polyps, patient is 18 years of age or over.

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., allergist, immunologist, otolaryngologist, pulmonologist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be 6 months for initial, 12 months for renewal

Other Criteria:

- D. Patient has a diagnosis of IgE-mediated food allergy AND ALL of the following:
 - i. Patient is using the requested agent for the reduction of allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one or more foods AND
 - ii. IgE-mediated food allergy has been confirmed by an allergy diagnostic test (e.g., skin prick test, serum specific IgE test, oral food challenge) AND
 - iii. Patient will avoid known food allergens while treated with the requested agent AND
 - iv. The requested agent will NOT be used for the emergency treatment of allergic reactions, including anaphylaxis AND
- 2. Patient will NOT be using the requested agent in combination with Dupixent or an injectable Interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication AND
- 3. The requested dose is within FDA labeled dosing for the requested indication

Criteria for renewal approval require ALL of the following:

- 1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND
- 2. ONE of the following:
 - A. Patient has a diagnosis of moderate to severe persistent asthma AND ALL of the following:
 - i. Patient's weight is within the FDA indicated range for their age (i.e., 20 kg to 150 kg for patients age 6 to less than 12 years and 30 kg to 150 kg for patients 12 years of age or over) AND
 - ii. Patient has had clinical benefit with the requested agent AND
 - iii. ONE of the following:
 - a. Patient is currently being treated with AND will continue asthma control therapy (e.g., ICS, ICS/LABA, LRTA, LAMA, theophylline) in combination with the requested agent OR
 - b. Patient has an intolerance, FDA labeled contraindication, or hypersensitivity to an asthma control therapy OR
 - B. Patient has a diagnosis of chronic idiopathic urticaria AND the following:

- a. Patient has had clinical benefit with the requested agent OR
- C. Patient has a diagnosis of nasal polyps AND the following:
 - a. Patient has had clinical benefit with the requested agent OR
- D. Patient has a diagnosis of IgE-mediated food allergy AND ALL of the following:
 - a. Patient is using the requested agent for the reduction of allergic reactions (Type I), including anaphylaxis, that may occur with accidental exposure to one or more foods AND
 - b. Patient has had clinical benefit with the requested agent AND
 - c. Patient will avoid known food allergens while treated with the requested agent AND
 - d. The requested agent will NOT be used for the emergency treatment of allergic reactions, including anaphylaxis AND
- 3. Patient will NOT be using the requested agent in combination with Dupixent or an injectable interleukin 5 (IL-5) inhibitor (e.g., Cinqair, Fasenra, Nucala) for the requested indication AND
- 4. The requested dose is within FDA labeled dosing for the requested indication

Prior Authorization Group Description:

Zepatier PA

Drug Name(s)

Zepatier

Indications:

All Medically-Accepted Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for approval require ALL of the following:

1. Patient has a diagnosis of hepatitis C confirmed by serological markers AND
2. Prescriber has screened the patient for current or prior hepatitis B viral (HBV) infection and if positive, will monitor the patient for HBV flare-up or reactivation during and after treatment with the requested agent AND
3. The requested agent will be used in a treatment regimen and length of therapy that is supported in FDA approved labeling or AASLD/IDSA guidelines for the patient's diagnosis and genotype AND
4. The requested dose is within FDA labeled dosing or supported in AASLD/IDSA guideline dosing for the requested indication AND
5. If genotype 1, the patient's subtype has been identified and provided AND
6. If genotype 1a, the prescriber has tested the patient for the presence of virus with NS5A resistance-associated polymorphisms AND
7. ONE of the following:
 - A. There is evidence of a claim that the patient is currently being treated with the requested agent within the past 90 days OR
 - B. Prescriber states the patient is currently being treated with the requested agent within the past 90 days OR
 - C. Patient has an FDA labeled contraindication or hypersensitivity to TWO preferred agents: Epclusa and Harvoni for supported genotypes OR
 - D. Prescriber has provided information based on FDA approved labeling or AASLD/IDSA guidelines supporting the use of the non-preferred agent for the patient's diagnosis and genotype over TWO preferred agents: Epclusa and Harvoni for supported genotypes

Age Restriction:**Prescriber Restrictions:**

Prescriber is a specialist in the area of the patient's diagnosis (e.g., gastroenterologist, hepatologist or infectious disease) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Duration of therapy: Based on FDA approved labeling or AASLD/IDSA guideline supported

Other Criteria:

Prior Authorization Group Description:

Zokinvy PA

Drug Name(s)

Zokinvy

Indications:

All FDA-Approved Indications.

Off-Label Uses:**Exclusion Criteria:**

FDA labeled contraindications to the requested agent

Required Medical Information:

Criteria for initial approval require the following:

1. ONE of the following:

A. BOTH of the following:

- i. Patient has a diagnosis of Hutchinson-Gilford progeria syndrome (HGPS) AND
- ii. Genetic testing has confirmed a pathogenic variant in the LMNA gene that results in production of progerin OR

B. Patient has a diagnosis of processing-deficient progeroid laminopathy AND ONE of the following:

- i. Genetic testing has confirmed heterozygous LMNA mutation with progerin-like protein accumulation OR
- ii. Genetic testing has confirmed homozygous or compound heterozygous ZMPSTE24 mutations

Criteria for renewal approval require ALL of the following:

1. Patient has been previously approved for the requested agent through the plan's Prior Authorization criteria AND

2. Patient has a diagnosis of ONE of the following:

- A. Hutchinson-Gilford progeria syndrome (HGPS) OR
- B. Processing-deficient progeroid laminopathies with either: heterozygous LMNA mutation with progerin-like protein accumulation OR homozygous or compound heterozygous ZMPSTE24 mutations AND

3. Patient has had clinical benefit with the requested agent

Age Restriction:

Patient is within the FDA labeled age for the requested agent

Prescriber Restrictions:

Prescriber is a specialist in the area of the patient's diagnosis (e.g., cardiologist, geneticist) or the prescriber has consulted with a specialist in the area of the patient's diagnosis

Coverage Duration:

Approval will be for 12 months

Other Criteria: