

Print

Program Name: BadgerCare Plus and Medicaid

Handbook Area: Pharmacy

Prior Authorization : Preferred Drug List

Topic #16217

Cytokine and Cell Adhesion Molecule Antagonist Drugs

Note: The Preferred Drug List Quick Reference provides the most current list of preferred and non-preferred drugs in this drug class.

Clinical PA is required for non-preferred cytokine and CAM antagonist drugs.

PA requests for non-preferred cytokine and CAM antagonist drugs will only be approved for use to treat these identified clinical conditions:

- Alopecia areata
- Ankylosing spondylitis
- Crohn's disease
- DJRA
- ERA
- Giant cell arteritis
- GPP
- Hidradenitis suppurativa
- JIA and systemic JIA
- NMOSD
- NOMID
- nt-axSpA
- PMR
- Psoriasis
- Psoriatic arthritis
- RA
- SSc-ILD
- Ulcerative colitis
- Uveitis

Otezla, a preferred drug in the cytokine and CAM antagonist drug class, is the only cytokine and CAM antagonist drug indicated for the clinical condition of oral ulcers associated with Behcet's disease. Preferred drugs do not require PA.

PA requests for cytokine and CAM antagonist drugs will only be approved for **one cytokine and CAM antagonist drug per member**. ForwardHealth does not cover treatment with more than one cytokine and CAM antagonist drug.

PA requests will not be considered for subcutaneous dosage forms of cytokine and CAM antagonist drugs that will be administered in a medical office or medical facility.

Supporting clinical information and a copy of the member's current medical records must be submitted with all PA requests for non-preferred cytokine and CAM antagonist drugs. The supporting clinical information and medical records must document the following:

- The member's medical condition being treated
- Details regarding previous medication use
- The member's current treatment plan

Initial PA requests for non-preferred cytokine and CAM antagonist drugs may be approved for up to 183 days.

Renewal PA requests for non-preferred cytokine and CAM antagonist drugs may be approved for up to 365 days. Renewal PA requests for non-preferred cytokine and CAM antagonist drugs must include supporting clinical information and copies of the member's current medical records demonstrating that the member had a significant reduction in symptoms compared to their baseline prior to the initiation of the non-preferred cytokine and CAM antagonist drug.

All renewal PA requests require the member to be adherent with the prescribed treatment regimen.

Information is available about general ForwardHealth policy for drugs that require PA approval. This includes what may **not** be considered criteria to support the need for a drug.

Cytokine and Cell Adhesion Molecule Antagonist Biosimilar Drugs

All cytokine and CAM antagonist biosimilar drugs will be added to the cytokine and CAM antagonist drug class as non-preferred drugs (unless specifically identified or noted) until the next scheduled drug class review by the Wisconsin Medicaid Pharmacy PA Committee. The cytokine and CAM antagonists drug class is typically reviewed at the November PDL review each year.

ForwardHealth will use a blanket "xxxx" placeholder after the biosimilar generic name when referring to non-preferred biosimilar drug products in the clinical PA criteria for the appropriate clinical condition. The Preferred Drug List Quick Reference data table will list the brand names with the complete generic names of the biosimilar drug products in the cytokine and CAM antagonists drug class.

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Alopecia Areata

Leqselvi, Litfulo, and Olumiant are non-preferred drugs used to treat alopecia areata.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat alopecia areata are **both** of the following:

- The member has severe alopecia areata with at least 50% scalp hair loss (SALT score of greater than or equal to 50). The member's SALT score must be documented.
- The prescription is written by a dermatologist or through a dermatology consultation.

Clinical documentation and medical records must be submitted with the PA request to support the member's condition of alopecia areata and outline the member's current treatment plan for alopecia areata.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Alopecia Areata

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat alopecia areata must be submitted using Section VI (Clinical Information for Drugs With Specific Criteria Addressed in the ForwardHealth Online Handbook) of the PA/DGA form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed PA/DGA form and a completed PA/RE to ForwardHealth.

PA requests used to treat alopecia areata may be submitted on the Portal, by fax, or by mail (but **not** using the STAT:PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Ankylosing Spondylitis

Cimzia, Cyltezo, Enbrel, Humira, Simponi subQ, and Xeljanz are preferred drugs used to treat ankylosing spondylitis. Preferred drugs do not require PA.

Adalimumab-xxxx, Bimzelx, Cosentyx subQ, Rinvoq, Taltz, and Xeljanz XR are non-preferred drugs used to treat ankylosing spondylitis.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat ankylosing spondylitis are **all** of the following:

- The member has ankylosing spondylitis.
- The prescription is written by a rheumatologist or through a rheumatology consultation.
- The member has taken **two** preferred cytokine and CAM antagonist drugs for **at least three** consecutive months each and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction.

(Note: ForwardHealth will only consider use of Cyltezo or Humira as one of the preferred cytokine and CAM antagonist drugs.)

- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Non-Preferred Adalimumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira. The clinical information must document why the member cannot use Cyltezo and Humira, including why it is medically necessary that the member receive a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira.

Xeljanz XR

The prescriber must submit detailed clinical justification for prescribing Xeljanz XR instead of Xeljanz. The clinical information must document why the member cannot use Xeljanz, including why it is medically necessary that the member receive Xeljanz XR instead of Xeljanz.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Ankylosing Spondylitis

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat ankylosing spondylitis must be submitted using the Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Ankylosing Spondylitis form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Ankylosing Spondylitis form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat ankylosing spondylitis may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Crohn's Disease

Cimzia, Cyltezo, and Humira are preferred drugs used to treat Crohn's disease. Preferred drugs do not require PA.

Adalimumab-xxxx, Entyvio subQ, Omvoh subQ, Rinvoq, Skyrizi subQ, Stelara subQ, Tremfya subQ, ustekinumab-xxxx subQ, and Zymfentra are non-preferred drugs used to treat Crohn's disease.

Note: Omvoh, Skyrizi, Stelara, and ustekinumab-xxxx will require an IV induction prior to initiating treatment with the subQ. A PA request for the IV induction must be approved before ForwardHealth will consider PA for the subQ. PA for the IV induction may be obtained through the physician-administered drug PA process. An IV induction for Tremfya is optional prior to initiating treatment with the subQ.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat Crohn's disease are **all** of the following:

- The member has Crohn's disease.
- The prescription is written by a gastroenterologist or through a gastroenterology consultation.
- The member has taken **two** preferred cytokine and CAM antagonist drug for **at least three** consecutive months and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction. (Note: ForwardHealth will only consider use of Cyltezo or Humira as one of the preferred cytokine and CAM antagonist drugs.)
- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Non-Preferred Adalimumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira. The clinical information must document why the member cannot use Cyltezo and Humira, including why it is medically necessary that the member receive a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira.

Non-Preferred Ustekinumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred ustekinumab-xxxx drug instead of Stelara. The clinical information must document why the member cannot use Stelara, including why it is medically necessary that the member receive a non-preferred ustekinumab-xxxx drug instead of Stelara.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Crohn's Disease

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat Crohn's disease must be submitted using the Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Crohn's Disease and Ulcerative Colitis form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Crohn's Disease and Ulcerative Colitis form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat Crohn's disease may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Deficiency of Interleukin-1 Receptor Antagonist

Kineret is a non-preferred drug used to treat DIRA.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat DIRA are **both** of the following:

- The member has DIRA.
- The prescription is written by or through consultation with a DIRA specialist (for example, an immunologist or a rheumatologist).

Clinical documentation and medical records must be submitted with the PA request to support the member's condition of DIRA and outline the member's current treatment plan for DIRA.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Deficiency of Interleukin-1 Receptor Antagonist

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat DIRA must be submitted using PA/DGA form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed PA/DGA form and a completed PA/RF to ForwardHealth.

PA requests used to treat DIRA may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Enthesitis-Related Arthritis

Cosentyx subQ is a non-preferred drug used to treat ERA.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat ERA are **both** of the following:

- The member has ERA.
- The prescription is written by a rheumatologist or through a rheumatology consultation.

Clinical documentation and medical records must be submitted with the PA request to support the member's condition of ERA and outline the member's current treatment plan for ERA.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Enthesitis-Related Arthritis

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat ERA must be submitted using Section VI (Clinical Information for Drugs With Specific Criteria Addressed in the ForwardHealth Online Handbook) of the PA/DGA form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed PA/DGA form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat ERA may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system.)

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Giant Cell Arteritis

Tyenne subQ is a preferred drug used to treat giant cell arteritis. Preferred drugs do not require PA.

Actemra subQ and Rinvoq are non-preferred drugs used to treat giant cell arteritis.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat giant cell arteritis are **all** of the following:

- The member has giant cell arteritis.
- The prescription is written by a rheumatologist or through a rheumatology consultation.
- The member has taken Tyenne subQ for **at least three** consecutive months and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction.
- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Giant Cell Arteritis

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat giant cell arteritis must be submitted using Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Giant Cell Arteritis and nr-axSpA form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Giant Cell Arteritis and nr-axSpA form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat giant cell arteritis may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Non-Preferred Cytokine and Cell Adhesion Molecule Antagonist Drugs for Generalized Pustular Psoriasis

Spevigo subQ is a non-preferred drug used to treat GPP.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat GPP are **all** of the following:

- The member has GPP.
- The prescription is written by a dermatologist or through a dermatology consultation.

Clinical documentation and medical records must be submitted with the PA request to support the member's clinical condition of GPP and outline the member's current treatment plan for GPP.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Generalized Pustular Psoriasis

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat GPP must be submitted using Section VI (Clinical Information for Drugs With Specific Criteria Addressed in the ForwardHealth Online Handbook) of the PA/DGA form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed PA/DGA form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat GPP may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Additional Dosage Form for Spevigo

The intravenous dosage form of Spevigo is a physician-administered drug, and it uses HCPCS procedure code J1747 (Injection, spesolimab-sbzo, 1 mg). HCPCS code J1747 will also have a diagnosis restriction of L40.1 for GPP.

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Hidradenitis Suppurativa

Cyltezo and Humira are preferred drugs used to treat hidradenitis suppurativa. Preferred drugs do not require PA.

Adalimumab-xxxx, Bimzelx, and Cosentyx subQ are non-preferred drugs used to treat hidradenitis suppurativa.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat hidradenitis suppurativa are **all** of the following:

- The member has hidradenitis suppurativa.
- The prescription is written by a dermatologist or through a dermatology consultation.
- The member has taken **one** preferred cytokine and CAM antagonist drug for **at least three** consecutive months and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction.
- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Non-Preferred Adalimumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira. The clinical information must document why the member cannot use Cyltezo and Humira, including why it is medically necessary that the member receive a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Hidradenitis Suppurativa

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat hidradenitis suppurativa must be submitted using the Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Hidradenitis Suppurativa form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Hidradenitis Suppurativa form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat hidradenitis suppurativa may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Neuromyelitis Optica Spectrum Disorder

Enspryng is a non-preferred drug used to treat NMOSD.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat NMOSD are **all** of the following:

- The member has NMOSD.
- The prescription is written by a neurologist or through a neurology consultation.
- The member is anti-aquaporin-4 antibody positive.

Clinical documentation and medical records must be submitted with the PA request to support the member's clinical condition of NMOSD and outline the member's current treatment plan for NMOSD.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Neuromyelitis Optica Spectrum Disorder

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat NMOSD must be submitted using Section VI (Clinical Information for Drugs With Specific Criteria Addressed in the ForwardHealth Online Handbook) of the PA/DGA form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed PA/DGA form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat NMOSD may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system.)

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Neonatal Onset Multisystem Inflammatory Disease

Kineret is a non-preferred drug used to treat NOMID.

Clinical criteria for approval of a PA request for Kineret used to treat NOMID are **both** of the following:

- The member has NOMID.
- The prescription is written by a rheumatologist or through a rheumatology consultation.

Clinical documentation and medical records must be submitted with the PA request to support the member's condition of NOMID and outline the member's current treatment plan for NOMID.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Neonatal Onset Multisystem Inflammatory Disease

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat NOMID must be submitted using Section VI (Clinical Information for Drugs With Specific Criteria Addressed in the ForwardHealth Online Handbook) of the PA/DGA form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed PA/DGA form and a completed PA/RF to ForwardHealth.

PA requests for Kineret used to treat NOMID may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Non-Radiographic Axial Spondyloarthritis

Cimzia is a preferred drug used to treat nr-axSpA. Preferred drugs do not require PA.

Bimzelx, Cosentyx subQ, Rinvoq, and Taltz are non-preferred drugs used to treat nr-axSpA.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat nr-axSpA are **all** of the following:

- The member has nr-axSpA.
- The prescription is written by a rheumatologist or through a rheumatology consultation.
- The member has taken Cimzia **at least three** consecutive months and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction.
- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Non-Radiographic Axial Spondyloarthritis

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat nr-axSpA must be submitted using the Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Giant Cell Arteritis and nr-axSpA form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Giant Cell Arteritis and nr-axSpA form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat nr-axSpA may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Polymyalgia Rheumatica

Kevzara is a non-preferred drug used to treat PMR.

Clinical criteria for approval of a PA request for non-preferred drugs used to treat PMR are **all** of the following:

- The member has PMR.
- The prescription is written by or through consultation with a PMR specialist.
- The member has taken corticosteroids and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction.

Clinical documentation and medical records must be submitted with the PA request to support the member's condition of PMR and outline the member's current treatment plan for PMR.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Polymyalgia Rheumatica

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat PMR must be submitted using Section VI (Clinical Information for Drugs With Specific Criteria Addressed in the ForwardHealth Online Handbook) of the PA/DGA form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed PA/DGA form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat PMR may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Psoriasis

Cimzia, Cyltezo, Enbrel, Humira, and Otezla are preferred drugs used to treat psoriasis. Preferred drugs do not require PA.

Adalimumab-xxxx, Bimzelx, Cosentyx subQ, Siliq, Skyrizi subQ, Sotyktu, Stelara subQ, Taltz, Tremfya, and ustekinumab-xxxx subQ are non-preferred drugs used to treat psoriasis.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat psoriasis are **all** of the following:

- The member has psoriasis.
- The prescription is written by a dermatologist or through a dermatology consultation.
- The member has taken **two** preferred cytokine and CAM antagonist drugs for **at least three** consecutive months each and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction. (Note: ForwardHealth will only consider use of Cyltezo or Humira as one of the preferred cytokine and CAM antagonist drugs.)
- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Non-Preferred Adalimumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira. The clinical information must document why the member cannot use Cyltezo and Humira, including why it is medically necessary that the member receive a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira.

Non-Preferred Ustekinumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred ustekinumab-xxxx drug instead of Stelara. The clinical information must document why the member cannot use Stelara, including why it is medically necessary that the member receive a non-preferred ustekinumab-xxxx drug instead of Stelara.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Psoriasis

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat psoriasis must be submitted using the Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Psoriasis form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Psoriasis form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat psoriasis may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Psoriatic Arthritis

Cimzia, Cyltezo, Enbrel, Humira, Orencia subQ, Otezla, Simponi subQ, and Xeljanz are preferred drugs used to treat psoriatic arthritis. Preferred drugs do not require PA.

Adalimumab-xxxx, Bimzelx, Cosentyx subQ, Rinvoq, Rinvoq LQ, Skyrizi subQ, Stelara subQ, Taltz, Tremfya subQ, ustekinumab-xxxx subQ, and Xeljanz XR are non-preferred drugs used to treat psoriatic arthritis.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat psoriatic arthritis are **all** of the following:

- The member has psoriatic arthritis.
- The prescription is written by a dermatologist or rheumatologist or through a dermatology or rheumatology consultation.
- The member has taken **two** preferred cytokine and CAM antagonist drugs for **at least three** consecutive months each and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction.

(Note: ForwardHealth will only consider use of Cyltezo or Humira as one of the preferred cytokine and CAM antagonist drugs.)

- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Non-Preferred Adalimumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira. The clinical information must document why the member cannot use Cyltezo and Humira, including why it is medically necessary that the member receive a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira.

Non-Preferred Ustekinumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred ustekinumab-xxxx drug instead of Stelara. The clinical information must document why the member cannot use Stelara, including why it is medically necessary that the member receive a non-preferred ustekinumab-xxxx drug instead of Stelara.

Xeljanz XR

The prescriber must submit detailed clinical justification for prescribing Xeljanz XR instead of Xeljanz. The clinical information must document why the member cannot use Xeljanz, including why it is medically necessary that the member receive Xeljanz XR instead of Xeljanz.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Psoriatic Arthritis

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat psoriatic arthritis must be submitted using the Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for RA, JIA, and Psoriatic Arthritis form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for RA, JIA, and Psoriatic Arthritis form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat psoriatic arthritis may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Rheumatoid Arthritis

Cimzia, Cyltezo, Enbrel, Humira, Orencia subQ, Simponi subQ, Tyenne subQ, and Xeljanz are preferred drugs used to treat RA. Preferred drugs do not require PA.

Actemra subQ, adalimumab-xxxx, Kevzara, Kineret, Olumiant, Rinvoq, and Xeljanz XR are non-preferred drugs used to treat RA.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat RA are **all** of the following:

- The member has RA.
- The prescription is written by a rheumatologist or through a rheumatology consultation.
- The member has taken **two** preferred cytokine and CAM antagonist drugs for **at least three** consecutive months each and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction. (Note: ForwardHealth will only consider use of Cyltezo or Humira as one of the preferred cytokine and CAM antagonist drugs.)
- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Non-Preferred Adalimumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira. The clinical information must document why the member cannot use Cyltezo and Humira, including why it

is medically necessary that the member receive a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira.

Xeljanz XR

The prescriber must submit detailed clinical justification for prescribing Xeljanz XR instead of Xeljanz. The clinical information must document why the member cannot use Xeljanz, including why it is medically necessary that the member receive Xeljanz XR instead of Xeljanz.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Rheumatoid Arthritis

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat RA must be submitted using the Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for RA, JIA, and Psoriatic Arthritis form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for RA, JIA, and Psoriatic Arthritis form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat RA may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Juvenile Idiopathic Arthritis

Cimzia, Cyltezo, Enbrel, Humira, Orencia subQ, Tyrenne subQ, and Xeljanz are preferred drugs used to treat JIA. Preferred drugs do not require PA.

Actemra subQ, adalimumab-xxxx, Kevzara, Rinvoq, Rinvoq LQ, and Xeljanz Oral Solution are non-preferred drugs used to treat JIA.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat JIA are **all** of the following:

- The member has JIA.
- The prescription is written by a rheumatologist or through a rheumatology consultation.
- The member has taken **two** preferred cytokine and CAM antagonist drugs for **at least three** consecutive months each and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction. (Note: ForwardHealth will only consider use of Cyltezo or Humira as one of the preferred cytokine and CAM antagonist drugs.)
- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Non-Preferred Adalimumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira. The clinical information must document why the member cannot use Cyltezo and Humira, including why it is medically necessary that the member receive a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira.

Xeljanz Oral Solution

The prescriber must submit detailed clinical justification for prescribing Xeljanz Oral Solution instead of Xeljanz. The clinical information must document why the member cannot use Xeljanz, including why it is medically necessary that the member receive Xeljanz Oral Solution instead of Xeljanz.

Clinical Criteria for Systemic Juvenile Idiopathic Arthritis

Tyenne subQ is a preferred drug used to treat systemic JIA. Preferred drugs do not require PA.

Actemra subQ is a non-preferred drug used to treat systemic JIA.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat systemic JIA are **all** of the following:

- The member has systemic JIA.
- The prescription is written by a rheumatologist or through a rheumatology consultation.
- The member has taken Tyenne subQ for **at least three** consecutive months and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction.
- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Juvenile Idiopathic Arthritis and Systemic Juvenile Idiopathic Arthritis

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat JIA and systemic JIA must be submitted using the Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for RA, JIA, and Psoriatic Arthritis form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for RA, JIA, and Psoriatic Arthritis form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat JIA and systemic JIA may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Systemic Sclerosis-Associated Interstitial Lung Disease

Actemra subQ is a non-preferred drug used to treat SSc-ILD.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat SSc-ILD are **both** of the following:

- The member has SSc-ILD.
- The prescription is written by or through consultation with an SSc-ILD specialist.

Clinical documentation and medical records must be submitted with the PA request to support the member's condition of SSc-ILD and outline the member's current treatment plan for SSc-ILD.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Systemic Sclerosis-Associated Interstitial Lung Disease

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat SSc-ILD must be submitted using Section VI (Clinical Information for Drugs With Specific Criteria Addressed in the ForwardHealth Online Handbook) of the PA/DGA form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed PA/DGA form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat SSc-ILD may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system.)

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Ulcerative Colitis

Cyltezo, Humira, Simponi subQ, and Xeljanz are preferred drugs used to treat ulcerative colitis. Preferred drugs do not require PA.

Adalimumab-xxxx, Entyvio subQ, Omvoh subQ, Rinvoq, Skyrizi subQ, Stelara subQ, Tremfya subQ, ustekinumab-xxxx subQ, Xeljanz XR, and Zymfentra are non-preferred drugs used to treat ulcerative colitis.

Note: Omvoh, Skyrizi, Stelara, Tremfya, and ustekinumab-xxxx will require an IV induction prior to initiating treatment with the subQ. A PA request for the IV induction must be approved before ForwardHealth will consider PA for the subQ. PA for the IV induction may be obtained through the physician-administered drug PA process.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat ulcerative colitis are **all** of the following:

- The member has ulcerative colitis.
- The prescription is written by a gastroenterologist or through a gastroenterology consultation.
- The member has taken **two** preferred cytokine and CAM antagonist drugs for **at least three** consecutive months each and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction. (Note: ForwardHealth will only consider use of Cyltezo or Humira as one of the preferred cytokine and CAM antagonist drugs.)
- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Non-Preferred Adalimumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira. The clinical information must document why the member cannot use Cyltezo and Humira, including why it is medically necessary that the member receive a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira.

Non-Preferred Ustekinumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred ustekinumab-xxxx drug instead of Stelara. The clinical information must document why the member cannot use Stelara, including why it is medically necessary that the member receive a non-preferred ustekinumab-xxxx drug instead of Stelara.

Xeljanz XR

The prescriber must submit detailed clinical justification for prescribing Xeljanz XR instead of Xeljanz. The clinical information must document why the member cannot use Xeljanz, including why it is medically necessary that the member receive Xeljanz XR instead of Xeljanz.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Ulcerative Colitis

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat ulcerative colitis must be submitted using the Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Crohn's Disease and Ulcerative Colitis form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Crohn's Disease and Ulcerative Colitis form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat ulcerative colitis may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).

Clinical Criteria for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Uveitis

Cyltezo and Humira are preferred drugs used to treat uveitis. Preferred drugs do not require PA.

Adalimumab-xxxx is a non-preferred drug used to treat uveitis.

Clinical criteria for approval of a PA request for non-preferred cytokine and CAM antagonist drugs used to treat uveitis are **all** of the following:

- The member has uveitis.

- The prescription is written by an ophthalmologist or through an ophthalmology consultation.
- The member has taken **one** preferred cytokine and CAM antagonist drug for **at least three** consecutive months and experienced an unsatisfactory therapeutic response or experienced a clinically significant adverse drug reaction.
- The prescriber has indicated the clinical reason(s) why a non-preferred cytokine and CAM antagonist drug is being requested.

Non-Preferred Adalimumab-xxxx Drugs

The prescriber must submit detailed clinical justification for prescribing a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira. The clinical information must document why the member cannot use Cyltezo and Humira, including why it is medically necessary that the member receive a non-preferred adalimumab-xxxx drug instead of Cyltezo and Humira.

Submitting PA Requests for Cytokine and Cell Adhesion Molecule Antagonist Drugs for Uveitis

PA requests for non-preferred cytokine and CAM antagonist drugs must be completed, signed, and dated by the prescriber. PA requests for non-preferred cytokine and CAM antagonist drugs used to treat uveitis must be submitted using the Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Uveitis form.

The PA form must be sent to the pharmacy where the prescription will be filled. The prescriber may send the PA form to the pharmacy, or the member may carry the PA form with the prescription to the pharmacy. The pharmacy provider will use the completed PA form to submit a PA request to ForwardHealth. Prescribers should **not** submit the PA form to ForwardHealth.

Pharmacy providers are required to submit the completed Prior Authorization Drug Attachment for Cytokine and CAM Antagonist Drugs for Uveitis form and a completed PA/RF to ForwardHealth.

PA requests for non-preferred cytokine and CAM antagonist drugs used to treat uveitis may be submitted on the Portal, by fax, or by mail (but **not** using the STAT-PA system).