

## Assignment Questions on Advanced Linear Regression

**Q1** What is the optimal value of alpha for ridge and lasso regression? What will be the changes in the model if you choose to double the value of alpha for both ridge and lasso? What will be the most important predictor variables after the change is implemented?

### Answer

Optimal value of alpha for ridge: **10**

The optimal value of alpha for ridge: **100**

After making the double alpha for ridge and lasso i.e. 20 and 200

For Ridge: Coeff values are increasing as alpha will increase.  $r^2$  score of train data is also dropped from **.807 to 0.45**

For Lasso: As the alpha value increased more features were removed from the model. But  $r^2$  score is also

dropped by 1% in both test and train data

Top Features: Neighborhood\_NoRidge, Neighborhood\_Nridge, OverallQual, overallQual  
Neighborhood

**Q2** You have determined the optimal value of lambda for ridge and lasso regression during the assignment. Now, which one will you choose to apply and why?

### Answer

We will choose Lasso as it gives feature selection option. It has removed unwanted features from the model without affecting the model accuracy. Which makes our model generalized and simple and accurate.

**Q3** After building the model, you realised that the five most important predictor variables in the lasso model are not available in the incoming data. You will now have to create another model excluding the five most important predictor variables. Which are the five most important predictor variables now?

### Answer

The top predictor variables are: 'GrLivArea', 'OverallQual', 'TotalBsmtSF', 'OverallCond', 'YearBuilt'

GrLivArea 0.128692

OverallQual 0.065629

TotalBsmtSF 0.048505

OverallCond 0.043846

YearBuilt 0.043804

After removing them, the predictor variables are:

GrLivArea 0.146928

MSZoning\_RL 0.083526

MSZoning\_RM 0.065606

TotalBsmtSF 0.056324

GarageCars 0.043311

The predictor variables remain the same, with a slightly different order for Lasso applied to the RFE created variables.

GrLivArea 0.136432

MSZoning\_RL 0.083509

TotalBsmtSF 0.073250

MSZoning\_RM 0.069320

GarageCars 0.041585

It is noticeable that the top predictor variables are rather in line with the variables that I would intuitively consider important for prediction of Sales Price. Without them, the intuition I had does not carry to the new variables.

**Q4** How can you make sure that a model is robust and generalizable? What are the implications of the same for the accuracy of the model and why?

### **Answer**

Generalisation is important and test accuracy needs to be higher than the training score. However, the difference should not be exceedingly high. The model should generalise during training, but if the score is very high in training, and lower in testing, it means that the model has memorised the data, meaning that it is overfitting. Overall, there should not be large differences between the results. Low test scores could result from splitting the data set too early in the preprocessing step, so that some steps may be missed on the test data. Robustness of a model is generally not solely based on high test scores, but also depends on the assumption that the train scores are higher than the test scores. Both scores have to be high enough to be acceptable for the specific business case and expectations of the model.

It is also important to consider the values obtained for train and test, so that the model will perform well on unseen data. This means that the data should retain some outliers to help with predictions. As demonstrated in the assignment, accuracy of the model will vary, depending on the way data is processed and how features are selected. There may be no perfect model, but different steps are available to ensure that the model developed is fit for purpose for the specific context and the uniqueness of the business case.

This is in line with Occam's razor, that is, the model to be chosen should not be more complex than it needs to be

