

## HW 5

Robin Baldeo

February 8, 2019

```
library(purrr)
library(tidyr)
library(knitr)
library(invgamma)
```

### 2.3

```
set.seed(6)
```

```
b<- seq(0,1, by = .0001)
```

```
a_ <- 0
```

```
b_ <- 0
```

```
#checking a range of values to get the  
#appropriate a and b that meets conditions
```

```
for(i in 1:length(b)){  
  a = (100*b[i]^2)  
  med = qgamma(.5,a,b[i])  
  if(med%>%round(0) == 75){  
    a_ = a;b_ = b[i]  
    break  
  }  
}
```

```
#condition 1, var = 100
```

```
var<- a_/b_^2  
var
```

```
## [1] 100
```

```
#condition 2 median = 75
```

```
med_<- rgamma(10000, a_, b_)%>%median()  
med_%>%round()
```

```
## [1] 75
```

The prior that satisfy both conditions is  $\text{Gamma}(56.175025, 0.7495)$

## 2.5

*#information given in exercise*

```
c= c(1,1,2,2)
```

```
a<- c(.1,1,.1,1)
```

*#matrix to store values*

```
data<- matrix(.0,nrow = 4, ncol = 9)
```

```
data[,1]<- c
```

```
data[,2]<- a
```

```
data[,3]<- a
```

*#function to build determine the sse*

```
sse <- function(){  
  x2=15;x=-2;n= 20  
  ((1/(n-1))* x2- (x/n)^2)%>%return()  
}
```

*#function for the mean*

```
mean<- function(a, b){  
  n = 20  
  ((sse() + b)/(n + 2*a - 2))%>%return()  
}
```

*#calculating the std*

```
for(i in 1:nrow(data)){  
  #post A  
  data[i,4]= (20/2) + data[i,3]  
  #post B  
  data[i,5] = (data[i,3] + sse())  
  #P(sigma>c)  
  data[i,6] = 1- pinvgamma(data[i,1], data[i,4], data[i,5])  
  # post mean  
  data[i,7] = mean(data[i,3], data[i,3])  
  # post std  
  data[i,8] = (sse()/(20-1))^.5  
  #adding random a and b to see what effect the prior had on the posterior  
  data[i,9] = mean(10, 10)
```

```
}
```

```
df2<- as.data.frame(data)
```

```
names(df2)<- c("c", "a", "b", "A", "B", "postProb", "postMean", "postStd",  
"priorTest")
```

```
df2%>%kable(caption = "Results Of Posterior")
```

### Results Of Posterior

c	a	b	A	B	postProb	postMean	postStd	priorTest
1	0.1	0.1	10.1	0.8794737	0.0e+00	0.0483227	0.2025461	0.2836704
1	1.0	1.0	11.0	1.7794737	2.8e-06	0.0889737	0.2025461	0.2836704
2	0.1	0.1	10.1	0.8794737	0.0e+00	0.0483227	0.2025461	0.2836704
2	1.0	1.0	11.0	1.7794737	0.0e+00	0.0889737	0.2025461	0.2836704

The results for  $P(\theta > c)$  are in column postProb.

```
#P(sigma>c | c= 1, a=b=0.1) / P(sigma>c | c=1, a=b=1)
```

```
r1 <- data[1,6]/data[2,6]
```

```
#P(sigma>c | c= 2, a=b=0.1) / P(sigma>c | c=2, a=b=1)
```

```
r2 <- data[3,6]/data[4,6]
```

The ratio when  $c = 1$  the ratio is 0.0095544 and when  $c = 2$  the ratio is 0.0118331. Also, the posterior is very sensitive to the prior. This is shown in column priorTest where the prior mean changes greatly when I plug in any value for rate and shape.