

INTRODUCTION TO ELEMENTARY ANALYTIC NUMBER
THEORY

Masum Billal

INTRODUCTION TO ELEMENTARY ANALYTIC NUMBER THEORY

Masum Billal

Introduction to Elementary Analytic Number Theory

MASUM BILLAL

September 30, 2021

Contents

Notations	v
1 Arithmetic Functions	1
2 Dirichlet Convolution and Generalization	3
3 Bertrand to Tchebycheff	5
4 A Modest Introduction to Sieve Theory	7
4.1 Glossary	9

Notations

$\gcd(a, b)$ Greatest common divisor of a and b

$\text{lcm}(a, b)$ Least common multiple of a and b

$\varphi(n)$ Euler's totient function of n

$\tau(n)$ Number of divisors of n

$\sigma(n)$ Sum of divisors of n

$\omega(n)$ Number of distinct prime divisors of n

$\Omega(n)$ Number of total prime divisors of n

$\lambda(n)$ Liouville function of n

$\mu(n)$ Möbius function of n

$\vartheta(x)$ Tchebycheff function of the first kind

$\psi(x)$ Tchebycheff function of the second kind

$\zeta(s)$ Zeta function of the complex number s

Chapter 1

Arithmetic Functions

In this chapter, we will discuss some generalized arithmetic functions and their asymptotic behavior. We will skip discussing the basic definitions since they are common in most introductory number theory texts. But for the sake of completeness, check Möbius function in section 4.1

Summatory function. For an arithmetic function f , the *summatory function* of f is defined as

$$F(n) = \sum_{d|n} f(d)$$

Note that the number of divisor function $\tau(n)$ is the summatory function of the unit function $u(n) = 1$. The sum of divisor function $\sigma(n)$ is the summatory function of the invariant function $f(n) = n$. An interesting property that we will repeatedly use is that

$$\begin{aligned} \sum_{i=1}^n F(i) &= \sum_{i=1}^n \sum_{d|i} f(d) \\ &= \sum_{i=1}^n \left\lfloor \frac{n}{i} \right\rfloor f(i) \end{aligned}$$

Here, the last equation is true because there are $\lfloor n/i \rfloor$ multiples of i not exceeding n . We can

Generalized sum of divisor. The *generalized sum of divisor* function can be

defined as

$$\sigma_k(n) = \sum_{d|n} d^k$$

We can write $\tau(n) = \sum_{ab=n} 1$.

Generalized number of divisor. The *generalized number of divisor* function is defined as

$$\tau_k(n) =$$

We will focus on the following function for now.

$$\sum_{n \leq x} \tau_k(n)$$

THEOREM 1. *A test theorem*

This is a theorem¹. Now I am citing Apostol²

¹This is a theorem

²Tom M. Apostol: *Introduction to analytic number theory*. In: *Undergraduate Texts in Mathematics* (1976). DOI: 10.1007/978-1-4757-5579-4.

Chapter 2

Dirichlet Convolution and Generalization

In this chapter, we will discuss Dirichlet convolution and its generalization, use Dirichlet derivative to prove the Selberg identity, establish some results using generalized convolution and finally, prove the fundamental identity of Selberg.

Dirichlet product. For two arithmetic functions f and g , the *Dirichlet product* or *Dirichlet convolution* of f and g is defined as

$$f * g = \sum_{d|n} f(d)g\left(\frac{n}{d}\right)$$

It is not so easy to see how the idea of Dirichlet convolution originates even though it is highly used in number theory. We can connect its origin with the zeta function.

$$\begin{aligned}\zeta(s) &= \frac{1}{1^s} + \frac{1}{2^s} + \dots \\ &= \sum_{i \geq 1} \frac{1}{i^s}\end{aligned}$$

4 *CHAPTER 2. DIRICHLET CONVOLUTION AND GENERALIZATION*

Chapter 3

Bertrand to Tchebycheff

Definition. Tchebycheff function of the first kind or *Tchebycheff's theta function* is defined as

$$\vartheta(x) = \sum_{p \leq x} \log p$$

Definition. Tchebycheff function of the second kind or *Tchebycheff's psi function* is defined as

$$\begin{aligned} \psi(x) &= \sum_{n \leq x} \Lambda(n) \\ &= \sum_{p^e \leq x} \log p \\ &= \sum_{p \leq x} \left\lfloor \frac{\log x}{\log p} \right\rfloor \log p \end{aligned}$$

Chapter 4

A Modest Introduction to Sieve Theory

4.1 Glossary

Möbius function $\mu(n)$ is a very important function in number theory.

Index

Dirichlet convolution, 3

Generalized number of divisor, 2

Generalized sum of divisor, 1

Zeta function, 3