

Bordism Homology and Cohomology

Paul Jin Robaschik

Geboren am 25. Juni 2004 in Köln

21. Mai 2025

Bachelorarbeit Mathematik

Betreuer: Prof. Dr. Markus Hausmann

Zweitgutachterin: Dr. Elizabeth Tatum

MATHEMATISCHES INSTITUT

MATHEMATISCH-NATURWISSENSCHAFTLICHE FAKULTÄT DER
RHEINISCHEN FRIEDRICH-WILHELMS-UNIVERSITÄT BONN

Contents

0	Introduction/Motivation	2
1	Basic Definitions	2
2	Bordism	3
2.1	Unoriented bordism	3
2.1.1	Definitions	3
2.1.2	The Eilenberg-Steenrod Axioms	6
2.1.3	relative bordism	7
2.1.4	Bordism homology	8
2.1.5	Calculations	11
2.2	oriented bordism	11
3	Cobordism	11
3.1	Cobordism	11
3.1.1	Definitions	11
3.1.2	The Eilenberg-Steenrod axioms	13
3.1.3	Bordism cohomology	13
3.1.4	Calculations	13

0 Introduction/Motivation

Recall the definition of homotopy groups $\pi_n(X)$ as the set of homotopy classes of maps from the n -sphere S^n to a space X . The problem with these groups is that they are generally hard to compute, even for simple spaces. For example, the homotopy groups of spheres are not known in general. The general idea of bordism is to replace the n -sphere with a manifold of dimension n and to consider the homotopy classes of maps from this manifold to a space X .

Testing citations:[Ati61],[BD70],[Tho54],[Lee13],[Hat02],[Die08],[Lüc05]

1 Basic Definitions

Here, Examples still need to be added.

Definition 1.1 (Topological manifold [Lee13, pp.2-3]). An n -dimensional **topological manifold** is a topological space M such that:

- M is Hausdorff, (i.e. any two distinct points can be separated by disjoint open sets),
- M is second countable, (i.e. there exists a countable basis for the topology of M),
- M is locally Euclidean (i.e. every point in M has a neighbourhood homeomorphic to an open subspace of \mathbb{R}^n).

We will often write M^n for an n -dimensional manifold.

Remark. One could replace the condition of being second countable with the condition of being paracompact (i.e. every open cover of M admits a locally finite refinement). The following equivalence holds:

second countable \iff paracompact and countably many connected components

Definition 1.2 ((Smooth) Atlas [Lee13, p.12]). Let M be a topological manifold. A **(smooth) atlas** \mathcal{A} on M is a collection of smooth charts $(U_\alpha, \varphi_\alpha)$ such that:

- the $\{U_\alpha\}$ cover M ,
- the charts are pairwise smoothly compatible (i.e. the transition functions $\varphi_j \circ \varphi_i^{-1} : \varphi_i(U_i \cap U_j) \rightarrow \varphi_j(U_i \cap U_j)$ are smooth)

Definition 1.3 (Equivalence of atlases). Two atlases \mathcal{A} and \mathcal{A}' (on a fixed topological manifold) are said to be **equivalent**, if their union is still an atlas.

This is an equivalence relation. — Sadly, Lee does this over maximal atlases. So I need a different citation.

Definition 1.4 (Smooth manifold [Lee13, p.13]). A **smooth manifold** $M = (M, [\mathcal{A}])$ consists of

- a topological manifold M ,
- an equivalence class $[\mathcal{A}]$ of smooth atlases on M .

Remark. While being a topological manifold is just a property of the topological space M , being a smooth manifold gives the manifold extra structure.

Definition 1.5 (Manifold with boundary [Lee13, p.25]). To get a definition of a (smooth or topological) **manifold with boundary**, replace the condition of the manifold being locally Euclidean with the condition that every point has a neighbourhood homeomorphic to an open subspace of $\mathbb{H}^n := \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_1 \geq 0\}$ (the half space).

Remark. In this thesis, with manifold we will always mean a smooth manifold with boundary.

Definition 1.6 (Boundary [Lee13, p.25]). A point $x \in M^n$ is called a **interior point** if it admits a neighbourhood homeomorphic to \mathbb{R}^n . Otherwise, it is called a **boundary point**. The set of boundary points is denoted by ∂M and is called the **boundary** of M . If a manifold is compact and has empty boundary, it is called a **closed manifold**.

Example (Standard n -disk). *The **standard n -disk** is the set of points in \mathbb{R}^n such that $|x| \leq 1$. The boundary of the standard n -disk is the standard $(n-1)$ -sphere. The standard n -disk is denoted by D^n and the standard $(n-1)$ -sphere is denoted by S^{n-1} .*

Remark. The boundary of an n -dimensional manifold is an $(n-1)$ -dimensional submanifold.

Maybe I will make a theorem out of this and add a proof.

Non-example. *Line with two origins*

2 Bordism

The problem with working with manifolds is that they are hard to classify up to homeomorphism or diffeomorphism. Bordism is a way to classify manifolds up to a weaker equivalence relation, which is easier to work with.

2.1 Unoriented bordism

2.1.1 Definitions

Examples need to be added here, too.

Definition 2.1 (Singular manifold [BD70, II, Definition 1.1]). Let X be a topological space. An n -dimensional **singular manifold** in X is a pair M, f of a compact manifold M^n and a continuous map $f : M \rightarrow X$.

The **boundary** of a singular manifold is $\partial(M, f) := (\partial M, \partial f) := (\partial M, f|_{\partial M})$.

Definition 2.2 (Nullbordant [BD70, II, Definition 1.2]). Let (M, f) be a singular manifold in X . We say that (M, f) is **nullbordant**, if there exists a singular manifold (B, F) , such that $\partial(B, F) = (M, f)$.

B, F is then called a **nullbordism** of (M, f) .

Example.

Definition 2.3 (Bordant [BD70, II, Definition 1.3]). Let (M, f) and (N, g) be singular manifolds in X . We say that (M, f) and (N, g) are **bordant**, if their sum $(M, f) + (N, g) := (M + N, (f, g)) := (M \amalg N, f \amalg g)$ is nullbordant.

A nullbordism of $(M, f) + (N, g)$ is called a **bordism** between (M, f) and (N, g) .

We will sometimes refer to this relation as **bordism relation**.

Remark.

$$(M, f) + (\emptyset, g) \text{ are bordant } \iff (M, f) \text{ is nullbordant}$$

Example. *Cylinder, ...*

Non-example.

Proposition 2.4. [BD70, II, Satz 1.4] *Being bordant is an equivalence relation on the set of singular manifolds.*

Proof. [BD70]

- Symmetry: Follows from the symmetry of the disjoint union. If (M, f) and (N, g) are bordant, then there exists a nullbordism of $(M, f) + (N, g) = (M \amalg N, f \amalg g) = (N \amalg M, g \amalg f) = (N, g) + (M, f)$. So, a bordism between (M, f) and (N, g) is also bordism between (N, g) and (M, f) .
- Reflexivity: Cylinder
- Transitivity: Draw a picture

□

Check smooth structure!

Definition 2.5 (bordism group [BD70, II, Definition 1.5]). The equivalence classes of the bordism relation are called **bordism classes** and are denoted by $[M, f]$. The set of bordism classes of n -dimensional singular manifolds in X is denoted by $\mathfrak{N}_n(X)$ and is called the **n -th bordism group of X** .

$$\mathfrak{N}_n = \{\text{singular } n\text{-manifolds in } X\} / \text{bordism}$$

Observe the similarity with the definition of singular homology groups.

Theorem 2.6. [BD70, II, Satz 2.1] The bordism groups are abelian groups with the operation defined in 2.3:

$$[M_1, f_1] + [M_2, f_2] = [M_1 + M_2, (f_1, f_2)]$$

Every element in this group has order at most 2, making $\mathfrak{N}_n(X)$ a \mathbb{F}_2 -vector space.

Proof. [BD70] The neutral element is the bordism class of the empty manifold (i.e. the class of all nullbordant manifolds).

„+“ is associative and commutative, because the disjoint union is associative and commutative.

It is well-defined: by + of the two bordisms.

Since being bordant is a reflexive, every element is its own inverse. \square

Let X, Y be two topological spaces. Then we define a product map

$$\cdot : \mathfrak{N}_p(X) \times \mathfrak{N}_q(Y) \rightarrow \mathfrak{N}_{p+q}(X \times Y)$$

as

$$((M, f), (N, g)) \mapsto (M \times N, f \times g)$$

Remark. Check: Well-definedness, bilinearity

For $X = \{\text{pt}\}$, we write $\mathfrak{N}_n(X)$ as \mathfrak{N}_n and for elements of \mathfrak{N}_n , we will omit the map from the notation ($[M] = [M, f]$).

Definition 2.7 (graded bordism ring [BD70, Satz 2.2]).

$$\mathfrak{N}_* := \bigoplus_{n \in \mathbb{Z}} \mathfrak{N}_n$$

is a graded ring over \mathbb{F}_2 and is called the **bordism ring**.

Remark. As \mathbb{F}_2 is a field, \mathfrak{N}_* is a graded vector space over \mathbb{F}_2 .

Maybe do a quick check.

Definition 2.8 (graded bordism module [BD70, Satz 2.3]).

$$\mathfrak{N}_*(X) := \bigoplus_{n \in \mathbb{Z}} \mathfrak{N}_n(X)$$

is a graded module over \mathfrak{N}_* via

$$[M] \cdot [N, f] = [M \times N, f \circ \text{pr}_2]$$

where $\text{pr}_2 : M \times N \rightarrow N$ is the projection onto the second factor.

As a \mathbb{Z} -graded module over \mathfrak{N}_*

2.1.2 The Eilenberg-Steenrod Axioms

The Eilenberg-Steenrod axioms are a set of axioms that characterize the homology and cohomology theories.

Definition 2.9 (Homology theory [Lüc05, Definition 1.1]). A **homology theory** $\mathcal{H}_* = (\mathcal{H}_*, \partial_*)$ with coefficients in R -modules is a covariant functor

$$\mathcal{H}_* : \text{TOP}^2 \rightarrow \mathbb{Z}\text{-graded } R\text{-modules}$$

together with a natural transformation

$$\partial_* : \mathcal{H}_* \rightarrow \mathcal{H}_{*-1} \circ I$$

- **Homotopy invariance**

Let $f, g : (X, A) \rightarrow (Y, B)$ be homotopic maps. Then for all $n \in \mathbb{Z}$, we have

$$\mathcal{H}_n(f) = \mathcal{H}_n(g) : \mathcal{H}_n(X, A) \rightarrow \mathcal{H}_n(Y, B)$$

- **Long exact sequence**

Let (X, A) be a pair of spaces. Then for all $n \in \mathbb{Z}$, we have the long exact sequence of homology groups:

$$\dots \xrightarrow{\partial_{n+1}(X, A)} \mathcal{H}_n(A) \xrightarrow{\mathcal{H}_n(i)} \mathcal{H}_n(X) \xrightarrow{\mathcal{H}_n(j)} \mathcal{H}_n(X, A) \xrightarrow{\partial_n(X, A)} \mathcal{H}_{n-1}(A) \rightarrow \dots$$

- **Excision**

Let $A \subset B \subset X$ be subspaces of X such that $\overline{A} \subset B^\circ$. Then the inclusion $i : (X \setminus B, A \setminus B) \rightarrow (X, A)$ induces an isomorphism of homology groups for all $n \in \mathbb{Z}$:

$$\mathcal{H}_n(i) : \mathcal{H}_n(X \setminus A, B \setminus A) \xrightarrow{\cong} \mathcal{H}_n(X, B)$$

Sometimes one adds the following axioms:

- **Disjoint union axiom**

Let $\{X_i\}_{i \in I}$ be a family of topological spaces. Let $j_i : X_i \rightarrow \coprod_{i \in I} X_i$ be the inclusion. Then for all $n \in \mathbb{Z}$, we have a bijection:

$$\bigoplus_{i \in I} \mathcal{H}_n(j_i) : \bigoplus_{i \in I} \mathcal{H}_n(X_i) \xrightarrow{\cong} \mathcal{H}_n\left(\coprod_{i \in I} X_i\right)$$

- **Dimension axiom**

For the point space pt , we have

$$\mathcal{H}_n(\text{pt}) \cong \begin{cases} R & n = 0 \\ \{0\} & n \neq 0 \end{cases}$$

For homology theories, the abbreviation \mathcal{H}_* is often used to refer to $\mathcal{H}_*(\{\text{pt}\}, \emptyset)$.

Before we can say that bordism defines a homology theory, we need to define relative bordism.

2.1.3 relative bordism

Definition 2.10 (relative bordism [Die08, pp.524-525]). For a pair of topological spaces (X, A) , we call a $(M, f) = (M, \partial, f)$ a **singular manifold in X** , if $f : (M, \partial M) \rightarrow (X, A)$ is a continuous map of pairs.

Two singular manifolds (M_0, f_0) and (M_1, f_1) in X are called **bordant**, if there exists a singular manifold (B, F) in X , such that:

- (B, F) is a $(n + 1)$ -dimensional compact manifold with boundary,
- $\partial B = \partial_0 B \cup \partial_1 B \cup \partial_2 B$ such that $\partial(\partial_2 B) = \partial(\partial_0 B) \amalg \partial(\partial_1 B)$ and for $i \in \{0, 1\}$, $\partial_i B \cap \partial_2 B = \partial(\partial_i B)$
- There are diffeomorphisms $g : (M_i, f_i) \xrightarrow{\cong} (\partial_i B, \partial(\partial_i B))$ for $i \in \{0, 1\}$ such that $\partial F \circ g_i = \partial f_i$
 $\partial f_i, \partial g_i, \partial F_i$ are as defined in 2.1.
- $F(\partial_2 B) \subseteq A$

As before, B, F will be called a **bordism** between (M_0, f_0) and (M_1, f_1) and a **nullbordism** of M_0, f_0 if $M_1 = \emptyset$.

Example. *Disks on sphere*

Theorem 2.11. [Die08, p.525] *Relative bordism is an equivalence relation on the set of singular manifolds in X .*

Maybe do a proof or just say that it is similar to the proof of the bordism relation.

relative bordism groups are graded modules in the same way. They are denoted by $\mathfrak{R}_*(X, A)$. If $A = \emptyset$, the definition coincides with the definition of bordism groups.

Lemma 2.12. [Zha23, Lemma 5.1] *Let $[M, f] \in \mathfrak{R}_n(X, A)$ and V a embedded submanifold, such that $[V, f|_V] \in \mathfrak{R}_n(X, A)$ and $f(M \setminus N) \subseteq A$. Then $[M, f] = [V, f|_V]$ in $\mathfrak{R}_n(X, A)$.*

Proof. [Zha23] We need to show that (M, f) and $(N, f|_N)$ are bordant.

Let $B = M \times I$ the cylinder. $\partial B = \underbrace{M \times \{0\}}_{M_0} \cup \underbrace{M \times \{1\}}_{M_1} \cup \partial M \times I$. Define $g : B \rightarrow X$ as

$$g(p, t) = f(p).$$

Writing $M_1 = (M_1 \setminus N_1) \cup N_1$, we get that $M \times I$ is a bordism between M and N , because

$$g(\partial B \setminus (M_0 \cup N_1)) = g((M_1 \setminus N_1) \cup (\partial M \times I)) = f(M \setminus N) \cup f(\partial M) \subseteq A$$

□

2.1.4 Bordism homology

Lemma 2.13. [BD70, II, Satz 3.2] *Relative bordism is a covariant functor*

$$\mathfrak{N}_* : \text{TOP}^2 \rightarrow \text{graded } \mathfrak{N}_* \text{ modules}$$

Proof. [BD70] Let $(X, A) \in \text{Ob}(\text{TOP}^2)$, we already saw

$$(X, A) \xrightarrow{\mathfrak{N}_*} \mathfrak{N}_*(X, A)$$

For a map $\text{Mor}(\text{TOP}^2) \ni f : (X, A) \rightarrow (Y, B)$, we take the induced map on the bordism groups:

$$f_* := \mathfrak{N}_*(f) : \mathfrak{N}_*(X, A) \rightarrow \mathfrak{N}_*(Y, B)$$

given by $f_*[M, g] = [M, f \circ g]$ for $[M, g] \in \mathfrak{N}_n(X, A)$ and $n \in \mathbb{N}$.

Then we get that $\mathfrak{N}_*(\text{id}_{(X,A)}) = \text{id}_{\mathfrak{N}_*(X,A)}$ and for $f : (X, A) \rightarrow (Y, B)$, $g : (Y, B) \rightarrow (Z, C)$, we have for any $[M, h] \in \mathfrak{N}_*(X, A)$:

$$(g \circ f)_*[M, h] = [M, g \circ f \circ h] = g_*[M, f \circ h] = g_* \circ f_*[M, h]$$

□

Need to add: Naturality of the boundary map

Lemma 2.14 (Homotopy invariance). [BD70, II, Satz 3.1] *\mathfrak{N}_* is homotopy invariant.*

Proof. [BD70] Let $f, g : (X, A) \rightarrow (Y, B)$ be homotopic maps. Let $F : X \times I \rightarrow Y$ be a homotopy between f and g . Then we have a bordism between $f_*[M, h]$ and $g_*[M, h]$ by $(M \times I, F \circ (h \times \text{id}_I))$. □

Maybe do an example here. (Special case Cylinder done before showing that bordism is an equivalence relation)

Lemma 2.15 (Long exact sequence [Die08, Proposition 21.1.9]). *\mathfrak{N}_* satisfies the long exact sequence axiom.*

Compare with [Zha23]!

Proof. [Die08] Let i, j be the inclusion maps $i : A \rightarrow X, j : X = (X, \emptyset) \rightarrow (X, A)$.

Claim: The sequence

$$\dots \xrightarrow{\partial} \mathfrak{N}_n(A) \xrightarrow{i_*} \mathfrak{N}_n(X) \xrightarrow{j_*} \mathfrak{N}_n(X, A) \xrightarrow{\partial} \mathfrak{N}_{n-1}(A) \xrightarrow{i_*} \dots$$

is exact.

- **Exactness at $\mathfrak{N}_n(A)$** From the definitions, $i_* \circ \partial = 0$.

If (B, F) is a nullbordism of $f : M \rightarrow A$ in X , then $\partial[B, F] = [M, f]$

- **Exactness at $\mathfrak{N}_n(X)$** Let $[M, f] \in \mathfrak{N}_n(A)$. Choose $V = \emptyset$ and use 2.12 to get $[M, f] = 0$ in $\mathfrak{N}_n(X, A)$, so $j_* \circ i_* = 0$.
Now let $[M, f] \in \mathfrak{N}_n(X)$ such that $j_*([M, f]) = 0$. A nullbordism of $[M, f]$ in X, A is a bordism between M, f and N, g with $g(N) \subset A$. Then $i_*[N, g] = [M, f]$ follows.
- **Exactness at $\mathfrak{N}_n(X, A)$** Again, $\partial \circ i_* = 0$ follows from the definitions.
Let $\partial[M, f] = 0$, $[B, F]$ be a nullbordism of $\partial M, f|_{\partial M}$. Identify (M, f) and (B, f) along ∂M . Call the resulting singular manifold (C, g) . Now with 2.12, get $j_*[C, g] = [M, f]$.

□

Theorem 2.16 (Sard's theorem [Lee13]). *The set of critical values has measure zero.*

We only need to check the excision axiom now to see that bordism is a homology theory. But to see the excision property, we need some preliminary lemmas.

Lemma 2.17 ([Zha23, Lemma 5.6]). *Let $K, L \subseteq M$ be disjoint closed subsets of a compact manifold M . Then there exists a closed submanifold with boundary $N \subseteq M$ with $K \subseteq N, L \cap N = \emptyset$.*

Proof. [Zha23]

□

In the above (nonexistent yet, may look into Connor,Floyd) proof, a smooth α with regular value r is constructed.

Lemma 2.18 ([Zha23, Lemma 5.8]). *Let K, L, M, N be as in 2.17. Then,*

$$\partial N \subseteq \partial M \cup \alpha^{-1}(r) \subseteq \partial M \cup ((M \setminus K) \cap (M \setminus L))$$

Lemma 2.19 (Excision axiom [Zha23, Theorem 5.10]). *Let X, A, Z be a triple of topological spaces satisfying $\overline{Z} \subseteq \overset{\circ}{A}$. Then the inclusion map $i : (X \setminus Z, A \setminus Z) \hookrightarrow (X, A)$ induces an isomorphism of bordism groups:*

$$i_* : \mathfrak{N}_n(X \setminus Z, A \setminus Z) \xrightarrow{\cong} \mathfrak{N}_n(X, A)$$

Proof. Surjectivity: Let $[M, f] \in \mathfrak{N}_n(X, A)$. Then the preimages $K = f^{-1}(X \setminus \overset{\circ}{A})$ and $L = f^{-1}(\overline{Z})$ are disjoint and closed subsets of M . By 2.17, there exists a closed submanifold with boundary $N \subset M$ such that $K \subseteq N$ and $L \cap N = \emptyset$.

From $L \cap N = \emptyset$, it follows that $f(N) \subseteq X \setminus \overline{Z}$. By 2.18, we have $\partial N \subseteq \partial M \cup ((M \setminus K) \cap (M \setminus L))$. So, for any $p \in \partial N$, we have either $p \in \partial M$, implying $f(p) \in A$, or $p \in (M \setminus K)$, implying $f(p) \in \overset{\circ}{A}$. In any case, we get $f(\partial N) \subseteq A \setminus \overline{Z}$, so $[N, f|_N] \in \mathfrak{N}_n(X \setminus Z, A \setminus Z)$.

As $f^{-1}(X \setminus \overset{\circ}{A})$, we get $f(M \setminus N) \subseteq \overset{\circ}{A}$. By 2.12, we get $i_*[N, f|_N] = [M, f]$

Injectivity: Take $[M, f] \in \mathfrak{N}_n(X \setminus Z, A \setminus Z)$ such that $i_*[M, f] = 0$ in $\mathfrak{N}_n(X, A)$. Then there exists an $(n+1)$ -manifold B and $g : B \rightarrow X$, such that M is an embedded

submanifold of ∂B , $g(\partial B \setminus M) \subseteq A$ and $g|_M = f$.

Again, let $K = g^{-1}(X \setminus \overset{\circ}{A})$, $L = g^{-1}(\overset{\circ}{Z})$. Then we have an embedded submanifold $W \subseteq B$ such that $K \subseteq W$, $W \cap L = \emptyset$. So $[\partial W, g|_{\partial B}] \in \mathfrak{N}_n(X \setminus Z, A \setminus Z)$ nullbordant.

Claim: $M \cap \partial B = M \cap B$. „ \subseteq “ is clear. For „ \supseteq “, take $p \in M \cap B$. Then $p \in \partial W$, so there exists a chart $\varphi : U \rightarrow \mathbb{H}^{n+1}$ such that $\varphi(p) \in \partial \mathbb{H}^{n+1}$. So, $p \in \partial B$.

Then $M \cap \partial B$ is an embedded submanifold of M , because

$$M \cap \partial B = M \cap B = (\alpha|_M)^{-1}([0, r]) \quad (1)$$

and we can assume that r is a regular value, because of 2.16. Now we can see that $(M \cap \partial B, f|_{M \cap \partial B}) \in \mathfrak{N}_n(X \setminus Z, A \setminus Z)$: $f(M \cap \partial B) \subseteq f(M) \subseteq X \setminus Z$ and $f(\partial(M \cap \partial B)) \subseteq A \setminus Z$ because of 1 and 2.18, as $\partial(M \cap \partial B) \subseteq \partial M \cup (K^c \cap L^c)$.

Claim: $[M, f] = [M \cap B, f|_{M \cap B}] \in \mathfrak{N}_n(X \setminus Z, A \setminus Z)$. By 2.12, it is enough to show that $f(M \setminus (M \cap B)) = f(M \setminus \partial(M \cap B)) \subseteq A \setminus Z$. Let $p \in M \setminus \partial B = M \setminus B$, then $f(p) \in X \setminus Z$ because $p \in M$, and $f(p) \in \overset{\circ}{A}$, because $p \notin B$. The claim follows.

By the same argument, $[M \cap B, f|_{M \cap B}] = [\partial B, g|_{\partial B}] \in \mathfrak{N}_n(X \setminus Z, A \setminus Z)$. Since $M \cap B$ is an embedded submanifold of M , it is also an embedded submanifold of ∂B . Now, it is enough to show that $g(\partial B \setminus (M \cap B)) = g(\partial B \setminus M) \subseteq A \setminus Z$. We know $\partial B \subseteq \partial W \cup (K^c \cap L^c)$. Let $p \in \partial B \setminus M$, then either $p \in (K^c \cap L^c) \setminus M$ or $p \in \partial W \setminus M$. In the first case, we are done. In the second case, we know $p \notin L$, because B, Q are disjoint. So, $g(p) \notin \overset{\circ}{Z}$. By assumption, we have $g(\partial W \setminus M) \subseteq A$, so $g(p) \in A \Rightarrow g(p) \in A \setminus Z$. In total, we have shown that $(M, f), (M \cap B, f|_{M \cap B}), (\partial B, g|_{\partial B})$ are bordant, so nullbordant, since the last one is nullbordant in $\mathfrak{N}_n(X \setminus Z, A \setminus Z)$, so $[M, f] = 0$ in $\mathfrak{N}_n(X \setminus Z, A \setminus Z)$. \square

Now it already follows that bordism is a homology theory. Let let's take a look at the other axioms too.

Lemma 2.20 (Disjoint union axiom [Zha23, Theorem 5.3]). *The disjoint union axiom holds for bordism.*

Proof. [Zha23] We need to show that

$$\bigoplus_{i \in I} \mathfrak{N}_n(j_i) : \bigoplus_{i \in I} \mathfrak{N}_n(X_i) \rightarrow \mathfrak{N}_n\left(\coprod_{i \in I} X_i\right)$$

is an isomorphism.

Injectivity:

Surjectivity:

\square

Observation. *Bordism does not satisfy the dimension axiom.*

Now we can finally conclude:

Theorem 2.21. *Bordism defines a homology theory satisfying the disjoint union axiom.*

Proof. This follows directly from the lemmas 2.13, 2.14, 2.15, 2.19 and 2.20. The only axiom that does not hold is the dimension axiom. □

This theorem should probably be done after cobordism.

Theorem 2.22. *There is a natural equivalence of homology theories*

$$\mathfrak{N}_*(-) \xrightarrow{\cong} \mathcal{H}_*(-, \mathbf{MO})$$

2.1.5 Calculations

We will now calculate the bordism groups.

Of points

Of spheres

2.2 oriented bordism

Definition 2.23 (vector bundle).

Definition 2.24 (orientation).

orientation bundle

Definition 2.25 (bordant).

Definition 2.26 (oriented bordism group).

$$\Omega_n(X)$$

3 Cobordism

Now that we have seen that bordism is a homology theory, we can ask the question whether there is a dual theory, giving rise to a cohomology theory. The answer is yes, as we will see now.

I should give an explanation of how $O(n)$ (real bundles) and $SO(n)$ (oriented real bundles) come into play.

3.1 Cobordism

3.1.1 Definitions

Maybe I will not need this

Definition 3.1 (Classifying space). Let G be a topological group. The **classifying space** BG of G is the

Observation.

$$\begin{aligned} BO(n) &= \mathbb{G}_{k,\infty} \\ BSO(n) &= \tilde{\mathbb{G}}_{k,\infty} \end{aligned}$$

I might adjust this to be more like [Tho54]’s definition.

Definition 3.2 (Thom space[BD70]). Let $\xi : E \rightarrow B$ be a real k -dimensional vector bundle over a compact manifold B . Then the Thom space of ξ is defined as

$$M(\xi) = E^c$$

the one-point compactification of the total space of ξ , the added point serving as the base point.

Definition 3.3 ($MO(n), MSO(n)$ [BD70]).

$$MO(n) := M(\xi_{n,\infty})$$

with $\xi_{n,\infty}$ being the universal real vector bundle over $\mathbb{G}_{k,\infty}$.

$$MSO(n) :=$$

Maybe it is enough to just say that these are called this way because the Grassmannians are the classifying spaces of $O(n), SO(n)$ instead of defining classifying spaces.

Theorem 3.4 (Suspension sequence). *Let X, Y be pointed spaces. We denote by $[X, Y]^\circ$ the set of homotopy classes of pointed maps $X \rightarrow Y$. Then we have the **suspension sequence**:*

$$[X, Y] \rightarrow [\Sigma X, \Sigma Y] \rightarrow \cdots \rightarrow [\Sigma^n X, \Sigma^n Y]$$

Theorem 3.5 (Freudenthal suspension theorem).

No proof.

Lemma 3.6. [Tho54] *The natural map*

$$\Sigma\{MSO(n)\} \rightarrow MSO(n+1)$$

induces isomorphisms of homotopy groups π_{n+1} for n large.

Corollary 3.7 ([Ati61, p.201]). *For a finite CW-complex X with basepoint,*

$$[X, \Sigma\{MSO(n)\}] \rightarrow [X, MSO(n+1)]$$

is bijective for n large.

Lemma 3.8 ([Ati61, p.201]). *Let Y be a subcomplex of a CW-complex X . Then from 3.4 and 3.6, we get a map*

$$[\Sigma^{n-k}(X/Y), MSO(n)] \rightarrow [\Sigma^{n+1-k}(X/Y, MSO(n+1))]$$

Definition 3.9 (relative oriented cobordism group [Ati61, p.201]). Let (X, Y) be a pair of spaces, then for $k \in \mathbb{Z}$, the **k -th oriented cobordism group** is

$$MSO^k(X, Y) = \lim_{n \rightarrow \infty} [\Sigma^{n-k}(X/Y), MSO(n)]$$

with respect to the above map.

3.1.2 The Eilenberg-Steenrod axioms

To get a definition for cohomology theories, intuitively, we „reverse all arrows“ in the previously defined axioms for homology theories in 2.1.2.

Definition 3.10 (Cohomology theory [Lüc05, Definition 5.2]). A **cohomology theory** $\mathcal{H}^* = (\mathcal{H}^*, \partial^*)$ with coefficients in R -modules is a contravariant functor

$$\mathcal{H}^* : \text{TOP}^2 \rightarrow \mathbb{Z}\text{-graded } R\text{-modules}$$

together with a natural transformation

$$\partial^* : \mathcal{H}^* \circ I \rightarrow \mathcal{H}^{*+1}$$

satisfying the following axioms:

- **Homotopy invariance**
- **Long exact sequence**
- **Excision**

Sometimes one adds the following axioms:

- **Disjoint union axiom**
- **Dimension axiom**

3.1.3 Bordism cohomology

Again, we will check the axioms one by one.

3.1.4 Calculations

Thom isomorphism, Thom class, Thom isomorphism theorem

References

- [Ati61] M. F. Atiyah. „Bordism and cobordism“. In: Proc. Cambridge Philos. Soc. 57 (1961), pp. 200–208. ISSN: 0008-1981. DOI: 10.1017/s0305004100035064. URL: <https://doi.org/10.1017/s0305004100035064>.
- [BD70] Theodor Bröcker and Tammo tom Dieck. Kobordismentheorie. Vol. Vol. 178. Lecture Notes in Mathematics. Springer-Verlag, Berlin-New York, 1970, pp. xvi+191.
- [Die08] Tammo tom Dieck. Algebraic topology. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2008, pp. xii+567. ISBN: 978-3-03719-048-7. DOI: 10.4171/048. URL: <https://doi.org/10.4171/048>.
- [Hat02] Allen Hatcher. Algebraic topology. Cambridge University Press, Cambridge, 2002, pp. xii+544. ISBN: 0-521-79540-0.
- [Lee13] John M. Lee. Introduction to smooth manifolds. Second. Vol. 218. Graduate Texts in Mathematics. Springer, New York, 2013, pp. xvi+708. ISBN: 978-1-4419-9981-8.
- [Lüc05] Wolfgang Lück. Algebraische Topologie. Homologie und Mannigfaltigkeiten. German. Vieweg Stud. Wiesbaden: Vieweg, 2005. ISBN: 3-528-03218-9.
- [Tho54] René Thom. „Quelques propriétés globales des variétés différentiables“. In: Comment. Math. Helv. 28 (1954), pp. 17–86. ISSN: 0010-2571,1420-8946. DOI: 10.1007/BF02566923. URL: <https://doi.org/10.1007/BF02566923>.
- [Zha23] Zhuo Zhang. „A Geometric View on Bordism Homology“. English. In: (Dec. 2023).