
BECOMING A DOCTOR OF

MARRIAGE AND FAMILY THERAPY

DOCTORAL TEXT AND ADVANCED PROFESSIONAL REFERENCE



TRIUNE BIBLICAL UNIVERSITY GLOBAL EXTENSION USA

BECOMING A DOCTOR OF MARRIAGE AND FAMILY THERAPY

BECOMING A DOCTOR OF

Marriage and Family Therapy

A Doctoral Text and Advanced Professional Reference on Family Systems Theory, Couple and Family Assessment, Clinical Practice, Supervision, and Research in Marriage and Family Therapy

TRIUNE BIBLICAL UNIVERSITY GLOBAL EXTENSION USA

First Edition · 2026

Becoming a Doctor of Marriage and Family Therapy

First Edition, 2026

Published by TRIUNE BIBLICAL UNIVERSITY GLOBAL EXTENSION USA

© 2026 Triune Biblical University Global Extension USA. All rights reserved.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means — electronic, mechanical, photocopying, recording, or otherwise — without the prior written permission of the publisher, except for brief quotations embodied in critical articles, reviews, and scholarly works with full attribution.

ISBN (print): to be assigned
ISBN (electronic): to be assigned

Academic notice. This book is a doctoral-level academic text in psychology, psychotherapy, and the behavioural sciences. It is written for study, research, and professional education in marriage and family therapy. It is not clinical supervision, diagnosis, or treatment, and it does not by itself confer a doctoral or any other academic qualification. Doctoral standing depends on enrolment in a programme, supervised original research, and successful examination.

Professional notice. Nothing in this book authorises or supports practice beyond the reader's training, competence, licensure, and lawful scope of practice, or practice that is contrary to the ethical codes and legal requirements of the jurisdiction concerned. Readers must obtain the current editions of the codes, statutes, and standards referred to in these pages; the law and professional regulation described are stated as understood at the date of writing (2026).

Research notice. Studies, effect sizes, and findings are reported in summary form to support theoretical analysis. Readers who rely on a particular finding must consult the original publication, its methods, its sample, and any subsequent replication or critique.

Hypothetical material. Cases, families, and scenarios marked as hypothetical are fictitious and are used exclusively for teaching and examination purposes. Any resemblance to actual persons or families is coincidental. Where case material derives from a published source, the source is cited.

Illustrations. Diagrams and figures were prepared for this edition. Photographs marked "illustrative image (AI-generated)" were created for teaching purposes and do not depict any identifiable person, family, clinic, or agency.

Typeset in EB Garamond and Source Sans 3. Trim size 6 × 9 inches.

DEDICATION

To the researchers who test what practitioners believe, and to the practitioners who test what researchers propose —

and to the families whose difficulties have taught this discipline almost everything it knows.

Contents

Foreword ix
Preface xi
Acknowledgements xiii
About This Book xv
Abbreviations xvii
List of Figures and Illustrations xix
List of Tables xxiii
Introduction: Theory, Practice, and the Doctoral Study of Family Systems xxv
How to Use This Book xxvii

PART I Foundations of Systemic Theory and Practice 1

1 Advanced Theories of Family Systems 3
Introduction 3
1.1 The Systemic Turn: Cybernetics, Systems Theory, and Epistemology 5
1.2 Structural Family Theory 12
1.3 Strategic, Brief, and Milan Systemic Approaches 17
1.4 Intergenerational and Contextual Theories 20
1.5 Experiential, Communication, and Humanistic Theories 25
1.6 Attachment, Emotion, and Neurobiological Theories 28
1.7 Social Constructionist and Narrative Theories 33
1.8 Cognitive-Behavioural, Integrative, and Evidence-Based Synthesis 36

PART II Research, Evidence, and Clinical Decision-Making 49

2 Research Methods in Marriage and Family Therapy 51
Introduction 51
2.1 From Clinical Concern to Researchable Question 53
2.2 Paradigms and the Choice of Method 56
2.3 Research Designs, Sampling, and Inference 59
2.4 Measurement and the Relational Unit of Analysis 62
2.5 Evaluating Interventions and Causal Claims 64
2.6 Research Ethics, Safety, and Data Governance 67
2.7 Analysis, Reporting, and Evidence Synthesis 69
2.8 From Evidence to Practice and Implementation 71
2.9 Constructing a Doctoral Research Proposal 73
Summary 74
Doctoral Research Questions 75
Further Reading 76
Selected Methodological References 76
End of Chapter 2 78

PART III Culture, Context, and Relational Practice 79

3 Multicultural Issues in Family Therapy 81
Introduction 81
3.1 Culture, Identity, and the Context of Family Life 83
3.2 Cultural Humility, Positionality, and Power 86
3.3 Culturally Responsive Assessment and Formulation 89

3.4 Language, Communication, and Meaning . . . 93
3.5 Kinship, Migration, and Transnational Family Life . . . 94
3.6 Race, Ethnicity, and the Effects of Racialisation . . . 98
3.7 Gender, Sexuality, Disability, and Socioeconomic Context . . . 99
3.8 Adaptation, Evidence, and Organisational Practice . . . 102
3.9 Supervision, Documentation, and Continuing Development . . . 104
Summary . . . 106
Doctoral Research Questions . . . 107
Further Reading . . . 107
End of Chapter 3 . . . 108

PART IV Clinical Reasoning, Assessment, and the Therapeutic Relationship 109

4 Advanced Clinical Practice in Marriage Therapy . . . 111
Introduction . . . 111
4.1 The Advanced Couple Therapist's Clinical Stance . . . 113
4.2 Engagement, Contracting, and the First Sessions . . . 115
4.3 Comprehensive Couple Assessment . . . 117
4.4 Safety, Coercion, and Suitability for Conjoint Work . . . 120
4.5 Formulation: From Assessment to a Shared Map . . . 123
4.6 Treatment Planning and Selecting an Approach . . . 124
4.7 Managing the Therapeutic Process . . . 127
4.8 Monitoring Progress, Referral, and Ending . . . 129
4.9 Integrating the Clinical Arc . . . 131
Summary . . . 133
Doctoral Research Questions . . . 134
Further Reading . . . 134
End of Chapter 4 . . . 135

A Genogram Notation . . . 137
A.1 Household structure . . . 138
A.2 Descent and vital events . . . 139
A.3 Migration, culture, and social context . . . 139
A.4 Relationship notation and the cultural genogram . . . 139
A.5 Recording conventions for doctoral work . . . 139

B Comparative Grid of Family Systems Theories . . . 141
B.1 Theories compared on core analytical dimensions . . . 141
B.2 Therapist stance, session composition, and typical duration . . . 143
B.3 A checklist for evaluating a theory . . . 144

C The Doctoral Programme and the Study Pathway for MFT. 901 . . . 145
C.1 The programme . . . 145
C.2 Relation of this volume to the programme . . . 146
C.3 A nine-week study pathway for MFT. 901 . . . 147
C.4 Assessment design . . . 148

Selected Bibliography . . . 151
1. Founding texts of systemic theory and cybernetics . . . 151
2. Histories, handbooks, and systematic overviews of the field . . . 152
3. Structural family theory . . . 152
4. Strategic, brief, MRI, and Milan systemic approaches . . . 152
5. Intergenerational and contextual theories . . . 153
6. Communication, experiential, and humanistic theories . . . 154
7. Attachment, emotion, and neurobiological accounts . . . 154
8. Cognitive-behavioural, behavioural, and parenting models . . . 155
9. Multisystemic, multidimensional, and family-based treatment models . . . 156
10. Social constructionist, narrative, and solution-focused approaches . . . 156
11. Culture, gender, power, and critique . . . 157
12. Integration, common factors, and the evidence base . . . 158
13. Reference works, journals, and methodological resources . . . 159
14. Research methods, measurement, and evidence synthesis . . . 159
15. Multicultural, intersectional, and culturally responsive practice . . . 160
16. Professional and regulatory sources . . . 161

Foreword

Marriage and family therapy is a young discipline with an old subject. Its subject — the ways in which people who are bound to one another by kinship, partnership, and care shape each other's conduct, emotion, and health — is as ancient as human society. Its disciplinary identity, however, was consolidated only in the middle decades of the twentieth century, when clinicians and researchers in the United States, Britain, Italy, and elsewhere began to treat the family not as the background of individual pathology but as a system with properties of its own.

This book is the first volume in a curriculum-based series written for doctoral study in marriage and family therapy. Each volume addresses one course in the doctoral programme of Triune Biblical University Global Extension USA, and each is designed to stand alone as a text that can be read, taught, and examined in its own right. The present volume addresses the foundational course of the programme, *MFT. 901 Advanced Theories of Family Systems*.

The purpose of the series is not to replace the primary literature. Doctoral candidates must read the original texts — Bateson's essays on communication, Minuchin's structural analyses, Bowen's papers on differentiation, Johnson's attachment-based treatment manuals, and the empirical studies on which contemporary practice rests. What a text of this kind can do is to provide a disciplined map: to explain where the ideas came from, what they claim, how they relate to one another, what evidence supports them, where they have been criticised, and how a doctoral researcher can work with them. The series therefore treats theory as a set of tools that must be understood in their historical and cultural context and evaluated on their merits, not as doctrine to be recited.

The book is written for three audiences. The first is the doctoral candidate who is encountering family systems theory at an advanced level and must be able to compare, apply, and critique competing formulations. The second is the practising clinician who wishes to ground practice in theoretical understanding rather than in habit. The third is the researcher and teacher who needs a consolidated account of the field's conceptual architecture, together with its unresolved questions.

Each chapter follows a consistent design: learning objectives; an introduction; numbered sections of analysis; illustrative case material, clearly identified as hypothetical or documentary; applied exercises; a summary; doctoral research questions; and annotated guidance for further reading. Diagrams and illustrations are placed at the points in the argument where they are needed, and are drawn or prepared for this edition.

The series is academic in method and professional in orientation. It engages evidence, including evidence that is inconvenient for preferred theories. It attends to culture, gender, power, and context, because a systems perspective that ignores them is not systematic at all. Above all, it treats the reader as a colleague whose task is not to memorise conclusions but to form independent, defensible judgements.

The Editorial Board
Doctoral Texts in Marriage and Family Therapy

Preface

This volume is a doctoral-level introduction to the theories that constitute the intellectual foundation of marriage and family therapy. It has been written for the first year of the doctorate, where candidates are expected to acquire a secure command of the field's theoretical inheritance, together with the critical apparatus needed to evaluate it.

Theory in this discipline is not decoration. A therapist who has no explicit theory still has one: an implicit set of assumptions about what causes distress, who is responsible for it, what counts as change, and whose voice should be heard. Implicit theory is not thereby wrong, but it is unaudited, and unaudited assumptions produce predictable clinical failures — blaming a parent for a child's symptom, escalating a couple's conflict in the name of honesty, treating a cultural difference as resistance, or abandoning a family because the therapist's preferred model has stopped working. The purpose of a doctoral course in family systems theory is to make theory explicit, comparable, and testable.

The chapters in this volume are organised around the major traditions of the field: the cybernetic and systems-theoretical foundations; structural theory; strategic and systemic approaches; intergenerational and contextual theories; experiential, communication, and humanistic traditions; attachment, emotion, and neurobiological accounts; social constructionist and narrative approaches; and cognitive-behavioural and integrative frameworks. Each tradition is presented in its own terms before it is evaluated, so that the reader understands what its authors were trying to do rather than only what later critics said about them. The chapters also examine the empirical record: which theories have generated research programmes, which have demonstrated effects, and where the evidence is thin, contested, or simply absent.

Three editorial commitments shape the text. First, the argument is presented in a plain academic register, with technical terms defined when they are introduced and used consistently thereafter. Second, the material is comparative: theories are placed against one another so that their commitments, predictions, and limits

become visible. Third, the text is clinically honest. It does not promise more than the evidence supports, and it identifies the points at which the field is still divided.

The volume is a text for study, not a substitute for supervised clinical training, personal therapy, or professional regulation. Nothing in it should be read as a guarantee of outcome, and no account of clinical technique in these pages relieves the practitioner of the obligations imposed by law, professional codes, ethical guidelines, and the local systems within which practice occurs.

A word on case material. Illustrative cases in this book are hypothetical unless they are attributed to a published source, and hypothetical cases are labelled. Where a case is constructed to demonstrate a theoretical point, the point is the point; the family is a teaching device. Readers should assume that any resemblance between a hypothetical family and their own clients is a coincidence of the kind that family systems theory itself predicts — the patterns are generic, the particulars are not.

Acknowledgements

A text of this kind is a synthesis of other people's work. The authors whose theories are examined here — Gregory Bateson, Don Jackson, Salvador Minuchin, Jay Haley, Mara Selvini Palazzoli, Murray Bowen, Ivan Boszormenyi-Nagy, Virginia Satir, Carl Whitaker, Paul Watzlawick, Sue Johnson, Leslie Greenberg, John Gottman, Norman Epstein, Donald Baucom, Andrew Christensen, Neil Jacobson, William Pinsof, Douglas Breunlin, Michael White, David Epston, Steve de Shazer, Insoo Kim Berg, Harlene Anderson, Harry Goolishian, Richard Schwartz, and many others — wrote primary texts that remain indispensable. Any summary is a simplification; the reader is urged throughout to consult the originals.

The editorial team thanks the faculty and examiners of Triune Biblical University Global Extension USA for their review of the curriculum design, and the doctoral candidates whose questions in seminars shaped the structure and emphasis of the chapters. Their persistent demand was for clarity: what does this theory actually claim, what evidence supports it, and what follows for practice? This volume attempts to answer those questions in that order.

The diagrams in this book were prepared for this edition. Photographs marked as illustrative images are AI-generated representations created for teaching purposes and do not depict identifiable persons, agencies, or events. Tables, exercises, and measurement instruments (where reproduced) are illustrative; readers requiring standardised measures must obtain them from the copyright holders or the public-domain sources indicated in the text.

Responsibility for the selection, interpretation, and any errors in what follows rests with the editorial team alone.

About This Book

Triune Biblical University Global Extension USA publishes this series as part of its doctoral curriculum in marriage and family therapy. The series translates each course of the doctoral programme into a full-length text: comprehensive in coverage, comparative in method, and written to the standard expected of doctoral work.

This volume corresponds to **MFT. 901 Advanced Theories of Family Systems**, the foundational theoretical course of the programme. It is designed to be used alongside the remaining volumes of the series, which address research methods, multicultural practice, clinical practice and assessment, psychopathology, ethics and law, human development, couple therapy, trauma, substance use, family life-cycle transitions, group work, neurobiology, supervision, mediation, parenting, integrative practice, technology, play therapy, spirituality in practice as an academic subject, advanced research methodology, and supervised practice.

Scope and level. The text assumes graduate-level preparation in the behavioural sciences or a related professional field. It does not assume prior training in family therapy, but it does not linger over elementary material. Terms of art are defined on first use and indexed thereafter. Statistical and methodological material is introduced where a theory's credibility depends on it.

Structure. The book opens with the historical and epistemological foundations of the systemic approach. It then works through the major theoretical traditions, devoting a section to each; compares them on common dimensions; examines the empirical evidence; and closes with the integrative frameworks that dominate contemporary practice and research. Appendices provide the standard notation for genograms, a comparative grid of theories, and a guide to using the volume in seminar and examination preparation.

Learning features. Each chapter contains:

- **Learning objectives**, stating what the reader will be able to do after study;
- **Numbered analytical sections**, each covering one tradition or theme;
- **Case illustrations**, hypothetical or documented, showing how theory appears in practice;

- **Key theoretical points**, summarising the central claim of each tradition;
- **Comparative perspectives**, placing theories side by side;
- **Applied exercises**, designed for seminars and supervision;
- **Assessment instruments**, describing measurement tools used in research on the theories discussed;
- **Doctoral research questions**, identifying live research problems; and
- **Further reading**, indicating where to begin with the primary literature.

A note on academic honesty. This book cites real works, summarises real research programmes, and identifies real controversies. It contains no invented quotations and no fictional attributions. Where a case is hypothetical, it is labelled as such. Where evidence is limited, the text says so. Where the field is divided, the text reports the division rather than resolving it silently.

Professional notice. This book is an academic text. It is not a clinical manual for unsupervised practice, it does not constitute clinical supervision or professional advice, and it does not by itself confer any academic qualification. Practice in marriage and family therapy is regulated differently across jurisdictions; readers must comply with the law and the professional requirements of the jurisdiction in which they practise, and must work within the limits of their competence.

Abbreviations

Abbreviation	Full form
ACA	Affordable Care Act (United States)
AAMFT	American Association for Marriage and Family Therapy
ACT	Acceptance and commitment therapy
AFT	Association for Family Therapy and Systemic Practice (United Kingdom)
ASIA	Attachment and Biobehavioral Catch-up (attachment-based parenting programme)
ATM	Adaptive therapy model *(see also IST)*
BCT	Behavioural couple therapy
CBCT	Cognitive-behavioural couple therapy
CFT	Couple and family therapy
DBT	Dialectical behaviour therapy
DSM	*Diagnostic and Statistical Manual of Mental Disorders*
ECR	Experiences in Close Relationships (attachment measure)
EFT	Emotionally focused therapy; also Emotionally focused therapy for couples (EFT-C)
FAD	Family Assessment Device
FACES	Family Adaptability and Cohesion Evaluation Scales
FIT	Feedback-informed treatment
FST	Family systems theory
IBCT	Integrative behavioural couple therapy
ICD	*International Classification of Diseases*
IFS	Internal family systems therapy
IPT	Integrative problem-centred therapy
IST	Integrative systemic therapy

Abbreviation	Full form
MFT	Marriage and family therapy; marriage and family therapist
MRI	Mental Research Institute, Palo Alto
MST	Multisystemic therapy
NICE	National Institute for Health and Care Excellence (United Kingdom)
ORS/SRS	Outcome Rating Scale / Session Rating Scale
RCT	Randomised controlled trial
RFT	Reflecting team
SFBT	Solution-focused brief therapy
TBCT	Traditional behavioural couple therapy
UKCP	United Kingdom Council for Psychotherapy

List of Figures and Illustrations

1.1 The historical development of family systems theory. The field's founding texts and research programmes cluster in three generations: the cybernetic founding, the clinical systems building of the 1970s, and the integrative and evidence-based period after 1990. 7

1.2 Linear and circular causality. The linear formulation (A) locates pathology in one member and reads events as a one-way sequence. The circular formulation (B) describes reciprocal influence: each act is both a response to the preceding act and a stimulus for the next, and the same cycle can be entered at any point. 9

1.3 Second-order practice: the observing system. In first-order practice the therapist observes the family from outside the system. In second-order practice the therapist, the team behind the one-way mirror, the recording equipment, and the institutional setting are all inside the system that is being observed and that is, in turn, observing itself. 11

1.4 Structural maps and boundary types. Panel (A) shows a family with clear generation and subsystem boundaries: the partners maintain a spousal subsystem, the siblings maintain a sibling subsystem, and authority is distributed across generations. Panel (B) shows structural pathology: the spousal boundary is diffuse, the mother–daughter relationship is enmeshed, a cross-generational coalition is forming, and overt marital conflict is displaced into the parental subsystem. 13

1.5 Structural practice: the enactment. The therapist directs family members to speak to one another in the session while observing — and intervening in — the pattern that emerges. The session becomes a site of observation and of change at the same time. *Illustrative image (AI-generated).* 14

1.6 The structural sequence of clinical reasoning. The therapist moves from a presenting symptom to an observational description of sequence, a functional hypothesis about the pattern, an intervention designed to change it, and an assessment of structural change. 15

1.7 The emotional triangle. Anxiety in a dyad is displaced onto a third element — a person, an activity, a substance, or a symptom — which stabilises the pair. Triangles are normal features of relationship systems; they become clinically significant when they are rigid, chronic, and reproduced across generations. 21

1.8 Multigenerational transmission. Bowen's account of the amplification of small differences in differentiation across generations: anxiety concentrated in one relationship in one generation produces a lower level of differentiation in the next, where the pattern intensifies, until a descendant presents with symptoms. 22

1.9 Constructing a genogram with a couple. The diagram is built collaboratively in the session, so that the process of construction — who remembers what, who 23

disagrees, which facts were never spoken — becomes part of the assessment. *Illustrative image (AI-generated).* . . . . .

1.10 The negative interaction cycle in emotionally focused therapy. The cycle is presented to the couple as a shared pattern rather than as the responsibility of either partner, which is the precondition for de-escalation and for the attachment-level work that follows. 30

1.11 The integrative problem-solving sequence. Integration is disciplined by a specified sequence: define the system and the problem, formulate across levels, select mechanisms of change with the best available warrant, sequence the treatment, and monitor and revise in the light of outcome data. 39

1.12 Comparative summary of the major theoretical traditions. The grid organises each tradition by its core construct, its mechanism of change, and the state of its evidence base, and is intended for use in seminar work as a working analytical instrument. 41

2.1 A research question is refined by aligning the clinical concern, population and setting, construct, design, evidence, and bounded conclusion. The arrows are iterative: feasibility and stakeholder input may require revision at each stage. 55

2.2 Three broad methodological routes answer different kinds of questions. Mixed-methods research requires an explicit integration step; it is not simply two studies placed side by side. 57

2.3 Designs differ in allocation, timing, and the strength and kind of inference they can support. The continuum is conceptual rather than a universal hierarchy: appropriateness depends on the question, implementation, and assumptions. 60

2.4 MFT observations may be nested and interdependent: members within dyads or families, families within therapists and clinics, and clinics within service systems. The analytic model should match the sampling and intervention structure. 63

2.5 An intervention may influence outcomes through several pathways, while baseline context and delivery affect both participation and observed results. The diagram is a planning aid: the arrows represent hypotheses that must be justified, not established facts. 66

3.1 A culturally responsive formulation holds family interaction, social position, institutions, and historical or material conditions in view at the same time; influence can move in every direction. 85

3.2 Therapist authority, client and family power, institutional rules, and the working alliance influence one another; ethical practice makes these relationships explicit and negotiable. 88

3.3 A culturally responsive clinical formulation links members' meanings and interactional sequences with social position, structural context, resources, and safety; hypotheses are revisited collaboratively. 92

3.4 A practice cycle for culturally responsive MFT: locate context, ask about meaning, formulate tentatively, negotiate goals and adaptation, act, and review impact. The cycle is iterative, not a certification checklist. 103

4.1 Advanced couple therapy proceeds through engagement, assessment and safety, shared formulation, negotiated intervention, outcome review, and a deliberate decision to continue, adapt, refer, or end. Review points loop back when new information emerges. 114

4.2 A balanced assessment moves among each partner's account, the relationship sequence, strengths and history, current context, and safety; the clinician selects depth according to relevance and risk. 120

4.3 A safety decision pathway begins with private inquiry and assessment of freedom and risk, then weighs suitability, expertise, and protections before conjoint work; risk is reviewed throughout rather than screened once. 122

4.4 A formulation-to-plan pathway links each partner's account and the interactional pattern to goals, model choice, agreed tasks, and review criteria; safety and context remain active at every stage. 126

4.5 Progress review combines the couple's own goals and accounts with observed interaction, safety, relevant measures, and contextual change; results guide a shared decision to continue, adapt, refer, or end. 130

A.1 Standard genogram notation. Structural symbols (upper panels) record family composition, descent, and vital events; relationship symbols (lower panel) record the quality of emotional bonds. Notation conventions vary slightly between published sources; records should state which convention is used. 138

List of Tables

B.1 B.1 — Core analytical dimensions of the major traditions. 141
B.2 B.2 — Stance, composition, duration, and characteristic risk. 143
C.1 C.1 — Course structure of the doctoral programme (each course 3 units). 145
C.2 C.2 — Theoretical prerequisites supplied by this volume. 146
C.3 C.3 — Suggested seminar sequence for MFT. 901. 147

Introduction: Theory, Practice, and the Doctoral Study of Family Systems

Marriage and family therapy occupies an unusual position among the applied psychological professions. It treats a unit — the couple, the family, the household, sometimes a wider network of kin and care — that no single discipline owns. It borrows from psychology, psychiatry, sociology, anthropology, communication studies, nursing, social work, education, and, increasingly, neurobiology and developmental science. It practises in a setting — the consulting room — through which streams of law, culture, economics, and health policy flow continuously. And it depends on a body of theory that was assembled, in a single generation, by people who disagreed with one another about almost everything except the premise that the family is a system.

That premise is the starting point of this book. A system, in the sense used throughout these volumes, is a set of elements that interact, such that the state of each element depends on the states of the others, and the behaviour of the whole cannot be predicted from the properties of the parts considered in isolation. Applied to families, the premise yields a distinctive set of questions: what maintains a pattern of interaction, rather than what causes it; how emotional intensity is distributed among members; how hierarchies, alliances, and boundaries organise daily life; how meaning is negotiated; and how change in one part of the system propagates to others.

The premise has consequences for research as well as for practice. It implies that variables are usually reciprocal, that observation is always participant observation, that the unit of analysis is contested (individual, dyad, triads, household, kin network, community), and that culture, class, gender, and migration history are not "context" to be added on but constitutive of the system itself. Doctoral work in the field therefore requires both theoretical fluency and methodological discipline, and the two are inseparable: a theory that cannot be operationalised cannot be tested, and a study that has not decided what the system is cannot measure change within it.

This volume addresses the theory. The chapters that follow move historically and comparatively. They begin with the intellectual foundations of systemic thought — cybernetics, general systems theory, and the shift from first-order to second-order cybernetics — and then examine the major clinical traditions: structural, strategic, systemic, intergenerational, contextual, experiential, communication-based, attachment and emotion-focused, constructionist and narrative, cognitive-behavioural, and integrative. Each tradition is presented through its own concepts, its own founding texts, its own clinical methods, and its own evidence base. Each is then subjected to the same questions: What is the unit of analysis? What is the mechanism of change? What does it predict? What evidence exists? Where has it been criticised — theoretically, empirically, culturally, and ethically?

The final sections of the volume address the two developments that define the present state of the field. The first is the integrative turn: the recognition that no single model is sufficient for the range of problems clinicians encounter, and the construction of frameworks — metaframeworks, common factors, and empirically supported principles — that allow different theories to be combined without collapsing into incoherence. The second is the evidence-based movement, which has changed what it means to hold a theory responsibly: not merely to find it persuasive or useful in one's own practice, but to specify what it predicts, to test it, and to abandon or revise it when the evidence turns against it.

Two cautions apply throughout. The first concerns the seductions of systems language. Circular formulations can explain everything and therefore predict nothing; a theory that can accommodate any outcome has abandoned empirical responsibility. The second concerns power. A systems perspective that treats all members as equally responsible for a pattern can obscure asymmetries of power, dependency, and harm — including violence, coercion, and abuse — that are not distributed symmetrically at all. Subsequent chapters return to both cautions repeatedly, and the doctoral reader should treat them as standing obligations of the discipline.

The volume closes with material designed for use: a comparative grid of theories on common dimensions, the standard notation for genograms, and guidance on constructing a theoretical framework for a doctoral study. These are working tools. Their value is measured by what the reader can do with them — in a seminar, in a viva, in a clinic, and in the design of original research.

How to Use This Book

For the doctoral candidate. Read each section before the seminar in which it is discussed, then re-read it after. The numbered sections are self-contained enough to be read in any order, but the volume is designed to be read in sequence: the later traditions assume the conceptual vocabulary established in the first section. Keep a theory-comparison notebook, recording for each tradition its unit of analysis, its account of problem maintenance, its mechanism of change, its principal methods, and its evidence. By the end of the volume you should be able to reproduce that grid from memory and defend its entries.

For the seminar leader. Each chapter contains an applied exercise designed for group work in ninety minutes, followed by doctoral research questions suitable for written assignments. Case illustrations can be role-played: assign the family positions, one observer to each of two competing theoretical models, and a third observer whose task is to record which model generated a testable formulation and which generated a description that no data could refute.

For the supervisor. The comparative material is intended for use in supervision when a trainee is stuck. The question "which of the traditions in the grid would formulate this case differently, and what would that formulation lead you to do next?" converts theoretical knowledge into clinical flexibility.

For the researcher. The evidence discussions identify where the empirical record is strong, where it is thin, and where it is contested. Those passages are the natural starting points for dissertation work; the doctoral research questions at the end of each chapter mark live problems rather than settled questions.

Conventions. British spelling is used throughout. Terms of art appear in **bold** at first definition and in the index thereafter. Figures are numbered by chapter. Tables are numbered by chapter and carry a caption above the table. Case material is hypothetical unless a published source is cited, and hypothetical exercises are labelled as such. Where an instrument is described, its copyright status is noted; standardised instruments cannot be reproduced freely, and their use is subject to the terms of the publisher or author.

A caution about scope. A volume of this kind can teach theory, evidence, and method. It cannot confer clinical competence. Competence is acquired through supervised practice, direct observation and feedback, personal development work, and compliance with the regulatory and ethical framework of the jurisdiction in which one works. Nothing in this book should be read as authorising practice beyond the reader's training, licensure, or lawful scope of practice.



PART I

Foundations of Systemic Theory and Practice

The intellectual foundations of marriage and family therapy: cybernetics and general systems theory, the major clinical traditions built upon them, the constructionist and evidence-based turns, and the integrative frameworks that define contemporary doctoral practice.

1 CHAPTER

Advanced Theories of Family Systems

LEARNING OBJECTIVES

After studying this chapter, the reader should be able to:

1. Explain the intellectual origins of family systems theory in cybernetics, general systems theory, and information theory, and identify the specific concepts each contributed to the clinical field.
2. Distinguish first-order from second-order cybernetics and analyse the consequences of that distinction for the position of the therapist, the status of the observer, and the conduct of research.
3. Compare the foundational clinical traditions — structural, strategic, systemic, intergenerational, contextual, experiential, communication, attachment-based, constructionist, cognitive-behavioural, and integrative — on the dimensions of unit of analysis, account of problem maintenance, mechanism of change, and evidentiary status.
4. Apply the major theoretical constructs of each tradition to the formulation of a presented clinical problem, and demonstrate how competing formulations generate different interventions.
5. Evaluate the principal theoretical criticisms directed at the systemic tradition, including the charges of circularity, unfalsifiability, neglect of power, and cultural insensitivity.
6. Assess the empirical record of family systems theories, distinguishing between theories with established treatment effects, theories with promising but limited evidence, and theories whose influence rests primarily on clinical or historical rather than empirical grounds.
7. Identify unresolved theoretical problems in the field that constitute viable subjects for doctoral research.

Introduction

Family therapy began as a heresy. Its founders were trained as psychiatrists, psychoanalysts, social workers, and paediatricians, and the proposition they advanced from the 1950s onwards contradicted the dominant clinical orthodoxy

of the period: that disturbance resides inside the individual, is formed in the individual's early history, and is properly treated in the individual's mind. Against this orthodoxy the systemic pioneers proposed that disturbance is located in the space between people, that it is maintained by current patterns of interaction rather than by historical events alone, and that it may legitimately be treated by changing those patterns — sometimes without treating any individual separately at all.

The heresy did not arise in a vacuum. It drew on intellectual resources developed outside clinical medicine during the Second World War and immediately afterwards: Norbert Wiener's cybernetics, Ludwig von Bertalanffy's general systems theory, Claude Shannon's mathematical theory of communication, and the group dynamics of Kurt Lewin and his successors. It also drew on a specifically American tradition of social investigation — Talcott Parsons's structural-functional account of the family as a system of roles, in which the nuclear family was analysed as a differentiated unit performing functions of socialisation and tension management. When these resources were carried into the consulting room by clinicians struggling with schizophrenia, delinquency, psychosomatic illness, and marital breakdown, the result was a new discipline with a new vocabulary.

That vocabulary is now so familiar within the field that its origins and its limits are easily forgotten. Terms such as *system*, *boundary*, *subsystem*, *feedback*, *homeostasis*, *triangle*, *sequence*, *punctuation*, *reframe*, *structure*, *hierarchy*, *enactment*, and *second-order change* are the common currency of clinical training. Each was borrowed from a technical discipline in which it carried a precise meaning, and each was modified — sometimes degrades — in the process of importation. One task of advanced theoretical study is to recover the original meanings and to ask what was gained and lost in translation. A concept such as *homeostasis*, for example, was developed to explain the regulation of physiological variables around set points; imported into family theory as an explanation of resistance to change, it was stretched to cover phenomena — loyalty, fear, cultural obligation, economic dependency — that have nothing to do with physiological regulation, and it carried with it a normative bias in favour of stability that later critics of family therapy have identified as politically loaded.

This chapter has two aims. The first is expository: to reconstruct, with precision, what each major tradition in the field actually claims. The second is critical: to evaluate those claims against theoretical objections, empirical evidence, cultural critique, and clinical experience. Both aims are essential to doctoral work. A candidate who knows what Bowen wrote but cannot say what would count as evidence against Bowen's theory is not yet a scholar of the field; a candidate who can recite the criticisms of structural family therapy but cannot specify

what Minuchin's account of hierarchy enables a clinician to do that a competing account does not is equally unqualified.

The chapter proceeds as follows. Section 1.1 examines the systemic turn: the intellectual sources of the field, the cybernetic concepts that organised its early thinking, and the first-order-to-second-order transition that reshaped its epistemology from the 1980s onwards. Sections 1.2 to 1.7 examine the major clinical traditions in roughly the order in which they developed: structural theory; strategic, brief, and Milan systemic approaches; intergenerational and contextual theories; experiential, communication, and humanistic approaches; attachment, emotion, and neurobiological accounts; and social constructionist and narrative theories. Section 1.8 addresses the cognitive-behavioural tradition, the integrative frameworks that have become the organising structure of contemporary practice, and the evidence-based synthesis that disciplines all of the preceding theoretical claims. The chapter closes with case illustrations, an applied exercise, a summary, doctoral research questions, and guidance for further reading.

Because this chapter necessarily compresses a vast literature, the treatment is analytical rather than exhaustive. Each tradition is presented through a small number of conceptually decisive claims, the methods those claims generate, and the principal lines of criticism. Throughout, the reader should keep three questions in view. What is the unit of analysis that this theory asserts? What, according to this theory, would count as evidence that it is wrong? And what would a practitioner do differently — specifically, differently — if the theory were abandoned tomorrow?

1.1 The Systemic Turn: Cybernetics, Systems Theory, and Epistemology

Preconditions: the family as an object of scientific study

Systemic thinking about families did not appear from nowhere in 1952. It required three preconditions. The first was a tradition of treating the family as a social unit rather than merely a collection of individuals. Developed in sociology and social anthropology in the first half of the twentieth century — notably in the functionalist analyses of kinship and role differentiation, in the ethnographic studies of household organisation, and in the "family interaction" research produced at the University of Chicago and in the family studies movement of the 1920s and 1930s — this tradition gave clinicians a legitimate object: the family as a structure with roles, norms, and boundaries. Early interactional studies of families containing a member diagnosed with schizophrenia, carried out in the 1950s (notably the work on "pseudo-mutuality" and later on "communication deviance"), demonstrated that family conversation could be described systematically, and that the descriptions differed in interesting ways across families.

The second precondition was a language of interaction. This came mainly from cybernetics and information theory, whose vocabulary of feedback, input, output, signal, noise, channel, and code permitted clinicians to describe what passed between people in terms that were neither the language of inner psychic life (which they wished to avoid) nor the language of mechanical cause (which failed to capture reciprocal influence). The third precondition was a technology of observation. The emergence of one-way mirrors, audio recording, video recording, and later the coded observational laboratory — from the family interaction studies of the 1960s to the Gottman laboratory's observational protocols of the 1970s and beyond — made it possible to see and re-see what families actually did, rather than to rely on retrospective report. Each precondition remains active in the field today, and each carries a methodological obligation with it: the interactionist vocabulary must be operationalised, and the observational setting must be recognised as itself a system.



Figure 1.1 The historical development of family systems theory. The field's founding texts and research programmes cluster in three generations: the cybernetic founding, the clinical systems building of the 1970s, and the integrative and evidence-based period after 1990. *Source:* prepared for this edition.

Cybernetics: feedback, homeostasis, and deviation

Norbert Wiener's *Cybernetics*, published in 1948, defined the study of control and communication in the animal and the machine, and introduced the concept that became the field's organising metaphor: **feedback**. In a feedback loop, information about the current state of a system's output is returned to the system and modifies its subsequent behaviour. Where the returned information opposes the deviation — that is, where the system acts to restore a prior state — the loop is **negative** and the behaviour is regulatory. Where the returned information amplifies the deviation, the loop is **positive** and the behaviour is destabilising or escalatory. Walter Cannon's earlier concept of **homeostasis**, describing the physiological regulation of temperature, blood chemistry, and fluid balance around physiologically viable ranges, supplied the language of set points and regulation that family theorists subsequently adapted.

Two further cybernetic ideas were equally consequential. The first was **circularity**: in a feedback system, cause and effect are not arranged in a line but in a loop, so that any statement identifying a single cause is, strictly speaking, an artefact of where the observation is begun. The second was the distinction, drawn by W Ross Ashby and others, between the variety a system can generate and the variety it can absorb — a constraint later summarised as the law of requisite variety. Applied to families, the idea predicts that a system facing a degree of change greater than its regulatory capacity will either reorganise or fail.

The importation of cybernetics into clinical thinking was the work of a small group in Palo Alto. Gregory Bateson's research project, funded from 1952, assembled Don D Jackson, John Weakland, Jay Haley, and William Fry, and produced in 1956 the paper on the "double bind" hypothesis — an attempt to link schizophrenic communication deficits to pathological communicational contexts. The double bind hypothesis itself has not survived empirical scrutiny in its original form, and the causal claim that family communication causes schizophrenia was abandoned long ago. Its historical importance lies elsewhere: it established the practice of describing psychopathology in terms of communication within a relational system, and it produced the field's first generation of researchers and clinicians.




Figure 1.2 Linear and circular causality. The linear formulation (A) locates pathology in one member and reads events as a one-way sequence. The circular formulation (B) describes reciprocal influence: each act is both a response to the preceding act and a stimulus for the next, and the same cycle can be entered at any point. *Source:* prepared for this edition.

KEY THEORETICAL POINT

The concept of circular causality is the field's foundational epistemological commitment. It holds that in a stable interaction pattern there is no non-arbitrary first cause: the pattern is self-maintaining, and any causal narrative that begins with one member's behaviour is a *punctuation* of the sequence rather than an empirical discovery. This commitment generates the characteristic clinical question of systems practice — not "why does this person behave this way?" but "what maintains this pattern, and who benefits from its persistence?" — and it generates the characteristic research problem of the field, namely how to study reciprocal influence without collapsing it into a unidirectional claim.

General systems theory and open systems

In parallel with cybernetics, the biologist Ludwig von Bertalanffy developed **general systems theory**, which he presented as a set of principles applicable to entities that maintain themselves in exchange with their environments. Two of his emphases entered family theory directly. The first was the distinction between **closed** and **open** systems. A closed system tends toward entropy; an open system

imports energy and information from its environment, exports outputs, and maintains itself far from equilibrium — the biological and, by analogy, the familial condition of continuous exchange with surroundings. The second emphasis was **hierarchy and organisation**: systems are nested, so that each system is simultaneously a whole composed of parts and a part of a larger whole. Nested organisation supplied family therapy with a concept of levels — individual, dyad, household, kin network, community, society — and with a criterion of adequacy for theory: an account pitched at one level that ignores the levels above and below it is, by the standards of systems thinking, incomplete.

Later contributions enlarged the vocabulary. Magoroh Maruyama's analysis of deviation-amplifying processes distinguished morphostasis (pattern maintenance) from morphogenesis (pattern change), thereby escaping the connotation of universal conservatism attached to homeostasis. Information theory contributed the notion that patterns may be described economically as **rules** — regularities that constrain behaviour across situations — and that an observer may describe a system's rules even when no member could state them. Taken together, these ideas supplied the conceptual apparatus later chapters will examine in clinical form: subsystems, boundaries, hierarchy, feedback, rules, and levels of organisation.

From first-order to second-order cybernetics

The early systemic writings treated the family as an observable system and the therapist as an external agent who could observe it, map it, and intervene in it. This became known as **first-order cybernetics**. By the late 1970s this position was undermined from within, by two arguments. The epistemological argument, drawn from the biology of cognition developed by Humberto Maturana and Francisco Varela, held that living systems are *autopoietic*: they operate according to their own structure, and an external agent cannot determine what a system will do — it can only perturb it, with results determined by the system's own organisation. The observer, on this view, is not outside the observed system but part of the description, and the sharp distinction between objective observation and subjective involvement collapses. The clinical argument, developed by the Milan school and its successors and articulated influentially in the work of Lynn Hoffman and later by Harlene Anderson and Harry Goolishian, held that a therapist who claims a privileged position outside the family reproduces the very hierarchy of expertise that the systemic approach was supposed to question.

Second-order cybernetics therefore reconceived the therapist as a participant in the system under observation, whose own premises, culture, and emotional responses are part of the clinical picture. The clinical consequences are substantial. If the therapist cannot stand outside the system, then the ideal of neutrality — central to the Milan approach — must be reinterpreted as a discipline of

curiosity rather than a claim to objectivity; the therapist's hypotheses must be held tentatively, since the family will respond in ways the therapist's own structure permits it to respond; and the therapist's own reactivity becomes data to be monitored rather than noise to be eliminated. In research, the consequence is that the observer's presence and framing must be treated as part of the phenomenon, a methodological demand familiar from participant observation and from reflexive qualitative traditions.



Figure 1.3 Second-order practice: the observing system. In first-order practice the therapist observes the family from outside the system. In second-order practice the therapist, the team behind the one-way mirror, the recording equipment, and the institutional setting are all inside the system that is being observed and that is, in turn, observing itself. *Source:* prepared for this edition.

Illustrative image (AI-generated). This photograph was created for this edition for teaching purposes; it depicts no identifiable clinic, team, or family.

COMPARATIVE PERSPECTIVE

First-order and second-order positions compared. In a first-order frame the therapist's task is to obtain an accurate map of the family's structure and to intervene to change it; progress is assessed against the therapist's formulation; the family is the object of treatment. In a second-order frame the therapist's task is to participate in a conversation that generates new descriptions and possibilities; progress is assessed through the meanings that emerge in the system, including the therapist's contribution to them; and the appropriate stance is one of curiosity, tentativeness, and responsibility for one's own participation. Most contemporary practitioners work in a mixed frame. The distinction remains important because it determines what a therapist does when an intervention fails: in the first-order frame, a failed interven-

> tion suggests an inaccurate map; in the second-order frame, it is first an occasion to examine the therapist's own premises and position in the system.

The move to second-order cybernetics also changed the relationship between family therapy and its own history. If observation is always from within, then the field's theories cannot be treated as neutral discoveries about families; they are constructions produced by observers with particular locations, assumptions, and interests. That recognition opens the field to the social constructionist and narrative traditions examined in Section 1.7, and it makes the critical examination of power, culture, and gender — examined in Sections 1.7 and 1.8 — an internal requirement of systemic theory rather than an external reproach.

1.2 Structural Family Theory

The structural claim

Structural family theory, associated above all with Salvador Minuchin and his colleagues at the Philadelphia Child Guidance Clinic, holds that the central features of a family are its **structure**: the recurrent patterns of interaction that organise who relates to whom, in what way, and with what authority. Structure, in this usage, is not a hidden entity but an observable regularity. It is revealed by who sits where, who speaks for whom, who intervenes in whose conversations, how decisions are reached, and what happens when the family is under stress. Because structure is a pattern rather than a substance, it is described in relational terms — proximity, hierarchy, alliance, coalition, boundary — and it is understood to be both stable and changeable.

Three constructs carry most of the theoretical weight. The first is the **subsystem**: a component of the family defined by function, such as the parental subsystem, the sibling subsystem, or the spousal subsystem, each with its own tasks and its own rules of participation. The second is the **boundary**: the invisible line regulating the flow of information, affect, and contact between subsystems and between the family and its environment. Minuchin distinguished **clear** boundaries, which permit appropriate contact and appropriate separation; **diffuse** boundaries, which produce enmeshment and reduce autonomy; and **rigid** boundaries, which produce disengagement and reduce support. The third is **hierarchy**: the distribution of authority and responsibility, which in healthy families places the parental subsystem in an executive position with respect to younger children and requires that generational roles not be inverted. Structural pathology, on this account, appears as an inappropriate arrangement of these elements: weak executive authority, a **cross-generational coalition** in which one parent allies with a child against the other, an inverted hierarchy in which

the child's symptom regulates the marriage, or an enmeshed subsystem in which differentiation is impossible.




Figure 1.4 Structural maps and boundary types. Panel (A) shows a family with clear generation and subsystem boundaries: the partners maintain a spousal subsystem, the siblings maintain a sibling subsystem, and authority is distributed across generations. Panel (B) shows structural pathology: the spousal boundary is diffuse, the mother–daughter relationship is enmeshed, a cross-generational coalition is forming, and overt marital conflict is displaced into the parental subsystem. *Source:* prepared for this edition.

Method: enactment, joining, and restructuring

Structural theory is inseparable from its clinical method, and the method is inseparable from the theory: the therapist learns about the family's structure by attempting to change it and observing what happens. Minuchin described three overlapping phases. In **joining**, the therapist accommodates to the family's style — its language, pace, humour, and hierarchy — in order to gain access to the system without triggering its defences. Joining is not a technique of manipulation but a precondition of influence: a therapist who cannot be heard by the family's most influential member cannot restructure anything. In **enactment**, the therapist invites the family to talk to one another in the session rather than to the therapist, and then intervenes in the resulting interaction — increasing intensity, blocking a familiar move, redirecting a conversation from parent to child, or requiring a disengaged member to take a position. The third phase, **restructuring**, uses directives, boundary-making moves, and the deliberate redistribution of time


and attention to install a new pattern that the family can maintain without the therapist.



Figure 1.5 Structural practice: the enactment. The therapist directs family members to speak to one another in the session while observing — and intervening in — the pattern that emerges. The session becomes a site of observation and of change at the same time. *Illustrative image (AI-generated).*

Illustrative image (AI-generated), created for this edition for teaching purposes. It depicts no identifiable family, clinician, or agency, and the composition is stylised rather than representative of any particular service.

The theoretical significance of enactment has been under-appreciated in the literature. It is the field's clearest attempt to solve the problem of observation. Rather than asking a family to report on itself, or inferring structure from diagnostic assessment, the structural therapist generates structure in the room and intervenes on it in real time. The approach thereby treats the session as a microcosm — a move that presupposes, but does not demonstrate, that patterns displayed in the consulting room resemble patterns displayed at home. That presupposition is an empirical claim of exactly the kind later observational research on family interaction could test, and the results have been mixed: behaviour in the laboratory shows substantial stability for some variables and considerable reactivity for others.



›rder change alters intensity within an unchanged structure; second-order change alters the rules, boundaries, or pre that generated the sequence.

Figure 1.6 The structural sequence of clinical reasoning. The therapist moves from a presenting symptom to an observational description of sequence, a functional hypothesis about the pattern, an intervention designed to change it, and an assessment of structural change. *Source:* prepared for this edition.

Enmeshment, disengagement, and the empirical record

The enmeshment–disengagement dimension became one of the most widely used constructs in the field, and one of the most criticised. The critical literature identifies three problems. First, the concepts were originally formulated as poles of a continuum with the healthy range at the middle, but the clinical writing frequently treated enmeshment as a near-synonym for maternal over-involvement, which introduced a normative and gendered evaluation into what claimed to be a structural description. Second, the dimension overlaps with, but is not identical to, the closely related construct of family **cohesion** in the circumplex model of David Olson and colleagues, whose central hypothesis — that adaptive family functioning lies in the balanced middle range of cohesion and flexibility, not at either extreme — generated an extensive measurement programme using the Family Adaptability and Cohesion Evaluation Scales and thirty years of empirical debate about whether the relationship between cohesion and outcome is curvilinear. Third, the empirical work that has examined enmeshment directly has been conducted chiefly within the literature on adolescent development, where the relevant construct is **psychological control** and the evidence is that intrusive, autonomy-constraining parenting is associated with internalising and externalising difficulties, while warm and close, but autonomy-supportive, parenting is not. The structural distinction between clear and diffuse boundaries therefore survives in revised form: what matters is not closeness but control.

The treatment evidence for structural family therapy is more substantial than the theoretical controversy might suggest. The Philadelphia programme pro-

duced controlled studies of family therapy for adolescent substance misuse and delinquent behaviour (the family-therapy condition of the studies by Minuchin and colleagues and later by Howard Liddle and colleagues on multidimensional family therapy), and testable claims about the role of family interaction in the management of somatic conditions: the studies of family interaction and diabetes and asthma conducted with Maurizio and Dora Andolfi's group in Rome, and the well-known observations that symptom intensity in some children with brittle diabetes or severe asthma varies with family interactional conflict. Contemporary structural practice has been absorbed, in much of the literature, into broader integrative and multidimensional models — multidimensional family therapy, multisystemic therapy, and brief strategic family therapy — that retain the structural focus on hierarchy, alliance, and engagement in the family's context while adding interventions aimed at individual functioning, peer networks, school, and community.

Case illustration: structural formulation in action

::: casebox {title="Case Illustration 1.1 — The presenting symptom as an organisational requirement (hypothetical)"} **The referral.** A family is referred by the local school following a sudden decline in the academic performance of a fourteen-year-old boy, accompanied by episodes of asthma severe enough to require two hospital admissions in six weeks. The referring paediatrician notes that medical management has been optimised and suggests that family factors be assessed.

The family. Both parents work. The couple report that they have not discussed the state of their marriage "for years", and each describes the other as preoccupied. The mother manages all aspects of the son's medical care and communicates with the hospital directly; the father describes himself as "unable to do anything right" in relation to the illness and withdraws when the mother and son begin to discuss symptoms or schoolwork.

Structural formulation. From a structural standpoint the family presents an inverted hierarchy in the medical domain: the parental subsystem has no unified executive function. The mother and son form a strong subsystem whose organising theme is illness; the father occupies a disengaged position; the spousal subsystem is weak. The son's escalation of symptoms reliably produces a degree of parental coordination that is otherwise absent, and the sequence has the effect of suspending the marital distance by providing a shared (though conflictual) task. The symptom is not "caused" by the marriage in any linear sense; it is specified by the pattern that surrounds it.

Intervention. The therapist joins by taking the father's experience of exclusion seriously and by acknowledging the mother's burden without framing it as

over-involvement. Interventions focus on re-establishing a functioning parental subsystem with respect to health and schooling: a shared task assigned to the couple rather than to the mother alone; an enactment in which the parents negotiate a plan in the son's presence while the son is asked to comment only at the end; and the deliberate discussion of the couple's relationship, which the son is excused from early in treatment.

Methods note. This case is hypothetical. It illustrates how the same set of facts can be described as an individual illness trajectory, a systemic maintenance pattern, or a structural imbalance, and how the description chosen determines the treatment offered. :::

1.3 Strategic, Brief, and Milan Systemic Approaches

The strategic tradition

The strategic tradition grew from the same Palo Alto soil as the communication theory examined below, and it shares a characteristic emphasis: therapy is a planned attempt to resolve a defined problem, and techniques are evaluated by their effects rather than by their theoretical elegance. Jay Haley's *Strategies of Psychotherapy* (1963) and *Problem-Solving Therapy* (1976) defined the essentials: therapy has a social function as well as an analytical one; the therapist must establish a position of sufficient influence to direct change while remaining within the family's hierarchy; and the presenting problem is embedded in a sequence of interactions that can be interrupted. Haley distinguished the problem-maintaining sequence from its historical origins, which made possible a form of therapy that was deliberately brief and deliberately non-historical.

Cloe Madanes extended the strategic framework with an attention to metaphor and structure in symptom presentation, notably the argument that symptoms may serve as metaphors that render a family's situation comprehensible, and that interventions may be designed to shift the relational requirement the symptom satisfies. The strategic tradition's signature techniques include **directives** (tasks assigned for the interval between sessions), **reframing** (the deliberate re-description of a behaviour in terms that open a new response), **ordeals** (assigning a burdensome task that must be performed if the symptom is to continue, thereby increasing its cost), and — most controversially — **paradoxical** prescriptions, in which the therapist prescribes the symptom or its continuation on the assumption that the family will resist the prescription and thereby change.

Paradoxical techniques require extended comment because they carry consequential ethical risk. The theoretical rationale is that a system resistant to direct instruction may mobilise its own resistance against a therapeutic instruction, producing change; the empirical status of the technique is weak, with little controlled evidence; and the ethical objections are substantial where the technique

involves deception, where the family has not consented, where the presentation involves violence, self-harm, or abuse — conditions in which the prescriptive stance may be actively harmful — and where the therapist lacks the skill to manage an extreme response. Contemporary practice has largely absorbed reframing and directive tasks while abandoning paradox as a routine strategy, though it survives in some brief therapy traditions. That trajectory is instructive for doctoral readers: a technically elegant intervention may be abandoned not because its theory was refuted but because its risks were recognised as disproportionate to its plausible benefits.

MRI brief therapy

The Mental Research Institute, founded in Palo Alto in 1959 with Don Jackson as its first director, developed the most economical of the systemic models. In *Change: Principles of Problem Formation and Problem Resolution* (Watzlawick, Weakland, and Fisch, 1974), the MRI group advanced a theory of **problem formation**: difficulties of living are universal and usually resolve themselves, but a small proportion become chronic because the ways in which people attempt to solve them — more of the same of an ineffective solution, denial that a problem exists, or attempts to change a rule that cannot be changed at that level of the system — succeed only in maintaining them. Therapy, on this account, is directed at the attempted solution rather than at the original difficulty, and it is by design limited to a small number of sessions with the explicit goal of interrupting the vicious circle.

The MRI model's contribution to theory is its economy and its emphasis on the difference between the problem and the *problem about the problem*. Its contribution to practice is a set of interventions — attention to interactions around the problem rather than to the problem's content, negotiation of concrete and minimal goals, tasks framed to interrupt the attempted solution — that have been absorbed so widely that their origin is often unnoticed. Its principal vulnerability is the assumption that the presenting problem and the system's boundaries can be identified without controversy; a model configured around a single defined problem has difficulty with presentations that are diffuse, multiply defined, or constituted by legal or child-protection processes rather than by the family's own account.

The Milan group and systemic epistemology

The Milan group — Mara Selvini Palazzoli, Luigi Boscolo, Gianfranco Cecchin, and Giuliana Prata — began in the 1960s treating severe eating and psychiatric disorders with a psychoanalytic orientation and reworked their practice in the 1970s into the most epistemologically sophisticated of the early models. *Paradox and Counterparadox* (1978) described work with families in which the identified

patient's symptom was analysed as part of an ongoing "family game" characterised by obscured alliances, disqualification of messages, and covert coalitions, with the ritualised prescription of the symptom as the central intervention. In the 1980s, following internal disagreement and the group's division, the model was reformulated around **hypothesising, circularity, and neutrality** (1980) and then around a positive connotation of the family's behaviour rather than paradoxical prescription.

Circular questioning is the Milan contribution of greatest durability. The technique asks each family member about the relationships and beliefs of others — "how does your mother think your father feels when your sister refuses to eat?" — thereby generating information about difference, comparison, and relationship at the same time as it introduces, through the very form of the question, the idea that relationships are mutual. The intervention is thus inseparable from the assessment: the family learns a systemic view of itself in the process of being asked systemic questions. The related device of the **reflecting team**, developed by Tom Andersen in Tromsø in the late 1980s, moved the observing team's discussion into the family's presence, so that the family could hear multiple descriptions of its situation and choose among them — a practice that operationalises second-order principles by making the observing system transparent.

> **KEY THEORETICAL POINT**
> The Milan school's most durable theoretical claim is that the instruments of assessment and the instruments of intervention are the same: a question is already a perturbation. The claim has two consequences. First, it makes it impossible to regard assessment as a neutral phase that precedes therapy, since the assessment has already changed what it purports to measure. Second, it places a burden on the therapist to ask what the question itself does — whom it recruits as an observer, whose version it elevates, and whether the difference it installs is therapeutic or destructive.

Contemporary brief and systemic practice

The strategic and MRI traditions survive in two ways. First, their techniques are embedded in most contemporary brief work: task assignment between sessions, attention to interaction around the presenting problem, explicit goal negotiation, and the deliberate use of the session's end to consolidate change. Second, their intellectual descendants include solution-focused brief therapy (Section 1.7) and the various manualised brief models used in child and adolescent services. The empirical picture is uneven. Solution-focused and brief systemic approaches have moderate evidential support for several presentations, with the strength of evidence limited by short follow-up periods, heterogeneous outcome measures, and small samples in many studies; the specific claim that brevity itself is a mechanism of change — that fewer sessions produce better outcomes through the

intensification of focus — has not been established. Doctoral readers should note the general pattern: brief approaches accumulate evidence of non-inferiority with respect to longer treatments for mild and moderate presentations, and evidence of inferiority for severe, chronic, and high-risk presentations where extended work and risk management are required.

1.4 Intergenerational and Contextual Theories

Bowen family systems theory

Murray Bowen developed his theory over three decades from clinical work with families of people diagnosed with schizophrenia and from research on family interaction at the National Institute of Mental Health. *Family Therapy in Clinical Practice* (1978) collects the principal statements. Bowen's theory is distinguished from the traditions discussed above by its ambition: it offers a general account of emotional functioning in families across generations, and it treats the family as an emotional unit whose processes are subject to natural-scientific description.

Its central construct is **differentiation of self**: the capacity to distinguish thinking from feeling, to remain in emotional contact with significant others while holding one's own positions, and to tolerate anxiety without either fusing with others or cutting off from them. Differentiation is both an individual property and a property of the relationship system: Bowen distinguished the person's *basic level* of differentiation, reasonably stable over the lifespan, from *functional* differentiation, which fluctuates with the anxiety of the surrounding system. The theory's second construct is the **triangle**, discussed below. The third is the **nuclear family emotional system**, which describes the ways in which undifferentiation expresses itself in a couple: in marital conflict, in symptoms in one partner, in the projection of anxiety onto a child, or in emotional distance. The fourth is the **family projection process**, by which parental anxiety concentrates on a particular child and produces lower differentiation in that child; the fifth is **multigenerational transmission**, whereby small differences in differentiation between children are amplified over several generations, producing a descendant with significant impairment in a family that shows no other obvious pathology. **Emotional cutoff** — the management of unresolved attachment to parents by distancing, geographical separation, or refusal of contact — is the sixth.




Figure 1.7 The emotional triangle. Anxiety in a dyad is displaced onto a third element — a person, an activity, a substance, or a symptom — which stabilises the pair. Triangles are normal features of relationship systems; they become clinically significant when they are rigid, chronic, and reproduced across generations. *Source:* prepared for this edition.

Bowen's theory also makes an explicit claim about the relationship between family process and society: the **societal emotional process** extends the analysis of anxiety, reactivity, and regression to the wider social system. The claim is important for doctoral readers because it converts a clinical theory into a social theory, and thereby exposes it to a range of evidence — historical, sociological, and cross-cultural — that clinical research cannot address. The family's functioning, on Bowen's account, is not merely influenced by society; it reproduces the same emotional process at a larger scale.




Figure 1.8 Multigenerational transmission. Bowen's account of the amplification of small differences in differentiation across generations: anxiety concentrated in one relationship in one generation produces a lower level of differentiation in the next, where the pattern intensifies, until a descendant presents with symptoms. *Source:* prepared for this edition.

Assessment and method in the intergenerational tradition

The distinctive instrument of the intergenerational tradition is the **genogram** — a multigenerational family diagram recording structure (households, births, deaths, marriages, separations), relationships (closeness, conflict, cutoff), and functioning (occupations, migration, illness, substance use, violence, and mental health). Its theoretical function is to make visible the transmission pathways the theory predicts. Its clinical function is to reframe the presenting problem within a longer temporal frame, frequently reducing a family's sense of blame and introducing the possibility of curiosity about the previous generation. Genogram notation has become standard across the field and is reproduced in Appendix A.



Figure 1.9 Constructing a genogram with a couple. The diagram is built collaboratively in the session, so that the process of construction — who remembers what, who disagrees, which facts were never spoken — becomes part of the assessment. *Illustrative image (AI-generated).*

Illustrative image (AI-generated), created for this edition for teaching purposes. It depicts no identifiable family, clinician, or agency.

Method in Bowen's own version consists of **coaching** rather than treatment in the usual sense. Clients are taught the theory's basic concepts, encouraged to observe their own reactivity, and supported in undertaking a structured re-engagement with their families of origin: gathering factual information from relatives, identifying triangles, remaining in contact while holding a position, and reducing cutoff. Sessions may be conducted with one person alone, a couple, or several people, reflecting the theory's contention that the emotional process is the unit of treatment rather than the number of people in the room. That contention generated the useful concept of **isomorphism** — the appearance of the same pattern in different parts of a system, including the therapist's consulting team — which later became a central topic in supervision theory.

Critique of Bowen's theory

Bowen's theory attracts three lines of criticism. The first is empirical: differentiation of self is measured principally by self-report instruments developed within the tradition, notably the Differentiation of Self Inventory (Skowron and Friedlander), and the resulting research is largely correlational, prone to shared-method variance, and modest in effect size. Studies reporting associations between lower differentiation and anxiety, relationship distress, and symptom severity are numerous, but the causal claims they are used to support — that raising differentiation produces symptom relief — rest chiefly on clinical observation and single-

case reports rather than controlled trials. The second criticism is theoretical: the biological grounding Bowen claimed for the theory has not been substantiated, and the theory's core processes are difficult to falsify because their operation is inferred from the very symptoms they are invoked to explain. The third is normative: the theory's emphasis on thinking over feeling, and on a particular style of calm self-assertion, has been argued to privilege cultural and gendered modes of relating that are not universal, and to underestimate structural constraints — poverty, discrimination, forced migration, violence — that are not reducible to differentiation.

These criticisms have a constructive as well as a deflationary use. The multigenerational frame is one of the field's genuine contributions to clinical understanding: it disciplines the tendency to treat the present family as self-contained, and it generates formulations that competing models cannot easily produce, particularly in work with cut-off adults, with families marked by migration and separation, and with recurrent patterns of loss, addiction, and violence. The doctoral task is to specify which parts of the theory survive as clinical heuristics, which parts are empirically testable, and which parts belong to the history of ideas rather than to present-day science.

Contextual therapy and the ethics of relationship

Ivan Boszormenyi-Nagy developed contextual therapy in explicit opposition to what he regarded as the field's neglect of ethical accounting. His *Foundations of Contextual Therapy* (1987) and related writings argue that relationships are governed not only by emotional dynamics but by a ledger of fairness: between parent and child, between partners, and across generations, people keep track of what is given, withheld, exploited, and repaid. The theory's technical vocabulary includes **invisible loyalties** (unconscious commitments to the family of origin that constrain present choices), **entitlement** (the legitimate claim to have received and to receive fair care, and the destructive entitlement that develops when it is not), **legacy** (the obligation to pass on to the next generation what one has received), and **trustworthy giving**. Its therapeutic method is **multidirected partiality**: the therapist actively grants standing to each participant in turn, including absent and deceased members, without abandoning responsibility for the consequences of harmful conduct.

Contextual therapy has been comparatively neglected in the empirical literature, to which its concepts are difficult to operationalise, but its influence on practice has been considerable. Elements of multidirected partiality are recognisable in contemporary work with families affected by parental conflict, in reunification practice, and in approaches to intergenerational trauma. Its most important contribution to theory may be the claim that a purely descriptive or functional account of family process is inadequate to clinical reality, because

families organise themselves around questions of fairness and reliability — that is, around normative questions — and that therapy that treats those questions as mere content misses what the family is arguing about.

::: casebox {title="Case Illustration 1.2 — Transmission, loyalty, and the ledger of fairness (hypothetical)"} **The referral.** A woman of thirty-eight seeks help for chronic anxiety and for a pattern of intense conflict with her mother, which has intensified since the birth of her first child eight months ago. She describes her mother as controlling and intrusive and has considered ending contact. She reports no history of violence within the current household.

Intergenerational formulation. A genogram completed over two sessions shows a maternal line with three generations of separation: the client's maternal grandmother emigrated alone and left her eldest daughter (the client's mother) in the care of relatives for six years; the client's mother, in turn, worked away from home for extended periods during the client's early adolescence, and the client cared for two younger siblings. Emotionally, the family has managed separation by cutoff, each generation distancing rather than repairing. The current conflict is not merely an adult disagreement about infant feeding; it is the reactivation of an unpaid account. The client's mother is seeking to be the trusted caregiver she was never permitted to be, and the client is reproducing the position she occupied at thirteen — responsible for a child she did not bear.

Contextual elements. Multidirected partiality is essential: the grandmother's loss, the mother's deprivation, and the client's neglected adolescence all have standing, and the therapist's task is to facilitate acknowledgement between them rather than to arbitrate. Interventions include the reconstruction of family history with both women present, explicit acknowledgement of what each has given, and the negotiation of a specific arrangement for the mother's involvement with the grandchild that the client can trust.

Methods note. This case is hypothetical. It illustrates how intergenerational and contextual formulations produce a developmental and an ethical account of the same clinical presentation, and how those accounts generate interventions — history-taking in the presence of both parties, acknowledgement work, negotiated agreements — that differ sharply from the here-and-now interventions of strategic or structural practice. :::

1.5 Experiential, Communication, and Humanistic Theories

Communication theory and the pragmatics of human communication

Pragmatics of Human Communication (Watzlawick, Beavin, and Jackson, 1967) is the most systematically constructed statement of communication theory in the field, and it supplied a set of propositions — the "axioms" — that were widely

taught as the anatomy of family interaction. Five are conventionally listed and should be understood precisely, since they are frequently cited in loose form.

The first axiom states that one cannot not communicate: in a social field, every act and every abstention carries message value, so silence, absence, and immobility communicate as surely as speech. The second distinguishes the **report** and **command** aspects of any message: every communication conveys information and, at the same time, proposes a relationship, and it is in the command aspect that most relational difficulties arise. The third concerns **punctuation**: interactants punctuate the sequence of exchanges, one treating her behaviour as a response and the other treating his as the response, so that each holds a linear reading of what is in fact circular, and each holds the other responsible for the pattern. The fourth contrasts **digital** and **analogic** communication — the former discrete, conventional, and suited to content, the latter continuous, iconic, and suited to relationship — with the prediction that difficulties arise at the junction between them, where the analogic must be translated into the digital and meanings are lost or contested. The fifth distinguishes **symmetrical** from **complementary** interaction: relationships in which participants exchange equivalent behaviours are symmetrical, while those structured by a difference of position (parent and child, teacher and student, one partner's pursuit and the other's withdrawal) are complementary, and each form carries its characteristic pathologies — competitive escalation in symmetrical systems, and rigid, unchallenged inequality in complementary ones.

These propositions have proved productive and problematic in equal measure. Their productivity is evident in clinical practice: the concept of punctuation organises an enormous amount of couple work; the report–command distinction underpins the practice of listening for the relational message beneath content; the symmetrical–complementary distinction became the basis of the influential typology of couple relationships developed by Alan Sillars and others. Their problems are of two kinds. Empirically, the axioms are framed as logical necessities of communication rather than as contingent generalisations, so that they are difficult to test and were rarely tested. Ethically, the fifth axiom's account of complementary relationships treats inequality as a structural form rather than as a distribution of power to be evaluated — an omission that the feminist critics of family therapy (Section 1.7) identified as central, since it allows the subordination of one partner to be described in neutral terms as complementary interaction.

Virginia Satir and the humanistic tradition

Virginia Satir's *Conjoint Family Therapy* (1964) and *Peoplemaking* (1972) made family therapy intelligible to a wide public and established the humanistic strand of the field. Satir treated family problems as deficits in self-esteem and in congruence of communication, and she gave clinical prominence to the stances people

adopt under stress: **placating** (appeasing others at the cost of self), **blaming** (asserting self at the cost of others), **computing** (retreating into reasonableness at the cost of both), **distracting** (deflecting attention altogether), and **levelling** (congruent communication in which feeling, thought, and action align). Her method emphasised warmth, touch, structured communication exercises, **sculpting** (the arrangement of family members in physical space to represent relationships), **parts parties** (the dramatised conversation among internal parts), and family reconstruction (a guided re-enactment of the parents' family histories).

Satir's theoretical writing is less formal than Bowen's or Minuchin's, and her constructs have been studied less than they deserve; the empirical literature includes some support for communication-training components and for the association between congruence and relationship satisfaction. Her durable contributions are two. She insisted that affect — warmth, longing, hurt, hope — is not incidental to systemic process but constitutive of it, and she insisted that the therapist's own humanity is an instrument of change rather than a contaminant of technique. In that respect her work anticipates the experiential and emotion-focused traditions examined in Section 1.6.

Symbolic-experiential therapy

Carl Whitaker's symbolic-experiential therapy took the humanistic premise further. Whitaker treated therapy as an encounter between two systems — the family and the therapeutic team — and he privileged immediacy, spontaneity, and the absurd over technique. His use of **co-therapy**, often with a colleague of the opposite sex, was theoretically motivated: the co-therapy couple models a working relationship, provides each other with the stimulus of a second perspective, and allows the therapeutic system itself to become an object of observation. *The Family Crucible* (Napier and Whitaker, 1978) remains the field's most widely read exposition of process in family therapy, and it dramatises a claim that no manual can contain: that the therapist's presence, including the therapist's own emotional life, enters the system and must be accounted for.

Symbolic-experiential concepts — the **battle for structure** (the therapist's contest of initiative with the family's homeostatic pull), the **battle for initiative**, and the distinction between the family's fantasy of what the therapist will do and what is required — remain useful for describing the early phase of therapy when a family's organising anxiety is at its highest. The tradition's evidentiary base is thin, and its practices depend heavily on the personal qualities of the therapist; both facts are acknowledged by its own exponents, who have generally preferred to defend the approach as an art on the grounds that manualising it would destroy what makes it work. That defence is itself worth examining in a doctoral seminar: it makes a claim about the nature of therapeutic knowledge that empirical research can, at least in part, test.

Internal family systems and the multiplicity turn

Richard Schwartz's internal family systems (IFS) model (1995 and subsequent revisions) approaches the family from the opposite direction: it takes the language of systems and applies it to the mind, describing the person as composed of **parts** — protective managers, vulnerable exiles, and a core **Self** characterised by curiosity, compassion, and calm — organised into an internal system with the same logic of coalitions, conflicts, and burdens that family theorists describe between persons. Its clinical method proceeds by helping clients access the Self, negotiate with protective parts, and unburden exiled material. IFS has become widely used for trauma and is increasingly applied in couple therapy, where it offers a vocabulary for the reciprocation of parts between partners that maps closely onto the negative interaction cycles of emotion-focused therapy. Its evidence base consists of growing but still limited randomised trial data, with most published work on individual trauma presentations; claims for the model in couple and family work are, at present, more clinical than empirical.

1.6 Attachment, Emotion, and Neurobiological Theories

Attachment theory and its migration into couple work

John Bowlby's attachment theory proposed that the human infant is born with a species-typical need for a protective, responsive caregiver, that the quality of early care is internalised as an **internal working model** of self and other, and that disruptions of attachment produce characteristic patterns of distress and defence that persist into adult life. Mary Ainsworth's observational procedure — the Strange Situation — supplied the empirical method, identifying the organised patterns now described as secure, insecure-avoidant, and insecure-ambivalent (resistantly attached), and later the disorganised pattern described by Mary Main and Judith Solomon.

The migration of attachment theory into adult relationship research began with the argument by Cindy Hazan and Phillip Shaver (1987) that romantic love can be understood as an attachment process, and it was consolidated by two developments. The first was dimensional measurement. Kim Bartholomew and Leonard Horowitz distinguished four adult attachment categories — secure, dismissing, preoccupied, and fearful — along the orthogonal dimensions of model of self and model of other; Brennan, Clark, and Shaver's factor-analytic work on a large item pool produced the two-factor structure — **attachment anxiety** and **attachment avoidance** — that underlies the self-report instruments now used throughout the field, most prominently the Experiences in Close Relationships inventory. The second development was the theoretical extension by Mario Mikulincer and Phillip Shaver, who proposed that attachment-system dynamics

regulate distress across the lifespan and that secure attachment functions as a resource for affect regulation, exploration, and caregiving.

Two clarifications matter for doctoral work. First, attachment patterns in adults are not simply the childhood patterns grown up. Discontinuity is common, and the distribution of attachment classifications in adulthood is not a straightforward continuation of the infant distribution; longitudinal studies find both stability and substantial change, with change associated with intervening relationship experiences, loss, therapy, and social support. Second, self-report attachment measures and interview-based measures (notably the Adult Attachment Interview and its derivatives) do not measure the same construct reliably; correlations between them are modest. Research that treats attachment as a single variable measurable by any available instrument will produce findings that cannot be interpreted with confidence.

Emotionally focused therapy

Emotionally focused therapy (EFT) is the most theoretically integrated and empirically supported synthesis of attachment theory with systemic practice. Developed by Leslie Greenberg and Sue Johnson in the 1980s, and elaborated principally by Johnson, the model treats relationship distress as the product of a self-reinforcing **negative interaction cycle** driven by unacknowledged attachment panic: one partner, feeling that the bond is threatened, protests, criticises, and pursues; the other, overwhelmed and inadequate, withdraws and defends; each partner's behaviour confirms the other's worst expectation, and the cycle becomes the couple's enemy rather than either partner.




Figure 1.10 The negative interaction cycle in emotionally focused therapy. The cycle is presented to the couple as a shared pattern rather than as the responsibility of either partner, which is the precondition for de-escalation and for the attachment-level work that follows. *Source:* prepared for this edition.

The therapeutic process proceeds through three stages that are subdivided into nine steps. In **de-escalation**, the therapist tracks the cycle with the couple, accesses and deepens the underlying emotions, and reframes the problem as the cycle, so that both partners can recognise their positions as protective rather than malicious. In **restructuring**, each partner is helped to access the primary attachment need and to articulate it in the other's presence, and the other is helped to respond; incidents of **attachment injury** — betrayal, abandonment in a moment of need, infidelity — are processed in the same terms. In **consolidation**, new patterns are rehearsed, integrated into the couple's narrative, and prepared for maintenance.

The evidence base is the strongest of any couple therapy model. Multiple randomised trials report significant reductions in relationship distress and improvements in satisfaction, including trials with distressed couples, couples with a depressed partner, parents of chronically ill children, and couples in which one partner has experienced trauma, together with process research showing that in-session emotional deepening predicts outcome. Meta-analyses report large within-group effects and moderate between-group advantages over the control conditions used, with the caveat that control conditions have often been non-specific, that follow-up periods are relatively short, and that several trials were

conducted by the model's own developers or trainers. The methodological point is not to dismiss the evidence but to read it as a mature literature requires: effect sizes shrink when the investigators are independent, and the field still lacks large independent multi-site trials.

EFT has been extended to families (emotionally focused family therapy), and the underlying attachment framework has been applied to individual work with adults. The extension of an attachment-based model to whole families raises a theoretical question that remains open: if the couple's cycle is the unit of change in dyadic work, what is the corresponding unit in a family with several attachment relationships, and how is the therapist to hold multiple cycles in a single treatment?

Observational research on couple interaction

The most methodologically ambitious research programme on couple interaction is that of John Gottman and colleagues, who from the 1970s developed laboratory observation of couples' conflict discussions combined with physiological monitoring and mathematical modelling of the interaction. The programme's theoretical contributions include the identification of behavioural sequences that predict deterioration — patterns of escalating criticism, contemptuous expression, defensive response, and withdrawal (popularised as the "four horsemen"), low ratios of positive to negative interaction during conflict, and physiological arousal that interferes with empathic listening — and the modelling of relationship change in terms of attractors in an affective state space, developed with Robert Levenson as the **dynamic interactional model**.

Two features of this literature deserve the doctoral reader's attention. The first is the methodological contribution: by measuring behaviour rather than relying on report, the Gottman laboratory made it possible to test theories of couple interaction directly, and it demonstrated the extent to which self-reports of communication are reconstructions rather than records. Second, the predictive claims require care. The frequently cited figures for accuracy in predicting divorce from observational data derive from specific samples and prediction models, and their generalisability has been criticised on the grounds that the models incorporate base-rate and sample-specific information, so that the headline accuracy figures overstate what the models would achieve in new samples. Subsequent work by many independent groups has confirmed the broad finding that negative affect reciprocity and withdrawal during conflict predict deterioration, while the specific numerical claims are best treated as model-dependent.

The practical implication for the theory of change is significant: if destructive patterns are observable and measurable, then therapy can be evaluated on whether it changes them. That shift — from evaluating therapy by whether couples feel

better to evaluating it by whether measured interaction changes — is one of the field's most important methodological advances, and it has been extended to family interaction in the context of adolescent and child problems.

Affect regulation, psychophysiology, and the neurobiology of relationships

The last two decades have seen a substantial expansion of neuroscientific and psychophysiological research on close relationships, and the field's theoretical literature has absorbed its findings eagerly — sometimes more eagerly than the evidence warrants. Four lines of work are central.

The first concerns **co-regulation**. Tiffany Field's early studies of mother–infant interaction and Edward Tronick's still-face paradigm established that infants regulate affect through interaction, and that disruption of the interaction produces disorganisation within seconds. Tronick's mutual regulation model proposes that dyadic systems are continuously repaired rather than stably attuned, and that it is repair, not constant harmony, that builds regulatory capacity. This proposition is directly relevant to couple therapy, where it reframes the goal from the elimination of rupture to the improvement of repair.

The second concerns **social regulation of physiological threat response**. James Coan and colleagues demonstrated that holding a partner's hand attenuates threat-related neural responses in the hypothalamus and anterior cingulate during the anticipation of electric shock, with the magnitude of attenuation associated with marital quality; Coan's social baseline theory interprets such findings as evidence that the human brain expects access to close others as a regulatory resource. The third concerns hormonal and autonomic mechanisms, including oxytocin research. This literature is more fragile than its prominence suggests: early findings of oxytocin's effects on trust and cooperation involved small samples and specific procedures, and several have failed to replicate in larger studies, while the construct of **vagal tone** as a marker of social engagement remains empirically unsettled, and the polyvagal framework in which it has been popularised is not generally regarded as established in physiology.

The fourth line is the popular literature on "interpersonal neurobiology", which has been enormously influential in clinical training while resting chiefly on extrapolation from basic neuroscience rather than on clinical trials. The relevant warning for doctoral readers is the documented "seductive allure" effect: explanations that include neuroscientific content are rated more convincing than the same explanations without it, even when the neuroscience is irrelevant to the logic of the argument. In this field the warning has teeth, because claims about the brain are frequently used to justify clinical methods that have not been tested by clinical research. The defensible position is that neurobiological findings may

suggest mechanisms and generate hypotheses, but that they do not license treatment claims, which require clinical evidence.

> **PRACTICE NOTE**
>
> **Reading a neurobiological claim.** When a text asserts that an intervention "rewires the brain" or "releases oxytocin", ask four questions. First, what is the original study, and in what population and design was it conducted? Second, has the finding replicated in independent samples? Third, is the measure — neural, hormonal, autonomic, or behavioural — a valid proxy for the psychological construct being claimed? Fourth, does the argument move from correlation to causation, or from a laboratory effect to a clinical outcome, without warrant? Claims that survive these four questions are worth building on; claims that do not should be replaced by the behavioural or relational evidence that the intervention itself must generate.

1.7 Social Constructionist and Narrative Theories

The constructionist turn

By the mid-1980s the field's epistemology had shifted again, from the systems metaphors of cybernetics to the linguistic and interpretive resources of social constructionism. The shift was signalled by Kenneth Gergen's argument that psychological descriptions are not discoveries about a stable inner world but products of social interchange that acquire authority within particular communities. Applied to therapy, the argument implies that "the problem" is not an entity to be mapped but a description produced and maintained in language by particular people for particular purposes. The clinical corollaries were developed by Harlene Anderson and Harry Goolishian, whose **collaborative language systems** approach proposed that therapy is a conversation in which new meanings are generated, that the appropriate stance is **not-knowing** — a disciplined ignorance in which the therapist's understanding remains always open to revision — and that the client system is not the family as such but the **problem-determined system**: the set of people who are engaged in talking about the problem, which may include professionals, relatives, and neighbours, and which dissolves when the problem ceases to be talked about.

Narrative therapy

The most influential constructionist practice is narrative therapy, developed by Michael White and David Epston and presented in *Narrative Means to Therapeutic Ends* (1990) and elaborated by White's *Maps of Narrative Practice* (2007). Its central premise is that people make sense of their lives by telling stories, that problems arise when the stories available to a person or family are impoverished or dominated by a single negative account, and that therapy is the negotiated re-authoring of those stories. Two techniques are particularly important.

Externalising conversation separates the problem from the person: rather than asking a child to describe what is wrong with her, the therapist asks about the influence of "the worry" or "the anger" on her and on the family, and about the influence she has on it. The technique is more than linguistic politeness. It is designed to reduce the shame and blame that accompany identity-level descriptions, to open the way for collective action against the problem, and to make visible exceptions — occasions when the problem's influence was diminished — which provide material for a preferred story.

Re-authoring develops those exceptions into an alternative account: **unique outcomes** (moments that contradict the dominant story), the identification of the values and commitments that explain them, the thickening of the new account with detail and witnesses, and the performance of the new identity in the presence of others. White drew on anthropological work for the practice of **definitional ceremonies** — occasions on which an outsider-witness recounts how the person's account resonated with their own experience — and on **re-membering**, in which the role in one's identity of living and dead figures is deliberately revised rather than passively inherited.

Two critiques are standard and should be examined together. The first concerns evidence: narrative practice has a modest empirical literature, consisting mainly of case studies, qualitative research, and a small number of trials in specific applications (for example, work with children and adolescents and with families affected by chronic illness). This does not render the approach ineffective, but it means that claims for it must be made with corresponding caution. The second concern is the limits of conversation. Narrative practice developed largely in outpatient work with families presenting relationship difficulties, and its instruments are least suited to presentations involving risk — severe self-harm, violence, acute psychosis, child protection proceedings — where assessment, monitoring, and the management of legal duties cannot be replaced by re-authoring, and where an approach that externalises problems may be experienced by family members as minimising danger.

Solution-focused brief therapy

Solution-focused brief therapy, developed by Steve de Shazer and Insoo Kim Berg at the Brief Family Therapy Center in Milwaukee, applies the constructionist premise with unusual economy. It proposes that change is better served by elaborating solutions than by diagnosing problems, and it relies on a small set of conversational instruments: the **miracle question** (an invitation to describe in detail a future in which the problem no longer exists), **scaling questions** (rating progress on a defined scale to identify concrete next steps), the identification of **exceptions** (times when the problem did not occur), and the negotiation of small, observable goals. Later developments include the practice of beginning the first

session without an assessment of the problem's history, the deliberate elicitation of the client's resources, and the use of compliments as a form of feedback.

The literature on solution-focused brief therapy is larger than for narrative practice, with randomised trials in school, mental health, and substance-use settings and several meta-analyses. The overall picture is of small to moderate effects on a range of outcomes, with substantial heterogeneity, and with the recurring methodological problem of the control conditions: when the comparison is waiting list or treatment as usual, the situation is favourable to any structured intervention, and several trials measure outcomes on instruments developed within the approach itself. The theoretical questions worth pursuing in doctoral work concern specificity — whether the technique adds anything to the relationship factors examined in Section 1.8 — and durability, since short treatments are particularly exposed to the risk that measured gains decay without follow-up support.

Power, gender, and culture: the critical traditions

The constructionist turn did not by itself address power, and the field's most significant internal criticism came from feminist writers. Rachel Hare-Mustin's *A Feminist Approach to Family Therapy* (1978) charged the field with encoding traditional gender roles as structural health — the mother as the family's emotional manager, the father as the peripheral figure whose withdrawal was treated as normal — and with systematic double standards in the description of male and female behaviour. Later work developed the critique into a programme: the invisibility of women's relational and material labour, the way in which neutrality and circularity could be used to distribute responsibility for male violence across the whole family, and the frequency with which mothers were identified as the problem in child-focused referrals. The influential collection *The Invisible Web* (Walters, Carter, Papp, and Silverstein, 1988) and, from a psychoanalytic-systemic position, the work of Virginia Goldner on abusive relationships gave the field a vocabulary for describing couples in which conflict is not symmetrical and where the language of "complementary" interaction obscures coercion.

The cultural critique developed in parallel. The argument that family systems theory had been constructed on the experience of white, middle-class, Western nuclear households was pressed by a series of foundational texts: the comparative work of Monica McGoldrick, John Pearce, and Joseph Giordano on ethnicity and family therapy (first edition, 1982), Nancy Boyd-Franklin's analyses of therapy with black families across class and structure, and Celia Falicov's account of migration, acculturation, and the transition between cultural contexts as a central dynamic of "cultural borderlands". The **cultural genogram**, introduced by Kenneth Hardy and Tracey Laszloffy (1995), adapted the intergenerational instrument to the analysis of cultural identity, pride, shame, and the internalisation

of dominant-group evaluations. In the two decades since, decolonial and critical-race critiques have enlarged the field's obligations further, questioning the export of Western therapeutic models, the universalism of concepts such as differentiation and boundary, and the global asymmetry in which the discipline's theory is produced in a small number of high-income countries and applied everywhere else.

The theoretical consequences of these critiques are not confined to ethics. If power is constitutive of family relationships, then a theory that describes systems in terms of reciprocity and feedback is incomplete: it lacks a mechanism for asymmetric influence that does not depend on each person's contribution. If the family's structure is shaped by culture, migration, religion as a social identity, class, and legal status, then a universal developmental timetable — the family life cycle as a single sequence of stages — misrepresents the lives of families whose trajectory differs. If the therapist is inside the system, then the therapist's own cultural position, its privileges and blind spots, are clinical variables. Each of these propositions is now uncontroversial in the field's literature, and each remains incompletely implemented in its theories, its measures, and its research designs.

1.8 Cognitive-Behavioural, Integrative, and Evidence-Based Synthesis

Behavioural and cognitive-behavioural couple therapy

The behavioural tradition entered couple therapy through the social-exchange analysis of distressed relationships. Its founding proposition is that relationship satisfaction is a function of the ratio of rewarding to punishing exchanges, and that distressed couples are characterised by low rates of positive behaviour, high rates of negative behaviour, and reciprocal escalation. Traditional behavioural couple therapy (TBCT), systematised in the work of Robert Weiss, Gerald Patterson, and later Neil Jacobson and Gayla Margolin, therefore combined two components: **behaviour exchange**, in which partners negotiate specific changes in daily behaviour, and **communication and problem-solving training**, in which they are taught to express feelings in structured form (typically the speaker–listener technique), to separate issues from persons, and to negotiate solutions. The approach is notable for being the first in the field to be manualised and tested in controlled trials, and for evaluating outcome with observable measures as well as self-report.

Cognitive-behavioural couple therapy (CBCT), developed principally by Norman Epstein and Donald Baucom, added a cognitive level of analysis. Its central claim is that distress is mediated by partners' interpretations of each other's behaviour — selective attention to negative events, attribution of hostile intent, unrealistic standards and assumptions about relationships, and discrepant

expectations — and that changing those cognitions produces change in affect and behaviour. The therapy therefore includes identification and modification of attributions and standards, and the development of shared meaning. The model's manualised form (Epstein and Baucom, 2002) has been tested in randomised trials with effects comparable to behavioural approaches, and its analysis of attributional processes has generated a substantial research literature, including work on couples recovering from infidelity.

Integrative behavioural couple therapy (IBCT), developed by Andrew Christensen and Neil Jacobson, arose from two problems with the earlier models: their outcomes decayed over time, and their emphasis on negotiation and change encountered couples whose differences proved intractable. IBCT retains the behavioural framework but adds **acceptance** as a therapeutic goal, achieved through an emphasis on **empathic joining** (mutual understanding of each partner's soft emotions and of the differences between them), **unified detachment** (treating the couple's pattern as a shared problem to be observed rather than a dispute to be won), and **tolerance building** (deliberately generating understanding of differences without demanding change). Change-oriented techniques remain available and are targeted at behaviours amenable to compromise. The largest trial in the field compared IBCT with TBCT over two years: both produced comparable improvements, with IBCT showing a slower but more durable trajectory in the maintenance of gains.

The theoretical significance of IBCT is broader than the model itself. It demonstrated that acceptance and change are not mutually exclusive, contradicting the field's earlier tendency to treat them as opposed; and its findings on the durability of effects undercut the assumption, common to the brief-therapy traditions, that faster change is necessarily better change.

Behavioural parenting and psychoeducational models

The behavioural tradition also produced the most extensively evaluated family interventions for child and adolescent difficulties: parent management training, the Incredible Years programme, parent–child interaction therapy, and their derivatives. These models share a common theory of change: child behaviour problems are maintained by patterns of parental attention, reinforcement, and ineffective discipline, and change is produced by training parents in specific skills — attending, praising, giving clear instructions, using contingent consequences — and by modifying parental cognitions such as catastrophising and hostile attribution. The adaptations for high-risk families (the Oregon model of parent management training developed by Gerald Patterson and colleagues, and its dissemination as Parent Management Training – Oregon) address additional predictors including parental antisocial behaviour, maternal depression, and coercive family process.

These approaches stand apart from most of the field's traditions in two respects: their theory of change is specified at the level of observable behaviour, and their evidence base consists of large numbers of randomised trials and meta-analyses demonstrating effects on child behaviour that are maintained at follow-up, with effects on antisocial behaviour reported in long-term follow-up studies. They are also the models whose implementation has been most closely examined: the literature on transportability, therapist fidelity, and the "fidelity–adaptation" dilemma — whether evidence-based programmes retain their effects when adapted to local culture and resources — is now substantial, and the balance of evidence favours a degree of structured adaptation with fidelity to core mechanisms rather than either rigid replication or unconstrained localisation.

Multi-systemic and multi-dimensional models

Two further developments belong to this section because they constitute integrations in their own right. **Multisystemic therapy** (MST), developed by Scott Henggeler and colleagues for serious adolescent antisocial behaviour, treats the family as the primary agent of change but intervenes across the systems in which the adolescent is embedded — family, peers, school, neighbourhood, and juvenile justice — using techniques drawn from structural, strategic, cognitive-behavioural, and behavioural family therapy traditions according to the requirements of the case. **Multidimensional family therapy** (MDFT), developed by Howard Liddle, similarly combines individual and family work with attention to the adolescent's developmental context. Both models have substantial trial literatures with mixed results across replications: MST's effects in the original trials were large, while some independent replications found smaller or no effects, a pattern that has generated a vigorous methodological debate about the conditions under which complex family interventions transport across systems, populations, and fidelity regimes.

Integrative frameworks

The integrative movement addresses a problem that the field's early history created: a proliferation of models that disagree about mechanisms, methods, and even about what counts as the problem. Four integrating strategies can be distinguished.

Metaframeworks. Douglas Breunlin, Richard Schwartz, and Betty Mac Kune-Karrer proposed organising the field's theories into a small set of domains and levels — the organisation of the family (structure), sequences of interaction, development, culture, gender, and internal processes — on the argument that competing theories are describing different aspects of the same phenomenon rather than competing accounts of it. Integrative problem-centred therapy (Pinsof, 1995) and its successor **integrative systemic therapy** (Pinsof, Breunlin,

Russell, and Lebow, 2018) developed this into a full clinical method: the problem defines the treatment; assessment proceeds through a series of lenses; the therapist draws on the traditions best suited to the formulation at hand; and the therapy is planned, monitored, and revised in explicit stages. The metaframework approach is the framework most widely used in doctoral training, since it converts the task of learning many theories into the task of learning when each theory applies.




Figure 1.11 The integrative problem-solving sequence. Integration is disciplined by a specified sequence: define the system and the problem, formulate across levels, select mechanisms of change with the best available warrant, sequence the treatment, and monitor and revise in the light of outcome data. *Source:* prepared for this edition.

Common factors. The second strategy argues that the specific techniques of the field's models account for less of the variance in outcome than the factors they share: the alliance, the client's expectations and resources, the therapist's qualities, and the process of remoralisation. Scott Miller, Barry Duncan, and Mark Hubble pressed this argument in *Escape from Babel* (1997); Douglas Sprenkle, Adrian Blow, and later Sprenkle, Davis, and Lebow developed it specifically for couple and family therapy, identifying conceptual and practical common factors that include relational reframing, enlarging the direct treatment system, expanding the therapeutic alliance to multiple participants, and altering the family's interactional sequences. The common-factors position functions as an empirical hypothesis as well as an argument: it predicts that structured differences between models matter less than alliance quality within models, and it predicts that monitoring outcome will improve results whether or not a particular model is used. Support for the second prediction is stronger than for the first — the evidence that routine outcome monitoring and feedback improves outcomes, particularly for clients who are not progressing, is now reasonably robust across settings.

Empirically supported principles. A third strategy abandons the model as the unit of evaluation and asks instead which principles of change are supported by evidence across models. On this account, what is transferable is not a brand of therapy but a set of mechanisms — modifying coercive sequences, reducing expressed emotion and hostility, improving communication and problem-solving skills, changing attributions, strengthening attachment security, addressing motivational ambivalence — and the therapist's task is to apply the relevant principles rather than to implement a manual faithfully.

Assimilative integration. The fourth strategy accepts that therapists inevitably work within a principal orientation while incorporating techniques from others, and seeks to do so deliberately: the borrowed method is absorbed into the host theory rather than appended unreflectively.

	Core construct	Mechanism of change	Evidence base
Structural	Boundaries; hierarchy; alliances	Restructuring; enactment	Moderate–strong for adolescent and psychosomatic presentations
Strategic / MRI brief	Interactional sequence; function of symptom	Directives; reframing; paradox	Moderate; paradox carries greater risk
Milan systemic	Beliefs; circularity; family game	Circular questioning; neutrality; positive connotation	Limited; small trials and case series
Intergenerational (Bowen)	Differentiation; triangles; cut-off	Genogram; detriangulation; coaching	Moderate; largely correlational research
Contextual (Boszormenyi-Nagy)	Legacy; fairness; trustworthiness	Multidirected partiality; exoneration	Emerging; qualitative and small-scale studies
Experiential (Satir; Whitaker)	Self-worth; communication stances	Sculpting; immediacy; co-therapy	Limited; historically influential
Emotionally focused (Johnson)	Attachment; negative cycle	Emotion reprocessing; cycle de-escalation	Strong: multiple RCTs and meta-analytic support
Cognitive-behavioural (Epstein; Baucom)	Cognitions; behaviour exchange	Communication training; cognitive restructuring	Strong: extensive RCT literature

Figure 1.12 Comparative summary of the major theoretical traditions. The grid organises each tradition by its core construct, its mechanism of change, and the state of its evidence base, and is intended for use in seminar work as a working analytical instrument. *Source:* prepared for this edition.

Table: A comparative grid of the major traditions in family systems theory.

COMPARATIVE PERSPECTIVE

What the traditions agree and disagree about. Nearly every tradition in this chapter agrees on four propositions: that behaviour is patterned; that patterns are self-maintaining; that intervention should focus on present interaction as well as on history; and that the family's meaning system matters. They disagree on four others: *where* change should be pursued (structure, sequence, emotion, cognition, story, generational account); *how* the therapist should position herself (expert, consultant, participant, witness, coach); *how many people* should be in the room; and *what* should be treated as the index of success (symptom relief, relational satisfaction, observed interaction, narrative revision, or the client's own evaluation). For a doctoral candidate, the ability to state these agreements and disagreements precisely — rather than to describe the field as a set of interchangeable techniques — is the mark of theoretical competence.

Evaluating theory: criteria and the evidence base

The traditions discussed in this chapter can be assessed against a common set of criteria. The first is **empirical adequacy**: does the theory's account of maintenance and change survive measurement? On this criterion, the outcome literature is favourable to structural, behavioural, cognitive-behavioural, emotion-focused, multisystemic, and parenting models; weaker for strategic, Milan systemic, and symbolic-experiential approaches; and thin for narrative, contextual, and internal family systems approaches, particularly in couple and family applications. Systematic reviews and meta-analyses of couple and family therapy consistently report moderate-to-large effects on relationship distress and on a range of child and adult problems, but with the familiar limitations of heterogeneity, small samples, short follow-up, and investigator allegiance effects.

![Comparative grid of theories — see previous figure.] The second criterion is **specificity**: does the theory predict something that a rival theory does not? Much of the field's evidence supports family therapy as a class of treatment while failing to distinguish between its brands. The most informative studies for theory are therefore the comparative trials, the dismantling studies, the mediation analyses that test whether theorised mechanisms actually carry the effect, and the process research that links in-session events to outcome. There are far fewer of these than the field's theoretical confidence would suggest.

The third criterion is **falsifiability**, and it is here that systemic theory has historically been most vulnerable. Circular formulations, since they can account for any pattern after the fact, must be stated in a form that specifies in advance what would disconfirm them. The fourth is **scope**, on which the differences between the traditions are wide: Bowen's theory aspires to a general account of emotional functioning across generations and cultures; the behavioural models deliberately restrict themselves to observable interaction in defined problem areas. The fifth

is **clinical utility** — whether the theory helps a practitioner do something effective — which is distinct from truth and which, confusingly, may be enjoyed by theories that are false. The sixth is **ethical and cultural adequacy**: whether the theory's descriptions distribute responsibility fairly, whether it treats violence and abuse with the seriousness they require, whether it assumes a family form that is culturally or socially specific, and whether it can be used in ways that respect the dignity and agency of those it describes.

On the last criterion, the field's history supplies cautionary examples. The circular-causality axiom, applied without care, can convert an account of a violent pattern into a description in which the victim's responses are treated as half of the cycle. The concept of complementarity can describe a relationship of domination as a structural form. The language of structural health can pathologise families whose arrangements differ from the nuclear norm. A graduate of a doctoral programme in marriage and family therapy should be able to use the field's theories while refusing these uses — and should be able to explain, on theoretical as well as ethical grounds, why the refusal is correct.

::: casebox {title="Case Illustration 1.3 — Competing formulations of a single presentation (hypothetical)"} **The referral.** A couple, married eleven years with two children aged nine and six, present with escalating arguments about the division of household labour, the wife's resentment at the husband's long working hours, and her reported insomnia and low mood. The husband reports feeling criticised and "never able to win". They attended four sessions with a previous therapist and discontinued, each reporting privately that the therapist "took the other side".

Formulation 1 — structural. The hierarchy between the spousal and sibling subsystems has blurred: the eldest child has become an ally of the mother and a critic of the father, and family decisions are made around the children's preferences. The couple's arguments occur in the children's hearing and serve the function of maintaining an intimate connection that negotiation would require them to secure directly. Interventions: strengthen the spousal subsystem; establish an explicit boundary around parental conflict; require the parental coalition to resolve disagreements outside the children's earshot.

Formulation 2 — emotion-focused. A pursue–withdraw cycle is in operation. The wife's protest — criticism about tasks and absence — is a manifestation of attachment distress: she is asking whether she matters. The husband's withdrawal is a defence against a sense of inadequacy and the fear that any response will be used against him. Both partners' secondary and primary emotions are unexplored. Interventions: track and reflect the cycle; access underlying fear and longing; reframe the problem as the cycle; then restructure attachment-level responsiveness.

Formulation 3 — cognitive-behavioural. Attributional and standard-level cognitions maintain distress. The wife attributes the husband's overtime to a preference for work over family and holds a standard of partnership that equates love with equal task distribution; the husband attributes criticism to a persistent negative evaluation of him and holds assumptions about the meaning of his role as provider. Behavioural exchange is minimal and appears grudging to both. Interventions: modify attributions and increase accurate empathy; negotiate specific behaviour change; provide structured communication skills.

Formulation 4 — contextual. A ledger of fairness is unbalanced. The husband's family history includes a father who lost employment and was humiliated; his commitment to work is a legacy aimed at restoring a family dignity that was damaged before he was born, and his sense that his contribution is unacknowledged by his wife is not only interpersonal but historical. The wife's own family history includes caretaking that was never reciprocated, and her insistence on partner contribution is haunted by earlier unpaid accounts. Interventions: multidirected acknowledgement of each partner's invisible loyalty and of what each has contributed; negotiation of entitlement and responsibility; work with the couple's families of origin where possible.

Task for the reader. Which formulation most adequately accounts for the presentation, and by what criteria would you decide? What intervention would each formulation suggest first, and what would each predict about the likely failure modes of the other three? This exercise is the point of the chapter: theory is not decoration, and different theories lead a therapist to do different things, to ask different questions, to notice different information, and to risk different mistakes. :::

::: exercise {title="Applied Exercise 1: Theoretical pluralism under a single case"} **This exercise is entirely hypothetical. The family, its members, and their circumstances are fictitious; participants who wish to discuss real cases in supervision must do so in accordance with confidentiality obligations and institutional policy.**

Participants work in four groups. Each group receives the same written vignette and is assigned one theoretical tradition from the following list: structural; strategic or Milan systemic; intergenerational or contextual; emotion-focused or attachment-based; narrative or solution-focused; cognitive-behavioural. Each group has ninety minutes to complete five tasks.

1. **Formulate.** Write a one-page formulation of the case in the assigned tradition. Name the unit of analysis, the maintenance mechanism, and the hypothesised function of the presenting problem within it.
2. **Predict.** State three specific observations or data patterns that your formulation predicts, and one pattern that would falsify it.

3. **Plan.** Specify the first three interventions you would use, and state what each is intended to change.
4. **Anticipate.** Identify the failure mode of your own formulation: the kind of case or client for whom your approach would be unsuitable or harmful.
5. **Cross-examine.** Read another group's formulation and write three questions that would test it. In plenary, the groups exchange formulations and defend them without knowing who wrote them.

Debrief questions. Which of the six formulations were genuinely incompatible, and which differed only in emphasis? Did any formulation make a claim that no conceivable evidence could refute? If the case had involved an adult disclosing fear of a partner at the first session, how would each formulation have to change, and which would survive the change intact?

The vignette is: *a family of five, referred by a general practitioner, in which a seventeen-year-old daughter has been refusing school for ten weeks; the parents disagree about the seriousness of the difficulty and about whether to involve the school; an older brother has left home and does not communicate with the family; and both parents are second-generation migrants whose extended family history includes a contested inheritance, which is discussed only in the mother tongue and never with the children present.* :::

CHAPTER SUMMARY

This chapter has traced the intellectual construction of marriage and family therapy from its cybernetic foundations to its present condition as an integrative, evidence-conscious, and self-critical discipline. Seven conclusions deserve to be carried forward.

First, the systemic turn rested on borrowed concepts — feedback, homeostasis, open systems, hierarchy, information — whose precision was often lost in clinical translation, and the doctoral reader should recover that precision.

Second, the field's foundational commitment to circular causality generated both its characteristic clinical question (what maintains this pattern?) and its characteristic vulnerability (an account that can explain everything predicts nothing).

Third, each of the major traditions supplies something the others lack: structural theory, an account of hierarchy and boundary; strategic and Milan approaches, an account of sequence and of the constitutive role of the therapist's questions; intergenerational and contextual theories, a temporal and ethical frame; experiential and communication approaches, an account of affect and of the message beneath the content; attachment and emotion-focused work, a theory of bonding and of negative cycles with the field's strongest couple-therapy evidence; behavioural and cognitive-behavioural models, the field's most rigorous research machinery and its most scalable parenting interventions; narrative and solution-focused approaches,

a theory of meaning and a set of respectful, brief practices; and integrative frameworks, the discipline of deciding which theory to use, when, and with what warrant. Fourth, the second-order and constructionist turns permanently altered the position of the practitioner: the therapist is a participant in the system under observation, and the therapist's own premises, culture, and reactions are part of the clinical data. Fifth, the field's internal critiques of power, gender, and culture are not external moralising but theoretical corrections, without which systems concepts are descriptively and ethically incomplete.

Sixth, the evidence base is uneven, and knowing where it is strong and where it is thin is a professional competence in its own right. Couple and family therapy as a class of treatment is supported by substantial evidence; the relative merits of specific models are supported in fewer instances than the field's rhetoric implies; and several influential approaches rest on clinical tradition rather than trial evidence.

Seventh, the integrative movement is the mature response of a pluralistic field. Integration is disciplined when it specifies the problem, the levels at which it is formulated, the mechanisms selected, and the criteria by which progress will be judged — and it degenerates into eclecticism when it amounts to doing whatever the therapist happens to do.

DOCTORAL RESEARCH QUESTIONS

The following questions represent live theoretical problems suitable for doctoral investigation.

1. **Mechanism specificity.** Which of the change mechanisms claimed by the major traditions can be shown, through mediation and process research, to carry treatment effects? Design a study capable of discriminating between attachment-level change, cognitive change, behavioural change, and structural change in couple therapy.
2. **The transportability of systemic concepts.** Do the core constructs of structural and intergenerational theory describe families outside the cultural and socio-economic settings in which they were formulated, or do they require reformulation? Specify the constructs, the settings, and a defensible comparative method.
3. **Circularity and responsibility.** How should systemic formulations be constructed in cases involving violence, coercion, or child protection, such that circular description does not distribute responsibility symmetrically? Develop a formal account and test it against practitioners' formulations.
4. **Therapist effects and second-order claims.** If the therapist is part of the system under observation, what follows for outcome research? Design a study that treats therapist variables as systemic variables rather than as statistical nuisance.
5. **Neurobiological claims.** Which neurobiological findings about co-regulation and threat response are sufficiently replicated to warrant use in family therapy

theory, and what are the rules for importing a laboratory finding into a clinical rationale?

6. **Common factors versus model specificity.** Are the documented benefits of couple and family therapy better explained by shared relational processes or by model-specific mechanisms? Identify a design that can distinguish the two, including the necessary measurement of alliance across multiple participants.
7. **Cultural adaptation and fidelity.** When a manualised family intervention is adapted to a different cultural setting, which components may be altered without loss of effect, and which must be preserved? This question requires a mixed-methods design and a clear theory of the active ingredients.
8. **The historical record.** How did the field's founding concepts acquire their authority, and which of them were retained for reasons that were rhetorical and institutional rather than empirical? A rigorous historiographical study of one concept (homeostasis, enmeshment, or double bind) would constitute a valuable original contribution.

FURTHER READING

- Ludwig von Bertalanffy, *General System Theory: Foundations, Development, Applications* (revised edn, George Braziller 1968).
- Gregory Bateson, *Steps to an Ecology of Mind* (Chandler 1972; reprinted University of Chicago Press 2000).
- Paul Watzlawick, Janet Beavin Bavelas, and Don D Jackson, *Pragmatics of Human Communication: A Study of Interactional Patterns, Pathologies, and Paradoxes* (W W Norton 1967).
- Salvador Minuchin, *Families and Family Therapy* (Harvard University Press 1974).
- Jay Haley, *Problem-Solving Therapy* (2nd edn, Jossey-Bass 1987).
- Mara Selvini Palazzoli, Luigi Boscolo, Gianfranco Cecchin, and Giuliana Prata, *Paradox and Counterparadox* (Jason Aronson 1978).
- Murray Bowen, *Family Therapy in Clinical Practice* (Jason Aronson 1978).
- Ivan Boszormenyi-Nagy and Barbara R Krasner, *Between Give and Take: A Clinical Guide to Contextual Therapy* (Brunner/Mazel 1986).
- Virginia Satir, *Conjoint Family Therapy* (3rd edn, Science and Behavior Books 1983).
- Augustus Y Napier and Carl A Whitaker, *The Family Crucible* (Harper & Row 1978).
- Sue Johnson, *The Practice of Emotionally Focused Couple Therapy: Creating Connection* (3rd edn, Routledge 2019).
- John M Gottman and Nan Silver, *The Seven Principles for Making Marriage Work* (revised edn, Harmony 2015) — for the popular statement; consult Gottman and Levenson's journal articles for the research programme.

- Norman B Epstein and Donald H Baucom, *Enhanced Cognitive-Behavioral Therapy for Couples: A Contextual Approach* (American Psychological Association 2002).
- Andrew Christensen and Neil S Jacobson, *Reconcilable Differences* (revised edn, Guilford 2002).
- William M Pinsof, Douglas C Breunlin, and colleagues, *Integrative Systemic Therapy: Metaframeworks for Problem Solving with Individuals, Couples, and Families* (American Psychological Association 2018).
- Douglas H Sprenkle, Sean D Davis, and Jay L Lebow, *Common Factors in Couple and Family Therapy: The Overlooked Foundation for Effective Practice* (Guilford 2009).
- Michael White and David Epston, *Narrative Means to Therapeutic Ends* (W W Norton 1990).
- Steve de Shazer, *Clues: Investigating Solutions in Brief Therapy* (W W Norton 1988).
- Harlene Anderson, *Conversation, Language, and Possibilities: A Postmodern Approach to Therapy* (Basic Books 1997).
- Monica McGoldrick, Joseph Giordano, and Nydia Garcia-Preto (eds), *Ethnicity and Family Therapy* (3rd edn, Guilford 2005).
- Froma Walsh (ed), *Normal Family Processes: Growing Diversity and Complexity* (4th edn, Guilford 2012).
- Alan S Gurman, Jay L Lebow, and Douglas K Snyder (eds), *Clinical Handbook of Couple Therapy* (6th edn, Guilford 2023) — the standard reference for comparative model review.
- *Journal of Marital and Family Therapy*; *Family Process*; *Journal of Family Therapy*; *Family Relations* — for current empirical and theoretical work.



PART II

Research, Evidence, and Clinical Decision-Making

The design, conduct, interpretation, and ethical governance of research in marriage and family therapy, with particular attention to relational data, clinical complexity, and the translation of evidence into practice.

2 CHAPTER

Research Methods in Marriage and Family Therapy

LEARNING OBJECTIVES

After studying this chapter, the reader should be able to:

1. Formulate answerable research questions about relational processes, clinical interventions, service delivery, and lived experience.
2. Select and justify quantitative, qualitative, mixed-methods, single-case, and implementation designs appropriate to the question and setting.
3. Distinguish association, prediction, mediation, moderation, and causal inference, and identify the assumptions required for each claim.
4. Design studies that represent dyadic and family interdependence rather than treating related observations as independent.
5. Evaluate sampling, measurement, missingness, analytic strategy, reporting, and risk of bias in MFT research.
6. Apply ethical principles to recruitment, consent, confidentiality, safety, data governance, and researcher-clinician dual roles.
7. Translate research findings into cautious, culturally responsive clinical decisions and identify limits on generalisability.
8. Develop a defensible doctoral research proposal, including a reproducible analysis plan and a feasible dissemination strategy.

Introduction

Research methods in marriage and family therapy (MFT) are not a technical appendix to clinical work. They are the means by which the field tests its accounts of relational processes, distinguishes helpful interventions from plausible but ineffective ones, identifies who benefits and under what conditions, and detects the unintended consequences that clinical enthusiasm may overlook. A research question about a family is also a question about how the family is represented: whether the unit is an individual, a couple, a household, a kin network, or an interaction sequence; whose report is treated as authoritative; and which social

conditions are included in the model. Methods therefore carry theoretical and ethical commitments.

The distinctive challenge is interdependence. A partner's report is not independent of the other partner's behaviour; a parent's account of a child may be shaped by the child's account of the parent; siblings share environments; and a therapist's actions may vary in response to the family while the family changes in response to the therapist. Statistical procedures that assume independent observations can produce misleading standard errors and false precision when applied uncritically to such data. Qualitative procedures can also miss relational structure if each participant is interviewed and interpreted as though their account were isolated. MFT research must make its unit of analysis explicit and align its design, measurement, analysis, and conclusions with that unit.

A second challenge is the distance between efficacy and routine practice. A tightly supervised trial may establish that an intervention can produce change under specified conditions; it does not automatically establish that the intervention will be acceptable, accessible, safe, or equally effective in ordinary services. Conversely, a family's improvement in routine care cannot by itself identify which intervention caused it. Doctoral researchers must be able to reason across this continuum, neither dismissing controlled evidence as artificial nor treating it as a universal guarantee.

This chapter moves from question formation to design, measurement, analysis, ethics, and application. Section 2.1 develops the logic of research questions and the alignment of question, theory, design, and claim. Section 2.2 examines research paradigms and the complementary roles of quantitative, qualitative, and mixed-methods inquiry. Sections 2.3 and 2.4 address sampling, measurement, and the analysis of interdependent observations. Sections 2.5 and 2.6 consider intervention evaluation, causal inference, and research ethics. Sections 2.7 and 2.8 focus on evidence synthesis, implementation, and the construction of a doctoral proposal. A hypothetical case runs through the chapter to demonstrate how a clinically important problem can be translated into a sequence of answerable research questions.

The discussion assumes a secular, evidence-oriented academic frame. It does not treat any clinical model as self-validating. Theoretical coherence can guide a study, but only transparent methods and appropriately bounded claims can support conclusions. Readers should also remember that standards, data-protection law, ethics-review procedures, and professional regulation differ by jurisdiction; the researcher is responsible for checking the currently applicable requirements before recruitment or data access.

2.1 From Clinical Concern to Researchable Question

The question precedes the method

A common doctoral error is to begin with a favoured method — a survey, a randomised trial, a thematic analysis — and then search for a problem to which it can be attached. The more defensible sequence is the reverse: specify the uncertainty that matters, identify what kind of answer would reduce it, and select a method capable of producing that answer. This sequence protects the study from methodological fashion and makes the proposal intelligible to clinicians, participants, reviewers, and funders.

A researchable question has at least five properties. It identifies a population or context; it names the phenomenon or intervention; it states the outcome or experience of interest; it defines a time frame or process where relevant; and it implies an answer that could alter knowledge or practice. "Does family therapy work?" is not yet researchable: the population, intervention, comparator, outcome, context, and meaning of "work" are unspecified. "Among adolescents referred to outpatient services for recurrent school refusal, does a manualised family-based intervention, compared with usual care, improve attendance and family functioning over six months?" is more precise, although it still requires a specification of eligibility, treatment fidelity, outcome measures, and the meaning of usual care.

Research questions can be descriptive, explanatory, predictive, evaluative, or interpretive. A descriptive question estimates prevalence or characterises a process: how often do couples disengage before seeking care, or what forms of support do families report using? An explanatory question asks how or why a pattern arises, while recognising that explanation may be causal, mechanistic, or interpretive. A predictive question asks whether information available at one time forecasts a later outcome; predictive accuracy does not itself demonstrate a causal mechanism. An evaluative question estimates the consequences of an intervention, service, or policy. An interpretive question seeks to understand the meanings participants give to an experience and the social conditions in which those meanings are formed. These categories overlap, but naming the primary purpose helps prevent a design from being asked to answer a different question than the one it can support.

A useful discipline is to write three versions of the question. The clinical version preserves the concern in ordinary professional language; the scientific version makes the population, construct, and inference explicit; and the participant-facing version explains the study without jargon or pressure. If the three versions cannot be reconciled, the construct or proposed design may be confused.

Involving service users and community partners early can reveal that a technically measurable outcome is not the outcome families consider meaningful.

Theory, construct, and operational definition

MFT concepts such as alliance, differentiation, emotional safety, conflict escalation, co-parenting, family functioning, and relational satisfaction are not directly observable in the same way as a recorded appointment date. They are **constructs**: theoretical abstractions inferred from indicators. Operationalisation specifies how the construct will be represented in a study, without pretending that the measure exhausts the construct. For example, "family functioning" might be represented by scores on a validated self-report instrument, structured observations of problem-solving, or participants' descriptions of coordination during a transition. Each operationalisation captures different aspects and carries different assumptions.

Construct validity is not established by naming a familiar scale. Researchers ask whether the measure represents the intended domain, whether its structure and scores behave as expected, whether it is distinguishable from related constructs, and whether it functions comparably across the groups in which it is used. A scale validated in one language, age range, cultural setting, or service context may not have equivalent meaning elsewhere. Translation alone does not establish equivalence; cognitive interviewing, local consultation, and empirical testing may be required. Researchers should report the exact version, scoring, source, permission status, and relevant psychometric evidence rather than relying on the label "validated".

A framework for question alignment

For intervention questions, a structured framework such as population, intervention, comparator, outcome, and time (PICOT) can expose missing elements. For qualitative questions, frameworks such as population, phenomenon of interest, design, evaluation, and research type (often abbreviated PICO or SPIDER in different traditions) may help, but should not force an interpretive question into a trial template. The framework is a prompt, not a substitute for intellectual judgement.




Figure 2.1 A research question is refined by aligning the clinical concern, population and setting, construct, design, evidence, and bounded conclusion. The arrows are iterative: feasibility and stakeholder input may require revision at each stage. *Source:* prepared for this edition.

An alignment table is a compact way to test a proposal before data collection:

Element	Proposal question to answer	Example: hypothetical study of couple conflict
Clinical concern	What uncertainty matters to practice?	Whether brief video feedback changes escalation in routine couple care
Population and setting	To whom and where might the answer apply?	Adult couples referred to two community clinics; eligibility and exclusions stated
Construct	What is meant by the key term?	Escalation defined through coded interaction and participant reports
Design	What comparison or account is needed?	Pragmatic allocation to video feedback plus usual care versus usual care
Outcome	What change is meaningful and when?	Conflict behaviour, distress, acceptability, and adverse experiences at post-treatment and follow-up
Inference	What conclusion will the design permit?	An estimate of the added intervention's effect under the study conditions, not proof of a universal mechanism

The researcher should write the intended conclusion before finalising the design. If the intended claim is "the intervention causes improvement", a cross-sectional survey will not support it. If the intended claim is "participants describe the intervention as restoring a sense of shared understanding", a randomised trial's outcome scores may be insufficient. Design alignment is a form of intellectual honesty: it makes explicit what the study can and cannot know.

Feasibility and value

A scientifically interesting question can still be unfeasible or unethical. Feasibility includes recruitment rate, staff capacity, participant burden, intervention availability, language access, data quality, follow-up retention, secure storage, analytic expertise, and realistic doctoral time. A pilot study may test recruitment, procedures, or acceptability; it should not be presented as a definitive effectiveness trial when its sample is too small for that purpose. Value includes likely contribution, relevance to affected communities, and the possibility that findings could inform decisions. The strongest proposal is not necessarily the largest study; it is a study whose question, design, resources, and claims fit together.

2.2 Paradigms and the Choice of Method

Quantitative, qualitative, and mixed-methods purposes

Quantitative methods represent observations numerically to describe distributions, estimate associations, evaluate differences, test specified models, or quantify uncertainty. They are particularly useful when constructs can be measured with adequate quality and when the question requires estimates that can be compared across people, time, or conditions. Quantification does not make a study objective by itself: decisions about inclusion, measure selection, missing data, model specification, and interpretation remain human and theoretically situated.

Qualitative methods examine language, interaction, documents, and other forms of meaning-making in context. Interviews, focus groups, observation, recordings, diaries, and documents may each be appropriate, depending on the question. Approaches differ in their assumptions: reflexive thematic analysis, grounded theory, interpretative phenomenological analysis, discourse analysis, ethnography, and qualitative case study are not interchangeable labels for "reading transcripts". The chosen approach should define what counts as data, how interpretation is developed, what role theory plays, and how the researcher will make their analytic decisions visible. A qualitative study may generate a conceptual account or a nuanced description; it does not estimate population prevalence unless paired with an appropriate sampling and measurement strategy.

Mixed methods intentionally integrate quantitative and qualitative components so that one can illuminate, extend, or challenge the other. Merely placing a survey and interviews in the same project is not sufficient. The researcher must

specify the rationale for integration, the sequence (concurrent, explanatory sequential, exploratory sequential, or another design), the priority of components, the point at which findings are brought together, and the inference generated by the combined evidence. Integration can occur through connecting samples, building one phase on another, merging data for comparison, or embedding one method within another. A joint display can make convergence, complementarity, and discrepancy visible.




Figure 2.2 Three broad methodological routes answer different kinds of questions. Mixed-methods research requires an explicit integration step; it is not simply two studies placed side by side. *Source:* prepared for this edition.

Pragmatism and methodological pluralism

A pragmatic approach often asks which methods are best suited to the question and its consequences, without claiming that all methods share identical assumptions. This can be valuable in applied MFT research, where a service may need to know both whether outcomes changed and how families experienced the change. Pragmatism does not mean "anything goes". The researcher remains accountable for the validity of each component and for the logic that integrates them. Nor does pluralism require every study to use multiple methods; a well-designed single-method study can be more informative than an incoherent mixed-methods project.

Paradigmatic disagreement is most useful when made explicit. A post-positivist design may treat constructs as imperfectly measured phenomena and seek

estimates with uncertainty; a critical realist account may distinguish observed events from underlying mechanisms and contextual conditions; an interpretive design may treat meaning as situated and co-produced in research interaction. These positions can overlap in practice, but they make different claims about what evidence is and how conclusions should be warranted. A doctoral proposal should state its assumptions in language that shapes concrete decisions, not as a decorative paragraph.

Case Illustration: the referral question

CASE ILLUSTRATION

Hypothetical case — from referral concern to staged research. A community clinic reports that some adolescent referrals for school non-attendance involve escalating conflict between caregivers and young people. Clinicians believe that a brief family intervention may improve attendance, but their records are incomplete and families describe different priorities. A doctoral researcher should not begin by promising a randomised trial. An initial descriptive audit may establish referral characteristics and the quality of routine attendance data; confidential interviews with families may clarify what outcomes matter and identify barriers to engagement; a feasibility study may then test recruitment, acceptability, and data collection. Only after the intervention and procedures are sufficiently specified would an adequately powered comparative evaluation be justified. The staged sequence is not a weaker version of science; it is a rational response to uncertainty.

In this hypothetical example, three questions arise at different levels: What patterns are present in the clinic's referrals? How do adolescents and caregivers understand the conflict and the support they want? Can the service recruit, deliver, and evaluate the proposed intervention safely? The questions require different methods. Trying to answer all three with a single symptom questionnaire would reduce the problem to what that instrument happens to measure. A staged programme can preserve clinical relevance while making each inference appropriately modest.

The next methodological decision is the unit of analysis. If attendance is measured for each young person but family interaction is measured at the household level, the design must explain how those levels relate. If two caregivers provide separate reports, a disagreement is not simply error to be averaged away; it may itself be meaningful. These issues lead directly to the sampling and relational measurement problems examined in the following sections.

2.3 Research Designs, Sampling, and Inference

Design as a structure of comparison

A research design determines which observations will be made, when they will be made, under what conditions, and what comparisons can be interpreted. No design is universally strongest. A design is strong when it addresses a well-defined question while controlling or describing the most important alternative explanations within its ethical and practical limits.

Cross-sectional studies measure variables at one time. They can describe distributions and estimate associations, but temporal order is usually ambiguous: distress may contribute to conflict, conflict may contribute to distress, and both may be influenced by economic pressure or a recent transition. Longitudinal studies repeat observations and can establish temporal ordering more clearly. Repeated measurement does not, by itself, establish causality; time-varying confounding, measurement changes, attrition, and reciprocal effects remain possible. Prospective cohort studies follow a defined group over time, while retrospective studies reconstruct prior events from records or recollection and therefore face different biases.

Experimental designs manipulate or assign an intervention. A randomised controlled trial (RCT) uses random allocation to make treatment groups comparable in expectation, subject to successful implementation and follow-up. Randomisation does not guarantee balance in a small sample, correct measurement, or absence of bias after allocation. Allocation concealment, pre-specified outcomes, transparent reporting, treatment fidelity, and appropriate analysis are essential. In MFT, the trial may randomise couples or families, clinicians, clinics, or service sites; the level of allocation must match the level at which contamination and treatment are expected. If a therapist delivers both conditions, learning or allegiance effects may blur the contrast. If whole clinics are randomised, the number of clusters and intraclass correlation become central to power and analysis.

Quasi-experimental designs estimate effects without random allocation. Options include interrupted time series, regression discontinuity, matched comparison groups, difference-in-differences, and natural experiments. Each depends on assumptions that should be stated and examined. For example, a difference-in-differences interpretation assumes that, absent the intervention, the groups would have followed parallel trends; a regression-discontinuity design depends on the allocation threshold not being manipulated and on continuity around the cut-off. Quasi-experimental methods can be useful when randomisation is unavailable or inappropriate, but the name of a method does not remove the need to evaluate its assumptions.

Single-case experimental designs collect repeated observations from one person, couple, or family across phases or conditions. Designs such as A-B-A-B reversal, multiple baseline, changing criterion, or alternating treatments can help test whether change is temporally linked to an intervention. Reversal may be unsuitable where a treatment effect is irreversible or withdrawal would be harmful. A multiple-baseline design can offer stronger control without withdrawing a beneficial intervention, provided baselines and staggered intervention points are carefully planned. Single-case work is not equivalent to an anecdote: it requires systematic repeated measurement, a defensible design, and transparent analysis of level, trend, variability, immediacy, overlap, and clinical significance.




Figure 2.3 Designs differ in allocation, timing, and the strength and kind of inference they can support. The continuum is conceptual rather than a universal hierarchy: appropriateness depends on the question, implementation, and assumptions. *Source:* prepared for this edition.

Sampling and representativeness

Sampling begins with a target population and an accessible sampling frame. A clinic-based sample may represent people who reach that clinic, not all families experiencing the problem. Inclusion and exclusion criteria affect both safety and generalisability; researchers should state them with enough detail to allow readers to understand who was not represented and why. Probability sampling can support population estimates when the sampling frame and response processes are adequate. Convenience, purposive, snowball, and consecutive sampling are often appropriate for different applied and qualitative aims, but their limitations should be described rather than disguised.

In qualitative work, sample adequacy is argued in relation to the research purpose, analytic depth, diversity of relevant experience, and information rich-

ness. A fixed number cannot guarantee saturation or depth. In quantitative work, sample size should be justified by the primary estimand, expected effect or precision, variability, clustering, repeated measures, attrition, and multiplicity. "A large sample" is not a calculation. For a trial involving couples, an analysis that ignores dyadic dependence may make a nominally adequate sample misleading. Power calculations should identify the allocation unit, outcome scale, anticipated attrition, and assumptions; where uncertainty is substantial, sensitivity analyses are more honest than a single precise figure.

Recruitment procedures can shape the sample. Clinicians may preferentially refer families they believe will comply; families may decline because of work, transport, language, privacy, or fear of consequences. Recruitment logs should document numbers approached, eligible, consenting, declining, and lost to follow-up, with reasons where ethically and practically available. Researchers should consider compensation carefully: it should recognise time and costs without creating undue influence. Accessible formats, interpreters, childcare support, flexible scheduling, and remote options may improve inclusion, although each introduces confidentiality and data-quality considerations.

Threats to validity and transferability

Internal validity concerns whether the observed difference can reasonably be attributed to the exposure or intervention in the study. Threats include selection, history, maturation, regression to the mean, instrumentation change, attrition, contamination, differential attention, and selective reporting. Construct validity concerns whether the manipulation and measures represent the intended constructs. Statistical conclusion validity concerns whether the analysis supports the stated inference, including assumptions, precision, and multiplicity. External validity concerns the conditions under which results may generalise. These categories are diagnostic tools, not a checklist that can certify a study.

Qualitative quality is often evaluated through transparency, credibility, reflexivity, coherence between question and method, attention to divergent cases, and sufficient contextual detail for readers to judge relevance. The term *transferability* recognises that a reader may assess the relevance of a finding to another setting, rather than assuming statistical generalisation. Member reflection, peer discussion, audit trails, triangulation, and reflexive journals can serve different purposes; none is a ritual that automatically proves truth. Researchers should say what a procedure contributed and what it could not establish.

Hypotheses, outcomes, and preregistration

A confirmatory analysis tests hypotheses specified before the outcome data are examined; an exploratory analysis searches for patterns that may generate later hypotheses. Both are legitimate, but they should not be presented as the same

kind of evidence. Preregistration records the question, design, outcomes, sample, and analysis plan before results are known. It can reduce undisclosed analytic flexibility and selective outcome reporting, though it cannot prevent poor measurement or make a weak theory sound. Protocol amendments should be dated and explained, and exploratory analyses should be labelled as such. For qualitative research, transparent protocols can specify the analytic approach and reflexive commitments without pretending that interpretation is a mechanical algorithm.

2.4 Measurement and the Relational Unit of Analysis

Choosing outcomes that matter

An outcome should be relevant to the question, meaningful to affected people, sufficiently reliable for the intended use, feasible to collect, and interpretable over time. MFT studies frequently need more than one outcome domain: symptoms or distress, relational functioning, safety, quality of life, service engagement, and adverse experiences may all matter. A measure can be statistically sensitive yet clinically unimportant; conversely, a change important to a family may not be captured by a standardised scale. Researchers should distinguish primary from secondary outcomes, define time points, and explain how clinically meaningful change will be interpreted.

Self-report is efficient and captures subjective experience, but may be influenced by recall, social desirability, literacy, mood, and the relationship with the person collecting data. Clinician ratings add a professional perspective but may be affected by allegiance and knowledge of treatment allocation. Administrative outcomes such as attendance, referrals, or service use can be useful, but record completeness and the meaning of a recorded event must be checked. Observational coding can capture interaction directly, but requires a clear sampling procedure, trained raters, blinding where possible, and assessment of inter-rater agreement. Physiological measures may add information about arousal, but should not be treated as transparent indicators of a particular emotion or relational mechanism.

Measurement invariance is the question of whether a construct is measured comparably across groups or time. If a scale's items function differently by language, age, gender, or cultural context, apparent group differences may partly reflect measurement rather than the construct. Researchers may assess configural, metric, and scalar invariance in suitable samples, but statistical tests cannot replace qualitative inquiry into what questions mean to participants. For cross-cultural adaptation, a careful process can include forward and back translation, expert review, cognitive interviewing, pilot testing, and psychometric evaluation; the process should be reported rather than reduced to the statement "translated into X".

Dyads, families, and nested data

The **actor-partner interdependence model** (APIM) is one framework for analysing dyadic data. It distinguishes actor effects — the association between a person's predictor and that same person's outcome — from partner effects — the association between one person's predictor and the partner's outcome. APIM can be extended to distinguish distinguishable dyads, such as two roles that are meaningfully different, from indistinguishable dyads where labels are arbitrary. The assumptions, coding, and interpretation should be explained. A partner effect is not automatically a causal influence; it is an association conditional on the model and data.

For family data, observations may be nested within members, members within families, and families within therapists, clinics, or communities. Multilevel models can represent variation at these levels, provided the number of units at each level is adequate and the model is specified appropriately. A family-level intervention does not imply that every outcome should be collapsed to a single family score. Disagreement between members can be clinically meaningful, while an aggregated score may conceal who changed and who did not. The design should decide whether the estimand concerns individual members, dyads, families, or a distribution of effects across these units.




Figure 2.4 MFT observations may be nested and interdependent: members within dyads or families, families within therapists and clinics, and clinics within service systems. The analytic model should match the sampling and intervention structure. *Source:* prepared for this edition.

Repeated measures introduce another dependence: observations from the same person or family over time are correlated. Growth models, mixed-effects models, or other longitudinal approaches may represent trajectories, but model choice depends on the question and data structure. Two or three time points usually cannot capture a complex nonlinear process with precision. Researchers should specify whether time is treated categorically or continuously and explain the meaning of the estimated trajectory. When only pre- and post-treatment measures are available, a simple change score may be transparent but can be sensitive to measurement error and baseline differences; a model adjusting for baseline outcome may answer a different but often useful question.

Missing data and data quality

Missingness is common in clinical research and may be related to the outcome. A participant who stops attending because the intervention is distressing or because conflict has escalated is not equivalent to a participant who misses a questionnaire because of a scheduling error. Complete-case analysis can be biased and inefficient. Researchers should describe the extent, timing, and reasons for missingness; compare available characteristics of retained and lost participants; and consider methods such as multiple imputation or likelihood-based estimation when their assumptions are plausible. No statistical method recovers information that was never collected without assumptions. Sensitivity analyses should test how conclusions might change under different missing-data mechanisms.

Data cleaning should be planned, reproducible, and documented. Range checks, duplicate detection, coding rules, and a data dictionary reduce errors. Corrections should retain an audit trail. For observational coding, a proportion of material should be independently coded, with agreement assessed using a statistic appropriate to the data and design; percent agreement alone may be misleading when categories are imbalanced. Researchers should not delete inconvenient observations without a pre-specified reason. Outlier handling, transformations, and model changes need justification and reporting.

2.5 Evaluating Interventions and Causal Claims

From change to effect

Families often improve during treatment, but improvement during treatment is not identical to improvement caused by treatment. Symptoms may change through natural development, regression to the mean, a change in circumstances, concurrent services, measurement reactivity, or the family's own efforts. A causal effect is a comparison between what happened under an intervention and what would have happened to the same unit, at the same time, under a relevant alternative. Because both counterfactual outcomes cannot be observed for one family,

research design creates a comparison that approximates the missing counterfactual.

The intervention must be specified sufficiently for the contrast to be interpretable. A treatment manual may describe session structure, but delivery also includes therapist competence, responsiveness, supervision, dose, adaptations, and the surrounding service context. "Family therapy" is not a single intervention. If the study compares a named model with usual care, usual care should be described: who provides it, what services are available, whether families receive additional treatment, and how those services are recorded. Otherwise the comparison group is a label rather than a replicable condition.

Randomisation can support causal inference by balancing known and unknown baseline characteristics in expectation. The analysis should generally preserve participants in their assigned groups (intention-to-treat) to estimate the effect of assignment under the trial's conditions. Per-protocol analyses may address a different question about outcomes among those who received treatment as planned, but attendance and adherence are not randomly assigned; people who complete therapy may differ systematically from those who do not. Both analyses may be informative if clearly distinguished and appropriately qualified. Blinding therapists and participants to psychotherapy allocation is usually impossible, but outcome assessors, coders, and analysts can sometimes be blinded.

Mechanism, mediation, and moderation

A treatment outcome asks whether an intervention changes an outcome; a mechanism question asks how change occurs. A **mediator** is a variable through which an intervention is hypothesised to influence an outcome. A **moderator** identifies conditions or characteristics associated with variation in an intervention's effect. These concepts are often invoked loosely. To support a mediation claim, the mediator should be measured at a time that precedes the outcome, the theoretical causal structure should be specified, and confounding of mediator-outcome relations must be considered. Cross-sectional associations between alliance and outcome cannot show that alliance caused improvement: early improvement may strengthen alliance, or a third factor may affect both. Moderation should also be pre-specified where possible and tested with adequate power; post hoc subgroup findings are especially vulnerable to false positives.

A plausible causal diagram can expose assumptions. For example, therapist experience may influence treatment delivery and outcome; baseline severity may influence both attendance and outcome; and social adversity may affect both engagement and follow-up. A directed acyclic graph is not proof of a causal model, but it forces researchers to state which variables they consider causes, mediators, colliders, or consequences. Adjusting for every available variable is not a safe

strategy: adjustment for a collider or a post-treatment variable can introduce bias. The causal question should determine the adjustment set, not a software default.




Figure 2.5 An intervention may influence outcomes through several pathways, while baseline context and delivery affect both participation and observed results. The diagram is a planning aid: the arrows represent hypotheses that must be justified, not established facts. *Source:* prepared for this edition.

Fidelity, adaptation, and therapist effects

Treatment fidelity is not simply therapist compliance with a checklist. It can include adherence to core components, competence in delivery, differentiation from comparison conditions, and the quality of the therapeutic relationship. Fidelity measures should be chosen in relation to the intervention theory. In a relational treatment, rigid adherence that prevents responsive engagement may undermine the model's intended process. At the same time, unrecorded adaptation makes it difficult to know what was delivered. Studies should document planned adaptations, distinguish core functions from adaptable forms, and record deviations with reasons.

Therapists are part of the intervention and a source of variation. Therapist effects may be estimated when the number of therapists and allocation structure support such analysis, but small numbers make estimates unstable. Therapists' allegiance to a model can influence recruitment, delivery, and ratings; equipoise, training balance, supervision, and transparent reporting can help manage but not eliminate this risk. In pragmatic settings, implementation variation may be a substantive finding rather than mere noise. The researcher should decide whether the question concerns the efficacy of a model under controlled delivery, the

effectiveness of a service package in routine care, or the processes through which implementation succeeds or fails.

Clinical significance and harms

Statistical significance is not synonymous with clinical importance. Effect estimates should be accompanied by uncertainty intervals and interpreted in relation to a pre-defined meaningful difference, baseline risk, service costs, participant burden, and possible harms. Reliable change indices, minimal clinically important differences, responder definitions, and number-needed-to-treat can each be useful, but each depends on assumptions and should be selected before results are known where possible. Reporting only a p-value obscures magnitude and precision.

Harms in relational treatment may include increased conflict, disclosure-related risk, pressure to reconcile, breaches of privacy, distress from recordings, or consequences of involving a family member in a session. Monitoring should be prospective, with procedures for response and referral. A low rate of reported adverse events can reflect inadequate monitoring or a definition that excludes relational harms. Safety planning and separate opportunities for private disclosure are especially important where coercion or violence is suspected; research protocols should not assume that joint participation is safe for every couple or family.

2.6 Research Ethics, Safety, and Data Governance

Consent and participation in relational research

Informed consent is a continuing process, not a signature collected once. Participants need understandable information about purpose, procedures, foreseeable risks, alternatives, confidentiality limits, data retention, recording, withdrawal, and use of quotations or images. Consent must be voluntary and free of inappropriate pressure from clinicians, employers, schools, or family members. Where minors participate, researchers must follow applicable law and ethics guidance on parental permission, young people's assent, capacity, and circumstances in which confidentiality may be limited. Procedures should explain what will happen if a young person discloses abuse, imminent risk, or another safeguarding concern.

In family research, one member's decision may affect others. A researcher should not presume that one person can consent on behalf of the entire family to the collection or disclosure of another member's data. Each participant should receive a clear explanation of what data are individual, what will be shared, who may access recordings, and whether participants can review or withdraw their own contribution. Group interviews create limits: researchers can promise that the research team will protect confidentiality, but cannot guarantee that other participants will keep information private. This limitation must be communicated before participation.

Confidentiality and the limits of "anonymity"

Clinical and family data can be identifiable through combinations of details even after names are removed. Rare family structures, locations, occupations, migration histories, or unusual events may enable recognition. Pseudonymisation reduces direct identification but does not make data anonymous if a key or contextual clues remain. Reports should minimise unnecessary detail, alter non-essential characteristics where appropriate, and state when a composite or hypothetical case has been used. Any alteration must preserve the analytic point and should not mislead readers about the source of evidence.

Recordings and transcripts require security proportional to sensitivity. A data-management plan should identify what is collected, where it is stored, encryption, access permissions, transfer, backup, retention, destruction, and permitted secondary uses. Cloud services, transcription platforms, and generative tools may process data beyond the researcher's intended jurisdiction; they should not be used with identifiable or confidential material without institutional approval, legal review, and informed consent. Researchers should minimise collection: data that are not necessary should not be gathered simply because a platform makes them easy to obtain.

Dual roles, power, and reciprocal obligations

The clinician-researcher may have therapeutic, evaluative, and academic authority at once. Patients may believe that declining research will affect care; students or supervisees may believe refusal will affect assessment; staff may fear that criticism will reach managers. Separate recruitment from treatment decisions where possible, use an independent recruiter, and make clear that refusal or withdrawal will not alter care or professional standing. When the therapist is also the researcher, the protocol should address access to clinical records, disclosure of research findings during treatment, and management of distress or conflict.

Research with families must attend to power within the family as well as power between institution and participant. A joint interview can silence a member or create risk after the session. Researchers need private, safe ways for individuals to express concerns, and a pre-established response pathway for coercion, abuse, suicidal risk, or danger to others. Confidentiality cannot be promised without limits, and the limits should be described in plain language. Safety procedures should be developed with local safeguarding professionals and reflect jurisdictional obligations.

Equity and community partnership

Research samples often overrepresent people with time, transport, institutional trust, dominant-language fluency, and stable access to services. Inclusion is not achieved simply by adding a demographic table after recruitment. It requires

early attention to accessibility, appropriate language, disability accommodation, compensation, cultural safety, and the distribution of research burden and benefit. Community advisors can identify problems in terminology, recruitment, outcome choice, or dissemination that academic teams may miss. Partnership must be substantive: clarify decision rights, compensation, credit, data access, and how disagreement will be handled.

Ethical review is a minimum governance process, not a substitute for ongoing judgement. A protocol may be approved before circumstances change. Researchers should report amendments, adverse events, protocol deviations, and conflicts of interest through appropriate channels. International or cross-institutional research adds questions of data transfer, local review, ownership, and fair benefit sharing. The relevant standards and law must be checked in each jurisdiction; a generic statement of compliance does not establish that a project is lawful or ethically adequate.

2.7 Analysis, Reporting, and Evidence Synthesis

Analysis begins before data collection

An analysis plan should state the primary question, estimand or analytic objective, outcome variables, time points, predictors, covariates, handling of clustering and repeated measures, missing-data approach, multiplicity strategy, and sensitivity analyses. The plan should be detailed enough to distinguish the intended analysis from later choices made after seeing results. It need not anticipate every contingency; unforeseen problems should be documented with reasons and their status as confirmatory or exploratory made clear.

For quantitative studies, report descriptive statistics that allow readers to understand the sample and data distribution, not only test statistics. Estimate effects in interpretable units where possible and provide confidence or credible intervals. Model assumptions should be checked and reported, including distributional form, residual structure, independence where assumed, and influence of unusual observations. If several outcomes or subgroup analyses are tested, the report should explain how multiplicity is addressed or why results are exploratory. "No statistically significant difference" is not evidence that two interventions are equivalent; equivalence or non-inferiority requires a pre-specified margin, a design appropriate to that question, and sufficient precision.

In qualitative research, analysis should make visible how the researcher moved from material to themes, categories, cases, or theoretical claims. The report can describe familiarisation, coding or interpretive practices, development and revision of analytic ideas, attention to disconfirming material, and the influence of the researcher's position. A coding framework may be useful in some approaches, while in reflexive thematic analysis the aim is not necessarily to produce an

objective codebook or inter-rater reliability coefficient. Methods should fit the epistemology and analytic tradition. A quotation illustrates an interpretation; it does not independently prove that the interpretation is representative.

Mixed-methods analysis requires both component analyses and an account of integration. A table that juxtaposes outcome change with interview themes may reveal agreement, elaboration, or contradiction. Disagreement between methods is not a flaw to be hidden. Families might report improved communication while observational coders detect unchanged interruption rates; the discrepancy may reflect different contexts, different constructs, or different definitions of progress. The researcher should explore the discrepancy and state what it means for the combined inference.

Transparent reporting and reproducibility

Reporting guidelines support completeness but do not replace sound design. Researchers should select a guideline appropriate to the study, such as CONSORT for randomised trials, TREND for non-randomised evaluations, STROBE for observational studies, COREQ or SRQR for qualitative reports, and relevant mixed-methods guidance. The guideline should be used during planning as well as manuscript preparation. Protocols, registration records, analytic code, materials, and de-identified data may be shared when ethically, legally, and practically appropriate. Family-level data are unusually sensitive; open science does not require public release of data that could expose participants. Controlled access, data-use agreements, synthetic data, and detailed codebooks may support transparency while protecting privacy.

Research integrity includes accurate authorship, acknowledgement of community contributions, disclosure of funding and conflicts, and correction of errors. Selective reporting, p-hacking, HARKing (hypothesising after results are known while presenting hypotheses as prior), and suppression of null findings distort the evidence base. A null result may be informative if the study was adequately designed and precisely reported. Researchers should avoid overstating implications in abstracts and press releases, where qualifications are often lost.

Systematic reviews and meta-analysis

A systematic review uses explicit methods to identify, select, appraise, and synthesise relevant studies. It begins with a focused question, a reproducible search strategy, inclusion criteria, duplicate screening or verification procedures, and a plan for risk-of-bias assessment and synthesis. The review should include unpublished or grey literature where relevant and feasible, because publication bias can make an intervention appear more effective than it is. Search limits, language restrictions, and date cut-offs should be justified. A flow diagram documents study

selection, while an evidence table supports assessment of setting, population, intervention, comparator, outcome, and design.

Meta-analysis combines estimates statistically when studies are sufficiently comparable and the data permit. It is not automatically appropriate merely because several studies exist. Differences in treatment content, population, comparator, outcome measure, follow-up period, and risk of bias may make a pooled estimate difficult to interpret. A random-effects model allows that true effects may vary, but does not explain why they vary and does not make incompatible studies comparable. Heterogeneity statistics should be interpreted alongside clinical and methodological differences. Publication-bias tests are unreliable with few studies and should not be used as a reassurance ritual. Certainty-of-evidence frameworks can help distinguish the magnitude of an estimate from confidence in it.

For MFT, systematic synthesis should attend to intervention format and context: whether treatment included the identified patient alone, a couple, parents, or a wider family; how many sessions were delivered; provider training and supervision; cultural and service context; and the comparator. "Family-based treatment" is too broad to function as a single category without further specification. Reviewers should also consider whether outcome measures privilege individual symptom reduction over relational or participant-defined outcomes.

2.8 From Evidence to Practice and Implementation

Evidence-informed practice as a decision process

Evidence-informed clinical practice integrates the best available research with clinical expertise and the preferences, goals, circumstances, and values of clients. It is not the mechanical application of an average treatment effect to every family. A therapist should ask whether the study population resembles the family before them; whether the intervention can be delivered with adequate competence; whether the outcome is meaningful; whether risk and safety have been assessed; and whether the family has a genuine choice. Evidence may inform a decision without dictating it.

The distinction between efficacy, effectiveness, and implementation helps organise this reasoning. Efficacy asks whether an intervention can produce benefit under controlled conditions. Effectiveness asks whether benefit occurs in ordinary practice. Implementation research studies the strategies and conditions that support adoption, fidelity, adaptation, reach, and sustainment. A treatment may have efficacy evidence but poor reach because it is inaccessible; an effective local programme may not be sustainable without supervision and funding. Dissemination is not merely publishing findings: it includes communication to families, clinicians, service leaders, educators, and policymakers in usable forms.

Implementation frameworks can help specify determinants and outcomes. For example, the Consolidated Framework for Implementation Research organises contextual influences, while implementation outcome taxonomies distinguish acceptability, adoption, appropriateness, feasibility, fidelity, cost, penetration, and sustainability. These are not interchangeable outcomes. A programme can be acceptable but ineffective, effective but unaffordable, or feasible at one clinic but difficult to sustain. Researchers should select implementation outcomes in relation to the decision being made.

Adaptation without loss of meaning

Interventions may need adaptation for language, culture, disability, family structure, service resources, or local law. Adaptation should be documented rather than treated as a deviation that is either automatically good or automatically bad. A useful distinction is between a model's core functions — the mechanisms it is intended to activate — and its forms — the particular materials, examples, or session arrangements used to activate them. The distinction is theoretical and must be tested; researchers should not declare any inconvenient element "non-core" after the fact. Community partners, clinicians, and service users can help identify adaptations, while outcome and process evaluation can assess what changed.

Implementation research should include unintended effects and equity. A digital or manualised approach might increase standardisation while excluding families with limited connectivity or literacy. A fidelity target might improve consistency while narrowing therapist responsiveness. An intervention delivered in a school or court context may be experienced as compulsory even if the research consent form is voluntary. Monitoring who is reached, who declines, who completes treatment, and who benefits is essential to equitable translation.

Research-informed clinical humility

Evidence can discipline clinical confidence, but it cannot eliminate uncertainty. A practitioner should communicate what is known, what is not known, and how the family's own observations will be used to evaluate progress. Routine outcome monitoring and feedback can make treatment responsive, provided scores are discussed collaboratively and are not used as a substitute for listening. If outcome data suggest no improvement, the clinician can revisit goals, alliance, formulation, safety, fit, and external conditions rather than blaming the family or defending a preferred model.

Research translation should preserve the level of evidence. A small uncontrolled study may justify further investigation, not a claim that a treatment is proven. A statistically significant average effect may not mean that every family benefits. An association between therapist alliance and outcome may support

attention to alliance, but does not prove that a particular alliance-building technique will cause improvement. Precision in language is part of ethical practice.

2.9 Constructing a Doctoral Research Proposal

A doctoral proposal should tell a coherent story: a significant problem; what is already known; what remains uncertain; the specific question; why the proposed design can answer it; how participants will be protected; how data will be analysed; and what contribution is expected. The literature review should synthesise, not merely list studies. It should compare designs and populations, identify limitations, and show how the present question follows logically from the evidence. The theoretical framework should clarify constructs and proposed relationships without being treated as a conclusion in advance.

A practical proposal can be organised around the following elements:

1. **Title and abstract:** concise, accurate, and explicit about the population, phenomenon, intervention, or design.
2. **Problem statement:** define the clinical or service problem and its significance without exaggerating prevalence or harm.
3. **Literature synthesis:** evaluate relevant theory, measures, designs, findings, and gaps; distinguish absence of evidence from evidence of absence.
4. **Aim and questions:** state one primary aim, linked sub-questions, and hypotheses where appropriate.
5. **Conceptual model:** define constructs, specify proposed relations, and identify assumptions or rival explanations.
6. **Methods:** setting, eligibility, recruitment, intervention or phenomenon, comparator where relevant, procedures, measures, and time points.
7. **Sample and feasibility:** justify size and recruitment, including clustering, attrition, access, and inclusion strategy.
8. **Analysis plan:** specify the estimand or interpretive approach, data preparation, primary analysis, missingness, sensitivity checks, and integration if mixed methods.
9. **Ethics and governance:** consent, safety, privacy, data protection, dual roles, adverse-event response, and community partnership.
10. **Contribution and dissemination:** identify who may use the findings, what decisions may change, and how limitations will be communicated.

A proposal should include a timeline with dependencies. Ethics approval precedes recruitment; instrument permissions may take time; clinician training should precede intervention delivery; translation and accessibility work should not be left until the final month. A risk register can identify recruitment shortfall, staff turnover, incomplete records, technical failure, participant distress, or changes in service policy. Each risk should have a mitigation strategy and a threshold for

revising the design. Doctoral feasibility is a scholarly criterion: an elegant question that cannot be completed responsibly is not yet a good proposal.

Applied Exercise: design a staged study

> **APPLIED EXERCISE — HYPOTHETICAL CASE**
>
> **Hypothetical exercise.** A regional service wants to know whether adding a structured co-parenting intervention to usual child-focused care improves family coordination after separation. In a short proposal outline, define the target population and safety exclusions; distinguish the primary outcome from secondary outcomes; choose an initial feasibility or effectiveness design and justify the choice; identify how each caregiver's report will be analysed without assuming independence; describe how children will be included or protected; propose a plan for missing data; and state one conclusion the design could support and one it could not. Include a brief account of how families and clinicians would help interpret the findings.

The purpose of the exercise is not to produce one correct design. A researcher might begin with qualitative interviews and a feasibility study because the intervention is not yet specified; another service with a stable programme and adequate infrastructure might justify a pragmatic trial. Either choice can be defensible if the question, assumptions, safeguards, and claims are aligned. The quality of the proposal lies in the reasoning made visible.

Summary

Research methods in MFT begin with the uncertainty that matters, not with a preferred technique. A well-formed question defines population and context, clarifies constructs, and implies an answer that can alter knowledge or practice. The researcher aligns the question, theoretical account, design, measurement, analysis, and intended conclusion. Descriptive, explanatory, predictive, evaluative, and interpretive questions require different forms of evidence; quantitative, qualitative, and mixed-methods approaches have complementary purposes and distinct assumptions.

Design determines the comparisons available and the inferences that can reasonably be made. Cross-sectional association is not causal evidence; longitudinal order improves temporal description but does not eliminate confounding; randomisation supports causal inference under conditions that include sound allocation, follow-up, measurement, and analysis. Quasi-experimental and single-case designs can be valuable when their assumptions and limitations are explicit. Sampling shapes both precision and applicability. Feasibility, recruitment, language access, participant burden, and equity are methodological concerns, not administrative details.

Measurement is a theory-laden act. Instruments represent constructs imperfectly, and validity depends on evidence for the intended interpretation and use. Outcomes should matter to families and be sufficiently precise for the question. MFT research must represent interdependence: partner reports, family members, repeated observations, therapists, and clinics may be clustered or mutually influential. Actor-partner and multilevel approaches can model some of this structure, but they do not turn association into causation. Missing data, measurement invariance, fidelity, therapist variation, and adverse effects deserve planned attention.

Ethical research protects voluntary participation, safety, privacy, and fair inclusion. Relational research demands special care because one participant's disclosure can affect others; consent, confidentiality, recording, safeguarding, and dual roles must be addressed before data collection. Secure governance and meaningful community partnership are essential. Transparent analysis and reporting, appropriate registration, and honest distinction between confirmatory and exploratory results strengthen the credibility of findings. Systematic reviews and meta-analyses are only as trustworthy as their search, appraisal, comparability, and interpretation.

The translation of evidence into practice is a reasoned decision process rather than a prescription. Efficacy, effectiveness, and implementation answer related but different questions. Families' goals, circumstances, safety, preferences, and access must remain central. Doctoral researchers demonstrate methodological maturity when they state not only what their study may show, but what it cannot establish.

Doctoral Research Questions

1. How can MFT research design represent reciprocal influence between partners while distinguishing temporal prediction from causal effect?
2. Which outcomes do families identify as meaningful after couple or family treatment, and how do these compare with commonly used clinician- or researcher-selected measures?
3. Under what conditions do brief family interventions improve adolescent engagement with care, and what mechanisms are supported by temporally ordered evidence?
4. How do language, culture, socioeconomic conditions, disability, and service context affect measurement equivalence and treatment access in family research?
5. Which strategies improve recruitment and retention of diverse families without increasing burden or compromising voluntariness?
6. How should family-level adverse experiences be defined, monitored, and reported in couple and family intervention trials?

7. What is the added value of integrating observational interaction coding with each partner's account of relational change?
8. How do therapist effects and supervision practices contribute to variation in outcomes across routine MFT services?
9. Which forms of adaptation preserve an intervention's intended function while improving local acceptability and feasibility?
10. How can routine outcome monitoring be used collaboratively without converting measurement into a compliance or surveillance tool?

Further Reading

- Creswell JW and Plano Clark VL, *Designing and Conducting Mixed Methods Research* (3rd edn, SAGE 2018).
- Kazdin AE, *Research Design in Clinical Psychology* (5th edn, Cambridge University Press 2021).
- Laurenceau J-P and Bolger N, 'Using Diary Methods to Study Marital and Family Processes', *Journal of Family Psychology* (2005).
- McHale SM and Lindahl KM (eds), *COPARENTING: A Conceptual and Clinical Examination of Family Systems* (American Psychological Association 2011).
- Pinsof WM, Zinbarg RE, and Knobloch-Fedders LM, 'Factorial and Construct Validity of the Systemic Therapy Inventory of Change', *Psychotherapy Research* (2008).
- Raudenbush SW and Bryk AS, *Hierarchical Linear Models: Applications and Data Analysis Methods* (2nd edn, SAGE 2002).
- Tashakkori A and Teddlie C (eds), *SAGE Handbook of Mixed Methods in Social & Behavioral Research* (2nd edn, SAGE 2010).
- Yin RK, *Case Study Research and Applications: Design and Methods* (6th edn, SAGE 2018).

KEY THEORETICAL POINT

Methodological rule of thumb. Make the unit of analysis explicit; define the claim before choosing the method; preserve the distinction between association and causation; treat participant safety and data governance as design features; and report uncertainty without overstating what the evidence can bear.

Selected Methodological References

The following sources are starting points for methods discussed in this chapter. Researchers should consult current editions and reporting guidance, and verify that each standard is appropriate to their jurisdiction and study design.

- American Psychological Association, *Journal Article Reporting Standards for Quantitative, Qualitative, and Mixed Methods Research in Psychology* (APA, current edition).
- Campbell DT and Stanley JC, *Experimental and Quasi-Experimental Designs for Research* (Houghton Mifflin 1963).
- Cohen J, *Statistical Power Analysis for the Behavioral Sciences* (2nd edn, Lawrence Erlbaum 1988).
- CONSORT Group, *CONSORT 2010 Statement: Updated Guidelines for Reporting Parallel Group Randomised Trials* (2010; consult current extensions).
- Hernán MA and Robins JM, *Causal Inference: What If* (Chapman & Hall/CRC 2020).
- Hussey MA and Hughes JP, 'Design and Analysis of Stepped Wedge Cluster Randomized Trials', *Contemporary Clinical Trials* (2007).
- Little RJA and Rubin DB, *Statistical Analysis with Missing Data* (3rd edn, Wiley 2019).
- McHale SM and Lindahl KM (eds), *Coparenting: A Conceptual and Clinical Examination of Family Systems* (American Psychological Association 2011).
- Moher D, Liberati A, Tetzlaff J, Altman DG, and the PRISMA Group, 'Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement', *PLoS Medicine* (2009; consult current PRISMA guidance).
- Morgan DL, *Integrating Qualitative and Quantitative Methods: A Pragmatic Approach* (SAGE 2014).
- Muthén BO and Satorra A, 'Complex Sample Data in Structural Equation Modeling', *Sociological Methodology* (1995).
- Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, Griffey R, and Hensley M, 'Outcomes for Implementation Research: Conceptual Distinctions, Measurement Challenges, and Research Agenda', *Administration and Policy in Mental Health and Mental Health Services Research* (2011).
- Shadish WR, Cook TD, and Campbell DT, *Experimental and Quasi-Experimental Designs for Generalized Causal Inference* (Houghton Mifflin 2002).
- Tong A, Sainsbury P, and Craig J, 'Consolidated Criteria for Reporting Qualitative Research (COREQ): A 32-Item Checklist for Interviews and Focus Groups', *International Journal for Quality in Health Care* (2007).
- von Bertalanffy L, *General System Theory: Foundations, Development, Applications* (George Braziller 1968).

- Wilkinson MD et al, 'The FAIR Guiding Principles for Scientific Data Management and Stewardship', *Scientific Data* (2016).

End of Chapter 2

MFT. 902 — Research Methods in Marriage and Family Therapy



PART III

Culture, Context, and Relational Practice

A critical and practice-oriented examination of culture, social location, power, identity, and context in couple and family therapy, with methods for culturally responsive assessment, formulation, intervention, and evaluation.

3 CHAPTER

Multicultural Issues in Family Therapy

LEARNING OBJECTIVES

After studying this chapter, the reader should be able to:

1. Define culture as dynamic, relational, internally diverse, and shaped by historical and material conditions rather than as a fixed list of group traits.
2. Analyse how race, ethnicity, language, migration, class, gender, sexuality, disability, age, family structure, and geography intersect in family life and in access to care.
3. Identify how therapist and institutional power shape assessment, diagnosis, alliance, confidentiality, and intervention.
4. Conduct a culturally responsive assessment that elicits each member's meanings, explanatory models, resources, risks, and preferences without stereotyping.
5. Formulate relational problems in a way that attends to both interactional processes and structural conditions, including discrimination and economic constraint.
6. Adapt systemic interventions transparently while preserving safety, client choice, and the intervention's stated therapeutic purpose.
7. Evaluate cultural responsiveness in research, supervision, service design, and outcome monitoring.
8. Develop a reflexive plan for ongoing learning, accountability, and culturally responsive clinical practice.

Introduction

Culture is not an optional layer added to family therapy after a clinician has completed the "real" assessment. It shapes the language in which distress is described, the meanings attached to roles and obligations, the routes through which help is sought, the authority granted to clinicians, and the material conditions under which families can act on therapeutic recommendations. The therapist's own cultural location and the institution's rules also shape what can be said, what is recorded, and which forms of family life are treated as ordinary or problematic.

Multicultural competence therefore concerns the whole clinical encounter, not a specialised technique reserved for clients who appear different from the practitioner.

The phrase *multicultural issues* can suggest that cultures are bounded groups, each with a predictable set of values and behaviours. That model is inadequate. People participate in multiple cultural worlds, may identify with or distance themselves from categories assigned by others, and change their affiliations across time and setting. Within every named population there is disagreement, generational change, class difference, regional variation, political conflict, and individual choice. A family may share a language but not a migration history; siblings may interpret a tradition differently; partners may negotiate gender roles that diverge from those of their relatives; and a clinician may meet people whose identities are not visible in a first session. Culture is better approached as an evolving set of meanings, practices, relationships, institutions, and power arrangements than as a fixed property of a person.

Family therapy's systemic focus offers a particular opportunity and a particular risk. It can locate difficulty within interaction and context rather than assigning blame to an individual. At the same time, systemic language can obscure unequal power if it describes coercion as a reciprocal "cycle", or treats a family's response to discrimination as an internal dysfunction. A formulation that calls a young person's resistance "oppositional" without asking what the young person is resisting may reproduce institutional assumptions. A formulation that interprets a caregiver's caution as disengagement without considering prior experiences of surveillance or poor treatment may mistake a reasonable protective strategy for pathology. Cultural responsiveness requires curiosity about context and equal willingness to examine the therapist's hypotheses.

This chapter offers a practical and critical framework. Section 3.1 defines culture, identity, intersectionality, and social context. Section 3.2 considers cultural humility, therapist positionality, power, and alliance. Sections 3.3 and 3.4 address assessment, formulation, communication, and the role of language. Sections 3.5 to 3.7 examine clinical adaptation across racialised and ethnic identities, migration and transnational relationships, gender and sexuality, disability, age, and socioeconomic circumstances. Section 3.8 brings these questions together in supervision, research, service design, and a hypothetical case. The chapter's central proposal is that culturally responsive practice is a disciplined process of inquiry, negotiation, action, and review; it is not a certification that allows a clinician to assume mastery of another person's experience.

The terms used in this chapter are not interchangeable across countries or communities. Labels such as race, ethnicity, Indigenous identity, minority, migrant, and diaspora have different histories and legal meanings in different

jurisdictions. The clinician and researcher should use the language preferred by the people concerned, explain categories used in records or research, and avoid treating a national or ethnic label as a complete clinical description. The chapter is academic and secular in approach; it addresses social, relational, and institutional dimensions of culture without devotional instruction.

3.1 Culture, Identity, and the Context of Family Life

Culture as process, not catalogue

Culture includes shared and contested meanings, practices, languages, histories, forms of knowledge, and institutions through which people organise social life. It is learned and reproduced, but also revised. It is embodied in everyday interaction and in material arrangements: who is expected to provide care, whose work is paid, who can travel safely, which languages are available in a clinic, and whose account is accepted as credible. Culture is not synonymous with ethnicity, nationality, or race. Those categories can be important, but they do not specify the full cultural world of any individual or family.

An essentialist approach attributes a behaviour to a presumed group characteristic: "families from this background do not discuss emotion," "that community expects elders to decide," or "this culture is collectivist." Such claims may appear respectful because they recognise difference, yet they can constrain a person's options and make the clinician less curious. They convert a possible pattern into a rule and position the therapist as an authority on a culture to which the client may belong. Even broad comparative constructs such as individualism and collectivism can obscure within-group differences and changes across situations. They may be useful as sensitising concepts, but not as shortcuts from a label to a clinical conclusion.

A process-oriented approach asks how a specific expectation operates in this family, who endorses it, who contests it, how it has changed, what resources it provides, and what costs it imposes. It treats the family member as an interpreter of their own social world without assuming that any one member speaks for all. It also recognises that people may hold contradictory commitments: autonomy and interdependence, privacy and kin involvement, continuity and change. Contradiction is not automatically evidence of confusion; it may be a realistic way of negotiating multiple obligations.

Identity, social position, and intersectionality

Identity includes self-understandings and the categories through which institutions and other people classify individuals. Some dimensions are chosen, some inherited, some imposed, and many are negotiated. Social position refers to a person's location within systems of advantage and disadvantage, including but not limited to racialisation, ethnicity, language, class, gender, sexuality, age, dis-

ability, citizenship, migration status, and geography. These dimensions interact rather than simply accumulate. Intersectionality, developed in Black feminist scholarship and associated particularly with Kimberlé Crenshaw's analysis of how legal and social systems can fail to recognise overlapping forms of discrimination (Crenshaw, 1991), directs attention to the specific configurations of power that shape lived experience. It is not a demographic checklist and does not imply that all members of a category have the same experience.

In family work, intersectionality requires attention to differences within a household. A child may be racialised differently from a parent in a school context; a family member with a disability may depend on another member for access while also seeking privacy; a partner's immigration status may alter economic and relational power; an older relative's language preference may affect participation in a clinical meeting. The clinician should not assume that the family is a unified cultural subject. Nor should the therapist presume that visible agreement in a session means equal influence. Questions about who speaks first, who translates, who is interrupted, and whose goals become the treatment goals can reveal relational and institutional hierarchies.

Ecological levels and structural conditions

A family's interaction occurs within nested contexts: personal history, household relationships, extended kin, neighbourhood, school, workplace, health and legal systems, media, policy, and wider historical conditions. A systemic formulation that attends only to the household may be incomplete when the reported difficulty is shaped by unstable housing, unsafe working conditions, discriminatory discipline, inaccessible services, or a legal process. These conditions do not eliminate relational agency; they change the range and cost of available choices.

Structural competence describes the ability to recognise how social and economic structures shape health and clinical encounters, and to respond within the scope of one's role (Metzl and Hansen, 2014). It does not require a therapist to become a policy specialist or promise resources the service does not control. It does require distinguishing an individual or family process from a structural constraint, documenting relevant context accurately, and knowing how to connect families with appropriate advocacy or material support where available. "The couple cannot attend weekly at 3 p.m." may be a scheduling fact, not resistance. "A parent is vigilant in meetings with school staff" may be understandable in light of prior discriminatory treatment, not merely a communication deficit.




Figure 3.1 A culturally responsive formulation holds family interaction, social position, institutions, and historical or material conditions in view at the same time; influence can move in every direction. *Source:* prepared for this edition.

Culture across the family life course

Cultural meanings and practices change with development and transition. A young adult's movement towards greater privacy may be experienced by parents as healthy differentiation, disrespect, or both, depending on the family's history and current circumstances. Migration, a new partnership, divorce, disability, bereavement, parenthood, retirement, or changing work can reorganise roles and expectations. Generational difference is not identical to cultural conflict: age, cohort, class, education, language exposure, and access to digital networks may all contribute to disagreement. The clinician should ask how family members understand the transition and what they want to preserve or change, rather than assigning the disagreement to "traditional" and "modern" positions by default.

The family is not a single cultural unit

The therapist should assess both shared practices and meaningful differences. Members may use different names for the same relationship, disagree about who belongs in a family decision, or have different thresholds for privacy and disclosure. One member may want extended relatives involved while another experiences that involvement as intrusive. An adult child may value interdependence but reject an expectation that they provide unlimited care. These are negotiable relational questions, not evidence that one person is more or less culturally authentic.

The family's own terms for its structure and relationships should be documented when possible. A household, co-parenting network, kinship group, or chosen family may extend beyond legal or biological relationships. Conversely, the presence of extended kin does not imply that every relative should be invited into treatment. Consent, safety, confidentiality, and client preference remain central. Family membership is a clinical question to be negotiated, not inferred from a clinician's preferred definition.

3.2 Cultural Humility, Positionality, and Power

From mastery to accountable curiosity

The language of cultural competence can imply a set of facts to be mastered about distinct populations. Competence remains useful when it refers to practical knowledge and skills, but it becomes unsafe when it suggests that a clinician can become an expert in another person's culture through a brief training or checklist. Cultural humility, articulated by Tervalon and Murray-García in health-professions education, emphasises lifelong self-evaluation, attention to power imbalances, and institutional accountability. In MFT, humility is not passivity or an excuse for ignorance. It is an active stance: learn relevant history, ask rather than presume, acknowledge limits, repair misattunement, and remain accountable for how professional authority is used.

A therapist can ask open, concrete questions: "Are there parts of your background or community that you want me to understand as I work with you?" "What words would you like me to use for your family and relationships?" "Are there expectations from relatives or institutions that are important to this decision?" "Have previous services made it easier or harder for you to seek help?" These questions should be asked with a clear clinical purpose, not as an invitation to make the client educate the therapist. If a client does not wish to discuss identity, the therapist should respect that choice while remaining alert to context relevant to safety and care.

Positionality and reflexive practice

Positionality refers to how a practitioner's identities, experiences, institutional role, and social location shape what the practitioner notices, expects, and can do. It also concerns how clients may experience the therapist. A therapist's race or language may affect trust for some clients, but no identity guarantees a particular response. Similarity can support connection and can also create assumptions of shared experience; difference can be navigated respectfully and can also activate mistrust or concern. The clinician should not require clients to reassure the therapist or perform a role of cultural interpreter.

Reflexivity is the repeated examination of how one's assumptions and actions enter clinical work. It is more than a personal statement about identity. Useful

questions include: Which behaviour did I label as avoidance, aggression, enmeshment, or disengagement, and what alternative meanings are plausible? Whose account did I find easiest to believe? Did I mistake professional fluency or eye contact for cooperation? Did I apply a standard of "healthy boundaries" that fits my own cultural expectations rather than the family's goals and safety? What did I feel when a family challenged my formulation, and did I treat that reaction as evidence about them?

A brief reflexive record after sessions can identify assumptions, moments of uncertainty, actions taken, and questions to bring to supervision. It should protect confidentiality and avoid turning clients into objects of the therapist's self-development. Reflexivity is strongest when it changes observable clinical behaviour: clarifying an assumption, checking a term, revising a goal, arranging an interpreter, addressing a power imbalance, or acknowledging an error.

Power in the room and in the system

Therapists hold institutional power through diagnosis, record-keeping, access to services, referrals, and professional language. In some settings, a report can influence custody, immigration, education, or child protection decisions. The family may experience an invitation to therapy as compulsory even when the form says voluntary. Clinicians must explain their role, the purpose and limits of confidentiality, who can access records, and whether information may be shared outside treatment. Ambiguity around these matters undermines trust and may expose clients to risk.

Power is also present within families. A systemic emphasis on reciprocal influence should not flatten differences in capacity, coercion, or danger. In contexts of intimate partner violence, child abuse, elder abuse, exploitation, or coercive control, describing behaviour as a mutual communication cycle can obscure responsibility and increase risk. Assessment must distinguish disagreement from intimidation, and reciprocal conflict from a patterned restriction of another person's autonomy. Joint sessions are not automatically appropriate. Separate, private assessment and a safety plan may be needed before inviting members into the same room. Cultural context may shape how risk is disclosed, but must never be used to excuse violence or deny protection.




Figure 3.2 Therapist authority, client and family power, institutional rules, and the working alliance influence one another; ethical practice makes these relationships explicit and negotiable. *Source:* prepared for this edition.

Alliance as negotiated collaboration

The therapeutic alliance in MFT involves agreement on goals, agreement on tasks, and relational connection with each participant, while the therapist also manages the alliance among members. It is possible for the therapist to have a strong alliance with one person and a weak or unsafe alliance with another. Alliance should not be equated with compliance, warmth, or shared language. A client may disagree with a formulation and still be engaged; a client may appear agreeable because disagreement feels unsafe. Routine alliance questions can be helpful when responses are reviewed openly and not used as a score of cultural fit.

The therapist can explain the systemic frame in plain language and invite correction: "I am trying to understand how the pattern affects each of you, but I do not want that to imply that responsibility is equal. How does this description fit or fail to fit your experience?" This language both offers a hypothesis and preserves room for disagreement. When a rupture occurs, the clinician should name the specific action, listen to the impact, avoid defensive explanations, and negotiate a repair. If the therapist has used a harmful term or made a stereotyped assumption, a concise acknowledgement and changed practice are generally more useful than an elaborate apology that asks the client to comfort the therapist.

Supervision and organisational accountability

Cultural responsiveness cannot be delegated solely to an individual clinician. Supervisors and organisations shape caseloads, interpreter access, treatment

protocols, documentation, referral thresholds, and the time available for care. Supervisors can use case discussion to examine the therapist's power, divergent member perspectives, structural constraints, and the assumptions embedded in measures or service rules. They should also recognise that a supervisee may fear professional penalty for raising concerns about bias. A psychologically safe supervisory process is not one in which all actions are approved; it is one where actions can be examined honestly and repaired.

Organisations can audit who is referred, accepted, discharged, retained, and reported as improving; compare these patterns across groups where lawful and appropriate; and investigate differences rather than assuming that they reflect client deficits. Data should be interpreted carefully, with attention to small cell sizes, privacy, measurement equivalence, and the histories behind categories. Accountability includes changing systems when a pattern indicates exclusion, not merely offering staff another training module.

3.3 Culturally Responsive Assessment and Formulation

Assessment as inquiry and negotiation

Assessment is not a neutral collection of facts. The clinician chooses what to ask, which accounts to record, what is considered relevant, and how a description becomes a diagnosis or treatment plan. A culturally responsive assessment uses structured clinical skills while remaining open to meanings that the instrument or diagnostic system may not capture. It identifies the presenting concerns from each member's perspective, the family's own account of cause and desired change, strengths and resources, safety, developmental and relational history, and the external conditions affecting the problem.

A useful sequence begins with the family's reason for seeking care. The clinician can ask what led the family to attend now, what each person hopes will be different, and what they have already tried. Ask how the concern is named at home and in other settings, whether family members use different terms, and whether a particular term feels stigmatising or inaccurate. Explore the timeline and contexts in which the concern increases or decreases. The purpose is not to replace clinical judgement with the family's first explanation, but to understand their framework before imposing one of the therapist's own.

Questions about explanatory models can be framed in ordinary language: "What do you think is contributing to this?" "What worries you most about it?" "What kind of help would make sense to you?" "What would be a sign that treatment is useful?" The clinician can explore prior experiences with health, education, legal, or social-service systems and what the family wants done differently this time. Inquiry should be proportionate: not every session requires an

exhaustive cultural history. The therapist should explain why a question matters, allow a person to decline, and avoid treating disclosure as a test of readiness.

A multidimensional formulation

A clinical formulation should distinguish observations, interpretations, and hypotheses. It might describe a recurring sequence between caregiver and adolescent; note how housing instability and school exclusion intensify it; record that family members disagree about the meaning of the adolescent's withdrawal; and identify safety questions that remain unresolved. It should not convert a social identity into a cause or a group average into a prediction about an individual. Competing hypotheses should remain visible until there is enough evidence to revise them.

A useful formulation can organise information across six domains:

1. **Presenting concern and meanings:** each member's account, preferred language, goals, and level of distress.
2. **Interactional pattern:** sequence, triggers, responses, attempted solutions, exceptions, and consequences.
3. **Developmental and relational history:** transitions, attachment and caregiving experiences, kin relationships, and prior coping without assuming a single normative family pathway.
4. **Social position and context:** language, migration, racialisation, class, disability, gender, sexuality, age, geography, and relevant historical or institutional conditions.
5. **Resources and protective factors:** supportive relationships, skills, community connections, material resources, and prior successful adaptations.
6. **Risk, constraints, and uncertainty:** violence, coercion, safeguarding, access barriers, confidentiality limits, and questions requiring further assessment.

This is not a requirement to document every identity category. It is a prompt to include what is relevant and to identify what remains unknown. Formulations are provisional working tools, not authoritative narratives about a family. They should be reviewed with the people concerned when possible and updated when new information changes the picture.

Distinguishing distress from difference

Clinical systems can pathologise behaviour that differs from a clinician's expectations. Quietness, limited eye contact, emotional restraint, a large extended-family network, indirect disagreement, or a strong preference for privacy may have multiple explanations. None should be assigned a meaning solely from a cultural label. The clinician asks whether a behaviour causes distress, restricts chosen functioning, creates safety concerns, or conflicts with the person's own aims. At

the same time, cultural relativism should not become a reason to ignore coercion or harm. Respect for difference and protection from abuse are compatible obligations.

Diagnostic frameworks may help communicate patterns and support access to services, but diagnosis should not displace a relational and contextual account. A clinician can state that diagnostic criteria are met while also documenting the role of discrimination, trauma, poverty, language barriers, or family transition where relevant. Conversely, a social explanation does not rule out a mental-health condition that merits assessment and treatment. Avoid false choices between individual suffering and structural context: both may be real, and their relation may be complex.

Culture-bound concepts and idioms of distress

People express distress through locally meaningful language, bodily experiences, relational concerns, and metaphors. Clinical services may recognise some terms and overlook others. An idiom of distress is not necessarily a symptom category; it can communicate suffering in ways that are socially intelligible and useful to the person. The therapist should ask what the expression means to the speaker, who else uses it, and what help it implies. A clinician should not assume that a term has the same meaning across countries or even across families.

The DSM-5-TR Cultural Formulation Interview (CFI) is one structured aid for eliciting a person's definition of the problem, perceived causes and supports, coping, help-seeking, and preferences. It can improve consistency of inquiry when used flexibly and with training. It does not replace a comprehensive systemic assessment, establish cultural competence by itself, or compel disclosure. The CFI and related tools should be used in the version and language appropriate to the setting, with attention to licensing and current guidance. The clinician should explain the purpose and adapt pacing to the person's needs.

	Inquiry	Clinical use
Meaning	How does each member name the concern?	Goals and explanatory models
Interaction	What sequence maintains or changes it?	Tentative relational hypotheses
Context	Which identities and conditions matter here?	Access, history, structural constraints
Resources	What supports and adaptations are available?	Strengths, kin, community, practical help
Safety	What risks or confidentiality limits apply?	Separate assessment, protection, referral
Uncertainty	What is not yet known or agreed?	Review and revise formulation

Figure 3.3 A culturally responsive clinical formulation links members' meanings and interactional sequences with social position, structural context, resources, and safety; hypotheses are revisited collaboratively. *Source:* prepared for this edition.

Using measures responsibly

Standardised measures can support monitoring, but their use requires attention to literacy, language, cultural validity, accessibility, and the construct being measured. If a measure has not been validated for the relevant population, the clinician should not imply that its scores have established the same meaning as in a validation sample. Interpreters should not be asked to translate proprietary instruments spontaneously without considering standardisation and permission. If no adequate translation exists, the clinician may use qualitative inquiry or another appropriate measure and document the limitation.

A score that differs from a clinician's impression is an invitation to discuss perspectives, not a reason to dismiss one. A family member may interpret an item differently, or the measure may not capture their concern. Conversely, a reassuring score does not rule out risk when there are safety indicators. The therapist should ask how participants understood the questions and what the score means to them, particularly when the result may shape access, diagnosis, or discharge.

3.4 Language, Communication, and Meaning

Working with interpreters

Language access is an ethical and clinical requirement, not a courtesy. A professional interpreter can support accuracy, privacy, and participation; using a child, relative, or untrained staff member as interpreter can create coercion, role reversal, omissions, and confidentiality risks. Family members may choose to assist in some ordinary contexts, but should not be treated as the default interpreter for sensitive clinical conversations. The clinician should follow local law, organisational policy, and professional standards regarding qualified interpretation and translated consent materials.

Before the session, the therapist should brief the interpreter about the purpose, expected terminology, turn-taking, confidentiality, and any safety concerns. The therapist speaks directly to the family member in the first person, uses short clear segments, avoids idioms and layered questions, and allows time for interpretation. The interpreter is not a co-therapist or cultural representative for an entire community. Afterward, the clinician can debrief about terminology or process without asking the interpreter to make a diagnosis or disclose information outside their role. When dialect, gender, or community affiliation affects safety or comfort, ask participants about preferences where feasible.

Interpretation is relationally complex. A member may understand a language but prefer to discuss emotion in another; partners may disagree about which language should be used; and code-switching may signal intimacy, authority, or exclusion. The therapist can ask who wants what language for which parts of the meeting. Ensure that interpreters understand confidentiality and that the family understands the limits. Remote interpretation may increase access but can reduce non-verbal information and create privacy challenges in the home; these should be reviewed explicitly.

Communication styles and clinical assumptions

Communication is culturally patterned but not reducible to national styles. Directness, silence, overlap, gaze, gesture, emotional display, and deference can vary with context, relationship, power, neurotype, stress, and individual preference. A clinician should not treat eye contact as a universal indicator of honesty, treat emotional restraint as lack of engagement, or equate verbal fluency with insight. Nor should a therapist romanticise indirect communication or assume it is shared by all members. Ask how people prefer to communicate and whether the session format is making participation easier or harder.

Turn-taking in a family session can reproduce existing hierarchies. The therapist can invite each member to speak without forcing disclosure, notice who has not had a chance to contribute, and offer different ways of participating, such as

writing, drawing, interpreter-supported speech, or a private check-in. These adjustments should be offered rather than imposed. In sessions involving children, adolescents, older adults, or people who use augmentative communication, the therapist should adapt pacing and format while preserving the person's agency.

Confidentiality and disclosure across settings

Clients may worry that disclosure will affect immigration, custody, employment, school placement, benefits, or safety. The therapist should explain what is recorded, who can access it, whether diagnoses or reports may be shared, and the circumstances in which confidentiality has legal limits. Statements such as "everything is confidential" are misleading when exceptions apply. Avoid collecting sensitive identity information unless it is clinically or administratively justified, and explain why it is requested and how it will be used.

Family therapy creates a further confidentiality issue: the clinician may receive information from one member that changes the safety or meaning of joint work. The therapist should establish a clear policy at the outset about individual communications, secrets, records, and when information may be shared. Policies differ by jurisdiction and setting; no single "no-secrets" rule is appropriate for every family or every risk. The clinician should explain the policy before private disclosures occur, seek consultation when needed, and avoid promising a level of privacy that cannot be maintained.

Building a shared language for change

Therapist and family may use the same word differently. "Respect", "independence", "support", "discipline", "family", and "privacy" can carry distinct meanings. A useful practice is to ask for an example: "When you say respect, what would I see happening at home?" This moves conversation from abstract values to observable interaction without implying that one definition is correct. The family can then negotiate which changes are desired and which meanings should be preserved.

When clients correct a therapist's language, the response matters. The therapist should use the requested term, clarify if uncertain, and avoid asking a client to defend an identity. A correction is information about the relationship and the clinician's impact. Repeatedly returning to an unwanted label or name can damage trust even if the clinician believes the intention was neutral. Accuracy and repair are ongoing clinical tasks.

3.5 Kinship, Migration, and Transnational Family Life

Family structure and kinship networks

Many clinical systems presume that the relevant family consists of a co-resident couple and their dependent children. That pattern exists alongside single-parent

households, multigenerational households, separated co-parents, kinship care, foster and adoptive relationships, same-gender-parent families, chosen families, and networks whose members live across regions or countries. The therapist should ask who the family considers important, who provides care, who has decision-making authority, and who should participate. Legal status and relational significance may not align: a person may be central to care without legal guardianship, or legally related but not safe to involve.

Extended kin relationships can be a source of belonging, childcare, practical help, identity, and continuity. They can also involve conflict, pressure, unequal labour, or restrictions on autonomy. The clinical task is not to encourage or discourage extended-family involvement as a general principle, but to understand what it does in the particular network and what the family wants. Mapping kinship can reveal both support and burden. A genogram should be adaptable to chosen and culturally significant relationships, and the therapist should explain symbols rather than allow a conventional chart to erase family members who do not fit its categories.

Migration, displacement, and settlement

Migration is not one experience. Voluntary movement for work or education, forced displacement, family reunification, refugee resettlement, seasonal mobility, and internal migration involve different risks, resources, legal conditions, and meanings. A clinician should not presume trauma from the fact of migration, nor overlook the effects of violence, loss, separation, uncertainty, discrimination, or insecure status when they are present. Ask what happened, what is safe to discuss, what the family wants to make of the experience, and how current legal or economic conditions affect daily life.

Migration can reorganise family roles. A young person who learns the dominant language quickly may become an interpreter for parents; a parent may lose professional status; family members may be separated for years; remittances may create obligations; or caregiving may be distributed across borders. These shifts can produce new competence as well as strain. A child who translates at appointments may gain responsibility but also carry adult information; a parent may feel dependent on the child while still holding important authority in other domains. The therapist should identify the specific pattern and avoid the simplistic story that children "acculturate" while parents "resist".

Settlement stress can include housing insecurity, work exploitation, racism, uncertain documentation, restricted access to services, and isolation. Such pressures may aggravate couple conflict or parent-child disagreement without being reducible to family dysfunction. Clinical recommendations should be feasible under the family's actual constraints. A weekly daytime appointment, separate

transport, or a costly referral can be inaccessible. Where legal or practical expertise is needed, therapists should make appropriate referrals rather than offer unqualified advice about immigration or status.

Acculturation and cultural change

Acculturation refers to processes of cultural and psychological change associated with sustained contact between cultural groups. It is not a single scale on which one person moves from "heritage" to "host" culture. People may adapt language, work practices, parenting styles, or public presentation at different rates and in different contexts. They may maintain some practices, revise others, and reject still others. The process is shaped by power, discrimination, opportunity, social networks, and personal choice. A family's disagreement may concern a specific behaviour—dating, clothing, school participation, care of elders—rather than a global clash between cultures.

The therapist can explore how members describe change and what each fears losing or gaining. A parent may worry that a young person's new independence will weaken connection; the young person may experience the same movement as necessary for safety or belonging. Both perspectives deserve attention, while the clinician also assesses coercion, harm, and developmental rights. "Both sides" language should not erase asymmetries in power or responsibility. The aim is to make meanings and consequences discussable and to support negotiated choices where safe.

Transnational ties and distributed care

Family relationships may be maintained through calls, messaging, remittances, visits, and care coordination across distance. A household member in one country may participate in decisions about schooling, healthcare, or elder care from another. Time zones, connectivity, cost, surveillance, and language access influence this communication. A conventional in-room family session may not include the people who shape the family's decisions, but adding remote participants may compromise privacy or reinforce pressure. The therapist should ask what role distant relatives play and whether their involvement is desired and safe.

Transnational caregiving can be both sustaining and demanding. A family may coordinate care for an older relative across borders while managing employment and child-rearing locally. Remittances can express solidarity, but can also be a source of conflict if family members disagree about priorities. The clinician should avoid interpreting transnational obligation as pathology or as an uncomplicated virtue. The question is how the arrangement affects each person, what choices are available, and what forms of support could reduce strain.

Culture, trauma, and the danger of deficit narratives

Exposure to violence, persecution, displacement, or discrimination can affect family communication and regulation, but clinicians should not use a trauma label to define a family. Some people prefer not to recount events; forced retelling is not a prerequisite for care. Present-focused support, practical stabilisation, and relational safety may be more appropriate. Assessment should consider ongoing threats and systemic conditions, not only past events. Resilience should be recognised without using it to minimise harm or deny access to support.

Deficit narratives emerge when clinicians describe a community mainly through risk, dysfunction, and service need. Counter this by documenting competence, mutual aid, humour, adaptation, cultural knowledge, and aspirations alongside difficulties. Strengths are not an alternative to material support: praising resilience does not remedy unsafe housing or discrimination. The formulation should capture both the family's resources and the constraints that make those resources costly to sustain.

Hypothetical Case Illustration: a cross-generational transition

> **CASE ILLUSTRATION**
>
> **Hypothetical case — a family negotiating migration and autonomy.** A 16-year-old, their caregiver, and a younger sibling attend a clinic after repeated arguments about school attendance and social activities. The young person speaks the local language more fluently and has begun interpreting for the caregiver in public services. The caregiver describes the young person's new social circle as unsafe; the young person says that restrictions are isolating and that they are tired of translating adult matters. The clinician does not assume that the family represents a particular national pattern. In separate and joint conversations, the therapist asks what the caregiver fears, what the young person wants, what information is safe to share, how interpretation responsibilities developed, and what school or community pressures are involved. The formulation includes language burden, parent-child authority, safety, and the family's own goals. A professional interpreter is arranged for future sessions. The therapist helps the family negotiate specific limits and a safer way to share public-service tasks, while keeping the young person's privacy and the caregiver's concerns visible.

This hypothetical case illustrates why cultural formulation must be specific. The concern is not simply "acculturation conflict". It includes a role reversal in public institutions, a developmental negotiation, safety concerns, and access to professional language support. The clinician should not decide in advance that either the caregiver's caution or the young person's wish for autonomy is culturally determined. Each account is explored, safety is assessed, and the intervention is reviewed for whether it reduces burden without dismissing either person's agency.

3.6 Race, Ethnicity, and the Effects of Racialisation

Naming structural experience

Race is a social and political category with real consequences, not a biological explanation for family behaviour. Racialisation is the process through which people are assigned racial meanings and treated accordingly. Families may experience discrimination in healthcare, schools, housing, employment, policing, and other institutions; these experiences can shape stress, trust, vigilance, and decisions about disclosure. The therapist should be able to ask about discrimination without presuming it has occurred or requiring clients to provide proof before it can be considered.

A question such as "Have experiences of bias or unfair treatment affected how your family is approaching this problem or this service?" can open a discussion while leaving room for "no", "not sure", or "I do not want to discuss it". If a client describes discrimination, the therapist should not immediately reframe the account as a cognitive distortion, communication problem, or misunderstanding. Explore what happened, its effect, what support the client wants, and whether the service itself needs to respond. The clinician should distinguish validation of lived experience from making claims about facts outside the available information; uncertainty should not become reflexive disbelief.

Historical context and family meaning

Historical policies and collective histories can affect present-day trust, family narratives, and institutional contact. The relevance and form differ by country and community. Therapists should learn the histories that are relevant to the populations they serve, but should not treat a client as a representative of a history or assume that a particular event explains their current behaviour. Ask whether historical or community experiences matter to the family and how they understand their relevance. The clinician's task is to situate, not to impose a collective narrative.

Avoiding racialised diagnosis and unequal standards

Clinical labels can be influenced by expectations about threat, emotional expression, parenting, and authority. A therapist should examine whether the same behaviour would be described similarly if performed by a person from another social group. Terms such as "aggressive", "resistant", "non-compliant", "hostile", and "enmeshed" may encode judgement unless linked to specific observed behaviour and context. Documentation should distinguish what was observed from inference and avoid vague characterisations that could be harmful in later institutional decisions.

The clinician should also consider how systemic models have historically described families. Some theories have universalised middle-class, Euro-American

assumptions about household boundaries, gender roles, privacy, and autonomy. Their constructs can still offer useful questions, but must be evaluated rather than applied as cultural facts. A "clear boundary" may support safety in one context and sever essential care in another; closeness may be supportive, controlling, or both. The family should help define the functional meaning of a pattern, and the clinician should assess freedom, consent, safety, and consequences.

Community knowledge without tokenism

Community consultation can improve service design, terminology, recruitment, and referral pathways. It should not depend on one community member being expected to represent all people who share an identity. Define the purpose of consultation, compensate expertise, protect confidentiality, and communicate what changed as a result. When disagreement exists within a community, it should be represented rather than flattened into a single "cultural view". Researchers and organisations should share decision-making where possible and avoid extracting knowledge without reciprocal benefit.

3.7 Gender, Sexuality, Disability, and Socioeconomic Context

Gender, power, and relational roles

Gender is socially organised and personally experienced; it shapes expectations, opportunities, safety, and distributions of unpaid and paid labour. Couples may negotiate caregiving, income, emotional work, decision-making, and public identity under explicit and implicit gender norms. The therapist should assess how these arrangements function for each person, who has freedom to refuse or renegotiate, and what consequences follow. A claim of "equal contribution" should not be accepted without exploring invisible tasks such as planning, kin coordination, emotional monitoring, and care administration.

Gender diversity should be approached without pathologising identities or assuming that a person has a particular treatment goal. Ask for names and pronouns, use them consistently, and correct errors without making the client responsible for the therapist's embarrassment. In family work involving a transgender or gender-diverse member, clarify confidentiality and safety before disclosing information to relatives. A young person's identity should not be revealed to a caregiver without consent except where a clearly applicable safeguarding or legal obligation requires a different response; the therapist must check local law and policy and discuss limits of confidentiality in advance. The clinician should assess family responses, support, rejection, and risk without treating identity itself as a symptom.

Therapists should not infer relationship roles from gender presentation, legal status, or household labels. Ask how partners define their relationship, what commitments they recognise, and how they wish to be addressed. Where services

or records use binary categories, explain why information is needed, offer a way to describe identity in the person's own words, and avoid collecting details that are irrelevant. MFT's focus on relational context should not become a pretext for making one person's identity a family "problem" or requiring them to educate relatives in session.

Sexual orientation and family diversity

Couples and families include lesbian, gay, bisexual, queer, asexual, and other sexual identities, as well as people who do not use a label. The therapist should avoid heteronormative assumptions about partners, parenting, fidelity, family roles, or future plans. Ask open questions about significant relationships and chosen family. A client's decision to disclose an identity to relatives is theirs; therapy should not pressure disclosure as a universal route to authenticity or family repair. Consider how stigma and legal or social conditions affect safety, access, and trust, while recognising that clients differ in how they understand and prioritise these experiences.

For families with LGBTQ+ members, clinical work may address a particular relational rupture, ordinary life-cycle transition, or external stressor. Not every problem is caused by stigma, and not every family is in conflict about identity. Avoid both erasure ("we treat everyone the same") and overemphasis that makes identity the only relevant dimension. Tailor questions to the family's stated concern, acknowledge discriminatory conditions where relevant, and make services visibly safe through respectful intake forms, accurate referrals, and confidentiality practices.

Disability and neurodiversity

Disability can be understood through medical, social, and relational models. A medical account may attend to impairment and treatment; a social model highlights barriers created by inaccessible environments; relational approaches consider how disability, care, identity, and family organisation interact. These perspectives can be complementary. The clinician should ask how the person understands disability and what language they prefer; some identify as disabled, others prefer person-first language, and preferences vary. Do not presume that a diagnosis defines the person's capacity, goals, or family role.

Accessibility should be designed into practice: step-free access, clear sensory conditions, communication supports, accessible written materials, flexible pacing, and remote participation where suitable. Ask about needed accommodations rather than making assumptions. For a client who uses augmentative and alternative communication, allow enough time for responses and address the client directly. Do not treat a family member's interpretation of another person's communication as definitive; check the person's preferences and consent. Neurodi-

vergent people may have different sensory, social, or processing needs, and therapy should adapt without framing difference as lack of motivation or empathy.

Caregiving relationships deserve careful assessment. A relative may provide essential support while also holding substantial practical power; the person receiving assistance may want greater autonomy or a different allocation of tasks. The clinician should assess caregiver stress and the disabled person's own goals, rather than treating one as the sole client and the other as a resource. When capacity or supported decision-making is relevant, follow legal standards and professional consultation, and involve the person as fully as possible.

Age, development, and intergenerational power

Children, adolescents, and older adults are not homogeneous groups. Developmental stage, communication access, family authority, and legal status affect participation. A child may have a right to a private conversation in some settings, but the scope and limits depend on jurisdiction and circumstances. The therapist should explain confidentiality in developmentally appropriate language and avoid creating promises that cannot be kept. Parents' authority and responsibilities are important, but the young person's safety, dignity, and voice must also be represented.

Older adults may be positioned as decision-makers, dependants, caregivers, or all three at different times. Ageism can lead clinicians to discount distress, presume cognitive decline, or treat a person's preferences as subordinate to adult children's convenience. Conversely, a therapist should not overlook coercion or neglect in the name of respecting family hierarchy. Assess capacity in relation to the specific decision, use communication supports, and distinguish a family's preferred arrangement from the individual's informed choice.

Class, work, and material resources

Socioeconomic circumstances shape access to therapy and the daily options available to families. Insecure housing, transport costs, shift work, childcare, food insecurity, debt, and limited digital access can affect attendance and stress. A therapist who interprets missed sessions as resistance may overlook the service's scheduling assumptions. A treatment plan requiring private space, internet, paid time off, or a stable home may be unrealistic. Ask about practical barriers without shaming and consider whether the service can adapt timing, location, format, or referral pathways.

Class is not only income. It includes occupation, education, social capital, neighbourhood, and familiarity with professional institutions. Clinicians may unintentionally reward clients who speak in professional language, complete forms promptly, or display emotional styles that match the service's expectations. The therapist should examine how institutional convenience is being mistaken for

motivation. Where material conditions are central, relational work can still help family members coordinate, but should not substitute for advocacy or practical support when those are needed.

Culture and safety are not competing priorities

A culturally responsive therapist respects difference while maintaining clear standards against coercion, abuse, exploitation, and discriminatory treatment. The clinician must not impose their own preferred family structure under the name of safety, but must not excuse harm as cultural practice. Risk assessment should be individualised, confidential where possible, and attentive to the person's own account of danger and available resources. If joint sessions may increase risk, individual assessment, specialist consultation, or referral may be necessary. The decision should be documented, reviewed, and explained in a way that does not blame the person at risk.

3.8 Adaptation, Evidence, and Organisational Practice

Adapting a model with transparency

Intervention adaptation can involve language, examples, metaphors, session format, family participation, pacing, and the degree of directiveness. Adaptation should respond to a stated need and be discussed with participants. The clinician can distinguish the therapeutic function intended by an intervention from the form used to deliver it, but should not assume that a component is culturally neutral or universally essential. A change that improves fit for one member could intensify a power imbalance for another. Consultation and ongoing feedback help identify these effects.

When using a manualised approach, clinicians should document adaptations and consider whether they alter the model's hypothesised mechanism. Some adaptations are accommodation—such as translating materials or changing session time—while others alter intervention content. Both can be appropriate, but the consequences differ. Supervisors and researchers should record what changed, why, by whom, and with what response. Fidelity should not be equated with rigid sameness, and adaptation should not be a label that excuses unexamined practice.

Evidence and generalisability

Treatment evidence is produced in particular populations, languages, settings, and historical periods. A statistically effective intervention may not have been tested with the family before the therapist. A small number of studies in one group does not establish universal suitability, and an absence of subgroup evidence does not prove that a treatment is ineffective for that group. Researchers should report sample composition, recruitment, attrition, adaptations, language procedures, and subgroup analyses transparently. Subgroup effects require ade-

quate power and should not be inferred from post hoc comparisons without qualification.

Culturally responsive research asks whether outcomes were selected with affected people, whether measures function comparably, whose definitions of improvement were used, and whether the intervention was accessible. Qualitative process studies can explain how participants interpret treatment; quantitative studies can estimate reach and outcomes; mixed-methods studies can connect these findings if integration is planned. Community-engaged research can broaden relevance, but partnership requires equitable decision-making, compensation, and a clear account of data ownership and dissemination.

A practical cultural responsiveness cycle

A clinician can use a repeatable cycle: **locate** the presenting concern within relevant identities and contexts; **ask** each member what it means and what help is wanted; **formulate** interactional and structural hypotheses; **negotiate** goals, language, participation, and adaptations; **act** with safety and access in view; and **review** the effect with each participant. The cycle is not a linear checklist. New information may require returning to an earlier step. The clinician should record the rationale for significant adaptations and seek supervision when uncertainty or risk exceeds their competence.




Figure 3.4 A practice cycle for culturally responsive MFT: locate context, ask about meaning, formulate tentatively, negotiate goals and adaptation, act, and review impact. The cycle is iterative, not a certification checklist. *Source:* prepared for this edition.

Hypothetical Case Illustration: unequal participation in a couple

CASE ILLUSTRATION

Hypothetical case — work, language, and decision-making. A couple seeks therapy for recurring arguments about money and caregiving. One partner recently moved to the region, has limited confidence in the service's primary language, and works irregular hours; the other partner has stable employment and usually answers questions first. The therapist notices an apparent pattern of "withdrawal" by the first partner. Rather than concluding that this partner is avoidant or culturally deferential, the therapist asks about work schedules, language preference, financial access, and whether the partner wants to speak privately. A qualified interpreter is arranged for one member's preferred language; the therapist checks whether the couple wants interpretation in joint sessions or separate time. The assessment reveals that the quieter partner is managing extensive care coordination and does not feel safe disagreeing about a shared account. The therapist screens separately for coercion, discusses confidentiality limits, and postpones joint money exercises until safety is clearer. The formulation is revised from mutual miscommunication to a combination of unequal decision-making, practical strain, and a communication pattern that requires further assessment.

This hypothetical example shows why a therapist should not treat equal airtime as proof of equal power or silence as a cultural trait. Language access and income affect participation, but do not determine the interpretation. Separate safety assessment is essential because "money conflict" may conceal control. The case demonstrates how culturally responsive practice modifies the pace and structure of care without assuming that a single intervention is appropriate before risks are understood.

3.9 Supervision, Documentation, and Continuing Development

Recording culture and context responsibly

Clinical records should include information that is relevant to care, accurately attributed, and proportionate to the purpose of the record. Distinguish a client's account from a clinician's observation and from a clinical hypothesis. Avoid using broad labels—"traditional", "non-Western", "difficult", "resistant"—without specifying the behaviour, context, and source of the description. Do not record sensitive identity information merely because it is available. Where identity or context is clinically relevant, explain why it matters and use the person's preferred language where possible.

Documentation can travel beyond its original setting. Notes may be read by other clinicians, insurers, courts, schools, or clients themselves, depending on law

and policy. A clinician should consider whether a phrase could stigmatise a family, obscure discrimination, or imply certainty not supported by the assessment. If an interpreter was used, note the language and relevant process, while following privacy requirements. If the clinician's understanding changed, document the updated formulation rather than allowing an early assumption to persist unchallenged.

Supervision as a site of accountability

Supervision should examine how the therapist's identity, theoretical orientation, and institutional position affect the case. A supervisee can present one concrete interaction, the interpretation made at the time, alternative explanations, and the response of each family member. Supervisors can ask whose account is missing, what structural conditions are active, whether risk has been assessed privately, and whether the proposed intervention fits the family's stated goals. Supervision should neither demand disclosure of personal identity beyond its relevance nor treat the supervisor's own experience as a universal reference point.

A supervision plan can include targeted learning, consultation with a specialist, review of translated materials, analysis of service access, or a change in the session structure. Consultation with community experts should be respectful and compensated where appropriate; no individual should be asked to speak for an entire population. The therapist remains responsible for clinical decisions and should not transfer the burden of cultural judgement to a consultant.

Developing reflective expertise

Continuing development includes reading scholarship, learning relevant histories, studying local law and policy, seeking feedback from clients, reviewing service data, and reflecting on practice. Training is most useful when it connects knowledge to observed clinical decisions and creates opportunities for correction. A one-off workshop cannot guarantee culturally responsive care; nor does a clinician's good intention establish that care was experienced as respectful. Feedback should be gathered through safe channels and acted on visibly.

Applied Exercise: revise a formulation

APPLIED EXERCISE — HYPOTHETICAL CASE

Hypothetical exercise. A therapist writes: "The family is enmeshed; the grandmother interferes with the parents' authority; the adolescent is oppositional and unmotivated." Rewrite this note using observable descriptions and attributed perspectives. Identify at least three plausible alternative explanations for the interaction, including one structural factor and one safety question. Draft two questions that could clarify what each family member wants. Identify a change to the session

> format or service process that could improve participation without presuming the family's preferred arrangement.

A strong revision might state that the grandmother attends most school meetings at the parents' request, that the parents differ about her involvement, and that the adolescent left the last two meetings after being interrupted. The record would identify who reported each point, avoid asserting a single cause, and note that the therapist will ask each member privately about preferences and safety. The exercise illustrates a general principle: precision is not merely stylistic. It reduces the likelihood that an untested cultural assumption becomes an institutional fact.

Summary

Culture in MFT is dynamic, relational, and shaped by power, history, and material conditions. It cannot be reduced to ethnicity, nationality, or a catalogue of group traits. People occupy multiple social positions and may share, contest, or revise cultural meanings within the same family. Intersectional analysis helps explain how interacting systems of power shape experience without treating identity categories as fixed explanations.

Culturally responsive practice combines inquiry with accountability. The therapist examines their own positionality, explains institutional power and confidentiality, attends to unequal power within the family, and negotiates goals and tasks with each participant. Systemic thinking is useful when it reveals interaction and context; it is harmful when circular language disguises coercion or equalises responsibility where power is unequal. Safety assessment, including private inquiry where indicated, remains essential.

Assessment should elicit each member's account, explanatory model, preferred language, goals, resources, constraints, and concerns. Formulations should distinguish observation from inference, include structural conditions, recognise strengths, and remain provisional. Measures and diagnostic frameworks can be helpful when used with attention to validity, language, access, and local context. Interpreters should be qualified for the task, briefed appropriately, and not treated as cultural authorities. Confidentiality, documentation, and records require special care because clinical information may affect domains beyond treatment.

Migration, kinship, race, gender, sexuality, disability, age, class, and geography can shape family roles and service access, but none predicts a single clinical pattern. The clinician asks how these factors matter in this family and follows the answer with action: modifying access, pacing, communication, or intervention when needed. Adaptations should be transparent and evaluated. Research and organisations share responsibility for inclusion, meaningful outcome measurement, equitable access, and accountability for disparities.

Cultural responsiveness is therefore an iterative practice cycle: locate, ask, formulate, negotiate, act, and review. It is measured not by a therapist's claim of mastery but by the quality of inquiry, the fit and safety of care, willingness to revise, and the institution's capacity to learn.

Doctoral Research Questions

1. How do family members within the same household differ in their accounts of culturally meaningful roles, boundaries, and obligations, and how do those differences affect therapy engagement?
2. Which adaptations to systemic interventions improve accessibility or acceptability while preserving the intervention's hypothesised therapeutic functions?
3. How do interpreter-mediated couple and family sessions affect alliance, participation, disclosure, and perceived confidentiality?
4. How do clinicians distinguish relational conflict from coercive control in culturally diverse settings, and what training improves the accuracy and safety of that distinction?
5. Which structural barriers most strongly influence access, retention, and outcomes in community MFT services, and which service-level changes reduce them?
6. How do families define meaningful improvement when standardised outcomes do not adequately represent their priorities or relational arrangements?
7. What is the relationship between therapist reflexivity, client-reported cultural safety, alliance, and treatment outcomes, and how should these constructs be measured?
8. How do transnational caregiving networks influence couple and family stress, resource exchange, and treatment participation?
9. In what ways do diagnostic descriptions and clinical records affect later institutional decisions for families from racialised or marginalised groups?
10. How can community-engaged research distribute decision-making, authorship, data access, and benefit more equitably?

Further Reading

- American Psychiatric Association, *Diagnostic and Statistical Manual of Mental Disorders* (5th edn, text revision, 2022), Cultural Formulation Interview materials.
- Barkan E (ed), *A Cultural Approach to Interpersonal Conflict: A Guide for Working with Diverse Populations* (Routledge 2018).

- Crenshaw K, 'Mapping the Margins: Intersectionality, Identity Politics, and Violence against Women of Color', *Stanford Law Review* 43 (1991).
- Falicov CJ, *Latino Families in Therapy: A Guide to Multicultural Practice* (2nd edn, Guilford Press 2014).
- Hardy KV and Laszloffy TA, *The Cultural Genogram: Key to Training Culturally Competent Family Therapists* (1995 article and subsequent training resources).
- McGoldrick M and Hardy KV (eds), *Re-Visioning Family Therapy: Race, Culture, and Gender in Clinical Practice* (3rd edn, Guilford Press 2019).
- Sue DW, Arredondo P, and McDavis RJ, 'Multicultural Counseling Competencies and Standards: A Call to the Profession', *Journal of Counseling & Development* 70 (1992).
- Tervalon M and Murray-García J, 'Cultural Humility versus Cultural Competence: A Critical Distinction in Defining Physician Training Outcomes in Multicultural Education', *Journal of Health Care for the Poor and Underserved* 9 (1998).
- Williams DR and Mohammed SA, 'Racism and Health I: Pathways and Scientific Evidence', *American Behavioral Scientist* 57 (2013).

KEY THEORETICAL POINT

Practice principle. Ask how identity and context matter in this family; do not assume that a category supplies the answer. Make power, safety, language, and access discussable, then revise the formulation and intervention in response to what family members tell you and what the evidence shows.

End of Chapter 3

MFT. 903 — Multicultural Issues in Family Therapy



PART IV

Clinical Reasoning, Assessment, and the Therapeutic Relationship

Advanced couple and marriage therapy as a disciplined clinical practice: engagement, assessment, safety, formulation, intervention planning, management of complex process, and outcome-informed closure.

4 CHAPTER

Advanced Clinical Practice in Marriage Therapy

LEARNING OBJECTIVES

After studying this chapter, the reader should be able to:

1. Conduct an advanced, balanced assessment of couple distress, relationship strengths, individual factors, context, goals, and safety.
2. Distinguish ordinary relational conflict, high-intensity conflict, coercive control, and intimate partner violence, and select an appropriate service response.
3. Develop a provisional, systemic case formulation that links each partner's account, reciprocal interaction sequences, developmental and social context, and maintaining conditions.
4. Negotiate a treatment contract that defines the unit of care, goals, confidentiality limits, participation, review points, and referral boundaries.
5. Select and adapt an evidence-informed treatment approach in response to the couple's needs, preferences, risks, resources, and clinical progress.
6. Monitor alliance with both partners, manage imbalance and rupture, and respond competently to escalation, disengagement, disclosure, and impasse.
7. Evaluate progress using multiple indicators, distinguish change from symptom suppression or compliance, and plan continuation, referral, or ending collaboratively.
8. Demonstrate ethical, culturally responsive, reflective, and professionally accountable practice in complex couple therapy.

Introduction

Advanced clinical practice in marriage therapy is not defined by the number of techniques a clinician can name. It is demonstrated in the capacity to make sound decisions when partners offer competing accounts, goals shift, risk is uncertain, a familiar intervention is not helping, or the therapist's own responses begin to shape the work. The clinician must sustain attention to two people, the relationship between them, and the wider conditions that influence their options,

without reducing either partner to a symptom, a role, or a presumed cause of the other's distress.

The course title uses *marriage therapy*, but this chapter treats the term as a professional tradition rather than a restriction to legally married heterosexual couples. The clinical principles apply to diverse committed relationships and couple configurations; clinicians should use the language the partners prefer and remain attentive to jurisdictional differences in legal recognition, service eligibility, and professional scope. The focus here is advanced practice: clinical reasoning, assessment, formulation, treatment planning, risk management, alliance, process monitoring, and closure. Detailed protocol-level skills for particular intervention models are developed in the separate course on couples therapy techniques.

Couple therapy is a distinctive treatment context. Partners enter with different histories, explanatory models, degrees of hope, and accounts of what should change. One may seek restoration of closeness while the other is considering separation; one may view the problem as betrayal and the other as chronic disconnection; one may attend voluntarily while the other experiences pressure from a partner, court, family, or service. The clinician cannot assume that "the couple" has a single goal or that both members experience the treatment setting as equally safe. The first task is therefore not to impose a shared narrative but to create a process in which distinct accounts can be heard, differences can be assessed, and next steps can be negotiated responsibly.

This chapter moves through the clinical arc. Section 4.1 defines the advanced therapist's stance and the scope of couple care. Section 4.2 addresses engagement and the treatment frame. Sections 4.3 and 4.4 examine assessment, safety, and formulation. Section 4.5 develops collaborative treatment planning and model selection. Sections 4.6 and 4.7 address alliance, high-intensity process, rupture, and clinical impasse. Section 4.8 examines progress review, referral, and ending. Section 4.9 integrates these tasks in a hypothetical case and applied exercise. Across the chapter, the therapist is asked to keep four questions active: What is each partner asking for? What pattern is occurring between them? What contextual or safety conditions change the meaning of that pattern? What evidence would show that the work is helping?

The discussion is academic and professional, not devotional. Clinical decisions should be guided by current evidence, applicable ethics and law, professional competence, and the couple's goals and safety. No single model is suitable for all couples, and no stated commitment to neutrality removes the clinician's responsibility to respond when coercion, violence, abuse, or serious impairment is present.

4.1 The Advanced Couple Therapist's Clinical Stance

Holding two accounts without collapsing them

The couple therapist hears at least two accounts of the relationship, each shaped by memory, expectation, emotion, and present interaction. The accounts may contradict one another on observable events, but disagreement does not automatically indicate deception. Partners may encode events differently, remember different details, or assign different meanings. The clinician should neither force premature agreement nor treat both accounts as equally reliable in every domain. A balanced stance means procedural fairness—asking each person, checking understanding, and making reasoning transparent—not the suspension of judgement about safety or evidence.

A therapist can reflect a partner's experience without endorsing every interpretation: "You experienced the silence as a refusal to engage; I want to understand what it meant for you, and also how your partner understands that moment." This allows the clinician to stay curious while keeping the sequence observable. When partners dispute what happened, it can be helpful to shift from global accusation to specific event, context, action, impact, and response. This reduces the chance that the session becomes an adjudication of character and creates material for a relational formulation.

The therapist must also resist becoming the couple's judge, advocate for the more articulate partner, or messenger between partners outside the agreed frame. A partner may ask the therapist to confirm that the other is "the problem". A clinically useful response is to name the therapist's task: understand how each person's actions and the relationship context are connected, while not assuming equal responsibility where there is evidence of coercion or abuse. The clinician's position is neither automatic equivalence nor reflexive alliance with the first account heard.

The couple as a unit of care: a negotiated decision

Couple therapy often treats the relationship as the focus, but both partners remain individual people with distinct rights, needs, and safety concerns. The unit of care must be made explicit: Is the service treating the relationship, offering conjoint assessment, providing individual support to each partner, or coordinating couple therapy with other services? These arrangements have different implications for records, confidentiality, communication, and what the clinician can promise. The therapist should state the model of care plainly and revisit it if the work changes.


The partners may not share the same definition of success. One may wish to improve communication; another may want clarity about whether to stay. Some couples seek help to separate with less harm; others wish to repair after a major

rupture. The therapist should not assume that preservation of the relationship is the only legitimate outcome. Treatment goals may concern safety, respectful decision-making, reduced conflict, co-parenting, separation planning, or a mutually chosen attempt at reconciliation. The ethical task is to help partners make informed decisions, not to secure a particular relationship status.

Advanced practice as disciplined flexibility

Clinical flexibility is not improvisation without structure. It is the ability to maintain a clear rationale while adapting the pace, format, and intervention to the couple's responses. The therapist tracks several streams at once: each partner's emotional and behavioural response, the interaction between partners, the alliance with each person, the fit of the treatment model, the effect of the setting, and any risk indicators. These streams can compete for attention. A partner's escalating distress may require slowing down the exploration of a relational cycle; a disclosure may require pausing joint discussion to assess safety; a rupture with the therapist may need repair before further technique is useful.

Advanced clinicians make their uncertainty visible. A formulation is a working hypothesis, not a verdict. A treatment plan is a proposal that should be reviewed, not a guarantee. Where evidence is limited or the therapist's competence is insufficient, the appropriate action may be consultation, supervision, referral, or a transparent discussion of alternatives. Knowing when not to proceed with conjoint therapy can be as important as knowing how to conduct it.



A non-linear clinical arc: new information or risk may return the work to assessment or contracting.

Figure 4.1 Advanced couple therapy proceeds through engagement, assessment and safety, shared formulation, negotiated intervention, outcome review, and a deliberate decision to continue, adapt, refer, or end. Review points loop back when new information emerges. *Source:* prepared for this edition.

4.2 Engagement, Contracting, and the First Sessions

Engagement before intervention

The first contact begins treatment. Referral language, scheduling, intake forms, and contact with each partner can communicate whose perspective the service values. If one partner makes the enquiry, the clinician should obtain appropriate contact and consent from the other without allowing the initiating partner to control all communication. When there are privacy or safety concerns, individual contact procedures should be planned carefully and should not inadvertently disclose attendance or a private message. The service must explain its communication channels and limits.

An initial appointment should establish what brings the couple now, what each partner wants from the meeting, and what would make the process safe enough to begin. The clinician describes their role, training, approach to confidentiality, record-keeping, cancellation, fees or service access, communication between sessions, and how urgent risk is handled. These matters should be conveyed in accessible language and checked for understanding. If partners have different language needs, arrange qualified interpretation and clarify how each person can participate. Do not rely on one partner to interpret sensitive material for the other.

The therapist can invite each person to speak without requiring equal time as a mechanical rule. Useful questions include: "What is most important that I understand today?" "What would you want to be different if these meetings were useful?" "What worries you about starting therapy?" and "Is there anything about meeting together that does not feel safe?" The final question should not be asked only in the presence of a partner if there is reason to suspect coercion. Clinicians should create a clear route for confidential, separate communication and explain how such information will be handled within the treatment contract.

A workable treatment contract

The treatment contract clarifies what the therapist can and cannot provide. It should address: the intended focus of treatment; the relationship between conjoint and individual meetings; who is considered a client; the approach to secrets and private communications; limits of confidentiality; the record and access to it; how safety concerns will be managed; participation and decision-making; expected frequency and review points; fees or service conditions; and circumstances for referral or ending. Legal requirements vary, and the contract must be adapted to local law, professional standards, and the service setting.

A "no-secrets" policy is sometimes used in couple therapy, but it is not a universal ethical solution. A rigid policy can discourage disclosure of safety concerns or create pressure to reveal information prematurely. A completely unstructured

promise of individual confidentiality can create a covert alliance or make conjoint treatment impossible if clinically important information is withheld. The therapist should select and explain a policy appropriate to their role, competence, and jurisdiction before private disclosures occur. If the policy changes or becomes untenable, the clinician should revisit the contract and consider consultation or referral rather than improvising a response after a secret has been disclosed.

The clinician should also discuss how therapy will respond if partners decide to separate. The therapist may support decision-making and reduce harmful conflict, but should not assume the role of mediator, custody evaluator, legal adviser, or advocate for either partner unless specifically trained, contracted, and authorised to do so. Multiple roles can compromise trust and impartiality. Clarify referral options for legal, financial, parenting, or individual mental-health needs.

Managing hope, ambivalence, and unequal motivation

Partners often begin with unequal motivation. One may be hopeful, one sceptical, one ambivalent, or one primarily complying with a demand. The therapist should not treat reluctance as pathology or pressure the less willing partner into a shared goal. Explore what each would consider a worthwhile use of time, even if their broader aims differ. An early session can establish a modest, observable aim —such as being able to discuss a difficult topic without intimidation or leaving with a clearer decision process—while the partners consider whether longer-term therapy is appropriate.

Hope should be realistic. Avoid guarantees that therapy will "save" a marriage, resolve betrayal, or prevent separation. The clinician can explain that treatment may help partners understand patterns, make decisions, improve selected behaviours, or coordinate a safe separation, but outcomes vary. If one partner has decided to leave, a discernment-oriented process may be more suitable than standard conjoint treatment; therapists should practise only within their competence and explain the distinction. If partners have a shared wish to stay and work on the relationship, treatment can focus more directly on relationship change.

Accessibility and inclusion in engagement

Engagement is affected by transport, work schedules, childcare, disability access, cost, digital access, immigration or legal concerns, and prior experience of services. A missed appointment may reflect any of these factors, or a decision not to continue; the therapist should ask rather than infer. Offer reasonable accommodations where possible, including interpretation, flexible scheduling, accessible rooms, remote options, or referral to a more suitable service. Remote conjoint work has privacy and safety limitations that must be assessed, particularly when partners are joining from a shared home.

The first session also establishes whether the clinician can provide competent care. If the couple's needs involve violence, acute psychiatric instability, active substance-related risk, severe trauma, or another condition outside the therapist's scope, the therapist should consult, coordinate, or refer as indicated. A referral is not a rejection. It should be explained as a step toward appropriate and safe support, with attention to continuity rather than abrupt abandonment.

4.3 Comprehensive Couple Assessment

Multiple perspectives and the relationship timeline

A comprehensive assessment integrates each partner's account with observation of interaction, relevant history, current context, and available records or collateral information where consent and law permit. Neither a joint interview nor separate interviews alone are sufficient for every case. Joint conversation shows how partners engage each other in the present, but may inhibit disclosure. Separate conversations may improve privacy and safety, but can create challenges around information-sharing and perceived alliances. The therapist should explain the purpose and boundaries of each format, and avoid promising that a private session will remain confidential if the treatment contract does not allow it.

Start with the relationship's current concern and the path to treatment. Ask how long the difficulty has been present, what preceded the referral, what makes it better or worse, and why help is being sought now. Elicit each partner's account of key events, including where their accounts converge and diverge. The timeline may include courtship, commitment, cohabitation, marriage or other chosen milestones, parenting, illness, work transitions, relocation, caregiving, betrayal, bereavement, and previous treatment. These events do not mechanically cause current distress; they provide context for understanding the couple's meanings, expectations, and adaptations.

Assess relationship strengths as carefully as problems. Ask what the partners appreciate, how they have coped with previous adversity, what still works, and when they feel more connected or coordinated. Strengths should be described concretely rather than used to minimise harm. A history of mutual support does not negate a current safety concern, just as severe conflict does not erase a couple's capacities. Elicit the social network, extended-family relationships, community resources, and practical constraints that may support or limit change.

Domains to assess

A systematic assessment can consider the following domains, adjusting depth to the referral question and clinical need:

- **Presenting problems and priorities:** distress, relationship satisfaction, recurring disputes, individual concerns, and each partner's preferred outcomes.
- **Interactional process:** how disagreement begins, what each partner notices, emotional and behavioural responses, escalation or withdrawal, repair attempts, and the consequences of the sequence.
- **Commitment and decision status:** hopes for change, ambivalence, possible separation, and any external pressure to attend.
- **Attachment, development, and learning history:** experiences that may shape expectations of closeness, trust, conflict, and care, explored without treating history as destiny.
- **Sexual and affectionate relationship:** only as relevant and with consent; ask about desire, intimacy, pain, safety, boundaries, and differences without presuming a normative frequency or form.
- **Parenting and caregiving:** co-parenting, division of labour, child impact, fertility or reproductive decisions where relevant, and care for older or disabled relatives.
- **Individual and health factors:** mood, anxiety, trauma, neurodevelopmental differences, medical conditions, medication, substance use, sleep, and functional impairment, with referral when assessment exceeds the therapist's scope.
- **Social and cultural context:** identity, language, discrimination, migration, economic circumstances, family structure, and institutional involvement.
- **Risk and protection:** violence, coercion, self-harm, harm to others, child or elder safety, stalking, access to weapons where clinically relevant, and protective resources.
- **Treatment history and preference:** previous therapy, what helped or harmed, desired approach, accommodations, and concerns about confidentiality.

Assessment should not become a questionnaire marathon. The therapist prioritises domains according to the couple's concerns and risk profile, explaining why sensitive questions are asked. Standardised measures can support baseline description and monitoring, but should be selected for validity, accessibility, and relevance. A measure does not replace a direct safety interview or a careful discussion of the partners' goals.

Observing interaction without staging a test

The therapist learns from how partners speak to one another in the session, but the observation is shaped by the setting. A couple may behave differently when being watched, when one partner controls transport or finances, or when an audi-

ence is present. The therapist should not deliberately provoke an intense conflict simply to observe it. Instead, invite discussion of a manageable topic and monitor whether each partner can pause, listen, express disagreement, and respond to cues. Observations should be described in behavioural terms: interruption, topic shift, raised voice, silence, checking for agreement, or repair. Interpretations such as "contemptuous" or "avoidant" should be tested with the partners and considered alongside context.

A sequence map can focus clinical attention: what happened immediately before the conflict, what each person did, what the other inferred, what response followed, and how the interaction ended. Ask each partner what they noticed internally and what they hoped the other would understand. This creates a shared object of inquiry without asserting that both are equally responsible for the pattern. If one partner is afraid, coerced, or constrained, the interaction must be interpreted in relation to that power difference.

Individual and relational assessment

Couple distress and individual mental-health concerns can influence one another. Depression, anxiety, trauma symptoms, chronic pain, sleep disruption, or substance use may affect communication, availability, and conflict; relationship distress may also exacerbate individual symptoms. The clinician should assess whether an individual condition requires concurrent or prior treatment, and avoid attributing it solely to relationship dynamics. When an individual assessment or diagnosis is needed, use appropriate professional procedures and make clear whether the information is part of couple treatment or a separate clinical service.

A therapist must be alert to the limits of their expertise. Screening tools can identify the need for further evaluation but are not diagnoses. Acute suicidal risk, mania, psychosis, severe substance withdrawal, cognitive impairment, or serious medical concerns may require urgent assessment by qualified professionals. The couple therapist should know local referral and emergency pathways and should not attempt to contain a crisis solely within conjoint sessions.

	Clinical question	Evidence to gather
Each partner	What matters to each person?	Individual goals, meanings, hopes, concerns
Interaction	What happens in the recurring sequence?	Specific actions, interpretations, responses, repair
History	What has changed across the relationship?	Transitions, treatment history, exceptions
Individual	What health or functioning factors matter?	Symptoms, health, substance use, referral needs
Context	What shapes options and access?	Work, caregiving, culture, resources, institutions
Safety	Can both participate freely?	Private inquiry, coercion, risk, protective plan

Figure 4.2 A balanced assessment moves among each partner's account, the relationship sequence, strengths and history, current context, and safety; the clinician selects depth according to relevance and risk. *Source:* prepared for this edition.

4.4 Safety, Coercion, and Suitability for Conjoint Work

Conflict, violence, and coercive control

Not all couple conflict is equivalent. Partners may engage in mutual disagreement, high-intensity arguments, situational violence, or patterned coercive control. A clinician must distinguish these phenomena as carefully as available information allows. **Coercive control** involves a pattern of domination that restricts another person's autonomy through intimidation, surveillance, isolation, threats, economic control, or other tactics; it is not defined simply by disagreement or by a single heated exchange. Intimate partner violence can include physical, sexual, psychological, and other forms of harm, and may occur in any gender or relationship configuration.

A symmetrical "negative cycle" formulation is unsafe if it converts coercion into reciprocal miscommunication or implies equal responsibility. The person using coercive behaviour is responsible for that behaviour. The therapist should assess the pattern, impact, fear, freedom to disagree, and consequences of disclosures, rather than relying only on whether both partners report "arguments". Direct but non-leading questions may include: "Has either of you felt afraid of the other?" "Has anyone been prevented from leaving, contacting others, or accessing

money?" "What happens after a disagreement?" "Would it be safe for you to speak honestly here, and what might happen after the session?" Questions should be asked privately where possible, with an explanation of confidentiality limits.

Screening and safety planning

A written instrument may support but cannot replace a private clinical conversation and professional judgement. The therapist should clarify whether the information will be kept in a couple record, who may access it, and what circumstances require action. Screen for immediate danger, escalation, threats, stalking, strangulation, sexual coercion, child exposure, access to lethal means where relevant, and the person's own assessment of risk. These are sensitive questions that require training; the clinician should consult current, jurisdiction-specific guidance and specialist services rather than assume a brief checklist is sufficient.

Safety planning is collaborative and should not prescribe a single path. It may include a safe contact method, emergency contacts, access to documents or medication, a plan for leaving a session safely, specialist advocacy, child protection consultation, or referral. Do not leave written information or voicemail in a way that could increase danger. The therapist should document objectively, consult as needed, and follow applicable law and safeguarding duties. The clinician should not confront a suspected perpetrator with confidential information without considering foreseeable risk.

Deciding whether to proceed conjointly

Conjoint therapy may be inappropriate or unsafe when one partner cannot speak freely, there is active coercive control, severe violence, fear of retaliation, or a significant imbalance that the treatment setting cannot safely manage. Other factors—active psychosis, uncontrolled substance-related impairment, acute suicidality, or severe instability—may require individual stabilisation or specialised care before couple work. These are not automatic exclusions based solely on a diagnostic label; the decision depends on current severity, safety, consent, treatment focus, available expertise, and coordination with other providers.

When conjoint work is not indicated, explain the reasoning without blaming the person at risk. Offer referral to specialist services, individual treatment, advocacy, or other appropriate support. If conjoint therapy proceeds despite a history of harm, it should do so only under a defensible plan, with specialised competence, ongoing private safety assessment, clear limits, and a willingness to stop or change course. A therapist should not promise that the office is a safe space simply because it is professionally supervised.




Figure 4.3 A safety decision pathway begins with private inquiry and assessment of freedom and risk, then weighs suitability, expertise, and protections before conjoint work; risk is reviewed throughout rather than screened once. *Source:* prepared for this edition.

Safety in diverse relationships

Clinicians should not assume that a particular gender, body, or relationship role determines who is at risk or who uses violence. They should also recognise that legal protections, community resources, and social credibility may be unevenly available. A partner may fear disclosure because of prior discrimination, outing, immigration consequences, or mistrust of authorities. Ask about the client's own concerns and seek specialist consultation without treating identity as evidence of risk. Use inclusive language and attend to the interactional pattern, access to resources, and concrete behaviour.

Risk assessment must include child and dependent-person safety when relevant. Children may witness conflict or be drawn into adult disputes. The therapist should ask how children are affected, what they understand, and what protective steps are needed, while following local safeguarding duties. A child should not be used as a messenger or mediator between parents. If the service is court-involved or mandated, clarify the clinician's role and limits; couple therapy is not a substitute for forensic assessment.

4.5 Formulation: From Assessment to a Shared Map

A systemic formulation with individual specificity

A clinical formulation organises relevant information into a provisional account of how the difficulty arose, what maintains it, what protects against it, and what might support change. It is not a diagnosis of the relationship, a summary of every fact, or a technique-selection chart. A sound formulation integrates each partner's account, the interactional sequence, life history and current transitions, individual and health factors, cultural and structural context, resources, and safety. It makes uncertainty explicit and identifies what information is still needed.

The clinician can develop the formulation through five linked questions:

1. **What is the problem as each partner defines it?** Identify shared concerns, distinct priorities, and disagreements over the meaning of events.
2. **What happens between partners when the problem appears?** Describe observable sequences, triggers, interpretations, responses, and short- and long-term consequences.
3. **What individual, developmental, and contextual factors influence the sequence?** Consider learning history, health, work, caregiving, social position, and transitions without treating any one factor as a complete cause.
4. **What resources and exceptions are available?** Identify times when the couple handles the issue differently, supportive relationships, shared values, practical capacities, and individual strengths.
5. **What risks or constraints change the treatment options?** Clarify coercion, violence, acute mental-health needs, access barriers, confidentiality, and limits of competence.

The map should capture partner-specific meanings. One person may escalate to obtain a response; the other may withdraw to reduce overload; each response can intensify the other's expectation. This cycle may be a useful account when both partners recognise it and there is no coercive pattern that makes symmetry misleading. If one partner threatens, monitors, or punishes the other, the formulation must name those behaviours and their effects rather than translating them into a mutually maintained "cycle".

Differentiating precipitating, perpetuating, and protective factors

A practical formulation distinguishes **precipitating factors**—events associated with the recent onset or worsening—from **perpetuating factors** that sustain the present difficulty and **protective factors** that reduce harm or support recovery. These categories are not always cleanly separable. A job loss may precipitate conflict, economic uncertainty may perpetuate it, and a supportive relative may protect the couple from isolation. The therapist should avoid presenting correlations as established causes. Formulation is a hypothesis to guide observation and

intervention; its value is judged by whether it helps the couple understand the problem, identifies feasible change, and remains open to correction.

The clinician should include the couple's preferred account of meaning alongside professional concepts. If the therapist uses terms such as pursue-withdraw, attachment threat, negative attribution, or demand-avoid, explain them in accessible language and ask whether they fit. Technical language can help organise clinical reasoning, but can also make the therapist's interpretation seem incontestable. A formulation should be concise enough to guide action and specific enough to distinguish the case from a generic description of couple distress.

Sharing a formulation without imposing it

A collaborative formulation is offered as a proposal: "I have a possible map of what happens, and I want to check it with each of you." The therapist can state what is observed, identify the hypothesised sequence, and ask each partner what fits or does not. Disagreement can refine the account. The clinician should not insist that partners accept a formulation because it is theoretically elegant. Nor should the therapist avoid offering a clinical perspective altogether; collaboration includes the professional's responsibility to provide a clear, reasoned view.

If partners do not agree on a joint formulation, treatment may still proceed with multiple parallel hypotheses and shared behavioural goals. For example, one partner may see a conflict as a response to neglect and the other as an effort to prevent criticism. They may nevertheless agree to pause an argument when voices rise and return to the topic at a set time. A shared plan does not require identical explanations, provided that the plan is voluntary and safe.

4.6 Treatment Planning and Selecting an Approach

From goals to a treatment contract

Treatment planning translates the assessment and formulation into goals, tasks, time frames, and criteria for review. Goals should be meaningful to both partners, but need not be identical. Distinguish **relationship goals** (such as improved emotional connection), **process goals** (such as discussing conflict without intimidation), and **decision goals** (such as clarifying whether to continue the relationship). Where separation is possible, state this without treating it as treatment failure. Goals should be specific enough to guide sessions and flexible enough to adapt as circumstances change.

A treatment plan can identify: the principal concerns; each partner's goals; the agreed focus of couple work; any individual or specialist services required; the initial model or intervention rationale; session frequency and likely duration; indicators of progress; safety procedures; and the date or conditions for review. The plan should also state what would prompt a change—worsening safety, declining engagement, symptom escalation, no meaningful progress, new information, or

a change in the couple's decision status. The therapist should obtain informed agreement and document revisions.

A treatment plan is not a promise of a fixed number of sessions or an obligation for partners to remain together. It is a provisional agreement about how therapy will be conducted. Review the goals periodically, and check privately where appropriate whether each partner remains willing and able to participate. A partner's continued attendance does not necessarily mean consent to the process if there is pressure or fear.

Evidence-informed model selection

Several couple-therapy traditions have credible clinical frameworks and varying degrees of empirical support, including behavioural and cognitive-behavioural approaches, integrative behavioural couple therapy, emotionally focused couple therapy, insight-oriented approaches, and systemic or integrative models. No approach should be selected solely because it is the therapist's preferred school. The clinician considers the couple's goals, presenting concerns, risk, treatment history, preferences, developmental and cultural context, access needs, and the evidence base relevant to the population and outcome. A therapist may use an integrative plan, but should be able to explain its rationale and avoid incoherent eclecticism.

Behavioural and cognitive-behavioural approaches may be appropriate where partners seek structured work on interaction, problem-solving, communication, behavioural exchange, and interpretations. Emotionally focused approaches may be relevant where partners identify recurring attachment-related disconnection and both can participate safely in emotionally engaged work. Insight-oriented work may help partners examine patterns linked to personal history and relational expectations. Systemic approaches can focus on interaction sequences, wider context, and how attempted solutions maintain difficulty. These descriptions are broad, not manual instructions; model-specific procedures require training and supervision.

Evidence-informed practice combines research, clinical expertise, and the couple's preferences and circumstances. Average treatment effects do not guarantee an outcome for a particular couple. The therapist should be transparent about what is known, discuss reasonable alternatives, and use outcome monitoring to identify whether the selected approach is helping. If the couple's situation falls outside the evidence base, the clinician should acknowledge uncertainty rather than present a model as proven for that case.

Sequencing and dose

A common sequence is engagement and assessment, shared formulation and goal setting, active intervention, review and consolidation, and closure or transition. It is not rigid. Safety may require delaying conjoint exploration; new disclosures

may return the therapist to assessment; a lack of progress may require revising the formulation; a couple may complete a decision-focused process earlier than expected. The number and spacing of sessions should match need, accessibility, service conditions, and evidence where available. Too little treatment may not allow meaningful change; an indefinite schedule without review can foster dependency or avoid difficult decisions.

Session structure can support fairness and focus. A clinician may begin with a brief check-in from both partners, identify the agreed agenda, work on one interactional issue, and end with a summary and feedback. This format is not universal, but helps prevent sessions from being taken over by the faster or more forceful speaker. The therapist can adapt use of written agendas, turns, pause signals, or separate check-ins if these improve safety and participation. Any structure should be explained and evaluated with the couple.




Figure 4.4 A formulation-to-plan pathway links each partner's account and the interactional pattern to goals, model choice, agreed tasks, and review criteria; safety and context remain active at every stage. *Source:* prepared for this edition.

Individual sessions within couple treatment

Separate meetings can support assessment, safety, or focused individual work, but they change the clinical frame. Before offering them, clarify their purpose, duration, information-sharing policy, and how the therapist will prevent a perceived alliance with one partner. If private sessions reveal information that cannot be held within the agreed policy or that makes joint work unsafe, the therapist may need to pause, renegotiate, or refer. The clinician should not invite a disclosure without a defensible plan for responding to it.

Where a partner needs individual therapy, the couple therapist may coordinate with a separate provider with appropriate consent. Dual roles can complicate con-

fidentiality, objectivity, and treatment goals. The same clinician providing both individual treatment and couple therapy may be inappropriate in some circumstances or jurisdictions. If unavoidable, the clinician should seek supervision and use a clear written agreement, while considering whether separate practitioners better protect each person's interests.

4.7 Managing the Therapeutic Process

Alliance with each partner and between partners

Alliance in couple treatment is multidimensional. The therapist needs a working relationship with each partner and a workable process between them. A strong alliance with one person can coexist with mistrust from the other; a couple may be warm toward the clinician while avoiding the agreed work. Therapists should monitor each partner's sense of being heard, the clarity of goals and tasks, and whether the session structure feels safe. Short, direct feedback questions may help: "Did today's conversation feel balanced?" "Was there a point where I moved too quickly?" "Is there something you could not say in this format?" Responses should lead to discussion, not be filed away as a score.

Therapist impartiality is not identical to treating every action as equivalent. The clinician can respect both partners while naming a specific harmful behaviour and protecting the person affected. Avoiding "taking sides" should not become a reason to avoid responsibility or safety. If there is no coercion or abuse, the therapist can help partners recognise their respective contributions to a reciprocal pattern. If there is coercion, the therapist should not use symmetry as a clinical shortcut.

De-escalation and emotional regulation

When interaction intensifies, the clinician's immediate task may be to regulate the session rather than pursue insight. Indicators include rapid turn-taking, repeated interruption, raised voices, contemptuous language, visible shutdown, or a partner's inability to remain oriented to the conversation. The therapist can pause, name what is happening without accusation, check each person's state, and re-establish the agreed ground rules. A break is not automatically a failure; it can prevent harm and create a chance to return more deliberately.

De-escalation should not require the less powerful partner to soothe the more powerful one or accept a compromise to restore calm. The therapist should assess whether the response is fear-based, whether a pause is safe after the session, and whether joint work remains indicated. When appropriate, a structured turn-taking exercise can help partners articulate one concern at a time, but it should not be imposed when a partner cannot speak freely. Clinical techniques are always subordinate to safety and consent.

Rupture, misattunement, and repair

Ruptures can occur when a partner feels blamed, unseen, exposed, or pressured; when the therapist aligns too quickly with one account; or when an intervention misses the couple's meaning. A rupture may be explicit—"You are taking their side"—or indirect, through withdrawal, sarcasm, reduced participation, or missed appointments. The clinician should consider whether the rupture reflects a misunderstanding, a mismatch in goals, a harmful clinical action, or a legitimate challenge to the therapist's frame.

A repair process can include noticing the shift, naming the specific moment, asking how it affected each person, taking responsibility for the therapist's contribution where appropriate, and renegotiating the next step. Defensive explanations often deepen the rupture. At the same time, the therapist need not agree with every accusation; they can acknowledge impact while clarifying intention and inviting further discussion. Repair is demonstrated by changed practice, not by the therapist's apology alone.

Stalemate and impasse

Couples may repeat the same argument in therapy, refuse a proposed exercise, disagree about the problem, or appear to make progress and then return to an earlier pattern. The clinician should treat impasse as information, not automatically as resistance. Possible explanations include an inaccurate formulation, incompatible goals, unaddressed safety, practical barriers, a treatment model mismatch, therapist reactivity, external stress, or a pace that is too fast or too slow. The therapist can ask each partner what makes the next step difficult and what they fear would happen if change occurred.

A structured case review can help: restate the treatment goal; check whether both partners still consent; review progress and risk; identify the therapist's assumptions; consider alternative formulations; consult supervision; and decide whether to continue, adapt, pause, or refer. Consultation is especially important when the therapist feels pressured to prove a theory, rescue the relationship, punish a partner, or avoid a difficult disclosure. Self-awareness is not sufficient on its own; use supervision and objective information where possible.

Therapist use of self, bias, and supervision

The therapist's emotional responses can be clinically informative, but they are not direct evidence about either partner. A wish to rescue one person, irritation with another, boredom during repeated conflict, or urgency to prove that the treatment model works may signal something about the interaction, the clinician's history, fatigue, or the service context. The therapist should pause before converting a reaction into a hypothesis. Ask what was happening immediately before the response, which partner's account was being heard, whether a safety issue was

minimised, and what alternative explanations are possible. Use supervision to test the formulation rather than recruit a supervisor to confirm a preferred side.

Supervision is especially important when there is unequal power, risk, legal involvement, strong model allegiance, or therapist uncertainty about the confidentiality contract. A useful presentation includes the referral question, each partner's stated goals, a brief sequence map, safety assessment, the therapist's intervention and response, and the specific decision that needs consultation. Where session recording is considered, obtain informed consent from both partners, explain access and storage, and allow either person to decline without penalty where feasible. Recordings should be used only for the purpose agreed and managed under applicable data-protection rules.

Therapists should monitor their own process without making clients responsible for it. If the clinician notices a strong pull to align with one partner, the response is not to force artificial balance but to examine the basis of the pull and return to observable behaviour, impact, and risk. If the therapist has acted unfairly, repair should be specific. If the clinician cannot maintain a useful position despite consultation, transfer or referral may be preferable to continuing in a compromised role. Professional accountability includes recognising when the therapist's presence is no longer helpful.

Rupture after a significant disclosure

A disclosure of infidelity, undisclosed debt, sexual boundary violation, or another major event can reorganise the therapeutic frame. The clinician should assess what the disclosure means to each partner, whether there is current risk, what each person wants to know or not know, and whether the existing confidentiality agreement applies. The therapist should not use a disclosure as a dramatic intervention or pressure a partner into immediate forgiveness, confrontation, or separation. If disclosure changes the feasibility of conjoint work, explain why and discuss alternatives.

Some couples seek therapy after a breach of trust. Treatment may need to address the meaning of the event, the injured partner's safety and agency, the accountable partner's responsibility, and the conditions under which rebuilding trust is possible. "Forgiveness" or reconciliation should not be imposed as a required outcome. The therapist should not present trust repair as a simple sequence that guarantees restoration. Partners may choose different outcomes, and therapy should support informed, safe decision-making.

4.8 Monitoring Progress, Referral, and Ending

Multiple indicators of change

Progress is not a single score or a therapist's impression. It can include reduced distress, safer disagreement, improved repair, greater clarity, increased emotional

accessibility, changed division of labour, more effective co-parenting, or a decision made with less coercion. Partners may define success differently. A clinician should identify baseline indicators with each person and review them regularly. Brief validated measures can track relationship distress or individual symptoms when appropriate, but behavioural examples and client feedback add context.

A review can ask: What has changed since we began? What is still difficult? What did each person do that contributed? Are there goals that no longer fit? Has safety changed? Is therapy helping enough to justify continuing? Measure results should be discussed in plain language and not treated as proof that the couple is or is not trying. If one partner reports improvement and the other does not, the divergence itself is clinically relevant. The clinician can explore whether goals differ, effects are uneven, or a measure misses an important domain.

Routine progress monitoring can support responsive care when it includes feedback, shared interpretation, and the option to revise treatment. It can become counterproductive if used as surveillance, a condition of service, or a substitute for dialogue. If progress is limited, revisit fit, alliance, formulation, dose, external constraints, and treatment goals before concluding that the couple is resistant or untreatable. Consider whether individual or specialist care is needed.




Figure 4.5 Progress review combines the couple's own goals and accounts with observed interaction, safety, relevant measures, and contextual change; results guide a shared decision to continue, adapt, refer, or end. *Source:* prepared for this edition.

Referral and coordination

Referral is indicated when the couple needs expertise or support outside the clinician's scope, when risk cannot be managed in the current setting, when individual stabilisation is needed, or when the therapist cannot maintain an appropriate therapeutic position. A referral should be specific and collaborative: explain the reason, identify an appropriate service, obtain consent for communication where required, and make a plan for continuity. If urgent risk exists, follow local emergency and safeguarding procedures.

Coordination with individual therapists, medical providers, legal professionals, or social services requires clear consent and attention to role boundaries. Couple therapy must not become an informal forensic assessment, custody evaluation, or legal opinion. If a court or employer requests information, the therapist should clarify the request, review law and policy, seek consultation, and provide only information within the scope of the role. Do not promise absolute confidentiality where legal exceptions apply.

Ending therapy

Ending may occur because goals have been met, partners have decided to separate, the work is no longer useful, a referral is needed, the service ends, or one or both partners withdraw. Planned ending includes summarising what has changed, identifying strategies and resources the couple can use, discussing foreseeable stressors, and clarifying how to re-access support if needed. The therapist should ask what ending means to each partner and whether there are unresolved safety or practical concerns.

A couple may have mixed feelings about ending. The clinician can acknowledge these without fostering dependency or abruptly disengaging. If treatment is ending despite lack of progress, review the reasons honestly and identify alternatives. A therapist should not extend treatment merely to avoid disappointment or end prematurely because the couple's choices differ from the clinician's hopes. When partners decide to separate, therapy may focus on respectful communication, co-parenting, or transition planning if safe and within competence; otherwise refer to appropriate services.

4.9 Integrating the Clinical Arc

Hypothetical Case Illustration: pursuing, withdrawing, and the need for reassessment

> **CASE ILLUSTRATION**
>
> **Hypothetical case — couple distress and changing safety information.** Two partners in a long-term relationship seek therapy after repeated arguments about

finances and time together. In the first joint session, one partner speaks rapidly and asks for a plan; the other is quiet and agrees to attend. The therapist initially notices a pursue-withdraw sequence: one partner presses for discussion, the other becomes silent, and the first partner escalates. During separate safety check-ins, the quieter partner reports that disagreement about money can lead to threats to restrict access to a shared account and that they fear what might happen after the session. The therapist does not proceed with a communication exercise or describe the pattern as mutually maintained. They review confidentiality and safety, consult a specialist supervisor, and arrange a safe referral and follow-up plan. The more vocal partner is not confronted with private information without a risk plan. The clinician documents the specific account, the steps taken, and the limits of the current service. Conjoint work is paused because the conditions for free participation are not established.

The hypothetical case shows why formulation must remain provisional. The initial interactional hypothesis was plausible but incomplete. New information changed the clinical question from "How can the couple communicate more effectively?" to "Can each partner participate freely and safely, and what response is required?" The therapist's task is not to defend the original model but to revise practice when evidence changes. The case also demonstrates that a referral can be the appropriate clinical outcome of an assessment, not a failure to provide therapy.

Applied Exercise: construct and review a treatment plan

APPLIED EXERCISE — HYPOTHETICAL CASE

Hypothetical exercise. A couple reports recurring conflict about caregiving and household labour. One partner wants more appreciation and the other wants a fairer distribution of tasks. They each report that the other interrupts, and there is no disclosure of fear or violence during the initial meeting. Draft a one-page provisional formulation that includes both accounts, one observable interaction sequence, one contextual factor, at least two strengths, and the safety questions that remain. Then write three measurable goals, a proposed initial treatment approach, one progress indicator each partner can report, and two conditions that would require reassessment or referral. State how you would check whether the session process is balanced and safe.

The exercise has no single correct model. A behavioural plan may focus on observable task allocation and collaborative problem-solving; an emotion-focused plan may explore the meaning of care and appreciation; a systemic plan may examine how work schedules, kin expectations, and attempted solutions shape the cycle. A defensible plan specifies why it fits, how it will be reviewed, and what evidence would prompt a change. It does not confuse a mutually negotiated goal with a therapist's preferred outcome.

Summary

Advanced marriage therapy requires disciplined attention to two individuals, the relational process, and the context in which the couple lives. The therapist holds competing accounts without collapsing them, defines the unit of care and the confidentiality frame, and avoids treating relationship preservation as the sole measure of success. Engagement begins before the first intervention: referral, contact, access, informed consent, and contracting all shape whether partners can participate freely.

Assessment integrates each partner's perspective, relationship history, interactional sequences, strengths, individual and health factors, social context, and safety. It is paced and purposeful rather than exhaustive. Observations should be described concretely and interpreted provisionally. Standardised measures can support monitoring, but cannot replace conversation, private safety assessment, or clinical judgement. Couple distress may coexist with individual mental-health needs, requiring consultation or coordinated care.

Safety is a threshold issue, not a later treatment module. Ordinary conflict, high-intensity disagreement, situational violence, and coercive control require careful differentiation. A symmetrical systemic formulation can be useful for reciprocal patterns but dangerous when it obscures coercion or unequal responsibility. Private assessment, clear confidentiality limits, safety planning, and specialist consultation are essential when indicated. Conjoint therapy should proceed only when both partners can participate with sufficient freedom and the clinician can manage risk within their competence.

A formulation links each partner's meanings with observable interaction, relevant history, context, strengths, and risk. It remains a revisable hypothesis and should be shared in language that invites correction. Treatment planning converts the formulation into goals, tasks, time frames, progress indicators, and conditions for review. Model selection considers evidence, therapist competence, couple preference, safety, access, and clinical fit. Flexibility requires a coherent rationale, not unstructured eclecticism.

During treatment, the clinician monitors alliance with each person and the process between them. De-escalation, rupture repair, and impasse review are core clinical tasks. Feedback and outcome measures are useful when interpreted collaboratively; change should be assessed across several domains, including safety, distress, relational functioning, decision clarity, and the couple's own goals. Referral, coordination, and ending are deliberate parts of treatment. The clinician should revise or stop conjoint work when new information changes the risk or fit of the service.

Doctoral Research Questions

1. Which assessment practices best detect differences in partners' goals and willingness to participate without increasing pressure or alliance imbalance?
2. How do clinicians distinguish reciprocal high-conflict interaction from coercive control, and what training or consultation pathways improve decision quality?
3. Which dimensions of couple-therapy alliance predict engagement and outcome for each partner, and how do partners' alliance ratings diverge over time?
4. How do treatment goals change when couples begin therapy with unequal commitment or uncertainty about continuing the relationship?
5. Which progress-monitoring strategies help therapists adapt treatment without making clients feel surveilled or judged?
6. How do individual and structural stressors—such as depression, chronic illness, caregiving burden, or economic instability—interact with couple distress and treatment response?
7. What service adaptations improve access and retention for couples with language, disability, transport, childcare, or shift-work barriers?
8. How do couples define a successful ending when their chosen outcome is respectful separation rather than relationship continuation?
9. What clinical conditions and safeguards are associated with safe and effective conjoint treatment after a history of violence?
10. How do therapists' model allegiance, supervision, and reflective practice influence formulation, intervention choice, and referral decisions?

Further Reading

- Christensen A and Jacobson NS, *Reconcilable Differences* (revised edn, Guilford Press 2002).
- Christensen A and Doss BD, *Integrative Behavioral Couple Therapy: A Therapist's Guide to Creating Acceptance and Change* (W W Norton 2017).
- Gurman AS, Lebow JL, and Snyder DK (eds), *Clinical Handbook of Couple Therapy* (6th edn, Guilford Press 2023).
- Johnson SM, *The Practice of Emotionally Focused Couple Therapy: Creating Connection* (3rd edn, Routledge 2019).
- Lebow JL, Chambers AL, Christensen A, and Johnson SM, 'Research on the Treatment of Couple Distress', *Journal of Marital and Family Therapy* 38 (2012).

- McGoldrick M, Gerson R, and Petry S, *Genograms: Assessment and Intervention* (3rd edn, W W Norton 2008).
- Snyder DK, Castellani AM, and Whisman MA, 'Current Status and Future Directions in Couple Therapy', *Annual Review of Psychology* 57 (2006).

KEY THEORETICAL POINT

Advanced practice principle. Formulate tentatively, assess safety privately when needed, negotiate goals with both partners, make the treatment frame explicit, and review whether the work is helping. When new information changes the risk or fit, change the plan rather than defending the original formulation.

End of Chapter 4

MFT. 904 — Advanced Clinical Practice in Marriage Therapy

APPENDIX A

Genogram Notation

The genogram is the standard multigenerational diagram used in marriage and family therapy for recording family structure, relationships, and functioning across at least three generations. Its notation is largely conventional rather than theoretically derived, and it has been codified most influentially in the work of Monica McGoldrick and colleagues. The symbols below follow that convention. Where practice varies between published sources — particularly in the notation of emotional relationships, where different texts use single, double, and triple lines with differing conventions — the variation is noted, and the governing principle is stated: notation must be defined for the record in which it is used, applied consistently, and explained to the family when the diagram is constructed with them.




Figure A.1 Standard genogram notation. Structural symbols (upper panels) record family composition, descent, and vital events; relationship symbols (lower panel) record the quality of emotional bonds. Notation conventions vary slightly between published sources; records should state which convention is used. *Source:* prepared for this edition.

A.1 Household structure

The household is drawn as an enclosing line around the persons who live together, with the household's identifying details recorded beside or beneath it. A person who has left the household is drawn inside the boundary with a diagonal line through the box, or outside the boundary with the date of departure recorded, according to the convention adopted in the record. Where members of the extended family live in the same compound or building, a dashed outer boundary may be used to indicate a wider domestic unit.

A.2 Descent and vital events

Birth order is recorded from left to right, with the eldest child nearest the couple line. Where children have different parents, the descent lines are drawn separately from the relevant couple lines. Deaths are marked with a diagonal cross through the symbol and the age or date of death recorded above it. Multiple births are shown by lines converging from the descent point to each child. Pregnancy, miscarriage, stillbirth, and termination are recorded by standard symbols and should be included where the family reports them, since their omission distorts the generational record and is frequently experienced by families as a silencing of significant losses.

A.3 Migration, culture, and social context

Because structure alone omits what often matters most, the record should carry information on migration and displacement (places and dates of origin, moves, separations of family members during migration, language use within the household), legal and immigration status where the family raises it and where recording is safe, occupation and socio-economic change across generations, and religious or community affiliation as a social identity. Serious caution is required: written records of legal status, ethnicity, or religion create risks for some families and must be kept in accordance with data-protection law and professional confidentiality standards, and in some jurisdictions must not be created in written form at all.

A.4 Relationship notation and the cultural genogram

Emotional relationships are recorded along the line connecting two persons: close and harmonious; fused or enmeshed; distant; conflictual; cutoff; and combinations such as close but conflictual, or distant and conflictual. Arrows may be added to indicate the direction of an asymmetric attachment, caretaking relationship, or hostility. The **cultural genogram** applies the same instrument to cultural identity, recording the family's experience of cultural pride, shame, discrimination, assimilation, and the internalisation of dominant-group evaluations of the family's group. It is constructed with the family and used as a conversational instrument rather than an assessment completed by the clinician alone.

A.5 Recording conventions for doctoral work

1. **State the convention.** Every genogram presented in a dissertation, article, or clinical report must be accompanied by a key naming the symbols used, including any local conventions.
2. **State the source of the information.** Record who supplied each cluster of information. Discrepant accounts from different family members are data, not errors to be resolved silently: noting that a mother and daughter disagree

about whether a grandfather left or was sent away is more informative than choosing one version.

3. **Record dates and ages, not impressions.** "Left home at 17" is data; "was in a hurry to escape" is an interpretation and belongs in the narrative, not the diagram.
4. **Do not over-diagram.** A diagram with ninety relationships marked is unreadable and functions as a display of thoroughness rather than an instrument of analysis. Record the relationships that carry the hypothesis.
5. **Anonymise.** In teaching, publication, and examination material, genograms must be anonymised: initials or labels replace names, identifying occupations and locations are generalised, and composite families are used where a single family would be identifiable.
6. **Review the diagram with the family.** Families frequently correct, dispute, and enlarge a diagram, and those corrections are among the most clinically informative moments of the assessment. A genogram that has never been shown to a family is an artefact of the clinician's research, not a family's history.

APPENDIX B

Comparative Grid of Family Systems Theories

The tables in this appendix are working instruments. They summarise the traditions examined in Chapter 1 on dimensions that allow systematic comparison, and they are intended to be used in seminar preparation, in the construction of formulations, and in the design of doctoral research. Two warnings accompany them.

First, any summary of a tradition compresses it. The cell entries are claims about what a tradition *characteristically* holds; individual theorists within a tradition differ, and later writings frequently revise earlier ones. Where a grid entry is contested within the tradition itself, the entry states the dominant version and the reader should consult the primary sources.

Second, the grid is not a menu. A formulation selected by matching a presentation to a cell in a table is not a formulation. The tables are aids to comparison; competence consists in using them to interrogate one's own reasoning.

B.1 Theories compared on core analytical dimensions

Table B.1 B.1 — Core analytical dimensions of the major traditions.

Tradition	Unit of analysis	Core construct	Account of problem maintenance	Mechanism of change
Structural	Household subsystem organisation	Boundary; hierarchy; alliance	Rigid or diffuse boundaries; weak executive authority; cross-generational coalition	Restructuring through enactment, boundary-making, realignment
Strategic and brief (MRI)	The interactional sequence around a defined problem	Attempted solution; feedback loop	Ineffective attempted solutions are repeated and amplified	Interrupting the attempted solution; task design; reframing
Milan systemic	The family's belief system and premises	Circularity; the family game; hypothesising	Covert coalitions, disqualification, and	Circular questioning; positive connota-

Tradition	Unit of analysis	Core construct	Account of problem maintenance	Mechanism of change
			premises that exclude alternatives	tion; reflecting team
Intergenerational (Bowen)	The multigenerational emotional unit	Differentiation of self; triangles; cutoff	Anxiety management through fusion, conflict, projection, or cutoff	Coaching; detriangulation; re-engagement with the family of origin
Contextual	The relational ledger across generations	Fairness; legacy; invisible loyalty; entitlement	Unbalanced give-and-take; destructive entitlement; unacknowledged merit	Multidirected partiality; acknowledgement; exoneration work
Experiential and communication	The here-and-now encounter and the message system	Congruence; report and command; punctuation	Incongruent communication; rigid punctuation; symmetrical escalation	Sculpting; immediacy; congruent communication practice
Attachment and emotion-focused	The dyadic attachment bond	Attachment security; the negative interaction cycle	Pursue–withdraw cycle driven by unacknowledged attachment panic	Emotion reprocessing; cycle de-escalation; restructuring responsiveness
Cognitive-behavioural	Observable exchange and cognition within the dyad or parent–child system	Reinforcement ratio; attribution; standards	Low reward-to-cost ratio; negative attributions; unrealistic standards	Behaviour exchange; communication skills; cognitive restructuring; acceptance work
Social constructionist and narrative	The meaning system and the conversational network	Story; problem-saturated account; language	Dominant stories that totalise identity; problem talk sustained by the network	Externalising; re-authoring; unique outcomes; solution elaboration
Integrative (metaframeworks, common factors)	Problem-defined system across levels	Lens selection; common factors	The problem is maintained across several levels simultaneously	Sequenced interventions selected by formulation; alliance and outcome monitoring

B.2 Therapist stance, session composition, and typical duration

Table B.2 B.2 — Stance, composition, duration, and characteristic risk.

Tradition	Therapist stance	Who attends	Typical duration	Characteristic risk
Structural	Active, directive, accommodating during joining	Whole household; subsystems seen separately as needed	Weeks to months	Imposing the therapist's map; insufficient attention to violence
Strategic and brief	Directive, problem-focused, influential	Those relevant to the defined problem	One to ten sessions	Manipulation; inadequate assessment; paradox misapplied
Milan systemic	Curious, neutral in discipline, hypothesising	Family plus reflecting team	Months	Neutrality misread as passivity or evasion in the face of harm
Intergenerational (Bowen)	Coach; teacher of the theory	One person, a couple, or a family; family of origin work between sessions	Months to years	Intellectualisation; underestimation of structural constraint
Contextual	Advocate for all parties in turn	Whoever can be present; absent and deceased members named	Variable; often medium-term	Acknowledging harm may be heard as excusing it
Experiential and communication	Personal, spontaneous, use of self	Family, often with co-therapy team	Variable	Dependence on therapist charisma; weak manualisation
Attachment and emotion-focused	Attuned, process-guiding, actively shaping emotion	Couple (or family in family versions)	Eight to twenty sessions typically	Decentring practical and structural problems; suitability with current violence
Cognitive-behavioural	Structured, collaborative, data-oriented	Couple, parent(s), or family depending on focus	Ten to twenty sessions typically	Rigidity; underattention to affect and to systemic context
Constructionist and narrative	Curious, decentred, influential	Those who define the problem, including professionals	Variable; often brief	Limits under risk; claims exceeding the evidence base

Tradition	Therapist stance	Who attends	Typical duration	Characteristic risk
Integrative	Formulation-driven; stance selected deliberately	Determined by the formulation	Planned and monitored, then revised	Eclecticism without a controlling framework

B.3 A checklist for evaluating a theory

1. **Unit of analysis.** What does the theory take as its system, and how are its boundaries drawn?
2. **Observables.** What would an observer measure in order to see the theory's constructs in operation?
3. **Prediction.** What does the theory predict that a rival theory does not?
4. **Disconfirmation.** What result would count as evidence against the theory?
5. **Mechanism.** By what causal pathway does change in the theory's terms produce change in the presenting problem?
6. **Evidence.** What is the state of the outcome, process, and mediation evidence, and how independent is it of the theory's proponents?
7. **Scope and limits.** For which presentations, ages, structures, and settings has the theory been tested, and for which has it not?
8. **Power and harm.** How does the theory describe relationships in which power is unequal or harm is occurring?
9. **Culture.** Whose family form does the theory implicitly assume? What happens to its constructs when applied across cultures?
10. **Utility.** What does a competent practitioner do differently because of this theory, and would clients notice?

These ten questions can be applied to any formulation, model, or published claim. A doctoral candidate who can answer them fluently about any tradition in the field has achieved what this course is designed to teach.

APPENDIX C

The Doctoral Programme and the Study Pathway for MFT. 901

This appendix situates the present volume within the doctoral programme in marriage and family therapy and sets out a study pathway for the course to which it corresponds, **MFT. 901 Advanced Theories of Family Systems**.

C.1 The programme

The doctoral programme comprises twenty-four required courses (72 units) distributed across three years, together with a pool of elective courses (each 3 units). The courses are as follows.

Table C.1 C.1 — Course structure of the doctoral programme (each course 3 units).

Year	Code	Course title
1	MFT. 901	Advanced Theories of Family Systems
1	MFT. 902	Research Methods in Marriage and Family Therapy
1	MFT. 903	Multicultural Issues in Family Therapy
1	MFT. 904	Advanced Clinical Practice in Marriage Therapy
1	MFT. 905	Psychopathology in Families
1	MFT. 906	Ethical and Legal Issues in Family Therapy
1	MFT. 907	Child and Adolescent Development
1	MFT. 908	Couples Therapy Techniques
2	MFT. 909	Trauma-Informed Care in Family Therapy
2	MFT. 910	Substance Abuse and Family Dynamics
2	MFT. 911	Family Life Cycle and Transitions
2	MFT. 912	Group Therapy in Family Contexts
2	MFT. 913	Neurobiology of Attachment and Relationships
2	MFT. 914	Supervision and Consultation in Therapy
2	MFT. 915	Assessment and Diagnosis in Family Therapy
2	MFT. 916	Conflict Resolution and Mediation
3	MFT. 917	Parenting and Family Dynamics
3	MFT. 918	Integrative Approaches in Therapy

Year	Code	Course title
3	MFT. 919	Family Therapy with LGBTQ+ Families
3	MFT. 920	Technology in Therapy Practice
3	MFT. 921	Advanced Play Therapy Techniques
3	MFT. 922	Spirituality in Family Therapy
3	MFT. 923	Advanced Research Methodology
3	MFT. 970	Dissertation
Elective	MFT. 924	Community Resources and Family Support
Elective	MFT. 925	Advanced Communication Skills for Therapists
Elective	MFT. 926	Narrative Therapy Techniques
Elective	MFT. 927	Psychodynamic Approaches to Family Therapy
Elective	MFT. 928	Family Therapy and Public Policy
Elective	MFT. 929	Advanced Practicum in Family Therapy
Elective	MFT. 930	Evaluation and Outcome Measurement in Therapy
Elective	MFT. 931	Advanced Pedagogy

C.2 Relation of this volume to the programme

MFT. 901 is the theoretical foundation on which subsequent courses build. The following table indicates, for each later course, the sections of this volume that provide its theoretical prerequisites.

Table C.2 C.2 — Theoretical prerequisites supplied by this volume.

Later course	Prerequisite material in this volume
MFT. 902 Research Methods	Section 1.1 (epistemology of observation); Section 1.8 (criteria for evaluating theory and evidence)
MFT. 903 Multicultural Issues	Section 1.7 (power, gender, and culture); Section 1.4 (genogram and cultural genogram)
MFT. 904 Advanced Clinical Practice	Sections 1.2, 1.3, 1.6, 1.8 (methods and integrative planning)
MFT. 905 Psychopathology in Families	Section 1.2 (symptom in context); Section 1.6 (affect regulation and attachment)
MFT. 906 Ethics and Legal Issues	Section 1.3 (paradox and risk); Section 1.7 (responsibility, power, and neutrality)
MFT. 907 Child and Adolescent Development	Section 1.5 (communication and self-esteem); Section 1.6 (attachment and co-regulation)
MFT. 908 Couples Therapy Techniques	Section 1.6 (attachment and emotion-focused models); Section 1.8 (cognitive-behavioural models)

Later course	Prerequisite material in this volume
MFT. 913 Neurobiology of Attachment	Section 1.6 (psychophysiology and the limits of neurobiological claims)
MFT. 914 Supervision and Consultation	Section 1.1 (second-order cybernetics); Section 1.4 (isomorphism)
MFT. 915 Assessment and Diagnosis	Section 1.2 (structural assessment); Section 1.4 (genograms); Appendix A
MFT. 918 Integrative Approaches	Section 1.8 (metaframeworks; common factors; monitoring)
MFT. 923 Advanced Research Methodology	Section 1.8 (specificity, mediation, and the evidence base)
MFT. 926 Narrative Therapy	Section 1.7 (constructionism and narrative practice)
MFT. 970 Dissertation	Entire volume: theoretical framework construction and critique

C.3 A nine-week study pathway for MFT. 901

The following pathway assumes a fifteen-week semester with a weekly three-hour seminar. Weeks 1–9 cover the analytical content; weeks 10–12 are devoted to integrative work; weeks 13–15 to assessment and consolidation.

Table C.3 C.3 — Suggested seminar sequence for MFT. 901.

Week	Reading focus	Seminar work
1	Section 1.1; Bateson (1972), selections; von Bertalanffy (1968)	Map the origins of systemic concepts; identify what was lost in translation
2	Section 1.2; Minuchin (1974)	Construct structural maps of two recorded or role-played families
3	Section 1.3; Watzlawick, Weakland, and Fisch (1974); Selvini Palazzoli et al (1978)	Practise circular questioning; debate the ethics of paradoxical technique
4	Section 1.4; Bowen (1978), selections; Boszormenyi-Nagy and Krasner (1986)	Build genograms; practise multidirected partiality in a role-play
5	Section 1.5; Watzlawick, Beavin, and Jackson (1967); Satir (1983)	Analyse session transcripts for punctuation, report, and command
6	Section 1.6; Johnson (2019); selected observational research	Track a negative cycle from a transcript; evaluate a neurobiological claim
7	Section 1.7; White and Epston (1990); de Shazer (1988); McGoldrick et al (2005)	Externalising practice; critique a formulation for cultural assumptions

Week	Reading focus	Seminar work
8	Section 1.8; Epstein and Baucom (2002); Christensen and Jacobson (2002)	Design a behaviour-exchange and an acceptance intervention for one case
9	Section 1.8 (integrative frameworks); Pinsof et al (2018); Sprenkle et al (2009)	Apply the integrative sequence (Figure 1.9) to the week 2 cases
10	Review of all sections	Comparative grid exercise: complete Tables B.1 and B.2 from memory
11	Evidence base review; meta-analyses and systematic reviews	Appraise three studies against the checklist in Appendix B.3
12	Critical traditions and ethics	Seminar debate: circularity, responsibility, and safety in a violent case
13	Applied Exercise 1	Group presentations and cross-examination
14	Examination preparation	Mock viva on competing formulations
15	Consolidation	Written examination or assignment submission

C.4 Assessment design

Three assessments are recommended for this course, in ascending order of independence.

Assignment 1 (2,000 words): concept analysis. Select one theoretical construct from Chapter 1 (for example, homeostasis, triangulation, enactment, punctuation, differentiation, or externalising). Trace its origin, describe its migration into family therapy, and evaluate the extent to which its clinical use departs from its original technical meaning.

Assignment 2 (3,000 words): comparative formulation. Using a case vignette supplied by the course leader — or a documented case from the literature, fully referenced — construct formulations of the same presentation from three traditions, state what each predicts, and defend a choice of treatment with reasons that address the alternatives.

Assignment 3 (5,000 words): theoretical framework for a research proposal. Develop the theoretical framework for a prospective doctoral study: state the research problem, specify the theories that frame it, justify their selection against at least two rival frameworks, and state the study's key constructs and their proposed operationalisation. This assignment is designed to feed directly into the dissertation stage.

Examiners should require candidates to state, for any theoretical claim they advance, what evidence would count against it. A candidate who cannot do so has not yet completed the course in the sense in which this volume defines it.

Selected Bibliography

This bibliography lists the principal works on which this volume relies and the works recommended for advanced study. It is deliberately weighted toward primary sources: the founding texts of each tradition, the research programmes that tested them, and the critical literature that has disciplined them. Where a work exists in multiple editions, the edition most useful for doctoral study is given. Where an item is a journal article, the citation convention is author, title, year, and journal; readers should consult the source for pagination and for the full bibliographic details required by their institution's citation standard.

1. Founding texts of systemic theory and cybernetics

- Bateson G, *Steps to an Ecology of Mind* (Chandler 1972; reprinted University of Chicago Press 2000).
- Bateson G, Jackson DD, Haley J, and Weakland J, 'Toward a Theory of Schizophrenia', *Behavioral Science* (1956).
- Bertalanffy L von, *General System Theory: Foundations, Development, Applications* (revised edn, George Braziller 1968).
- Cannon WB, *The Wisdom of the Body* (W W Norton 1932).
- Maturana HR and Varela FJ, *Autopoiesis and Cognition: The Realization of the Living* (D Reidel 1980).
- Maturana HR and Varela FJ, *The Tree of Knowledge: The Biological Roots of Human Understanding* (Shambhala 1987).
- Watzlawick P, Beavin JH, and Jackson DD, *Pragmatics of Human Communication: A Study of Interactional Patterns, Pathologies, and Paradoxes* (W W Norton 1967).
- Watzlawick P, Weakland JH, and Fisch R, *Change: Principles of Problem Formation and Problem Resolution* (W W Norton 1974).
- Wiener N, *Cybernetics: Or Control and Communication in the Animal and the Machine* (MIT Press 1948).

2. Histories, handbooks, and systematic overviews of the field

- Goldenberg H, Goldenberg I, and Peluso PR, *Family Therapy: An Overview* (9th edn, Cengage 2017).
- Gurman AS and Kniskern DP (eds), *Handbook of Family Therapy* (Brunner/Mazel 1981; later volumes with different editors).
- Gurman AS, Lebow JL, and Snyder DK (eds), *Clinical Handbook of Couple Therapy* (6th edn, Guilford 2023).
- Hoffman L, *Foundations of Family Therapy: A Conceptual Framework for Systems Change* (Basic Books 1981).
- Nichols MP and Davis SD, *Family Therapy: Concepts and Methods* (12th edn, Pearson 2020).
- Sexton TL and Lebow JL (eds), *Handbook of Family Therapy: The Science and Practice of Working with Families and Couples* (Routledge 2016).
- Walsh F (ed), *Normal Family Processes: Growing Diversity and Complexity* (4th edn, Guilford 2012).

3. Structural family theory

- Minuchin S, *Families and Family Therapy* (Harvard University Press 1974).
- Minuchin S and Fishman HC, *Family Therapy Techniques* (Harvard University Press 1981).
- Minuchin S, Montalvo B, Guerney BG, Rosman BL, and Schumer F, *Families of the Slums: An Exploration of Their Structure and Treatment* (Basic Books 1967).
- Minuchin S, Rosman BL, and Baker L, *Psychosomatic Families: Anorexia Nervosa in Context* (Harvard University Press 1978).
- Minuchin S, Nichols MP, and Lee W-Y, *Assessing Families and Couples: From Symptom to System* (Pearson 2007).
- Minuchin S, Reiter MD, and Borda C, *The Craft of Family Therapy: Challenging Certainties* (Routledge 2014).
- Szapocznik J and Kurtines WM, *Breakthroughs in Family Therapy with Drug-Abusing and Problem Youth* (Springer 1989).
- Szapocznik J, Hervis O, and Schwartz S, *Brief Strategic Family Therapy for Adolescent Drug Abuse* (National Institute on Drug Abuse 2003).

4. Strategic, brief, MRI, and Milan systemic approaches

- Boscolo L, Cecchin G, Hoffman L, and Penn P, *Milan Systemic Family Therapy: Conversations in Theory and Practice* (Basic Books 1987).

- Boscolo L and Bertrando P, *Systemic Therapy with Individuals* (Karnac 1996).
- de Shazer S, *Clues: Investigating Solutions in Brief Therapy* (W W Norton 1988).
- de Shazer S, *Words Were Originally Magic* (W W Norton 1994).
- Fisch R, Weakland JH, and Segal L, *The Tactics of Change: Doing Therapy Briefly* (Jossey-Bass 1982).
- Haley J, *Strategies of Psychotherapy* (Grune & Stratton 1963).
- Haley J, *Problem-Solving Therapy: New Strategies for Effective Family Therapy* (2nd edn, Jossey-Bass 1987).
- Madanes C, *Strategic Family Therapy* (Jossey-Bass 1981).
- Selvini Palazzoli M, Boscolo L, Cecchin G, and Prata G, *Paradox and Counterparadox: A New Model in the Therapy of the Family in Schizophrenic Transaction* (Jason Aronson 1978).
- Selvini Palazzoli M, Boscolo L, Cecchin G, and Prata G, 'Hypothesizing–Circularity–Neutrality: Three Guidelines for the Conductor of the Session', *Family Process* (1980).
- Tomm K, 'Interventive Interviewing: Intended Clinical Consequences of Hypothesizing, Circularity, and Neutrality', *Family Process* (1987).
- Andersen T, *The Reflecting Team: Dialogues and Dialogues about the Dialogues* (W W Norton 1991).

5. Intergenerational and contextual theories

- Bowen M, *Family Therapy in Clinical Practice* (Jason Aronson 1978).
- Boszormenyi-Nagy I and Spark GM, *Invisible Loyalties: Reciprocity in Intergenerational Family Therapy* (Harper & Row 1973).
- Boszormenyi-Nagy I and Krasner BR, *Between Give and Take: A Clinical Guide to Contextual Therapy* (Brunner/Mazel 1986).
- Boszormenyi-Nagy I, Grunebaum H, and Ulrich D, 'Contextual Therapy', in Gurman and Kniskern (eds), *Handbook of Family Therapy* (Brunner/Mazel 1981).
- Framo JL, *Family-of-Origin Therapy: An Intergenerational Approach* (Brunner/Mazel 1992).
- Kerr ME and Bowen M, *Family Evaluation: An Approach Based on Bowen Theory* (W W Norton 1988).
- McGoldrick M, Gerson R, and Petry S, *Genograms: Assessment and Intervention* (4th edn, W W Norton 2020).

- Skowron EA and Friedlander ML, 'The Differentiation of Self Inventory: Development and Initial Validation', *Journal of Counseling Psychology* (1998).
- Titelman P (ed), *Clinical Applications of Bowen Family Systems Theory* (Haworth 1998).

6. Communication, experiential, and humanistic theories

- Napier AY and Whitaker CA, *The Family Crucible* (Harper & Row 1978).
- Papp P, *The Process of Change* (Guilford 1983).
- Satir V, *Conjoint Family Therapy* (3rd edn, Science and Behavior Books 1983).
- Satir V, *Peoplemaking* (Science and Behavior Books 1972).
- Satir V, Banmen J, Gerber J, and Gomori M, *The Satir Model: Family Therapy and Beyond* (Science and Behavior Books 1991).
- Schwartz RC, *Internal Family Systems Therapy* (2nd edn, Guilford 2021; first published 1995).
- Whitaker CA and Keith DV, 'Symbolic-Experiential Family Therapy', in Gurman and Kniskern (eds), *Handbook of Family Therapy* (Brunner/Mazel 1981).

7. Attachment, emotion, and neurobiological accounts

- Ainsworth MDS, Blehar MC, Waters E, and Wall S, *Patterns of Attachment: A Psychological Study of the Strange Situation* (Erlbaum 1978).
- Bartholomew K and Horowitz LM, 'Attachment Styles among Young Adults: A Test of a Four-Category Model', *Journal of Personality and Social Psychology* (1991).
- Bowlby J, *Attachment and Loss*, vol 1: *Attachment* (Basic Books 1969).
- Bowlby J, *A Secure Base: Parent–Child Attachment and Healthy Human Development* (Basic Books 1988).
- Brennan KA, Clark CL, and Shaver PR, 'Self-Report Measurement of Adult Attachment: An Integrative Overview', in Simpson and Rholes (eds), *Attachment Theory and Close Relationships* (Guilford 1998).
- Coan JA, Schaefer HS, and Davidson RJ, 'Lending a Hand: Social Regulation of the Neural Response to Threat', *Psychological Science* (2006).
- Coan JA and Sbarra DA, 'Social Baseline Theory: The Social Regulation of Risk and Effort', *Current Opinion in Psychology* (2015).
- Gottman JM, *Marital Interaction: Experimental Investigations* (Academic Press 1979).

- Gottman JM, *The Marriage Clinic: A Scientifically Based Marital Therapy* (W W Norton 1999).
- Gottman JM and Levenson RW, 'The Timing of Divorce: Predicting When a Couple Will Divorce over a 14-Year Period', *Journal of Marriage and Family* (2000).
- Greenberg LS and Goldman RN, *Clinical Handbook of Emotion-Focused Therapy* (American Psychological Association 2019).
- Hazan C and Shaver P, 'Romantic Love Conceptualized as an Attachment Process', *Journal of Personality and Social Psychology* (1987).
- Johnson SM, *The Practice of Emotionally Focused Couple Therapy: Creating Connection* (3rd edn, Routledge 2019).
- Johnson SM, Hunsley J, Greenberg L, and Schindler D, 'Emotionally Focused Couples Therapy: Status and Challenges', *Clinical Psychology: Science and Practice* (1999).
- Main M and Solomon J, 'Discovery of an Insecure-Disorganized/Disoriented Attachment Pattern', in Brazelton and Yogman (eds), *Affective Development in Infancy* (Ablex 1986).
- Mikulincer M and Shaver PR, *Attachment in Adulthood: Structure, Dynamics, and Change* (2nd edn, Guilford 2016).
- Tronick EZ, 'Emotions and Emotional Communication in Infants', *American Psychologist* (1989).

8. Cognitive-behavioural, behavioural, and parenting models

- Baucom DH, Epstein NB, Kirby JS, and LaTaillade JJ, 'Cognitive-Behavioral Couple Therapy', in Gurman, Lebow, and Snyder (eds), *Clinical Handbook of Couple Therapy* (Guilford, various editions).
- Christensen A and Jacobson NS, *Reconcilable Differences* (revised edn, Guilford 2002).
- Christensen A, Atkins DC, Berns S, Wheeler J, Baucom DH, and Simpson LE, 'Traditional versus Integrative Behavioral Couple Therapy for Significantly and Chronically Distressed Married Couples', *Journal of Consulting and Clinical Psychology* (2004).
- Epstein NB and Baucom DH, *Enhanced Cognitive-Behavioral Therapy for Couples: A Contextual Approach* (American Psychological Association 2002).
- Jacobson NS and Margolin G, *Marital Therapy: Strategies Based on Social Learning and Behavior Exchange Principles* (Brunner/Mazel 1979).
- Kazdin AE, *Parent Management Training: Treatment for Oppositional, Aggressive, and Antisocial Behavior in Children and Adolescents* (Oxford University Press 2005).

- Kazdin AE, 'Evidence-Based Treatment and Practice: New Opportunities to Bridge Clinical Research and Practice, Enhance the Knowledge Base, and Improve Patient Care', *American Psychologist* (2008).
- Patterson GR, *Coercive Family Process* (Castalia 1982).
- Patterson GR, Reid JB, and Dishion TJ, *Antisocial Boys* (Castalia 1992).
- Sanders MR, *Positive Parenting and the Triple P System* (see Triple P programme literature; numerous trial reports).
- Weiss RL, Hops H, and Patterson GR, 'A Framework for Conceptualizing Marital Conflict, a Technology for Altering It, Some Data for Evaluating It', in Hamerlynck, Handy, and Mash (eds), *Behavior Change: Methodology, Concepts and Practice* (Research Press 1973).
- Webster-Stratton C, *The Incredible Years: A Troubleshooting Guide for Parents of Children Aged 3–8* (2nd edn, Incredible Years 2005).

9. Multisystemic, multidimensional, and family-based treatment models

- Henggeler SW, Schoenwald SK, Borduin CM, Rowland MD, and Cunningham PB, *Multisystemic Therapy for Antisocial Behavior in Children and Adolescents* (2nd edn, Guilford 2009).
- Liddle HA, *Multidimensional Family Therapy for Adolescent Cannabis Users* (Center for Substance Abuse Treatment 2002).
- Liddle HA, *Multidimensional Family Therapy: A Developmental-Ecological Treatment for Adolescent Behaviour Problems* (Routledge 2016).
- Alexander JF and Parsons BV, *Functional Family Therapy* (Brooks/Cole 1982).
- Sexton TL, *Functional Family Therapy in Clinical Practice: An Evidence-Based Treatment Model for Working with Troubled Adolescents* (Routledge 2011).

10. Social constructionist, narrative, and solution-focused approaches

- Anderson H, *Conversation, Language, and Possibilities: A Postmodern Approach to Therapy* (Basic Books 1997).
- Anderson H and Goolishian HA, 'Human Systems as Linguistic Systems: Preliminary and Evolving Ideas about the Implications for Clinical Theory', *Family Process* (1988).
- Gergen KJ, 'The Social Constructionist Movement in Modern Psychology', *American Psychologist* (1985).

- Freedman J and Combs G, *Narrative Therapy: The Social Construction of Preferred Realities* (W W Norton 1996).
- de Shazer S, Berg IK, Lipchik E, Nunnally E, Molnar A, Gingerich W, and Weiner-Davis M, 'Brief Therapy: Focused Solution Development', *Family Process* (1986).
- White M, *Maps of Narrative Practice* (W W Norton 2007).
- White M and Epston D, *Narrative Means to Therapeutic Ends* (W W Norton 1990).

11. Culture, gender, power, and critique

- Boyd-Franklin N, *Black Families in Therapy: Understanding the African American Experience* (2nd edn, Guilford 2003).
- Falicov CJ, *Latino Families in Therapy* (2nd edn, Guilford 2014).
- Goldner V, 'Feminism and Family Therapy', *Family Process* (1985).
- Goldner V, 'The Treatment of Violence and Victimization in Intimate Relationships', *Family Process* (1998).
- Goldner V, Penn P, Sheinberg M, and Walker G, 'Love and Violence: Gender Paradoxes in Volatile Attachments', *Family Process* (1990).
- Hardy KV and Laszloffy TA, 'The Cultural Genogram: Key to Training Culturally Competent Family Therapists', *Journal of Marital and Family Therapy* (1995).
- Hare-Mustin RT, 'A Feminist Approach to Family Therapy', *Family Process* (1978).
- Hare-Mustin RT, 'The Problem of Gender in Family Therapy Theory', *Family Process* (1987).
- Knudson-Martin C and Mahoney AR (eds), *Couples, Gender, and Power: Creating Change in Intimate Relationships* (Springer 2009).
- McGoldrick M, Giordano J, and Garcia-Preto N (eds), *Ethnicity and Family Therapy* (3rd edn, Guilford 2005).
- McGoldrick M and Hardy KV (eds), *Re-Visioning Family Therapy: Race, Culture, and Gender in Clinical Practice* (3rd edn, Guilford 2019).
- Tervalon M and Murray-García J, 'Cultural Humility versus Cultural Competence: A Critical Distinction in Defining Physician Training Outcomes in Multicultural Education', *Journal of Health Care for the Poor and Underserved* (1998).
- Walters M, Carter B, Papp P, and Silverstein O, *The Invisible Web: Gender Patterns in Family Relationships* (Guilford 1988).

12. Integration, common factors, and the evidence base

- Breunlin DC, Schwartz RC, and Mac Kune-Karrer B, *Metaframeworks: Transcending the Models of Family Therapy* (Jossey-Bass 1992).
- Carr A, 'The Evidence Base for Family Therapy and Systemic Interventions for Child-Focused Problems', *Journal of Family Therapy* (2014).
- Carr A, 'The Evidence Base for Couple Therapy, Family Therapy and Systemic Interventions for Adult-Focused Problems', *Journal of Family Therapy* (2018).
- Duncan BL, Miller SD, Wampold BE, and Hubble MA (eds), *The Heart and Soul of Change: Delivering What Works in Therapy* (2nd edn, American Psychological Association 2010).
- Lebow JL, Chambers AL, Christensen A, and Johnson SM, 'Research on the Treatment of Couple Distress', *Journal of Marital and Family Therapy* (2012).
- Miller SD, Duncan BL, and Hubble MA, *Escape from Babel: Toward a Unifying Language for Psychotherapy Practice* (W W Norton 1997).
- Miller SD, Hubble MA, Chow DL, and Seidel JA, 'The Outcome of Psychotherapy: Yesterday, Today, and Tomorrow', *Psychotherapy* (2013).
- Pinsof WM, *Integrative Problem-Centered Therapy: A Synthesis of Family, Individual, and Biological Therapies* (Basic Books 1995).
- Pinsof WM, Breunlin DC, Russell WP, and Lebow JL, *Integrative Systemic Therapy: Metaframeworks for Problem Solving with Individuals, Couples, and Families* (American Psychological Association 2018).
- Shadish WR and Baldwin SA, 'Meta-Analysis of MFT Interventions', *Journal of Marital and Family Therapy* (2003).
- Snyder DK, Castellani AM, and Whisman MA, 'Current Status and Future Directions in Couple Therapy', *Annual Review of Psychology* (2006).
- Sprenkle DH and Blow AJ, 'Common Factors across Theories of Marriage and Family Therapy: A Modified Delphi Study', *Journal of Marital and Family Therapy* (2004).
- Sprenkle DH and Piercy FP, 'Toward a Common Factors Approach to Couple and Family Therapy', *Family Process* (2005).
- Sprenkle DH, Davis SD, and Lebow JL, *Common Factors in Couple and Family Therapy: The Overlooked Foundation for Effective Practice* (Guilford 2009).
- Wampold BE, 'How Important Are the Common Factors in Psychotherapy? An Update', *World Psychiatry* (2015).

- Wampold BE and Imel ZE, *The Great Psychotherapy Debate: The Evidence for What Makes Psychotherapy Work* (2nd edn, Routledge 2015).

13. Reference works, journals, and methodological resources

- *Family Process* (Wiley) — the field's oldest specialist journal, founded 1962.
- *Journal of Marital and Family Therapy* (Wiley, for the American Association for Marriage and Family Therapy).
- *Journal of Family Therapy* (Wiley, for the Association for Family Therapy and Systemic Practice, United Kingdom).
- *Journal of Family Psychology* (American Psychological Association).
- *Family Relations* (Wiley, for the National Council on Family Relations).
- *Journal of Family Theory & Review* (Wiley) — theoretical and integrative scholarship.
- *Psychotherapy Research*; *Journal of Consulting and Clinical Psychology*; *Clinical Psychology Review* — for treatment outcome and process research.
- *Journal of Family Theory & Review* special issues on theory construction and evaluation.
- Hill AB, *The Environment and Disease: Association or Causation?* (Proceedings of the Royal Society of Medicine 1965) — still the clearest short statement of causal reasoning for clinicians.
- Kazdin AE, *Research Design in Clinical Psychology* (5th edn, Cambridge University Press 2021).
- Yin RK, *Case Study Research and Applications: Design and Methods* (6th edn, SAGE 2018).

14. Research methods, measurement, and evidence synthesis

- American Psychological Association Publications and Communications Board Task Force, 'Journal Article Reporting Standards for Quantitative, Qualitative, and Mixed Methods Research in Psychology', *Psychological Bulletin* 146(1) (2020).
- Campbell DT and Stanley JC, *Experimental and Quasi-Experimental Designs for Research* (Houghton Mifflin 1963).
- Creswell JW and Plano Clark VL, *Designing and Conducting Mixed Methods Research* (3rd edn, SAGE 2018).
- Hernán MA and Robins JM, *Causal Inference: What If* (Chapman & Hall/CRC 2020).
- Kazdin AE, *Research Design in Clinical Psychology* (5th edn, Cambridge University Press 2021).

- Kenny DA, Kashy DA, and Cook WL, *Dyadic Data Analysis* (Guilford Press 2006).
- Little RJA and Rubin DB, *Statistical Analysis with Missing Data* (3rd edn, Wiley 2019).
- Moher D, Liberati A, Tetzlaff J, Altman DG, and the PRISMA Group, 'Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement', *PLoS Medicine* 6(7) (2009).
- Page MJ et al, 'The PRISMA 2020 Statement: An Updated Guideline for Reporting Systematic Reviews', *BMJ* 372 (2021).
- Proctor E et al, 'Outcomes for Implementation Research: Conceptual Distinctions, Measurement Challenges, and Research Agenda', *Administration and Policy in Mental Health and Mental Health Services Research* 38 (2011).
- Raudenbush SW and Bryk AS, *Hierarchical Linear Models: Applications and Data Analysis Methods* (2nd edn, SAGE 2002).
- Shadish WR, Cook TD, and Campbell DT, *Experimental and Quasi-Experimental Designs for Generalized Causal Inference* (Houghton Mifflin 2002).
- Tashakkori A and Teddlie C (eds), *SAGE Handbook of Mixed Methods in Social & Behavioral Research* (2nd edn, SAGE 2010).
- Tong A, Sainsbury P, and Craig J, 'Consolidated Criteria for Reporting Qualitative Research (COREQ): A 32-Item Checklist for Interviews and Focus Groups', *International Journal for Quality in Health Care* 19(6) (2007).
- Wilkinson MD et al, 'The FAIR Guiding Principles for Scientific Data Management and Stewardship', *Scientific Data* 3 (2016).
- Yin RK, *Case Study Research and Applications: Design and Methods* (6th edn, SAGE 2018).

15. Multicultural, intersectional, and culturally responsive practice

- American Psychiatric Association, *Diagnostic and Statistical Manual of Mental Disorders* (5th edn, text revision, APA 2022), Cultural Formulation Interview materials.
- Crenshaw K, 'Mapping the Margins: Intersectionality, Identity Politics, and Violence against Women of Color', *Stanford Law Review* 43 (1991).
- Falicov CJ, *Latino Families in Therapy: A Guide to Multicultural Practice* (2nd edn, Guilford Press 2014).
- Hardy KV and Laszloffy TA, 'The Cultural Genogram: Key to Training Culturally Competent Family Therapists', *Journal of Marital and Family Therapy* 21(3) (1995).

- McGoldrick M and Hardy KV (eds), *Re-Visioning Family Therapy: Race, Culture, and Gender in Clinical Practice* (3rd edn, Guilford Press 2019).
- Metzl JM and Hansen H, 'Structural Competency: Theorizing a New Medical Engagement with Stigma and Inequality', *Social Science & Medicine* 103 (2014).
- Sue DW, Arredondo P, and McDavis RJ, 'Multicultural Counseling Competencies and Standards: A Call to the Profession', *Journal of Counseling & Development* 70 (1992).
- Tervalon M and Murray-García J, 'Cultural Humility versus Cultural Competence: A Critical Distinction in Defining Physician Training Outcomes in Multicultural Education', *Journal of Health Care for the Poor and Underserved* 9 (1998).
- Williams DR and Mohammed SA, 'Racism and Health I: Pathways and Scientific Evidence', *American Behavioral Scientist* 57 (2013).

16. Professional and regulatory sources

Members of the profession must work within the law, ethics codes, and training standards of the jurisdiction in which they practise. The following are representative sources rather than an exhaustive list:

- American Association for Marriage and Family Therapy, *Code of Ethics* (current edition).
- Association for Family Therapy and Systemic Practice (United Kingdom), *Code of Ethics and Practice* (current edition).
- American Psychological Association, *Ethical Principles of Psychologists and Code of Conduct* (current edition).
- World Health Organization, *International Classification of Diseases* (current edition), chapter on mental and behavioural disorders.
- American Psychiatric Association, *Diagnostic and Statistical Manual of Mental Disorders* (current edition).
- National Institute for Health and Care Excellence (United Kingdom), clinical guidelines relating to couple and family interventions.

Readers should verify the current edition and scope of every instrument, code, and classification cited in this volume before relying on it, and should note that some of the instruments discussed in the literature are proprietary and cannot be reproduced without permission from their publishers.