AuraHealth: first-aid guides for clinical review

AuraHealth is a free research prototype (simulation only, not a medical device). While someone waits for an ambulance, it shows these steps one at a time and reads them aloud, and it saves the time of each step for the doctor. The steps are fixed text written in plain English from the **International Federation of Red Cross and Red Crescent Societies (IFRC)**, *International first aid, resuscitation and education guidelines 2020*. No AI writes or changes them. For teenagers and adults. Babies and young children need different steps: tell the 112 operator the age. Medicine doses are left out on purpose (for example aspirin for chest pain). Ask the 112 operator. The emergency number shown is 112 (India).

Please check each step: is it correct, safe and clear for an ordinary person under stress? Tick OK, Change or Remove, and write the change in Comments. Anything missing that a bystander should do (or must not do), please add.

1. Someone collapsed: start here

When: You are not sure what is wrong. · **IFRC 2020 pages:** 128-139

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Make sure the place is safe for you, for example from traffic, fire or electricity. Go to the person only when it is safe.	☐ OK ☐ Change ☐ Remove	
2	Question	Shout their name and tap their shoulders. Do they respond? Answers: Yes, they respond / No response	☐ OK ☐ Change ☐ Remove	
3	Question	Look at their chest and listen. Are they breathing normally? Gasping, noisy or very slow breathing is NOT normal. Answers: Yes, normally / No, or only gasping / I'm not sure	☐ OK ☐ Change ☐ Remove	

Good to know

- If you are not sure whether they are breathing normally, start CPR. That is what the Red Cross guidelines advise.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Call 112 now, or ask someone to call. Shout and tap their shoulders. If they don't respond, look and listen: are they breathing normally? If not, start CPR: open the guide below. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

2. Not responding and not breathing normally

When: No response to voice or touch, and no breathing or only gasping. · **IFRC 2020 pages:** 134-146, 397

#	Type	What the app says (read aloud)	Your check	Comments
1	Call	Call 112 now, or ask someone nearby to call. Put the phone on speaker so your hands are free.	☐ OK ☐ Change ☐ Remove	
2	Step	Kneel beside them. Put the heel of one hand in the middle of the chest, the other hand on top, fingers locked together.	☐ OK ☐ Change ☐ Remove	
3	CPR beat	With straight arms, push hard, straight down, about 5 cm, then let the chest come all the way back up. Push on every beat: 100 to 120 times a minute.	☐ OK ☐ Change ☐ Remove	
4	Step	Keep going without stopping until help takes over, or they cough, open their eyes, move on purpose or breathe normally.	☐ OK ☐ Change ☐ Remove	

Do not

- Do not stop to look for a pulse. A pulse check is unreliable and wastes time.
- Do not stop compressions to wait for the ambulance.

Good to know

- If a defibrillator (AED) is nearby, ask someone to bring it, switch it on and follow its voice.
- If someone else can take over, swap so you don't get too tired, with as little pause as possible.
- If you are carrying the person to hospital, keep CPR going on a firm surface.

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

3. Not responding but breathing normally

When: No response to voice or touch, but normal, regular breathing. · **IFRC 2020 pages:** 128-133

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Roll them onto their side and gently tilt their head back. This keeps the airway open and stops them choking if they vomit.	☐ OK ☐ Change ☐ Remove	
2	Call	Call 112. Someone who does not respond needs help, even if they breathe.	☐ OK ☐ Change ☐ Remove	
3	Step	Stay with them and keep checking their breathing. If they stop breathing normally, start CPR.	☐ OK ☐ Change ☐ Remove	

Do not

- If you think they hurt their neck or back, do not roll them. Keep the airway open by tilting the head and lifting the chin.

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

4. Choking

When: Something is stuck in the throat. · **IFRC 2020 pages:** 170-174, 400

#	Type	What the app says (read aloud)	Your check	Comments
1	Question	Can they cough, speak or breathe? Answers: Yes / No	☐ OK ☐ Change ☐ Remove	
2	Step	Tell them to keep coughing. Stay with them and watch closely: it can get worse.	☐ OK ☐ Change ☐ Remove	
3	Step	Stand behind them, bend them forward, and give up to 5 sharp blows between the shoulder blades with the heel of your hand.	☐ OK ☐ Change ☐ Remove	
4	Step	Still blocked? Stand behind, put a fist between the belly button and the ribs, hold it with your other hand, and pull sharply in and up, up to 5 times.	☐ OK ☐ Change ☐ Remove	
5	Call	Still blocked? Call 112 now. Keep repeating 5 back blows and 5 abdominal thrusts.	☐ OK ☐ Change ☐ Remove	

Do not

- Do not sweep inside the mouth blindly with your fingers: it can push the object further down. Only take out something you can see.

Good to know

- Babies under one year need different steps. Tell the 112 operator the age.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Act now: if they cannot cough or speak, give up to 5 sharp blows between the shoulder blades, then up to 5 abdominal thrusts, and keep repeating. Ask someone to call 112. The guide below shows each step. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

5. Heavy bleeding

When: Blood is flowing fast from a wound. · **IFRC 2020 pages:** 187-194

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Press hard on the wound. Use a cloth, bandage or plastic between your hand and the blood. If they can, they can press it themselves.	☐ OK ☐ Change ☐ Remove	
2	Step	Help them lie down, so they do not faint and fall.	☐ OK ☐ Change ☐ Remove	
3	Call	Call 112, or ask someone to call while you keep pressing.	☐ OK ☐ Change ☐ Remove	
4	Step	If blood soaks through, put another cloth on top and press harder. Do not lift the first cloth.	☐ OK ☐ Change ☐ Remove	
5	Step	Only if pressing does not stop heavy bleeding from an arm or leg, and you have a tourniquet: put it on above the wound. The time is saved for the doctor.	☐ OK ☐ Change ☐ Remove	
6	Step	Keep them lying down and warm, for example covered with clothes or a blanket.	☐ OK ☐ Change ☐ Remove	

Do not

- If an object such as a knife is stuck in the wound, do not pull it out. Press around it.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Press hard on the wound with a cloth and keep pressing. Call 112, or ask someone to call while you press. Help them lie down. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

6. Chest pain

When: Pain or pressure in the chest that does not go away. · **IFRC 2020 pages:** 264-270

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Help them sit down in a comfortable position. If a doctor prescribed them heart medicine for this, help them take it.	☐ OK ☐ Change ☐ Remove	
2	Call	Call 112 now. Ask someone to bring a defibrillator (AED) if there is one nearby.	☐ OK ☐ Change ☐ Remove	
3	Step	Stay with them, keep them calm, and watch their breathing and whether they respond. If they stop responding and are not breathing normally, start CPR.	☐ OK ☐ Change ☐ Remove	

Good to know

- Women, older people and people with diabetes may have milder or unusual signs. If in doubt, treat it as serious.
- Medicine doses are not given here. Ask the 112 operator what to give.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Call 112 now. Help them sit down and rest, and keep them calm. Ask someone to bring a defibrillator (AED) if there is one nearby. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

7. Signs of a stroke

When: Sudden weak face or arm, or strange speech. · **IFRC 2020 pages:** 271-278

#	Type	What the app says (read aloud)	Your check	Comments
1	Question	Ask them to smile. Does one side of the face droop? Answers: Yes / No / Not sure	☐ OK ☐ Change ☐ Remove	
2	Question	Ask them to raise both arms. Does one arm drift down or not move? Answers: Yes / No / Not sure	☐ OK ☐ Change ☐ Remove	
3	Question	Ask them to repeat a simple sentence. Is their speech slurred or strange? Answers: Yes / No / Not sure	☐ OK ☐ Change ☐ Remove	
4	Call	Any of these signs? Call 112 now. Tell them when the signs started, or the last time the person was seen well. If you don't know, say so.	☐ OK ☐ Change ☐ Remove	
5	Step	Help them into a comfortable position and support them so they don't fall. Keep talking to them and watch their breathing.	☐ OK ☐ Change ☐ Remove	

Do not

- Do not give them anything to eat or drink: swallowing may not be safe.

Good to know

- Even if the signs go away, they still need a doctor quickly.
- Very high or low blood sugar can look similar. Tell the doctor about diabetes.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Call 112 now: these can be signs of a stroke. Note the time the signs started. Keep them comfortable and give nothing to eat or drink. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

8. Snakebite

When: Bitten by a snake, or you think so. · **IFRC 2020 pages:** 254-257

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Help them lie down and keep as still as possible. Moving makes the venom spread faster.	☐ OK ☐ Change ☐ Remove	
2	Call	Call 112. They need a hospital that has antivenom.	☐ OK ☐ Change ☐ Remove	
3	Step	Take off rings, bangles, watches and tight clothes near the bite before it swells. Move the limb as little as possible.	☐ OK ☐ Change ☐ Remove	
4	Step	If help is far away, keep the bitten arm or leg still with a cloth or non-stretchy bandage wrapped around it. Do not wrap it tight.	☐ OK ☐ Change ☐ Remove	
5	Step	Do not let them walk on a bitten leg. Carry them, for example two people making a chair with their arms.	☐ OK ☐ Change ☐ Remove	
6	Step	Keep watching their breathing and whether they respond. If it is safe, take a photo of the snake from a distance.	☐ OK ☐ Change ☐ Remove	

Do not

- Do not tie a tight band (tourniquet) around the limb.
- Do not cut the bite or suck out the venom.
- Do not put ice on it or rub it.
- Do not try to catch or kill the snake.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Call 112 now. Help them lie down and keep as still as possible. Do not tie the limb, cut the bite or suck out the venom. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

9. Seizure (fit)

When: Shaking, stiffening or not responding with twitching. · **IFRC 2020 pages:** 295-298

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Tap Done now: the seizure start time is saved for the doctor.	☐ OK ☐ Change ☐ Remove	
2	Step	Move hard or sharp things away. Do not move them unless they are in danger, for example on a road.	☐ OK ☐ Change ☐ Remove	
3	Step	Put something soft under their head. Take off glasses and loosen anything tight around the neck.	☐ OK ☐ Change ☐ Remove	
4	Call	Call 112 if: it is their first seizure, it lasts more than 5 minutes, they are hurt, they don't wake up after, it happens again, they are pregnant or diabetic, or you are not sure.	☐ OK ☐ Change ☐ Remove	
5	Step	When the shaking stops, tap Done: the stop time is saved.	☐ OK ☐ Change ☐ Remove	
6	Question	Now check their breathing. Are they breathing normally? Answers: Yes, normally / No, or only gasping	☐ OK ☐ Change ☐ Remove	

Do not

- Do not hold them down.
- Do not put anything in their mouth or between their teeth.
- Do not give them food, drink or medicine until they are fully awake.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Move hard or sharp things away and put something soft under their head. Do not hold them down or put anything in their mouth. Call 112 if it lasts more than 5 minutes, it is their first seizure, or they are hurt. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

10. Low blood sugar (diabetes)

When: Someone with diabetes is sweating, shaky, weak, confused or drowsy. · **IFRC 2020 pages:** 291-294

#	Type	What the app says (read aloud)	Your check	Comments
1	Question	Are they awake and able to swallow safely? Answers: Yes / No, very drowsy or not responding	☐ OK ☐ Change ☐ Remove	
2	Step	Help them sit down.	☐ OK ☐ Change ☐ Remove	
3	Step	Give something sugary now: their own glucose tablets if they have them, or a glass of fruit juice or a sugary (not diet) drink, or a few spoonfuls of sugar or some sweets.	☐ OK ☐ Change ☐ Remove	
4	Step	Stay with them. If they still feel unwell after 15 minutes, give something sugary again.	☐ OK ☐ Change ☐ Remove	
5	Call	Call 112 if they are not clearly better in about 30 minutes, or if they get more drowsy, have a fit or stop responding. Keep watching their breathing.	☐ OK ☐ Change ☐ Remove	
6	Step	When they feel better, give them something more filling, such as a slice of bread or a chapati, and check their blood sugar if they have a meter.	☐ OK ☐ Change ☐ Remove	

Do not

- Only give food or drink if they are awake and able to swallow: otherwise it can choke them.
- Do not give diet or sugar-free drinks or sweets: they do not raise blood sugar.

Good to know

- If you are not sure whether their sugar is low or high, treat it as low: sugar is unlikely to harm someone whose sugar is high, but not giving it to someone whose sugar is low can.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): If they are awake and can swallow, give something sugary now: their own glucose tablets, fruit juice or a sugary (not diet) drink, or sweets. Help them sit down. Call 112 if they cannot swallow safely, get more drowsy, have a fit or stop responding. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

11. Breathing difficulty or asthma attack

When: Short of breath, wheezing, or an asthma attack. · **IFRC 2020 pages:** 175-183

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Help them sit up, leaning forward with their arms resting on a table or on their knees. Stay calm and reassure them.	☐ OK ☐ Change ☐ Remove	
2	Step	If they have their own inhaler or medicine for this, help them use it. Loosen tight clothing and move them away from smoke or dust.	☐ OK ☐ Change ☐ Remove	
3	Call	Call 112 now if: there is no inhaler and it lasts more than a few minutes, the inhaler does not help within a few minutes, it is very hard to breathe or speak, their lips or fingers turn blue, they get confused or sleepy, or their breathing becomes slow and quieter as they tire.	☐ OK ☐ Change ☐ Remove	
4	Step	Stay with them, calm and sitting up, until their breathing is normal. They may use their inhaler again as their doctor prescribed.	☐ OK ☐ Change ☐ Remove	

Do not

- Do not make them lie flat.

Good to know

- Fast breathing from panic (with dizziness or tingling hands): take them somewhere quiet and ask them to breathe slowly into their cupped hands.
- If they stop responding, open "Someone collapsed: start here".

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Help them sit up, leaning forward with their arms resting on something, and loosen tight clothes. If they have an inhaler, help them use it. Call 112 now if it does not help within a few minutes or there is none, if they cannot speak, their lips turn blue, or they get confused or sleepy. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

12. Severe allergic reaction

When: Swelling of the lips, tongue or throat, hives, or trouble breathing after a food, a sting or a medicine. · **IFRC 2020 pages:** 279-287

#	Type	What the app says (read aloud)	Your check	Comments
1	Question	Is it severe: swelling of the lips, tongue or throat, trouble breathing or wheezing, or feeling faint? Answers: Yes / No, only an itchy rash or sneezing	☐ OK ☐ Change ☐ Remove	
2	Call	Call 112 now. Say it may be a severe allergic reaction (anaphylaxis).	☐ OK ☐ Change ☐ Remove	
3	Step	If they have their own adrenaline (epinephrine) auto-injector, help them use it now in the middle of the outer thigh. It can go through thin clothes. Tell the ambulance team when it was given.	☐ OK ☐ Change ☐ Remove	
4	Step	Help them lie down. If breathing is hard, help them sit up instead.	☐ OK ☐ Change ☐ Remove	
5	Step	Keep watching their breathing and whether they respond. If they are not better 5 to 10 minutes after the injection and have a second one, it may be used.	☐ OK ☐ Change ☐ Remove	
6	Step	For a mild reaction: move them away from what caused it, or rinse it off the skin, and help them take their own allergy medicine if they have it. Watch for any of the severe signs.	☐ OK ☐ Change ☐ Remove	

Do not

- Do not wait to see if it gets better when the lips, tongue or throat swell or breathing is hard.

Good to know

- A severe reaction can come back: watch them closely for the next three days.
- Someone who has had a severe reaction should always carry an allergy card.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): If the lips, tongue or throat swell, or breathing is hard, call 112 now: it may be a severe allergic reaction. If they have an adrenaline (epinephrine) auto-injector, help them use it in the middle of the outer thigh. Help them lie down, or sit up if breathing is hard. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

13. Poisoning

When: Swallowed medicines, a chemical, kerosene, a pesticide or a plant, or breathed in gas or smoke. · **IFRC 2020 pages:** 258-263

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Stop more contact with the poison. Take it away from them. With gas or smoke, get them into fresh air only if it is safe for you.	☐ OK ☐ Change ☐ Remove	
2	Step	Find out what it was, how much and when. Touch the packet only if it is safe.	☐ OK ☐ Change ☐ Remove	
3	Call	Call 112 and tell them what it was, how much and when. Follow their instructions.	☐ OK ☐ Change ☐ Remove	
4	Step	Help them into a comfortable position, on their left side if possible. Watch their breathing and whether they respond.	☐ OK ☐ Change ☐ Remove	
5	Step	Keep the bottles, packets or labels to give to the ambulance team.	☐ OK ☐ Change ☐ Remove	

Do not

- Do not make them vomit: it can damage the throat.
- If they swallowed a strong chemical (such as acid, bleach or drain cleaner), do not give water or milk to dilute it.

Good to know

- Poisoning can look like a fit, a stroke or low blood sugar.
- Keep medicines and chemicals locked away and out of reach of children.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Stop more contact with the poison: take it away, or get them into fresh air if it is safe for you. Call 112 and say what it was, how much and when. Do not make them vomit. Keep the bottle or packet. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

14. Burns

When: Burned by fire, hot water, oil, steam, a chemical or electricity. · **IFRC 2020 pages:** 213-220

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Make it safe first: stop the heat, for example cover a burning pot. With electricity, switch it off before you touch them.	☐ OK ☐ Change ☐ Remove	
2	Step	Cool the burn under clean running water for at least 10 minutes, best 20. Let the water flow over it gently. No running water? Use any cool, clean liquid.	☐ OK ☐ Change ☐ Remove	
3	Call	Call 112 if the burn is large or deep, on the face, mouth, throat or private parts, or caused by flames, electricity or a chemical. Any burn on a baby, an older person or someone with diabetes needs a doctor.	☐ OK ☐ Change ☐ Remove	
4	Step	Take off rings, bangles, watches and clothes on or near the burn, unless they are stuck to the skin.	☐ OK ☐ Change ☐ Remove	
5	Step	After cooling, cover it loosely with cling film, a clean wet cloth or a non-stick dressing. Cover it, do not wrap it tightly: it may swell.	☐ OK ☐ Change ☐ Remove	
6	Step	Keep the rest of the body warm. Watch their breathing and whether they respond, especially for a child or an older person.	☐ OK ☐ Change ☐ Remove	

Do not

- Do not put ice or ice-cold water on a burn.
- Do not pop blisters.
- Do not pull off clothing that is stuck to the burn.

Good to know

- A chemical on the skin or in the eye: rinse with plenty of clean running water until the pain eases.
- If the burn later gets more painful, red and warm, or they get a fever, see a doctor: it may be infected.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Cool the burn under clean running water now, for at least 10 minutes and best 20.

Take off rings and clothes near it unless they are stuck. Call 112 if the burn is large or deep, on the face, mouth, throat or private parts, or from flames, electricity or a chemical. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

15. Head injury (fall or blow to the head)

When: Fell, or hit their head. · **IFRC 2020 pages:** 233-237

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Help them stop what they were doing and rest. After a hard fall, keep their head and neck still.	☐ OK ☐ Change ☐ Remove	
2	Question	Any of these? Knocked out even for a few seconds; confused, slurred speech or behaving differently; a fit; weakness or tingling in an arm or leg; vomiting more than once; double or blurred vision; blood or clear fluid from the nose or ears; a bad headache for more than 2 hours. Answers: Yes, one or more / No	☐ OK ☐ Change ☐ Remove	
3	Call	Call 112 now. Keep their head and neck as still as possible: you can hold the head gently with your hands. Keep watching their breathing and whether they respond.	☐ OK ☐ Change ☐ Remove	
4	Step	A bump, a mild headache or feeling sick: let them rest and stay with them. Keep watching for the signs above for the next 24 hours, and call 112 if one appears.	☐ OK ☐ Change ☐ Remove	

Do not

- Do not let them drive, use machines or play sport until all signs have gone.

Good to know

- Signs of a serious head injury can appear hours, or even days, later.
- After a hard fall or blow, the neck may be hurt too.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Ask them to stop and rest, and watch them closely. Call 112 if they were knocked out even for a few seconds, are confused or behaving differently, vomit more than once, have weakness or tingling, double or blurred vision, a bad headache, or blood or clear fluid from the nose or ears. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

16. Feeling faint or fainting

When: Dizzy, pale, about to faint, or fainted and woke up quickly. · **IFRC 2020 pages:** 299-303

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Help them sit or lie down on the floor, in a safe place where they cannot fall.	☐ OK ☐ Change ☐ Remove	
2	Step	Ask them to cross their legs and squeeze their leg, belly and bottom muscles, or to squat. If they are lying down, you can raise their legs instead.	☐ OK ☐ Change ☐ Remove	
3	Question	Did they faint and are they still not responding? Answers: No, they are awake / Yes, not responding	☐ OK ☐ Change ☐ Remove	
4	Step	Stay with them and talk to them. Watch whether they respond and how they breathe. Give water if they are awake and can swallow, and ask what might have caused it (hunger, heat, standing for long).	☐ OK ☐ Change ☐ Remove	
5	Step	Let them get up slowly, only when they feel better: they may feel faint again.	☐ OK ☐ Change ☐ Remove	

Do not

- Do not let them stand up quickly.

Good to know

- A faint should last only seconds. If they do not wake up at once, it may be something more serious, such as a heart problem: call 112.
- If they fell, check for a head injury.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Help them sit or lie down where they cannot fall. If they lie down, raise their legs. Ask them to cross their legs and squeeze their leg, belly and bottom muscles. If they faint and do not wake up at once, call 112 and check their breathing. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

17. Fever (high temperature)

When: Temperature 38°C (100.4°F) or more, hot and sweaty, or shivering. · **IFRC 2020 pages:** 304-307

#	Type	What the app says (read aloud)	Your check	Comments
1	Step	Help them rest and dress lightly. Do not cover them with heavy blankets.	☐ OK ☐ Change ☐ Remove	
2	Step	Give water or other fluids to drink often, to replace what they lose by sweating.	☐ OK ☐ Change ☐ Remove	
3	Step	You can sponge them with lukewarm water if they like it. Never use cold water.	☐ OK ☐ Change ☐ Remove	
4	Step	If they feel really unwell, a fever medicine such as paracetamol can help: give only the dose written on the label or the prescription. Ask a doctor or pharmacist first for older people, or anyone with liver problems or on other medicines.	☐ OK ☐ Change ☐ Remove	
5	Call	Call 112 now if the fever comes with a rash, a change in how they think or behave (confused, very sleepy), trouble breathing, severe belly pain, eyes hurting in the light with vomiting, or cold, pale and clammy skin.	☐ OK ☐ Change ☐ Remove	
6	Step	See a doctor soon if they are over 65, have cancer or a weak immune system, if the fever lasts more than 3 days or does not come down with the medicine, or if anything else worries you.	☐ OK ☐ Change ☐ Remove	

Do not

- Do not use cold water or ice: it makes the body keep its heat in and causes shivering.
- Do not wrap them in thick blankets or many layers.

Good to know

- A fever is a temperature above 38°C (100.4°F). Above 39°C (102.5°F) it can be harmful.
- Where malaria is common, see a doctor for any fever, even without other signs.

Shown first in "Talk to Aura" (when someone describes this emergency in the chat, before they open the guide): Let them rest in light clothes and drink plenty of fluids. You can sponge them with lukewarm (not cold) water. Call 112 if the fever comes with a rash, confusion or a change in

behaviour, trouble breathing, severe belly pain, or eyes hurting in the light with vomiting. ☐ OK ☐ Change

Overall for this guide: ☐ Approve ☐ Approve with the changes above ☐ Do not use

Reviewer

Name	
Qualification / registration	
Date	
May we say on our demo video that you reviewed these steps? (Yes / No)	
Signature	