



Data 345

Applied Linear Algebra for Statistical Learning

Class 10 (Sep. 29, 2025)

Linear Transformations & Matrices

- Last time: we introduced a linear transformation $f_\theta: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ which took vectors in \mathbb{R}^2 and rotated them θ radians counterclockwise.
- We noted that determining a formula for f_θ boiled down to the effect of f_θ on basis elements of \mathbb{R}^2 .
- Let $T: V \rightarrow W$ be a linear transformation between vector spaces V and W , with $\dim(V) = n$ and $\dim(W) = m$. Let $B = (\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n)$ and $C = (\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_m)$ be two ordered bases for V and W .
- Since T sends vectors in V to vectors in W , then for any $j \in \{1, 2, \dots, n\}$, the vector $T(\mathbf{b}_j)$ may be expressed as a linear combination of basis elements from C , i.e.:

$$T(\mathbf{b}_j) = \alpha_{j1}\mathbf{c}_1 + \alpha_{j2}\mathbf{c}_2 + \dots + \alpha_{jm}\mathbf{c}_m$$

- These coefficients can be arranged into a matrix A_T , where $A_T[i, j] = \alpha_{ij}$.

Linear Transformations & Matrices

- The matrix A_T is called the **transformation matrix** of T with respect to B and C .
- Since linear transformations are entirely determined by their effect on basis elements, a transformation essentially just summarizes the effect of T on each individual basis element from B in terms of the basis C .
- Example: Define a linear mapping $T: \mathbb{R}^3 \rightarrow \mathbb{R}^4$ as:

$$T\left(\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}\right) = \begin{bmatrix} 3x_1 + 2x_2 + x_3 \\ x_1 + x_2 + x_3 \\ x_1 - 3x_2 \\ 2x_1 + 3x_2 + x_3 \end{bmatrix}$$

$$T\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 3 \\ 1 \\ 1 \\ 2 \end{bmatrix}$$

$$T\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 2 \\ 1 \\ -3 \\ 3 \end{bmatrix}$$


$$T\left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 1 \end{bmatrix}$$

$$A_T = \begin{bmatrix} 3 & 2 & 1 \\ 1 & 1 & 1 \\ 1 & -3 & 0 \\ 2 & 3 & 1 \end{bmatrix}$$


Visualizing Linear Transformations

- Since a linear transformation's behavior is entirely determined by its effect on basis vectors, one strategy for visualizing linear transformations (particularly from \mathbb{R}^2 to \mathbb{R}^2) is as follows:
- Draw two separate coordinate axes.
- Draw two arrows for the standard basis vectors $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 0 \\ 1 \end{bmatrix}$ in the first pair of coordinate axes.
- Draw a dense collection of points around the unit square in the first pair of coordinate axes.
- Draw $T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right)$ and $T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right)$ as vectors in the second pair of axes.
- Draw the images of the individual points from the unit square in the second pair of coordinate axes.


Visualizing Linear Transformations



For $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ where: $A_T = \begin{bmatrix} \cos \frac{\pi}{3} & -\sin \frac{\pi}{3} \\ \sin \frac{\pi}{3} & \cos \frac{\pi}{3} \end{bmatrix}$



For $S: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ where $A_S = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$



For a general linear transformation $\Phi: \mathbb{R}^2 \rightarrow \mathbb{R}^2$

Linear Transformations & Basis

- **Caution:** Matrices for linear transformations are always made with respect to an ordered basis for both the domain and the codomain. Choosing a different basis for V or W (or even a different ordering) leads to a different matrix.
- Example: Consider the transformation $S: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ that simply stretches in the horizontal direction by a factor of 2 and leaves the vertical direction unchanged. That is, $S\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = \begin{bmatrix} 2x_1 \\ x_2 \end{bmatrix}$.

- Letting $B = C = \left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right)$, we get:

$$\begin{aligned} S\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) &= \begin{bmatrix} 2 \\ 0 \end{bmatrix} = 2\begin{bmatrix} 1 \\ 0 \end{bmatrix} + 0\begin{bmatrix} 0 \\ 1 \end{bmatrix} \\ S\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) &= \begin{bmatrix} 0 \\ 1 \end{bmatrix} = 0\begin{bmatrix} 1 \\ 0 \end{bmatrix} + 1\begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{aligned} \longrightarrow A_S = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$$

- Letting $C = \left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix}\right)$, however, gives:


$$\begin{aligned} S\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) &= \begin{bmatrix} 2 \\ 0 \end{bmatrix} = 2\begin{bmatrix} 1 \\ 0 \end{bmatrix} + 0\begin{bmatrix} 1 \\ 1 \end{bmatrix} \\ S\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) &= \begin{bmatrix} 0 \\ 1 \end{bmatrix} = -1\begin{bmatrix} 1 \\ 0 \end{bmatrix} + 1\begin{bmatrix} 1 \\ 1 \end{bmatrix} \end{aligned} \longrightarrow \widetilde{A}_S = \begin{bmatrix} 2 & -1 \\ 0 & 1 \end{bmatrix}$$

Linear Transformation & Basis

- All this is to say that different matrices may represent the same linear transformation.
- This would be genuinely awful news under most circumstances because it means that the assignment of transformation matrices is not well-defined.
- Let $T: V \rightarrow W$ be a linear transformation between two vector spaces. Suppose that B, \tilde{B} and C, \tilde{C} are two ordered bases for V and W , respectively.

$$V \xrightarrow{T} W$$

Create a linear transformation $V \rightarrow V$ which simply “converts” the basis elements of \tilde{B} to basis elements of B . Its transformation matrix is P .



Create a linear transformation $W \rightarrow W$ which converts the basis elements of \tilde{C} to basis elements of C . Its transformation matrix is Q .

Equivalent Linear Transformations

- ▶ Let $A, B \in \mathbb{R}^{m \times n}$. Then A and B are said to be **equivalent** matrices if there exists invertible $Q \in \mathbb{R}^{m \times m}$ and invertible $P \in \mathbb{R}^{n \times n}$ such that $B = Q^{-1}AP$.
- ▶ Two matrices are equivalent if, and only if, they *represent the same linear transformation*.
- ▶ Under the equation $B = Q^{-1}AP$, the matrix A represents a linear transformation from V to W under one basis, while B represents the same linear transformation using a different basis. The matrices P and Q are called **change-of-basis matrices**.
- ▶ Now: **why should we care** about this? After all, each vector space has its own standard basis – why not just represent every transformation with respect to its standard bases?

Equivalent Linear Transformations

- Insisting on using the same bases every time makes some degree of sense for defining matrices, but it's incredibly restrictive and limits real solving power.
- Here's an extremely simple example of how change-of-basis speeds up computation.
- Consider a matrix $A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$ corresponding to a linear transformation of the standard basis vectors of \mathbb{R}^2 .
- Instead of the basis $B = \left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right)$, let's consider a different basis $\tilde{B} = \left(\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right)$.

$$\begin{aligned} \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix} &= \begin{bmatrix} 3 \\ 3 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} 1 \\ -1 \end{bmatrix} \\ \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ -1 \end{bmatrix} &= \begin{bmatrix} 1 \\ -1 \end{bmatrix} = 0 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 1 \begin{bmatrix} 1 \\ -1 \end{bmatrix} \end{aligned} \quad \rightarrow \quad \tilde{A} = \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix}$$

Equivalent Linear Transformations

- The big difference between A and \tilde{A} is that \tilde{A} is a **diagonal** matrix: its only nonzero entries lie on the main diagonal.
- If we wanted to repeatedly apply the transformation, we would chain together the transformation (i.e., multiply the matrices by themselves multiple times).

$$A^2 = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \cdot \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} (2 \cdot 2) + (1 \cdot 1) & (2 \cdot 1) + (1 \cdot 2) \\ (1 \cdot 2) + (2 \cdot 1) & (1 \cdot 1) + (2 \cdot 2) \end{bmatrix} = \begin{bmatrix} 4 + 1 & 2 + 2 \\ 2 + 2 & 1 + 4 \end{bmatrix} = \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix}$$

$$A^3 = \begin{bmatrix} 5 & 4 \\ 4 & 5 \end{bmatrix} \cdot \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} (5 \cdot 2) + (4 \cdot 1) & (5 \cdot 1) + (4 \cdot 2) \\ (4 \cdot 2) + (5 \cdot 1) & (4 \cdot 1) + (5 \cdot 2) \end{bmatrix} = \begin{bmatrix} 10 + 4 & 5 + 8 \\ 8 + 5 & 4 + 10 \end{bmatrix} = \begin{bmatrix} 14 & 13 \\ 13 & 14 \end{bmatrix}$$

$$A^4 = \begin{bmatrix} 14 & 13 \\ 13 & 14 \end{bmatrix} \cdot \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} (14 \cdot 2) + (13 \cdot 1) & (14 \cdot 1) + (13 \cdot 2) \\ (13 \cdot 2) + (14 \cdot 1) & (13 \cdot 1) + (14 \cdot 2) \end{bmatrix} = \begin{bmatrix} 28 + 13 & 14 + 26 \\ 26 + 14 & 13 + 28 \end{bmatrix} = \begin{bmatrix} 41 & 40 \\ 40 & 41 \end{bmatrix}$$

$$\tilde{A}^2 = \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} (3 \cdot 3) + (0 \cdot 0) & (3 \cdot 0) + (0 \cdot 1) \\ (0 \cdot 3) + (1 \cdot 0) & (0 \cdot 0) + (1 \cdot 1) \end{bmatrix} = \begin{bmatrix} 9 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\tilde{A}^3 = \begin{bmatrix} 9 & 0 \\ 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} (9 \cdot 3) + (0 \cdot 0) & (9 \cdot 0) + (0 \cdot 1) \\ (0 \cdot 3) + (1 \cdot 0) & (0 \cdot 0) + (1 \cdot 1) \end{bmatrix} = \begin{bmatrix} 27 & 0 \\ 0 & 1 \end{bmatrix}$$


Diagonal vs. Non-Diagonal

- If you look carefully at the entries in \tilde{A}^k , you can see that:

$$\tilde{A}^k = \begin{bmatrix} 3 & 0 \\ 0 & 1 \end{bmatrix}^k = \begin{bmatrix} 3^k & 0 \\ 0 & 1^k \end{bmatrix}$$

- In other words, to compute a high power of a diagonal matrix all you need to do is compute the exponentiated diagonals.
- If $A, B \in \mathbb{R}^{n \times n}$, then matrix multiplication – computing AB – is $\mathcal{O}(n^3)$. So to compute A^k (knowing that A is not a diagonal matrix) would have a time complexity of $\mathcal{O}(k \cdot n^3)$. For large powers of k , it would be extremely slow.
- On the other hand, for a diagonal matrix, one only needs to compute n separate exponentiated real numbers. So, if $D \in \mathbb{R}^{n \times n}$ is diagonal, then computing D^k is $\mathcal{O}(n \cdot \log k)$ instead.
- This is such a huge difference in time that even though it is slow and expensive to find a “diagonal” equivalent to a matrix it is **often worth the trade-off** by an order of magnitude.

Application – Image Processing



Applying a “blur” filter to an image requires a type of transformation called a **convolution**, which distributes pixel intensity values based on a sort of weighted average of pixel values of neighboring pixel values.

For an $N \times M$ image, “direct” convolution (i.e. using the standard basis elements) would take $\mathcal{O}(NMnm)$ time. Taking the time to diagonalize first changes the operation to $\mathcal{O}(NM \log NM)$ time instead.







Image & Kernel

- ▶ Let $T: V \rightarrow W$ be a linear transformation. The **image** or **range** of T is simply $T(V)$. The **kernel** or **null space** of T is the set of all vectors \mathbf{v} in V such that $T(\mathbf{v}) = \mathbf{0}_W$.
- ▶ In other words, the kernel of T is the set of vectors in V that get annihilated (sent to the zero vector) by the transformation T . The image of T is often written $\text{im } T$, while the kernel is often written $\text{ker } T$.
- ▶ For any linear transformation $T: V \rightarrow W$, we have that $T(\mathbf{0}_V) = \mathbf{0}_W$ so the null space / kernel is never empty.
- ▶ Because T is linear, the image of T is always a subspace of W , while the kernel of T is always a subspace of V .

Image & Kernel



The null space $\ker T$ is the part of the space V that is squashed into the zero vector.

The image $\text{im } T$ is the version of V that has been successfully “transferred” to W .