

Relative Equilibria and Diagonals

M. Shub

In the question of whether the relative equilibrium classes, \bar{R}_e , are finite, one of the first things to do seems to be to show that the Newtonian potential function

$$-\sum_{i \neq j} \frac{m_i m_j}{\|x_i - x_j\|}$$
 restricted to the sphere $\sum m_i x_i^2 = 1$ and $\sum m_i x_i = 0$ has no singularities in a neighborhood of the diagonals $x_i = x_j$, $i \neq j$, where the function is not defined. Here we let $x_i \in R^k$, $i = 1, \dots, n$, and $m_i \in R_+$, $i = 1, \dots, n$.

We proceed as follows: The derivative of $-\sum_{i \neq j} \frac{m_i m_j}{\|x_i - x_j\|}$ is

$$\sum_{i \neq j} \frac{m_i m_j \langle x_i - x_j, v_i - v_j \rangle}{\|x_i - x_j\|^3}. \text{ Let } x_1, \dots, x_{k_1}, x_{k_1+1}, \dots, x_{k_2}, \dots, x_*, \dots, x_n = X$$

be a point on the generalized diagonals where we have grouped the equal terms

$x_1 = \dots = x_{k_1}$, $x_{k_1+1} = \dots = x_{k_2}$, $x_* = \dots = x_n$. Let x_j^i denote the i^{th} component of x_j . Without loss of generality we may assume $k_1 \geq 2$ and $x_n^1 \neq x_1^1$. We represent a point in a neighborhood of X by $X + \delta$, where $\delta = (\delta_1, \dots, \delta_n)$. We will pick a vector $V(\delta) = (v_1(\delta), \dots, v_n(\delta))$ defined in this neighborhood so that:

(a) $\|V(\delta)\|$ is bounded

$$(b) \sum_{i \neq j} \frac{m_i m_j \langle x_i + \delta_i - (x_j + \delta_j), v_i(\delta) - v_j(\delta) \rangle}{\|x_i + \delta_i - (x_j + \delta_j)\|^3} \rightarrow +\infty$$

as $\delta_i - \delta_j \rightarrow 0$ for $i, j \leq k_1$ and $i \neq j$.

(c) $V(\delta)$ is in tangent space to the sphere $\sum_{i=1}^n m_i x_i^2 = 1$ and $\sum_{i=1}^n m_i x_i = 0$ if $X + \delta$ lies in the sphere.

We do this by solving the following system of linear equations continuously in a neighborhood of X .

$$(1) \sum_{i=1}^n m_i (x_i + \delta_i) v_i = 0$$

$$(2) \sum_{i=1}^n m_i v_i = 0$$

$$(3) v_1 - v_j = \delta_1 - \delta_j \text{ for } 2 \leq j \leq k_1$$

$$(4) v_{k_1+1} = v_{k_1+2} = \dots = v_{*-1} = 0$$

$$(5) v_* = v_{*+1} = \dots = v_n$$

If we have solved these equations then writing down the derivative

$$\sum_{\substack{i \neq j \\ i, j \leq k_1}} \frac{m_i m_j \langle x_i - x_j, v_i - v_j \rangle}{\|x_i - x_j\|^3}$$

we see that at $X + \delta$ if it is defined its value is

$$\geq \sum_{\substack{i \neq j \\ i, j \leq k_1}} \frac{m_i m_j \langle \delta_i - \delta_j, \delta_i - \delta_j \rangle}{\|\delta_i - \delta_j\|^3} + C \quad \text{where } C \text{ is a constant which may be chosen for}$$

the whole neighborhood if $\|\delta\|$ is small enough. Now

$$\sum_{\substack{i \neq j \\ i, j \leq k_1}} \frac{m_i m_j \langle \delta_i - \delta_j, \delta_i - \delta_j \rangle}{\|\delta_i - \delta_j\|^3} + C = \sum_{\substack{i \neq j \\ i, j \leq k_1}} \frac{m_i m_j}{\|\delta_i - \delta_j\|} + C \quad \text{which obviously tends to } +\infty$$

as any $\delta_i - \delta_j \rightarrow 0$. Which proves (b). Equations (1) and (2) prove c, and continuity insures that $\|V(\delta)\|$ is bounded.

It remains only to solve the equations for the v_i . We consider the family of linear maps

$$\begin{aligned} \sum m_i (x_i + \delta_i) v_i &: R^{kn} \rightarrow R^1 \\ \sum m_i v_i &: R^{kn} \rightarrow R^k \\ v_1 - v_i &: R^k \rightarrow R^k \quad \text{for } 2 \leq i \leq k_1 \end{aligned}$$

restricted to the linear subspace of R^{kn} defined by $v_{k_1+1} = \dots = v_{*-1} = 0$ and $v_* = \dots = v_n$. So we have a linear map from a $(k_1+1)k$ dimensional vector space to a $k_1 k + 1$ dimensional vector space. We will prove that it is surjective at $\delta = 0$. Then we will be able to solve the equations continuously in a neighborhood and we will be done. To prove the surjectivity we restrict to the $k_1 k + 1$ dimensional subspace determined by $v_n^j = 0$ for $j \neq 1$ and we will show that the map is an isomorphism by calculating the determinant of its matrix. Its matrix is:

$$\begin{pmatrix} m_1 x_1 & m_2 x_2 & \dots & m_{k_1} x_{k_1} & \dots & m_n x_n \\ m_1 I & m_2 I & \dots & m_{k_1} I & \dots & 0 \\ I & -I & & & & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \vdots \\ I & & & & -I & 0 \end{pmatrix}$$

where $\tilde{m} = \sum_{j=1}^n m_j$, I is the $k \times k$ identity matrix, and

$$\begin{pmatrix} \tilde{m} x_n^1 \\ \tilde{m} x_n^2 \\ \vdots \end{pmatrix}$$

is an ordinary

column of numbers. Let $m = \sum_{j=1}^{k_1} m_j$. Then after identifying $x_1 = \dots = x_{k_1}$ and column reducing we get:

$$\begin{pmatrix} mx_1 & m_2 x_2 & \dots & m_{k_1} x_{k_1} & \tilde{m} x^1 \\ mI & m_2 I & \dots & m_{k_1} I & \tilde{m} x^k \\ 0 & -I & & 0 & 0 \\ \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & & & -I & 0 \end{pmatrix}$$

so that the determinant is equal to the determinant of:

$$\begin{pmatrix} mx_1 & \tilde{m} x_n^1 \\ \vdots & \tilde{m} x_n^1 \\ mI & \vdots \\ 0 & 0 \end{pmatrix} \text{ which is } m^{k-1} \det \begin{pmatrix} mx_1^1 & \tilde{m} x_k^1 \\ m & \tilde{m} \end{pmatrix}$$

which is $m^{k-1}(\tilde{m} mx_1^1 - \tilde{m} mx_n^1) = m^k \tilde{m}(x_1^1 - x_n^1) \neq 0$ because by hypothesis $x_1^1 \neq x_n^1$.

Note that the calculation also gives that the neighborhood of the diagonals in which we may assume that there are no singularities varies continuously with the masses.

Brandeis University
WALTHAM, Mass.