Assessment and Differential Diagnosis

Chapter 17

In this chapter

Learning objectives 2
Introduction 2
17.1 Engagement and safety 2
17.2 Trauma history 3
17.3 Symptom assessment 3
17.4 Standardised instruments 3
17.5 Differential diagnosis 4
17.6 Dissociation and complex presentations 4
17.7 Cultural and linguistic considerations 5
17.8 Medical screening 5
17.9 Risk assessment 5
17.10 Formulation and feedback 6
17.11 Assessment of children and adolescents 6
17.12 Assessment in legal and occupational contexts 6
Synthesis 7
Composite case study 8
Review and discussion questions 8
Research assignments 8
Chapter summary 9
References 9

Central question. How should trauma-related conditions be assessed, and how can clinicians avoid the common errors of over- and under-diagnosis?

Learning objectives

- describe the components of a trauma-informed assessment;
- distinguish PTSD from common differential diagnoses;
- explain the role of standardised instruments;
- recognise the effects of culture, language and context on assessment;
- identify when medical investigation is indicated.

Introduction

Assessment is the foundation of good practice. It determines whether treatment is offered, which treatment, and whether the person is safe. It is also where many errors occur: trauma may be missed because it is not asked about, over-diagnosed because symptoms are attributed to trauma without considering alternatives, or misattributed to a medical condition that is not investigated.

This chapter sets out a structured approach. It covers history, symptom assessment, differential diagnosis, instruments, cultural considerations and medical screening. It emphasises that assessment is not a single event but a process that continues through treatment.

> **A note on asking.** *Many people do not volunteer trauma histories. They may not recognise what happened as trauma, may fear consequences, or may have been discouraged from speaking. Asking directly, privately and without pressure increases disclosure. The question should be followed by a response that is respectful and practical, not by immediate interrogation.*

17.1 Engagement and safety

Assessment begins with engagement. The person needs to know who you are, what you will do with the information, and what choices they have. Confidentiality and its limits should be explained.

Physical and emotional safety should be considered. Is the setting private? Is the person at risk at home? Are there immediate medical needs? These questions come before symptom inventories.

Assessment should be paced. A person who is highly dissociated or intoxicated may not be able to provide a reliable history, and the assessment should be resumed later rather than forced.

17.2 Trauma history

The trauma history should establish what happened, when, for how long, who was involved and what the consequences were. It should include childhood adversity, adult trauma, recent events and ongoing threats.

Questions should be specific enough to elicit information without being leading. Asking "have you ever been physically hurt by someone close to you" is more useful than asking "do you have trauma".

The history should also cover protective factors, supports and the person's own understanding of what happened. This informs formulation and engagement.

17.3 Symptom assessment

Core PTSD symptoms include intrusive memories, avoidance, negative alterations in cognition and mood, and alterations in arousal and reactivity. The assessment should establish frequency, intensity, duration and functional impact.

It should also assess dissociation, complex PTSD features, depression, anxiety, substance use, psychosis, suicidal ideation and self-harm. Comorbidity is the rule rather than the exception.

Symptoms should be linked to functioning: work, relationships, sleep, self-care and safety. A symptom count without functional context is of limited use.

Most adults report at least one traumatic event, but only a minority develop PTSD (Kessler et al., 1995).

17.4 Standardised instruments

Standardised instruments improve reliability and support monitoring. Commonly used measures include the PTSD Checklist for DSM-5, the Clinician-Administered PTSD Scale, the Impact of Event Scale-Revised and the Dissociative Experiences Scale.

Instruments are screening or severity measures, not diagnoses. They depend on the population, the language and the method of administration. Cut-off scores should be interpreted with knowledge of local validation.

Self-report instruments can be completed privately, which may increase disclosure. They can also be affected by literacy, language and cultural interpretation of items.

17.5 Differential diagnosis

Several conditions can mimic or co-occur with PTSD. Major depression shares low mood, sleep disturbance and concentration difficulties. Generalised anxiety shares hyperarousal and worry. Panic disorder shares autonomic arousal. Psychosis can involve intrusive experiences that resemble flashbacks. Traumatic brain injury can produce cognitive and emotional changes.

Medical conditions should be considered: thyroid disease, anaemia, sleep apnoea, chronic pain, medication effects and substance withdrawal. These can produce symptoms that overlap with trauma.

The goal is not to exclude trauma but to avoid attributing everything to it. A person can have PTSD and hypothyroidism, and treating one does not treat the other.

Diagnosis rests on clinical interview against established criteria rather than on screening scores alone (Yehuda et al., 2015).

17.6 Dissociation and complex presentations

Dissociation should be assessed specifically, including depersonalisation, derealisation, amnesia and identity confusion. Severe dissociation may require a modified treatment approach.

Complex PTSD, as defined in ICD-11, includes the core PTSD symptoms plus disturbances in self-organisation: affect dysregulation, negative self-concept and difficulties in relationships. Assessment should establish whether these features are present and how they affect functioning.

Personality disorder diagnoses should be made cautiously in the context of trauma, because many features overlap and because labelling can be stigmatising. Where a diagnosis is made, it should inform treatment rather than replace formulation.

The dissociative subtype and its assessment are reviewed by Lanius et al. (2010).

17.7 Cultural and linguistic considerations

Culture shapes how distress is expressed, what is considered traumatic, who is consulted and what help is acceptable. Somatic idioms, spiritual explanations and family-based coping are common and should be respected.

Language is central. Assessment in a second language may obscure nuance, and interpreters should be trained and briefed. Family members should not interpret for sensitive assessments.

Instruments developed in one culture may not be valid in another. Local validation and qualitative input are important.

17.8 Medical screening

Medical screening should be guided by the history and presentation. Indicated investigations might include thyroid function, full blood count, glucose, vitamin levels and, where relevant, toxicology.

Red flags—focal neurological signs, sudden cognitive decline, severe headache, seizures, unexplained weight loss—require urgent medical assessment. These should not be attributed to trauma without investigation.

The interface between trauma and medicine is often poorly managed. Coordinated care between mental health and primary care improves outcomes.

17.9 Risk assessment

Risk assessment should cover suicide, self-harm, harm to others, safeguarding and immediate danger. It should be documented and followed by a plan.

Asking about suicide does not increase risk. It provides an opportunity to intervene. Assessment should include intent, plan, means, protective factors and access to support.

Risk is not a static property of a person; it changes with circumstances. Assessment should be repeated at transitions and during crises.

17.10 Formulation and feedback

Formulation integrates the history, symptoms, context and strengths into an explanatory account that guides treatment. It should be shared with the person, checked for accuracy and revised as new information emerges.

Feedback improves engagement and outcomes. It also respects the person's expertise about their own life.

Formulation is not a diagnosis. It is a working model that can accommodate complexity, uncertainty and change.

17.11 Assessment of children and adolescents

Children rarely describe trauma in the way adults do. Younger children may show regression, clinginess, sleep disruption, somatic complaints, aggression or repetitive play; adolescents may show risk-taking, withdrawal, irritability or changes in school performance. A single informant is rarely sufficient, and caregivers, teachers and the young person should each contribute where possible.

Assessment should be developmentally calibrated. Questions should use the child's vocabulary, concrete examples and, for younger children, drawing or play-based methods. Forced-choice questions and repeated interrogation should be avoided.

Consent and confidentiality require care. A child may not understand the purpose of the assessment, and a caregiver may be the source of harm. Where safeguarding concerns arise, the clinician must follow local procedures and tell the child, in advance where possible, what will be shared and with whom.

Standardised instruments exist for trauma exposure and symptoms across age ranges, but they should be supplemented by clinical interview and, where appropriate, observation. Reports from school can reveal difficulties that are not apparent at home.

Assessment should identify strengths, relationships and resources, not only symptoms. A formulation that lists deficits without context is both inaccurate and demoralising for families.

17.12 Assessment in legal and occupational contexts

Clinicians are sometimes asked to assess trauma for courts, immigration tribunals, employers or insurers. These contexts differ from clinical care, and the differences should be made explicit.

The first issue is role clarity. A treating clinician is an advocate for the patient's wellbeing; an expert witness owes a duty to the court. The same person should not usually occupy both roles, and where they do, the dual role should be disclosed and managed.

The second issue is the standard of proof and the question asked. Legal questions are not clinical questions: "is this account consistent with the medical evidence" is different from "does this person meet criteria for PTSD". Reports should answer the question asked, state the limits of the assessment and avoid speculation.

The third issue is symptom validity. Effort and symptom exaggeration occur in some medico-legal assessments, but they are not the default, and no clinician can detect deception by impression. Validated symptom validity tests can be used where indicated, with attention to false positives and to the distress that testing can cause.

The fourth issue is documentation. Contemporaneous notes, verbatim quotes where appropriate, and clear reasoning are essential. Reports should distinguish what was reported, what was observed and what was inferred.

Finally, the clinician should consider the consequences of the assessment for the person. A report that overstates certainty can harm the person it is meant to help; one that understates genuine impairment can deny them justice. Accuracy, not advocacy in either direction, is the standard.

Synthesis

Table 17.1 summarises assessment components.

Table 17.1. Components of trauma-informed assessment

Component	Purpose	Common errors
Engagement	Safety, consent	Rushing to symptoms
Trauma history	Identify exposure	Not asking directly
Symptom assessment	Severity, function	Counting symptoms only
Instruments	Reliability, monitoring	Treating scores as diagnosis
Differential diagnosis	Avoid misattribution	Attributing all to trauma
Risk assessment	Safety planning	Avoiding the question
Formulation	Integrated explanation	Not sharing with the person

Composite case study

A patient presents with insomnia, irritability and intrusive images. Assessment identifies a traumatic event, but also untreated sleep apnoea and heavy alcohol use. The formulation integrates these factors. Treatment addresses sleep, alcohol and trauma in a sequenced plan, with medical follow-up. The clinician does not assume that trauma explains everything.

Review and discussion questions

How would you ask about trauma in a way that increases disclosure?

Why are screening instruments not diagnostic?

Which medical conditions should be considered in the differential diagnosis of PTSD?

How should interpreters be used in trauma assessment?

What are the risks of diagnosing personality disorder in trauma survivors?

Why should formulation be shared with the person?

Research assignments

- Compare two PTSD instruments, noting validation evidence and cultural limitations.

- Design an assessment protocol for a refugee clinic.
- Write a risk assessment template that is trauma-informed and legally defensible.

Chapter summary

Trauma-informed assessment integrates engagement, trauma history, symptoms, differential diagnosis, risk and formulation. Standardised instruments support but do not replace clinical judgement. Culture, language and medical conditions shape presentation and require attention. A good assessment avoids both under-recognition and over-attribution, and it results in a shared formulation that guides treatment and can be revised.

References

Kessler, R. C., Sonnega, A., Bromet, E., Hughes, M., & Nelson, C. B. (1995). Posttraumatic stress disorder in the National Comorbidity Survey. *Archives of General Psychiatry, 52*(12), 1048–1060. https://doi.org/10.1001/archpsyc.1995.03950240066012

Lanius, R. A., Vermetten, E., Loewenstein, R. J., Brand, B., Schmahl, C., Bremner, J. D., & Spiegel, D. (2010). Emotion modulation in PTSD: Clinical and neurobiological evidence for a dissociative subtype. *American Journal of Psychiatry, 167*(6), 640–647. https://doi.org/10.1176/appi.ajp.2009.09081168

Yehuda, R., Hoge, C. W., McFarlane, A. C., Vermetten, E., Lanius, R. A., Nievergelt, C. M., Hobfoll, S. E., Koenen, K. C., Neylan, T. C., & Hyman, S. E. (2015). Post-traumatic stress disorder. *Nature Reviews Disease Primers, 1*, Article 15057. https://doi.org/10.1038/nrdp.2015.57