

HOMEWORK 10

MATH 2001

SEBASTIAN CASALAINA

ABSTRACT. This is the first homework assignment. The problems are from Hammack [?, Ch. 1, §1.1]:

- **Chapter 12 Section 12.1**, Exercises: 4, 6. **Section 12.2**, Exercises: 5, 10. **Section 12.3**, Exercises: 1, 2.

CONTENTS

Chapter 12 Section 12.1	1
Ch.12, §12.1, Exercise 4	1
Ch.12, §12.1, Exercise 6	2
Ch.12, §12.2, Exercise 5	3
Ch.12, §12.2, Exercise 10	3
Ch.1, §1.1, Exercise 38	4
Ch.1, §1.1, Exercise 40	4

CHAPTER 12 SECTION 12.1

Ch.12, §12.1, Exercise 4. There are eight different functions $f : \{a, b, c\} \rightarrow \{0, 1\}$. List them all. Diagrams will suffice.

Date: April 12, 2020.

Solution to Ch.1, §1.1, Exercise 2.

$$f = \{(a, 0), (b, 0), (c, 0)\}$$

$$f = \{(a, 0), (b, 1), (c, 0)\}$$

$$f = \{(a, 0), (b, 0), (c, 1)\}$$

$$f = \{(a, 0), (b, 1), (c, 1)\}$$

$$f = \{(a, 1), (b, 0), (c, 0)\}$$

$$f = \{(a, 1), (b, 1), (c, 0)\}$$

$$f = \{(a, 1), (b, 0), (c, 1)\}$$

$$f = \{(a, 1), (b, 1), (c, 1)\}$$

□

Ch.12, §12.1, Exercise 6. Suppose $f : \mathbb{Z} \rightarrow \mathbb{Z}$ is defined as $f = \{(x, 4x + 5) : x \in \mathbb{Z}\}$. State the domain, codomain and range of f . Find $f(10)$.

Solution to Ch.12, §12.1, Exercise 6.

The domain of the function is $\{x : x \in \mathbb{Z}\}$.

The codomain of the function is $\{x : x \in \mathbb{Z}\}$.

The range of the function is $\{x : x = 4n, n \in \mathbb{Z}\}$.

□

Ch.12, §12.2, Exercise 5. A function $f : \mathbb{Z} \rightarrow \mathbb{Z}$ is defined as $f(n) = 2n + 1$. Verify whether this function is injective and whether it is surjective.

Solution to Ch.12, §12.2, Exercise 5.

Proposition. The function is injective but not surjective.

Step 1. Prove the function is injective. *Proof.* Assume $f(x) = f(y)$.

We have $2x + 1 = 2y + 1$, thus $x = y$.

Step 2. Prove the function is not surjective.

Proof. There exist element 2 which $f(x) = 2x + 1 \neq 2$ for every $x \in \mathbb{Z}$. The function is injective but not surjective. \square

Ch.12, §12.2, Exercise 10. Prove that the function $f : \mathbb{R} - 1 \rightarrow \mathbb{R} - 1$ defined by $f(x) = \left(\frac{x+1}{x-1}\right)^3$ is bijective.

Solution to Ch.12, §12.2, Exercise 10.

Proposition. The $f(x)$ is bijective.

Step 1. Prove $f(x)$ is injective.

Proof. Assume $f(x) = f(y)$, we have $\left(\frac{x+1}{x-1}\right)^3 = \left(\frac{y+1}{y-1}\right)^3$, taking the cubic root on both side get $\left(\frac{x+1}{x-1}\right) = \left(\frac{y+1}{y-1}\right)$. Therefor we have:

$$(x + 1) * (y - 1) = (y + 1) * (x - 1)$$

$$xy + y - x = xy + x - y$$

$$y - x = x - y$$

$$2y = 2x$$

$$x = y$$

Step 2. Prove $f(x)$ is surjective.

Proof. Suppose $f(x) = c$, we have $f(x) = \left(\frac{x+1}{x-1}\right)^3 = c$. Solve for x , we get $x = \frac{1+\sqrt[3]{c}}{1-\sqrt[3]{c}}, x \in \mathbb{R}$. It follows that $f(x)$ is surjective. The $f(x)$ is surjective and injective, so it is bijective. \square

Ch.1, §1.1, Exercise 38.


Solution to Ch.1, §1.1, Exercise 38. \square

Ch.1, §1.1, Exercise 40. Sketch the following set of points in the x, y -plane:

$$S = \{(x, y) : x \in [0, 1], y \in [1, 2]\}$$

Solution to Ch.1, §1.1, Exercise 40. For this problem I first sketched my own solution by hand. However, to implement my solution in L^AT_EX, I modified the tikz code from the webpage:

<https://tex.stackexchange.com/questions/140312/tikz-shading-region-bounded-by-s>



\square

UNIVERSITY OF COLORADO, DEPARTMENT OF MATHEMATICS, CAMPUS BOX
395, BOULDER, CO 80309-0395

Email address: casa@math.colorado.edu