

Linking early warning signals to the temporal epidemiology of measles in four Nigerian cities

Andrew T. Tredennick^{1,2*}, Pejman Rohani^{1,2}, Eamon O’Dea^{1,2}, Tobias Brett^{1,2}, & John M. Drake^{1,2}

¹ *Odum School of Ecology, University of Georgia, Athens, GA, USA*

² *Center for the Ecology of Infectious Diseases, University of Georgia, Athens, GA, USA*

Abstract

Forecasting the trajectory of infectious disease outbreaks over time is a fundamental challenge facing society.

1 Introduction

Theory shows that epidemic transitions can be anticipated by trends in the statistical properties of disease time series (AERO papers). The existence of statistical trends in the data that precede critical transitions, so-called ‘early warning signals’ (EWS), imply that we may be able to anticipate disease emergence and outbreaks. The end goal is a model-independent detection system, where statistical properties of disease surveillance data can trigger warnings of impending outbreaks without the need to fit mechanistic models of disease transmission (Han and Drake 2017).

However, there is currently a gap between the theoretical work, which has relied on knowing the underlying disease dynamics, and the eventual goal of applying EWS in real-world situations where the underlying disease dynamics may be unknown. Theoretical development of EWS has focused on anticipating when the population becomes supercritical, when $\mathcal{R}_0 > 1$, after which an outbreak is inevitable, perhaps with some bifurcation delay (Dibble et al. 2016). Knowing the value of \mathcal{R}_0 through time makes it possible to test the accuracy of EWS that are estimated from state variables alone. Empirical application of EWS does not require knowing the value of \mathcal{R}_0 through time, meaning that “tests” require making assumptions about when critical transitions occur. Whether EWS track and/or anticipate underlying dynamics of real disease time series remains unknown, and is a critical knowledge gap that must be filled before EWS can confidently be deployed.

To fill this gap we will fit a mechanistic model to incidence data of measles in Niger to estimate the temporal epidemiology of the disease, yielding the very same parameters that are known in data-free modeling studies. In particular, we are interested in the correlation between EWS and the time-varying reproductive ratio, known as the effective reproductive ratio (\mathcal{R}_E). If EWS and \mathcal{R}_E are significantly and positively correlated, then we have empirical evidence that EWS are applicable in real-world settings. If EWS and \mathcal{R}_E are negatively correlated or not significantly positively correlated, then we have evidence that EWS may not be applicable in certain settings.

*Corresponding author: atredenn@gmail.com

Materials and Methods

Data

We used weekly measles case report data from four Nigerien cities, collected over an 11 year period (1995-2005) (Figure 1). These data are ideal for stress testing EWS because each city has different population sizes, has different dynamics in terms of size of outbreaks and length of inter-epidemic periods, and each time series has different amounts of “noise” (though, the difference in variability is probably just reflective of population size differences). The data come from [somewhere/someone], and used here with permission from [somewhere/someone].




Figure 1: Time series of weekly measles case reports from four cities in Niger.

Stochastic *SEIR* model

The model is a discrete-time approximation of a continuous-time SEIR model with limited demography, specified as a set of difference equations,

$$S_{t+dt} = n_{S,t} - n_{E,t} \quad (1)$$

$$E_{t+dt} = n_{E,t} - n_{I,t} \quad (2)$$

$$I_{t+dt} = n_{I,t} + n_{O,t} - n_{R,t}, \quad (3)$$

where \mathbf{n}_t are random variables representing the number of individuals transitioning into or out of each class at each timestep $t \rightarrow t + dt$. n_S is the number of births, n_E is the number of newly infected individuals that have the disease but are not infectious, n_I is the number of newly infectious individuals, n_O is the number of imported infections, and n_R is the number of newly recovered individuals who are no longer infectious and have life-long immunity. The stochastic random variables are specified as follows:

$$n_{S,t} \sim \text{Poisson}(\mu_t N_t \times dt) \quad (4)$$

$$n_{E,t} \sim \text{Binomial}(\lambda_{E,t}, S_t) \quad (5)$$

$$n_{I,t} \sim \text{Binomial}(\lambda_{I,t}, E_t) \quad (6)$$

$$n_{O,t} \sim \text{Poisson}(\psi \times dt) \quad (7)$$

$$n_{R,t} \sim \text{Binomial}(\lambda_{R,t}, I_t), \quad (8)$$

where μ_t is the birth rate at time t , ψ is the rate of imported infections, and λ_E , λ_I , and λ_R are the probabilities of exposure, becoming infectious, and recovery, respectively. These probabilities reflect the processes of transmission, transition from the latent period to the infectious period, and recovery, which we model as:

$$\lambda_{E,t} = 1 - e^{-\frac{\beta_t I_t dt}{N_t}} \quad (9)$$

$$\lambda_{I,t} = 1 - e^{-\eta E_t dt} \quad (10)$$

$$\lambda_{R,t} = 1 - e^{-\gamma I_t dt}, \quad (11)$$

where β_t is time-varying rate of transmission, η is time-invariant rate from the exposed class to the infectious class, and γ is time-invariant recovery rate. We model rate of transmission as:

$$\beta_t = \beta \left(1 + \sum_{i=1}^6 q_i \xi_{i,t} \right) \Gamma_t. \quad (12)$$

β is the mean transmission rate, ψ accounts for measles infections from external sources that are not part of the local dynamics, and the term $\sum_{i=1}^6 q_i \xi_{i,t}$ is a B-spline to model seasonality in transmission. The B-spline bases ($\xi_{i,t}$) are periodic with a 1 year period. The transmission rate (β_t) is also subject to stochastic process noise at each time step, Γ_t , which we model as a gamma-distributed white (temporally uncorrelated) noise with mean 1 and variance σ^2 (Bretó and Ionides 2011).

We do not include a death process in the model because we expect death rates from the susceptible and infectious classes to be minimal relative to births and we are not interested in robust estimates of the recovered class. Excluding deaths means we can avoid making further assumptions about demographic rates – we are already making assumptions about birth rates (e.g., the rate is the same across cities, but with city-specific population size). We model demographic stochasticity in births and imported infections by drawing time-specific values from Poisson distributions. Transitions in the model are shown in Table 1. In this model, the effective reproductive ratio at time t is: $\mathcal{R}_{E(t)} = \beta_t / \gamma$.

We assume observed case reports (y) are drawn from a Negative Binomial distribution subject to a constant reporting fraction (ρ) and dispersion parameter τ ,

$$y_t \sim \text{Negative Binomial}(\rho I_t, \tau). \quad (13)$$

Table 1: Transitions in the SEIR model. We show the deterministic transmission rate for clarity, but our model uses the stochastic transmission rate.

Transition	$(\Delta S, \Delta E, \Delta I)$	Propensity
birth	$(1, 0, 0)$	$N_t \mu_t$
transmission (deterministic)	$(-1, 1, 0)$	$SI\beta_t/N_t$
transmission (stochastic)	$(-k, k, 0)$	$\frac{S}{k} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j+1} \tau_f^{-1} \ln(1 + (\beta_t I/N_t)) \tau_f (S - j)$
symptomatic (infectious)	$(0, -1, 1)$	$E\eta$
imported infections	$(0, 0, 1)$	ψ_t
recovery	$(0, 0, -1)$	$I\gamma$