
Supported backward leg lift

Hip extension / 6-10 each side



START / steady stance FINISH / small backward lift

MAIN: Gluteus maximus moves the thigh backward.

SECONDARY: Hamstrings assist hip extension.

BASE / SUPPORT: Stance-leg hip and ankle muscles and trunk control keep the pelvis level; hands assist balance on a fixed support.

01 / Check the setup

Begin without bands, holding a fixed counter or stable chair against a wall.

02 / Set the base and brace

Stand upright with feet under hips and knees soft. Keep ribs over the pelvis and toes facing forward.

03 / Move with control

Move one leg slightly backward from the hip while breathing normally, then return gently.

04 / Limit the range and return

Stop before the lower back arches, pelvis tips forward or the body leans. A small arc is enough.

WATCH FOR: Do not lift higher by bending the spine or swinging the leg. Keep the stance foot planted. EASIER: Use a smaller movement with more hand support, or omit if balance or symptoms limit control.

Soft joints, normal breathing. Colours describe exercise roles, not measured activation. Hidden muscles are named in text. No fixed resistance is prescribed.

Stop for pain, new numbness or weakness, dizziness, loss of grip, moving supports or damaged bands. Keep the recoil path clear. Follow manufacturer limits and individual clinical restrictions.

Sources: NHS - Strength exercises. Full source links: project100.fit/downloads/kinetics/index.html