

# Analysis 1 – Practicals

Lukas Prokop

October 2015 to January 2016

## Contents

<b>1</b>	<b>Exercise 1</b>	<b>6</b>
<b>2</b>	<b>Exercise 2</b>	<b>6</b>
<b>3</b>	<b>Exercise 3</b>	<b>6</b>
<b>4</b>	<b>Exercise 4</b>	<b>7</b>
<b>5</b>	<b>Exercise 5</b>	<b>8</b>
<b>6</b>	<b>Exercise 6</b>	<b>9</b>
6.1	Exercise 6a . . . . .	9
6.2	Exercise 6b . . . . .	9
<b>7</b>	<b>Exercise 7</b>	<b>9</b>
7.1	Exercise 7a . . . . .	9
7.2	Exercise 7b . . . . .	10
7.3	Exercise 7c . . . . .	10
<b>8</b>	<b>Exercise 8</b>	<b>10</b>
<b>9</b>	<b>Exercise 9</b>	<b>10</b>
9.1	Exercise 9a . . . . .	10
9.2	Exercise 9b . . . . .	11
<b>10</b>	<b>Reminder</b>	<b>11</b>
<b>11</b>	<b>Exercise 10</b>	<b>12</b>
11.1	Exercise 10a . . . . .	12
11.2	Exercise 10b . . . . .	12
<b>12</b>	<b>Exercise 11</b>	<b>13</b>
12.1	Exercise 11.1 . . . . .	13
12.2	Exercise 11.2 . . . . .	13

<b>13 Exercise 18</b>	<b>13</b>
<b>14 Exercise 19</b>	<b>14</b>
14.1 Exercise 19.1 . . . . .	14
14.2 Exercise 19.2 . . . . .	15
<b>15 Exercise 20</b>	<b>15</b>
<b>16 Exercise 21</b>	<b>16</b>
16.1 Exercise 21a . . . . .	16
16.2 Exercise 21b . . . . .	17
<b>17 Exercise 22</b>	<b>17</b>
<b>18 Exercise 23</b>	<b>18</b>
<b>19 Exercise 24</b>	<b>20</b>
<b>20 Exercise 25</b>	<b>20</b>
20.1 Exercise 25.a . . . . .	21
20.2 Exercise 25.b . . . . .	21
20.3 Exercise 25.c . . . . .	22
20.4 Exercise 25.d . . . . .	22
20.5 Exercise 25.e . . . . .	22
<b>21 Exercise 26</b>	<b>22</b>
21.1 Exercise 26.1: $\sum_{k=0}^n \binom{2n}{2k} = \sum_{k=1}^n \binom{2n}{2k-1}$ - approach 1 . . . . .	23
21.2 Exercise 26.1: $\sum_{k=0}^n \binom{2n}{2k} = \sum_{k=1}^n \binom{2n}{2k-1}$ - approach 2 . . . . .	24
21.3 Exercise 26.2: $\sum_{k=0}^n \binom{2n}{2k}$ . . . . .	24
<b>22 Exercise 27</b>	<b>24</b>
<b>23 Exercise 28</b>	<b>26</b>
<b>24 Exercise 29</b>	<b>27</b>
<b>25 Exercise 33</b>	<b>28</b>
<b>26 Exercise 34</b>	<b>28</b>
26.1 Exercise 34.1 . . . . .	28
26.2 Exercise 34.2 . . . . .	29
<b>27 Exercise 35</b>	<b>29</b>
27.1 Part 1 . . . . .	29
27.2 Part 2 . . . . .	30
<b>28 Exercise 36</b>	<b>30</b>

<b>29 Exercise 37</b>	<b>31</b>
<b>30 Exercise 38</b>	<b>32</b>
<b>31 Exercise 39</b>	<b>32</b>
31.1 Exercise 39a . . . . .	32
31.2 Exercise 39b . . . . .	32
<b>32 Exercise 40</b>	<b>33</b>
32.1 Exercise 40.a.1 . . . . .	33
32.2 Exercise 40.a.2 . . . . .	33
32.3 Exercise 40.b.1 . . . . .	33
32.4 Exercise 40.b.2 . . . . .	33
<b>33 Exercise 41</b>	<b>34</b>
<b>34 Exercise 42</b>	<b>34</b>
34.1 Exercise 42.a . . . . .	34
34.2 Exercise 42.b . . . . .	35
34.3 Exercise 42.c . . . . .	36
34.4 Exercise 42.d . . . . .	36
<b>35 Exercise 43</b>	<b>37</b>
35.1 Examples for $q = 1$ . . . . .	37
35.2 Proof for $q < 1$ . . . . .	37
<b>36 Exercise 44</b>	<b>38</b>
<b>37 Exercise 45</b>	<b>38</b>
<b>38 Exercise 46</b>	<b>39</b>
<b>39 Exercise 47</b>	<b>39</b>
39.1 Exercise 47.a . . . . .	40
39.2 Exercise 47.a, radical variant . . . . .	40
39.3 Exercise 47.b . . . . .	40
<b>40 Exercise 48</b>	<b>40</b>
<b>41 Exercise 49</b>	<b>41</b>
41.1 Shorter valid solution . . . . .	42
<b>42 Exercise 50</b>	<b>42</b>
42.1 Special case: $\mathbf{q} = \mathbf{0}$ implies a Cauchy sequence . . . . .	42
42.2 An observation . . . . .	42
42.3 Complete induction over $\mathbf{k} \in \mathbb{N}$ . . . . .	42

42.4	Selection of $\varepsilon$ . . . . .	43
42.5	Is $\varepsilon$ an appropriate constant? . . . . .	43
42.6	Does it also hold for $ x_{n+1} - x_n  <  x_n - x_{n-1} $ ? . . . . .	43
<b>43</b>	<b>Exercise 51</b>	<b>43</b>
<b>44</b>	<b>Exercise 52</b>	<b>44</b>
44.1	Proof direction $\Leftarrow$ . . . . .	44
44.2	Proof direction $\Rightarrow$ . . . . .	45
<b>45</b>	<b>Exercise 53</b>	<b>45</b>
45.1	Case 1: $n = 2k$ with $k \in \mathbb{N}$ . . . . .	45
45.2	Case 1.a: $k = 2l$ with $l \in \mathbb{N}$ . . . . .	46
45.3	Case 1.b: $k = 2l + 1$ with $l \in \mathbb{N}$ . . . . .	46
45.4	Case 2: $n = 2k + 1$ with $k \in \mathbb{N}$ . . . . .	46
45.5	Putting it together . . . . .	46
45.6	Limes . . . . .	46
<b>46</b>	<b>Exercise 54</b>	<b>46</b>
<b>47</b>	<b>Exercise 55</b>	<b>47</b>
47.1	Exercise 55.a . . . . .	47
47.1.1	Lathos' solution . . . . .	47
47.2	Exercise 55.b . . . . .	48
47.3	Exercise 55.c . . . . .	48
47.4	Exercise 55.d . . . . .	48
47.5	Exercise 55.e . . . . .	48
<b>48</b>	<b>Exercise 56</b>	<b>49</b>
<b>49</b>	<b>Exercise 57</b>	<b>49</b>
<b>50</b>	<b>Exercise 58</b>	<b>49</b>
<b>51</b>	<b>Exercise 59</b>	<b>50</b>
<b>52</b>	<b>Exercise 60</b>	<b>50</b>
52.1	Exercise 60.a . . . . .	50
52.2	Exercise 60.b . . . . .	50
<b>53</b>	<b>Exercise 61</b>	<b>50</b>
53.1	Exercise 61.a: $\sum_{n=1}^{\infty} \frac{1}{n^n} z^n$ . . . . .	50
53.2	Exercise 61.b: $\sum_{n=1}^{\infty} \frac{(n!)^2}{(2n)!} z^n$ . . . . .	50
53.3	Exercise 61.c: $\sum_{n=0}^{\infty} \frac{2^n}{n!} (z + i)^n$ . . . . .	50
53.4	Exercise 61.d: $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{z^{2n-1}}{(2n-1)!}$ . . . . .	50

<b>54 Exercise 62</b>	<b>50</b>
<b>55 Exercise 63</b>	<b>51</b>
<b>56 Exercise 64</b>	<b>51</b>
<b>57 Exercise 65</b>	<b>51</b>
<b>58 Exercise 66</b>	<b>51</b>
<b>59 Exercise 67</b>	<b>52</b>
59.1 Exercise 67.a: $ f $ is continuous, then $f$ is continuous . . . . .	52
59.2 Exercise 67.b: $f$ and $g$ are continuous, then $\max\{f, g\}$ is continuous . . . . .	52
59.3 Exercise 67.c: $fg$ is continuous, then $f$ and $g$ are continuous . . . . .	53
<b>60 Exercise 68</b>	<b>53</b>
<b>61 Exercise 72</b>	<b>53</b>
<b>62 Exercise 73</b>	<b>54</b>
<b>63 Exercise 74</b>	<b>54</b>
63.1 $f$ is continuous . . . . .	54
63.2 Find $f(x) = 0 = x^2 - 2$ . . . . .	54
63.3 Using the sequence theorem . . . . .	55
<b>64 Exercise 75</b>	<b>55</b>
64.1 Exercise 75.a: $\lim_{z \rightarrow 1} \frac{z^m - 1}{z^n - 1}$ with $m, n \in \mathbb{N}_+$ . . . . .	55
64.2 Exercise 75.b: $\lim_{x \rightarrow \infty} \sqrt{x - \sqrt{x}} - \sqrt{x}$ . . . . .	55
<b>65 Supporting sessions</b>	<b>55</b>
<b>66 Exercise 79</b>	<b>56</b>

## 1 Exercise 1

**Exercise 1.** Let  $p, q$  and  $r$  be statements. Prove the distributive law using the truth table:

$$p \wedge (q \vee r) \Leftrightarrow (p \wedge q) \vee (p \wedge r)$$

$p$	$q$	$r$	$(q \vee r)$	$(p \wedge (q \vee r))$	$(p \wedge q)$	$(p \wedge r)$	$(p \wedge q) \vee (p \wedge r)$
0	0	0	0	0	0	0	0
0	0	1	1	0	0	0	0
0	1	0	1	0	0	0	0
0	1	1	1	0	0	0	0
1	0	0	0	0	0	0	0
1	0	1	1	1	0	1	1
1	1	0	1	1	1	0	1
1	1	1	1	1	1	1	1

Therefore the truth table of both statements is equivalent. Two boolean statements are equivalent iff their truth table is equivalent.

## 2 Exercise 2

**Exercise 2.** Formalize the following colloquial combination of statements  $p, q$  and  $r$  in propositional calculus. Furthermore always create the negation:

- “Under the assumption, that  $p$  or  $q$  holds, we conclude that  $r$  cannot be true.”
- “It’s a requirement for  $r$ , that  $p$  and  $q$  hold.”
- “ $p$  or  $q$  holds, but  $p$  and  $q$  exclude each other”

- “Under the assumption, that  $p$  or  $q$  holds, we conclude that  $r$  cannot be true.”

$$(p \vee q) \rightarrow \neg r$$

Negation:  $(p \vee q) \wedge r$

- “It’s a requirement for  $r$ , that  $p$  and  $q$  hold.”

$$r \rightarrow (p \wedge q)$$

Negation:  $r \wedge (\neg p \vee \neg q)$

- “ $p$  or  $q$  holds, but  $p$  and  $q$  exclude each other”

$$(p \vee q) \wedge \neg(p \wedge q)$$

$$\Leftrightarrow (p \dot{\vee} q) \Leftrightarrow (p \oplus q)$$

Negation:  $p \leftrightarrow q$

## 3 Exercise 3

**Exercise 3.** Mister Travelmuch bought a Eurail ticket in August 1980 and has organized a large journey. When moving flats, he list his photo album, he tries to remember, which cities of Paris, Madrid and Rome he visited.

He remembers:

- If he was not in Madrid, then he was in Paris and Rome.
- If he was in Paris, he was not in Madrid and not in Rome.
- If he was not in Paris, he was also not in Rome.

Use appropriate variables for the statements and help Mister Travelmuch determining which cities (or city) he visited in 1980.

Let  $M$ ,  $P$  and  $R$  be visits to Madrid, Paris and Rome respectively. We formalize:

$$\begin{aligned}\neg M &\implies (P \wedge R) \\ P &\implies (\neg M \wedge \neg R) \\ \neg P &\implies \neg R\end{aligned}$$

As far as all three conditions need to be satisfied, we conjoint them:

$$[\neg M \rightarrow (P \wedge R)] \wedge [P \rightarrow (\neg M \wedge \neg R)] \wedge [\neg P \rightarrow \neg R]$$

We apply  $(a \rightarrow b) \Leftrightarrow (\neg a \vee b)$  to all three statements:

$$[\neg(\neg M) \vee (P \wedge R)] \wedge [\neg P \vee (\neg M \wedge \neg R)] \wedge [\neg(\neg P) \vee \neg R]$$

...and  $\neg(\neg A) \Leftrightarrow A$ :

$$[M \vee (P \wedge R)] \wedge [\neg P \vee (\neg M \wedge \neg R)] \wedge [P \vee \neg R]$$

...and the distributive law holds:

$$[(M \vee P) \wedge (M \vee R)] \wedge [(\neg P \wedge \neg M) \vee (\neg P \wedge \neg R)] \wedge [P \vee \neg R]$$

We reorder statements:

$$[(M \vee P) \wedge (M \vee R) \wedge (P \vee \neg R)] \wedge [(\neg P \wedge \neg M) \vee (\neg P \wedge \neg R)]$$

...and again the distributive law:

$$\begin{aligned}&[(M \vee P) \wedge (M \vee R) \wedge (P \vee \neg R) \wedge (\neg P \wedge \neg M)] \vee [(M \vee P) \wedge (M \vee R) \wedge (P \vee \neg R) \wedge (\neg P \wedge \neg R)] \\ &[(M \vee P) \wedge \neg P \wedge \neg M] \vee [(M \vee P) \wedge (M \vee R) \wedge (P \vee \neg R) \wedge (\neg P \wedge \neg R)]\end{aligned}$$

The left-hand side cannot be satisfied, but  $M \wedge \neg P \wedge \neg R$  holds for the right side. So,

- In 1980, he was in Madrid.
- In 1980, he was not in Paris.
- In 1980, he was not in Rome.

## 4 Exercise 4

**Exercise 4.** Let  $X$  be a set. Formalize the following colloquial combination of statements  $p(x)$ ,  $q(x)$ ,  $r(x)$  and  $s(x, y)$  with the help of quantifiers. Also create the negation:

1. "For all elements  $x$  of the set  $X$  for which  $p(x)$  holds, also  $q(x)$  or  $r(x)$  holds."
2. "For all  $x$  in  $X$ , there is one  $y$  in  $Y$  such that  $s(x, y)$  holds."
3. "If  $p(x)$  is not wrong for all  $x$  in  $X$ , then  $q(y)$  is true for at least one  $y$  in  $Y$ ."

1. “For all elements  $x$  of the set  $X$  for which  $p(x)$  holds, also  $q(x)$  or  $r(x)$  holds.”

$$\forall x \in X : p(x) \rightarrow q(x) \vee r(x)$$

$$\text{negation: } \exists x \in X : p(x) \wedge (\neg q(x) \wedge \neg r(x))$$

2. “For all  $x$  in  $X$ , there is one  $y$  in  $Y$  such that  $s(x, y)$  holds.”

$$\forall x \in X \exists y \in Y : s(x, y)$$

$$\text{negation: } \exists x \in X \forall y \in Y : \neg s(x, y)$$

3. “If  $p(x)$  is not wrong for all  $x$  in  $X$ , then  $q(y)$  is true for at least one  $y$  in  $Y$ .”

$$(\exists x \in X : p(x)) \rightarrow (\exists y \in Y : q(y))$$

$$\text{negation: } (\exists x \in X : p(x)) \wedge (\forall y \in Y : \neg q(y))$$

## 5 Exercise 5

**Exercise 5.** Prove in three ways (direct, indirect, by contradiction):

$$\forall x \in \mathbb{R} : x^3 + 2x > 0 \Rightarrow x > 0$$

Consider  $\phi$  to be given and  $\varphi$  to be our conclusion. Then the three ways of proving work as follows:

**Direct proof**  $\phi \Rightarrow \varphi$

**Indirect proof**  $\neg\varphi \Rightarrow \neg\phi$

Because  $\varphi \vee \neg\phi \Leftrightarrow \neg\phi \vee \varphi \Leftrightarrow \phi \rightarrow \varphi$ .

**Proof by contradiction**  $(\neg(\phi \Rightarrow \varphi) \Rightarrow \perp) \Rightarrow (\phi \Rightarrow \varphi)$

Because  $((\phi \rightarrow \varphi) \vee \perp) \rightarrow (\phi \rightarrow \varphi) \Leftrightarrow (\phi \rightarrow \varphi) \rightarrow (\phi \rightarrow \varphi)$ .

**Direct proof** Assume,

$$x(x^2 + 2) > 0$$

This requires that

- both factors are non-zero
- and
  - both factors are negative, or
  - both factors are positive

So,

$$(x \neq 0 \wedge (x^2 + 2) \neq 0) \wedge [(x < 0 \wedge (x^2 + 2) < 0) \vee (x > 0 \wedge (x^2 + 2) > 0)]$$

As far as a square cannot be negative,  $(x^2 + 2) < 0$  does not hold.

$$(x \neq 0 \wedge (x^2 + 2) \neq 0) \wedge [(x > 0 \wedge (x^2 + 2) > 0)]$$

Therefore it must hold that

$$(x \neq 0) \wedge (x^2 + 2 \neq 0) \wedge (x > 0) \wedge (x^2 + 2 > 0)$$

And so it holds that  $x > 0$ .

**Indirect proof** Assume  $x \leq 0$ . Then  $x \cdot x^2 \leq 0$ . And also  $x \cdot (x^2 + 2) \leq 0$ . Which is  $x^3 + 2x \leq 0$ .



**Proof by contradiction** Assume  $x(x^2 + 2) > 0 \implies x \leq 0$ .

$$\forall x \in \mathbb{R} : x \cdot \underbrace{(x^2 + 2)}_{\geq 2} > 0 \implies x \leq 0$$

$$\forall x \in \mathbb{R} : \underbrace{x}_{\Rightarrow \geq 0} \cdot \underbrace{(x^2 + 2)}_{\geq 2} > 0 \implies x \leq 0$$

$$\forall x \in \mathbb{R} : x > 0 \implies x \leq 0$$

$\downarrow$

$$\Rightarrow \forall x \in \mathbb{R} : x \cdot (x^2 + 2) > 0 \implies x > 0$$

## 6 Exercise 6

**Exercise 6.** Let  $p, q$  and  $r$  be statements. Show that

- $(p \rightarrow q) \iff \neg(p \wedge \neg q)$  “proof by contradiction”
- $[p \rightarrow (q \vee r)] \iff [(p \wedge \neg q) \rightarrow r]$

### 6.1 Exercise 6a

$$(p \rightarrow q) \iff \neg(p \wedge \neg q)$$

$$(\neg p \vee q) \iff (\neg p \vee q)$$

### 6.2 Exercise 6b

$$(p \rightarrow (q \vee r)) \iff ((p \wedge \neg q) \rightarrow r)$$

$$\neg p \vee (q \vee r) \iff \neg(p \wedge \neg q) \vee r$$

$$(\neg p \vee q) \vee r \iff (\neg p \vee q) \vee r$$

## 7 Exercise 7

**Exercise 7.** Let  $A, B, C$  and  $D$  be sets. Prove that

- $(A \setminus B) \cap (A \setminus C) = A \setminus (B \cup C)$
- $(A \setminus B) \cap (C \setminus D) = (A \setminus D) \cap (C \setminus B)$
- $B \subseteq A \implies B = A \setminus (A \setminus B)$

### 7.1 Exercise 7a

$$(A \setminus B) \cap (A \setminus C) = A \setminus (B \cup C)$$

Let  $a$  be a variable which is true if the considered element is contained in  $A$ .  $\neg a$  analogously means not contained. Same for  $b$  and  $c$ . Then:

$$(a \wedge \neg b) \wedge (a \wedge \neg c) = a \wedge \neg(b \vee c)$$

$$a \wedge \neg b \wedge a \wedge \neg c = a \wedge (\neg b \wedge \neg c)$$

$$a \wedge \neg b \wedge \neg c = a \wedge \neg b \wedge \neg c$$

$$\top = \top$$

## 7.2 Exercise 7b

$$\begin{aligned}
 (A \setminus B) \cap (C \setminus D) &= (A \setminus D) \cap (C \setminus B) \\
 (a \wedge \neg b) \wedge (c \wedge \neg d) &= (a \wedge \neg d) \wedge (c \wedge \neg b) \\
 a \wedge \neg b \wedge c \wedge \neg d &= a \wedge \neg b \wedge c \wedge \neg d \\
 \top &= \top
 \end{aligned}$$

## 7.3 Exercise 7c

$$B \subseteq A \Rightarrow B = A \setminus (A \setminus B)$$

$$\forall x \in X : (x \in B \rightarrow x \in A) \Rightarrow [x \in B \leftrightarrow x \in A \wedge \neg(x \in A \wedge x \notin B)]$$

$$\forall x \in X : (x \in B \rightarrow x \in A) \Rightarrow \left[ x \in B \leftrightarrow x \in A \wedge \underbrace{(x \notin A \vee x \in B)}_{\perp} \right]$$

$$\forall x \in X : (x \in B \rightarrow x \in A) \Rightarrow [x \in B \leftrightarrow x \in A \wedge x \in B]$$

$$\forall x \in X : (x \in B \rightarrow x \in A) \Rightarrow \left[ (x \in B \rightarrow x \in A \wedge \underbrace{x \in B}_{\top}) \wedge \underbrace{(x \in A \wedge x \in B \rightarrow x \in B)}_{\top} \right]$$

$$\forall x \in X : (x \in B \rightarrow x \in A) \Rightarrow (x \in B \rightarrow x \in A)$$

## 8 Exercise 8

**Exercise 8.** Let  $X$  be a set with  $X \neq \emptyset$  and  $X \neq \{\emptyset\}$ . Of which of the following sets is (a) the set  $X$ , (b) the set  $\{X\}$ , element of subset?

$\downarrow S$	op	$\rightarrow$	$x \in S$		$X \subseteq S$
$\{\{X\}, X\}$	$\checkmark$	2nd argument	$\times$	impossible to build	
$X$	$\times$	impossible to build	$\checkmark$	$X = X$	
$\emptyset \cap \{X\} = \emptyset$	$\times$		$\times$	unless $X = \emptyset$	
$\{X\} \setminus \{\{X\}\} = \{X\}$	$\checkmark$	1st argument	$\times$	impossible to build	
$\{X\} \cup X$	$\checkmark$	1st argument	$\checkmark$	$X = X$	
$\{X\} \cup \{\emptyset\}$	$\checkmark$	1st argument	$\times$	impossible to build	

## 9 Exercise 9

$(0, \infty)$  is the set  $\mathbb{R}_{>0}$

### 9.1 Exercise 9a

Prove in three ways the following statement:

$$\forall x \in (0, \infty) \forall y \in (0, \infty) : x \neq y \Rightarrow \frac{x}{y} + \frac{y}{x} > 2$$

**direct proof**

$$\begin{aligned} x &\neq y \\ x - y &\neq 0 \\ (x - y)^2 &\neq 0 \\ (x - y)^2 &> 0 \\ x^2 - 2xy + y^2 &> 0 \\ \frac{x^2}{xy} - \frac{2xy}{xy} + \frac{y^2}{xy} &> 0 \\ \frac{x}{y} - 2 + \frac{y}{x} &> 0 \\ \frac{x}{y} + \frac{y}{x} &> 2 \end{aligned}$$

**indirect proof**

$$\begin{aligned} \forall x \in (0, \infty) \forall y \in (0, \infty) : \frac{x}{y} + \frac{y}{x} \leq 2 &\Rightarrow x = y \\ \frac{x^2}{xy} + \frac{y^2}{xy} &\leq 2 \\ x^2 + y^2 &\leq 2xy \\ x^2 - 2xy + y^2 &\leq 0 \\ (x - y)^2 &\leq 0 \\ (x - y)^2 &= 0 \\ x - y &= 0 \\ x &= y \end{aligned}$$

**proof by contradiction**

$$\begin{aligned} \forall x \in (0, \infty) \forall y \in (0, \infty) : x \neq y &\Rightarrow \frac{x}{y} + \frac{y}{x} \leq 2 \\ x - y &\neq 0 \\ x^2 - 2xy + y^2 &\neq 0 \\ x^2 - 2xy + y^2 &> 0 \\ \frac{x^2}{xy} - 2 + \frac{y^2}{xy} &> 0 \\ \underbrace{\frac{x}{y}}_{>0} + \underbrace{\frac{y}{x}}_{>0} &> 2 \end{aligned}$$

## 9.2 Exercise 9b

Let  $z = \frac{x}{y}$  and illustrate the inequality geometrically.

$$\frac{x}{y} + \frac{y}{x} > 2 \Rightarrow z + z^{-1} > 2$$

## 10 Reminder

If  $n < m$ , then the *empty sum*  $\sum_{k=m}^n a_k$  has value 0, and the *empty product*  $\prod_{k=m}^n a_k$  has value 1.




Figure 1: Plot for  $z + z^{-1} > 2$  (Exercise 9b)

## 11 Exercise 10

### Exercise 9.

- Provide a concise definition of “n is an even number” and “n is an odd number” using the existence quantifier.
- Prove  $\forall n \in \mathbb{Z} : n \text{ is even} \Leftrightarrow n^2 \text{ is even}$   
**Hint:** Prove  $\Leftarrow$  using an indirect proof.

### 11.1 Exercise 10a

$$n \text{ is even} \Rightarrow \exists a \in \mathbb{Z} : n = 2a$$

$$n \text{ is odd} \Rightarrow \exists a \in \mathbb{Z} : n = 2a + 1$$

### 11.2 Exercise 10b

$$\forall n \in \mathbb{Z} \exists a_1 \in \mathbb{Z} \exists a_2 \in \mathbb{Z} : n = 2a_1 \Leftrightarrow n^2 = 2a_2$$

**Direction  $\Rightarrow$**

$$n = 2a_1 \Rightarrow n^2 = 4a_1^2$$

$$\text{Let } a_1 = \sqrt{\frac{a_2}{2}}.$$

$$n^2 = 4 \left( \sqrt{\frac{a_2}{2}} \right)^2 \Rightarrow n^2 = 2a_2$$

Such an  $a_2$  always exists. Proof finished.

**Direction  $\Leftarrow$**

$$n^2 \neq 2a_2 \Rightarrow n \neq 2a_1$$

Taking the square root preserves the parity of the value<sup>1</sup>.

$$n = \sqrt{2a_2 + 1}$$

So  $\sqrt{2a_2 + 1}$  gives an odd number. But this structure cannot match  $2a_1$ , which represents an even number. This shows a contradiction and  $n \neq 2a_1$  holds.

<sup>1</sup>Because an even number times an even number yields an even number. An odd number times an odd number yields an odd number.

## 12 Exercise 11

**Exercise 10.** For the following statement give

1. an indirect proof
2. a proof using Exercise 6b

$$\forall x \in \mathbb{R} \forall y \in \mathbb{R} : xy \notin \mathbb{Q} \Rightarrow x \notin \mathbb{Q} \vee y \notin \mathbb{Q}$$

### 12.1 Exercise 11.1

$$\forall x \in \mathbb{R} \forall y \in \mathbb{R} : x \in \mathbb{Q} \wedge y \in \mathbb{Q} \Rightarrow xy \in \mathbb{Q}$$

$$\forall x \in \mathbb{R} \forall y \in \mathbb{R} \exists p_0, p_1 \in \mathbb{R} \exists q_0, q_1 \in \mathbb{R} \setminus \{0\} :$$

$$\left( x = \frac{p_0}{q_0} \right) \wedge \left( y = \frac{p_1}{q_1} \right) \Rightarrow \left( \exists p_2 \in \mathbb{R} \exists q_2 \in \mathbb{R} \setminus \{0\} : xy = \frac{p_2}{q_2} \right)$$

$$xy = \frac{\overbrace{p_0 p_1}^{\in \mathbb{R}}}{\underbrace{q_0 q_1}_{\in \mathbb{R} \setminus \{0\}}} \Rightarrow xy = \frac{p_2}{q_2} \text{ for } p_2 = p_0 \cdot p_1 \text{ and } q_2 = q_0 \cdot q_1$$

### 12.2 Exercise 11.2

$$(p \Rightarrow (q \vee r)) \Leftrightarrow ((p \wedge \neg q) \Rightarrow r)$$

$$(xy \notin \mathbb{Q} \Rightarrow (x \notin \mathbb{Q} \vee y \notin \mathbb{Q})) \Leftrightarrow ((xy \notin \mathbb{Q} \wedge x \in \mathbb{Q}) \Rightarrow y \notin \mathbb{Q})$$

$$\forall x, y \in \mathbb{R} : \left( \left( \nexists p_2 \in \mathbb{R} \nexists q_2 \in \mathbb{R} \setminus \{0\} : xy = \frac{p_2}{q_2} \right) \wedge \left( \exists p_0 \in \mathbb{R} \exists q_0 \in \mathbb{R} \setminus \{0\} : x = \frac{p_0}{q_0} \right) \right) \Rightarrow \left( \nexists p_1 \in \mathbb{R} \nexists q_1 \in \mathbb{R} \setminus \{0\} : y = \frac{p_1}{q_1} \right)$$

Recognize  $p_2 = p_0 \cdot p_1$  and  $q_2 = q_0 \cdot q_1$ .

Therefore the conjunction yields the  $y \notin \mathbb{Q}$  because  $x \in \mathbb{Q}$ .

$$\forall x \in \mathbb{R} \forall y \in \mathbb{R} : \left( \nexists p_1 \in \mathbb{R} \exists q_1 \in \mathbb{R} \setminus \{0\} : y = \frac{p_1}{q_1} \right) \Rightarrow \left( \nexists p_1 \in \mathbb{R} \exists q_1 \in \mathbb{R} \setminus \{0\} : y = \frac{p_1}{q_1} \right)$$

This statement is true. The proof is complete.

## 13 Exercise 18

**Exercise 11.** Let  $n \in \mathbb{N}_+$ . Show that

$$\prod_{k=2}^n \left( 1 - \frac{1}{k} \right) = \frac{1}{n}.$$

**Induction base**  $n = 1$

$$\prod_{k=2}^1 \dots = 1 = \frac{1}{1} \quad \checkmark$$

**Induction step**  $n \rightarrow n + 1$

$$\begin{aligned}
 \prod_{k=2}^{n+1} \left(1 - \frac{1}{k}\right) &= \frac{1}{n+1} \\
 \prod_{k=2}^n \left(1 - \frac{1}{k}\right) \left(1 - \frac{1}{n+1}\right) &= \frac{1}{n+1} \\
 \frac{1}{n} \left(1 - \frac{1}{n+1}\right) &= \frac{1}{n+1} \\
 \frac{1}{n} \cdot \frac{n+1-1}{n+1} &= \frac{1}{n+1} \\
 \frac{n}{n} &= 1 \quad \checkmark
 \end{aligned}$$

Actually, can be rewritten as

$$\begin{aligned}
 &\prod_{k=2}^n \left(\frac{k-1}{k}\right) \\
 &= \frac{1}{2} \cdot \frac{2}{3} \cdot \frac{3}{4} \cdot \frac{4}{5} \cdots \frac{n-1}{n} \\
 &= \frac{1}{n}
 \end{aligned}$$

So this is the multiplication equivalent of telescoping sums.

## 14 Exercise 19

**Exercise 12.**  $X$  and  $Y$  are non-empty sets and  $f : X \rightarrow Y$  is a mapping. Furthermore let  $A \subseteq X$  and  $B \subseteq Y$ .

1. Prove that  $A \subseteq f^{-1}(f(A))$  and  $B \supseteq f(f^{-1}(B))$
2. Show (by providing counterexamples) that in the inclusions of (1) no equivalence is given.

### 14.1 Exercise 19.1

Show that,

$$a \in A \Rightarrow a \in f^{-1}(f(A))$$

So we take  $a$  and map it to the codomain:

$$f(a) \in f(A)$$

We denote the result as  $y$ :

$$y := f(a)$$

Because

$$f^{-1}(y) = \{x \in A \mid f(x) \in B\}$$

we know that  $a$  originates from:

$$a \in f^{-1}(f(A))$$

It is very important here to distinguish between *domain/codomain* and *function/inverse function*. Because an inverse function implies that the corresponding function is injective. Assuming this fact, the exercise is immediate. But we are talking about domains and co-domains here.

As second exercise we need to show that,

$$B \supseteq f(f^{-1}(B))$$

We need the definition that,

$$f^{-1}(B) = \{x' \in X \mid f(x') \in B\}$$

$$y' \in f(f^{-1}(B))$$

Does  $y' \in B$  hold? Yes, because ...

$$\begin{aligned} y' \in f(f^{-1}(B)) &\Rightarrow \exists x' \in f^{-1}(B) \\ &\Rightarrow y' \in B \end{aligned}$$

## 14.2 Exercise 19.2

Show that,

$$\exists f : A \subsetneq f^{-1}(f(A))$$

We use a surjective, but not injective function.

$$\begin{aligned} f : \{1, 2\} &\rightarrow \{a\} \\ 1 &\mapsto a \\ 2 &\mapsto a \end{aligned}$$

$$\begin{aligned} A &= \{1\} \\ f(A) &= \{a\} \\ f^{-1}(f(A)) &= \{1, 2\} \end{aligned}$$

Show that,

$$\exists f : A \subsetneq f(f^{-1}(A))$$

We use an injective, but not surjective function.

$$\begin{aligned} f : \{1\} &\rightarrow \{a, b\} \\ 1 &\mapsto a \end{aligned}$$

$$\begin{aligned} B &= \{b\} \\ f^{-1}(B) &= \emptyset \\ f(f^{-1}(B)) &= \emptyset \end{aligned}$$

## 15 Exercise 20

**Exercise 13.** Prove the following variant of Bernoulli's inequality: For  $x \in \mathbb{R}$  with  $0 < x < 1$  and  $n \in \mathbb{N}_+$  it holds that

$$(1 - x)^n < \frac{1}{1 + nx}.$$

$$\begin{aligned}
(1+x)^n &\geq 1+nx \\
\frac{(1+x)^n}{1+nx} &\geq \frac{1+nx}{1+nx} \\
\frac{(1+x)^n}{1+nx} &\geq 1 \\
\frac{(1-x)^n(1+x)^n}{(1-x)^n(1+nx)} &\geq 1 \\
\frac{(1-x)^n(1+x)^n}{(1+nx)} &\geq (1-x)^n \\
\frac{((1-x)(1+x))^n}{(1+nx)} &\geq (1-x)^n \\
\frac{(1-x^2)^n}{(1+nx)} &\geq (1-x)^n \\
\text{interval } (0,1) \\
\frac{\overbrace{(1-x^2)^n}^{\text{interval } (0,1)}}{(1+nx)} &\geq (1-x)^n \\
\frac{1}{(1+nx)} &> (1-x)^n
\end{aligned}$$

## 16 Exercise 21

**Exercise 14.**  $X$  and  $Y$  are nonempty sets and  $f : X \rightarrow Y$  is a mapping.

a) Show that the following holds: For all  $A, B \subseteq X$

$$f(A \cap B) \subseteq f(A) \cap f(B).$$

b) Show that the following statements are equivalent:

1.  $f$  is injective.
2. For all  $A, B \subseteq X$  it holds that  $f(A \cap B) \supseteq f(A) \cap f(B)$
3. For all  $A, B \subseteq X$  it holds that  $f(A \cap B) = f(A) \cap f(B)$

### 16.1 Exercise 21a

Let  $C = A \cap B$ . Case distinction:

$$A = B = C$$

$$\begin{aligned}
f(A \cap B) &= \{f(x) \mid x \in A\} \\
f(A) \cap f(B) &= f(A) \\
&= \{f(x) \mid x \in A\}
\end{aligned}$$

$$C = A \dot{\cup} C = B \text{ wlog. } C = A.$$

$$\begin{aligned}
f(A \cap B) &= f(A) \\
&= \{f(x) \mid x \in A\} \\
f(A) \cap f(B) &= \{f(x) \mid x \in A\} \cap \{f(x) \mid x \in B\} \\
&= \{f(x) \mid x \in A \wedge x \notin (B \setminus A)\} \\
&= \{f(x) \mid x \in A\}
\end{aligned}$$



$$C = \emptyset$$

$$\begin{aligned} f(A \cap B) &= f(\emptyset) \\ &= \emptyset \\ f(A) \cap f(B) &= \{f(x) \mid x \in A\} \cap \{f(x) \mid x \in B\} \end{aligned}$$

So,

$$C \neq \emptyset \Rightarrow f(A \cap B) = f(A) \cap f(B)$$

But if  $C = \emptyset$ , we get zero values on the left-hand side and zero to  $|A| + |B|$  values on the right-hand side. So,

$$C = \emptyset \Rightarrow f(A \cap B) \subseteq f(A) \cap f(B)$$

## 16.2 Exercise 21b

We prove 3 with 1:

Let  $C = A \cap B$ .  $f$  is injective, meaning

$$\forall x_1, x_2 \in X : x_1 \neq x_2 \Rightarrow f(x_1) \cap f(x_2) = \emptyset$$

Case distinction:

$$A = B = C$$

$$\begin{aligned} f(A \cap B) &= \{f(x) \mid x \in A\} \\ f(A) \cap f(B) &= f(A) \\ &= \{f(x) \mid x \in A\} \end{aligned}$$

$$C = A \vee C = B$$

wlog.  $C = A$  meaning  $A \subsetneq B$

$$\begin{aligned} f(A \cap B) &= f(A) \\ &= \{f(x) \mid x \in A\} \\ f(A) \cap f(B) &= \{f(x) \mid x \in A\} \cap \{f(x) \mid x \in B\} \\ &= \{f(x) \mid x \in A \wedge x \notin (B \setminus A)\} \\ &= \{f(x) \mid x \in A\} \end{aligned}$$

$$C = \emptyset$$

$$\begin{aligned} f(A \cap B) &= f(\emptyset) \\ &= \emptyset \\ f(A) \cap f(B) &= \emptyset \end{aligned}$$

Every element in  $A$  is distinct from values in  $B$ . Therefore  $\forall x_1 \in A, x_2 \in B : f(x_1) \neq f(x_2)$  because of injectivity. The intersection of all  $f(x_i)$  is therefore empty.

## 17 Exercise 22

**Exercise 15.** Let  $n \in \mathbb{N}$ . Use the following idea to derive an equation for the sum of powers of three.

$$\sum_{k=1}^n (k^4 - (k-1)^4)$$

This sum can be written in two different ways:

- As telescoping sum (the initial and trailing value will be left)

- (First resolve the parentheses.) As combination of sums of the third, second, first and zero-th power. With that (and known equations for sums of smaller powers) we can compute  $\sum_{k=1}^n k^3$ .

We look at the telescoping sum:

$$\begin{aligned}\sum_{k=1}^n (k^4 - (k-1)^4) &= (1^4 - (1-1)^4) + (2^4 - (2-1)^4) + (3^4 - (3-1)^4) \\ &\quad + \cdots + ((n-1)^4 - ((n-1)-1)^4) + (n^4 - (n-1)^4) \\ &= -0^4 + n^4 \\ &= n^4\end{aligned}$$

Then we use the combination of sums of lower powers.

$$\begin{aligned}\sum_{k=1}^n (k^4 - (k-1)^4) &= \sum_{k=1}^n (k^4 - (k^4 - 4k^3 + 6k^2 - 4k + 1)) \\ &= \sum_{k=1}^n (k^4 - k^4 + 4k^3 - 6k^2 + 4k - 1) \\ &= \sum_{k=1}^n (4k^3 - 6k^2 + 4k - 1) \\ &= \sum_{k=1}^n 4k^3 - \sum_{k=1}^n 6k^2 + \sum_{k=1}^n 4k - \sum_{k=1}^n 1 \\ &= \sum_{k=1}^n 4k^3 - 6 \frac{2n^3 + 3n^2 + n}{6} + 4 \frac{n(n+1)}{2} - n \\ &= \sum_{k=1}^n 4k^3 - 2n^3 - n^2\end{aligned}$$

Therefore,

$$\begin{aligned}n^4 &= \sum_{k=1}^n 4k^3 - 2n^3 - n^2 \\ \sum_{k=1}^n 4k^3 &= n^4 + 2n^3 + n^2 \\ \sum_{k=1}^n k^3 &= \frac{n^4 + 2n^3 + n^2}{4}\end{aligned}$$

## 18 Exercise 23

**Exercise 16.** Let  $n \in \mathbb{N}$ . Prove that

$$\sum_{k=0}^n \binom{n}{k} = 2^n$$

and if  $n \geq 1$ ,

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = 0$$

Binomial theorem with  $x = 1, y = 1$ :

$$\sum_{k=0}^n \binom{n}{k} 1^n 1^{n-k} = (1+1)^n$$

$$\sum_{k=0}^n \binom{n}{k} = 2^n$$

If  $n \geq 1$ ,

$$\begin{aligned} \sum_{k=0}^n (-1)^k \binom{n}{k} &= \sum_{k=0}^n (-1)^k \left( \binom{n-1}{k} + \binom{n-1}{k-1} \right) \\ &= \sum_{k=0}^n (-1)^k \binom{n-1}{k} + \sum_{k=0}^n (-1)^k \binom{n-1}{k-1} \\ &= (-1)^n \binom{n-1}{n} + \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} + \sum_{k=0}^n (-1)^k \binom{n-1}{k-1} \\ &= \underbrace{(-1)^n \binom{n-1}{n}}_0 + \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} + \sum_{k=1}^n (-1)^k \binom{n-1}{k-1} + \underbrace{(-1)^0 \binom{n-1}{-1}}_0 \\ &= \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} + \sum_{k=0}^{n-1} (-1)^{k+1} \binom{n-1}{k} \\ &= \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} - (-1) \sum_{k=0}^{n-1} (-1)^{k+1} \binom{n-1}{k} \\ &= \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} - \sum_{k=0}^{n-1} (-1)^{k+2} \binom{n-1}{k} \\ &= \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} - \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} \\ &= 0 \end{aligned}$$

## 19 Exercise 24

**Exercise 17.** Let  $k, n \in \mathbb{N}_+$  with  $k \leq n$ . Determine the number of vectors of length  $k$  with pairwise different entries from  $M_n = \{1, 2, \dots, n\}$ .

This question is covered by the field of combinatorics.

$$(a_0, a_1, a_2, \dots) \neq (a_0, a_2, a_1, \dots)$$

The order of elements is relevant. Therefore a variation, not combination, is given. The number of combinations without repetitions would be given by the binomial coefficient  $\binom{n}{k}$  (the number of ways to choose  $k$  of  $n$  elements disregarding their order). For variations the formula  $n^k$  holds to select  $k$  elements among  $n$  arbitrarily often (hence with repetition).

We model the given situation as

- “variation without repetition”
- i.e. “ $k$ -permutations of  $n$ ”
- i.e. the  $k$ -th falling factorial power  $n^{\underline{k}}$  of  $n$

The formula is given by,

$$P_k^n = \frac{n!}{(n-k)!}$$

We can estimate it in the following way: Consider a combination without repetition represented by the formula  $\binom{n}{k}$ :

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

So because we have a variation, not combination, the order of elements is relevant. Therefore given some combination, there are  $k!$  possible arrangements. Given the vector (and also combination)  $(1, 2, 3)$  there are  $k!$  possible arrangements (variations)  $\{(1, 2, 3), (1, 3, 2), (2, 1, 3), (2, 3, 1), (3, 1, 2), (3, 2, 1)\}$ . Indeed it holds that

$$\frac{3!}{(3-3)!} = \frac{6}{1} = 6$$

This argument explains why  $k!$  in the denominator is omitted for variations w/o repetitions.

combinations	variations					
(123)	(123)	(132)	(213)	(231)	(312)	(321)
(124)	(124)	(142)	(214)	(241)	(412)	(421)
(134)	(134)	(143)	(314)	(341)	(413)	(431)
(234)	(234)	(243)	(324)	(342)	(423)	(432)

Table 1: Combinations and variations for  $n = 4$  of  $k = 3$

## 20 Exercise 25

**Exercise 18.** Let  $K$  be a field and  $a, b, c \in K$ . Show (using the field axioms):

- (a)  $-(-a) = a$ .
- (b)  $(-a)(-b) = ab$ .
- (c)  $a + b = a + c \Rightarrow b = c$ .
- (d) From  $a \neq 0$  and  $ab = ac$  follows  $b = c$ .

(e) Is  $a \neq 0$ , then there is exactly one  $x \in K$  with  $ax + b = c$ .

The field axioms are defined as follows:

$$\mathbf{A1} \quad \forall a, b \in K : a + b = b + a$$

$$\mathbf{A2} \quad \forall a, b, c \in K : (a + b) + c = a + (b + c)$$

$$\mathbf{A3} \quad \exists 0 \in K \forall a \in K : a + 0 = a$$

$$\mathbf{A4} \quad \forall a \in K \exists \tilde{a} : a + \tilde{a} = 0$$

$$\mathbf{M1} \quad \forall a, b \in K : a \cdot b = b \cdot a$$

$$\mathbf{M2} \quad \forall a, b, c \in K : a \cdot (b \cdot c) = (a \cdot b) \cdot c$$

$$\mathbf{M3} \quad \exists 1 \in K : a \cdot 1 = a \forall a \in K \text{ (neutral element)}$$

$$\mathbf{M4} \quad \forall a \in K \setminus \{0\} \exists \hat{a} : \hat{a} \cdot a = 1$$

$$\mathbf{D} \quad \forall a, b, c \in K : a \cdot (b + c) = a \cdot b + a \cdot c$$

## 20.1 Exercise 25.a

$$A4 \Rightarrow \forall a \in K \exists -a : a + (-a) = 0$$

$$\text{equivalence} \Rightarrow a + (-a) - (-a) = 0 - (-a)$$

$$A1 \Rightarrow a + (-a) - (-a) = -(-a) + 0$$

$$A3 \Rightarrow a + (-a) - (-a) = -(-a)$$

$$A4 \Rightarrow a + 0 = -(-a)$$

$$A3 \Rightarrow a = -(-a)$$

## 20.2 Exercise 25.b

We have proven in the lecture: **M5**:  $-a = (-1) \cdot a$

First, we show **M7**

$$= a \cdot (-a)$$

$$M5 \Rightarrow a \cdot (-1) \cdot a$$

$$M1 \Rightarrow (-1) \cdot a \cdot a$$

$$\Rightarrow -(a \cdot a)$$

Secondly, we show (actually we have already shown that in the lecture) **M6**

$$D \Rightarrow \forall a, b, c \in K : a \cdot (b + c) = a \cdot b + a \cdot c$$

$$[\text{we choose } a := a, \quad b := a, \quad c := (-a)]$$

$$\Rightarrow a \cdot (a + (-a)) = a \cdot a + a \cdot (-a)$$

$$A3 \Rightarrow a \cdot 0 = a \cdot a + a \cdot (-a)$$

$$\text{previous theorem} \Rightarrow a \cdot 0 = a \cdot a + (-(a \cdot a))$$

$$A4 \Rightarrow a \cdot 0 = 0$$

Finally, we show

$$\text{previous theorem} \Rightarrow (-a) \cdot 0 = 0$$

$$A4 \Rightarrow (-a) \cdot (b + (-b)) = 0$$

$$D \Rightarrow (-a) \cdot b + (-a)(-b) = 0$$

$$\text{equivalence} \Rightarrow ab + (-a)b + (-a)(-b) = ab + 0$$

$$M1 \Rightarrow ab + (-a)b + (-a)(-b) = 0 + ab$$

$$A3 \Rightarrow ab + (-a)b + (-a)(-b) = ab$$

$$M6 \Rightarrow ab - ab + (-a)(-b) = ab$$

$$A4 \Rightarrow 0 + (-a)(-b) = ab$$

$$A3 \Rightarrow (-a)(-b) = ab$$

### 20.3 Exercise 25.c

$$a + b = a + c$$

$$\text{equivalence} \Rightarrow a + b + (-a) = a + c + (-a)$$

$$A1 \Rightarrow (a + (-a)) + b = (a + (-a)) + c$$

$$A4 \Rightarrow 0 + b = 0 + c$$

$$A3 \Rightarrow b = c$$

### 20.4 Exercise 25.d

$$a \neq 0 \wedge ab = ac$$

$$\text{equivalence} \Rightarrow aba^{-1} = aca^{-1}$$

$$M1 \Rightarrow aa^{-1}b = aa^{-1}c$$

$$M4 \Rightarrow 1b = 1c$$

$$M3 \Rightarrow b = c$$

### 20.5 Exercise 25.e

Proof by contradiction. Assume  $x_1, x_2 \in K$  with  $x_1 \neq x_2$  then  $\exists r \in K$ :

$$ax_1 = r \quad ax_2 = r$$

$$ax_1 = ax_2$$

$$\Rightarrow ax_1 = ax_2$$

$$\text{equivalence} \Rightarrow a^{-1}ax_1 = a^{-1}ax_2$$

$$M4 \Rightarrow 1x_1 = 1x_2$$

$$M3 \Rightarrow x_1 = x_2$$

This is a contradiction to our assumption  $x_1 \neq x_2$ . Therefore  $x$  is distinct.

## 21 Exercise 26

**Exercise 19.** Let  $n \in \mathbb{N}_+$ . Prove that

$$\sum_{k=0}^n \binom{2n}{2k} = \sum_{k=1}^n \binom{2n}{2k-1} = 2^{2n-1}.$$

**21.1 Exercise 26.1:**  $\sum_{k=0}^n \binom{2n}{2k} = \sum_{k=1}^n \binom{2n}{2k-1}$  - **approach 1**

*Proof.*

$$\begin{aligned} \sum_{k=0}^n \binom{2n}{2k} &= \sum_{k=1}^{n-1} \binom{2n}{2k} + 1 + 1 \\ &= \sum_{k=1}^{n-1} \left[ \binom{2n-1}{2k} + \binom{2n-1}{2k-1} \right] + 1 + 1 \\ &= \sum_{k=1}^{n-1} \binom{2n-1}{2k} + \sum_{k=1}^{n-1} \binom{2n-1}{2k-1} + 1 + 1 \\ &= \sum_{k=2}^n \binom{2n-1}{2(k-1)} + \sum_{k=1}^{n-1} \binom{2n-1}{2k-1} + 1 + 1 \\ &= \sum_{k=1}^n \binom{2n-1}{2k-2} + \sum_{k=1}^{n-1} \binom{2n-1}{2k-1} + 1 \\ &= \sum_{k=1}^{n-1} \binom{2n-1}{2k-2} + \sum_{k=1}^{n-1} \binom{2n-1}{2k-1} + \binom{2n-1}{2n-2} + 1 \\ &= \sum_{k=1}^{n-1} \left[ \binom{2n-1}{2k-2} + \binom{2n-1}{2k-1} \right] + \binom{2n-1}{2n-2} + 1 \\ &= \sum_{k=1}^{n-1} \binom{2n}{2k-1} + \left[ 1 + \binom{2n-1}{2n-2} \right] \\ &= \sum_{k=1}^{n-1} \binom{2n}{2k-1} + \left[ \binom{2n-1}{2n-1} + \binom{2n-1}{2n-2} \right] \\ &= \sum_{k=1}^{n-1} \binom{2n}{2k-1} + \binom{2n}{2n-1} \\ &= \sum_{k=1}^n \binom{2n}{2k-1} \end{aligned}$$

□

### 21.2 Exercise 26.1: $\sum_{k=0}^n \binom{2n}{2k} = \sum_{k=1}^n \binom{2n}{2k-1}$ - approach 2

*Proof.* Idea: Consider  $(1-1)^{2n}$  and split even/odd  $ks$ .

$$\begin{aligned}
 (1-1)^{2n} &= \sum_{k=0}^{2n} \binom{2n}{k} (-1)^k 1^{2n-k} && \text{[binomial theorem]} \\
 0 &= \sum_{k=0}^n \binom{2n}{2k} (-1)^{2k} + \sum_{k=1}^n \binom{2n}{2k-1} (-1)^{2k-1} \\
 &= \sum_{k=0}^n \binom{2n}{2k} + \sum_{k=1}^n \binom{2n}{2k-1} (-1) && [(-1)^{\text{even}} \text{ is } 1, (-1)^{\text{odd}} \text{ is } -1] \\
 &= \sum_{k=0}^n \binom{2n}{2k} - \sum_{k=1}^n \binom{2n}{2k-1} \\
 \sum_{k=1}^n \binom{2n}{2k-1} &= \sum_{k=0}^n \binom{2n}{2k}
 \end{aligned}$$

□

### 21.3 Exercise 26.2: $\sum_{k=0}^n \binom{2n}{2k}$

*Proof.* Idea: Consider  $(1+1)^{2n}$  and odd+even provides the factor 2 we need to divide with.

$$\begin{aligned}
 (1+1)^{2n} &= \sum_{k=0}^{2n} \binom{2n}{k} 1^k 1^{2n-k} && \text{[binomial theorem]} \\
 2^{2n} &= \sum_{k=0}^{2n} \binom{2n}{k} \\
 &= \sum_{k=0}^n \binom{2n}{2k} + \sum_{k=1}^n \binom{2n}{2k-1} && \text{[split even and odd]} \\
 &= \sum_{k=0}^n \binom{2n}{2k} + \sum_{k=0}^n \binom{2n}{2k} && \text{[from previous result]} \\
 &= 2 \sum_{k=0}^n \binom{2n}{2k} \\
 \frac{2^{2n}}{2} &= \sum_{k=0}^n \binom{2n}{2k} \\
 2^{2n-1} &= \sum_{k=0}^n \binom{2n}{2k}
 \end{aligned}$$

□

## 22 Exercise 27

**Exercise 20.** Let  $x \in \mathbb{R} \setminus \{0\}$ . Show: Let  $x + \frac{1}{x} \in \mathbb{Z}$ , then  $x^n + \frac{1}{x^n} \in \mathbb{Z}$  for all  $n \in \mathbb{N}$  (Remark: Consider  $(x + \frac{1}{x})^n$ .)

So we need to show that,

$$x \in \mathbb{R} \setminus \{0\} : \forall n \in \mathbb{N} : x + \frac{1}{x} \in \mathbb{Z} \implies x^n + \frac{1}{x^n} \in \mathbb{Z}$$



First we need to cover some fundamentals,

- $a, b \in \mathbb{Z} \implies (a + b) \in \mathbb{Z}$
- $a, b \in \mathbb{Z} \implies (a \cdot b) \in \mathbb{Z} \implies \forall n \in \mathbb{N} : a^n \in \mathbb{Z}$

*Proof.* **IB:**  $n = 0$

$$\begin{aligned}
 \forall x \in \mathbb{R} \setminus \{0\} : x + \frac{1}{x} &\in \mathbb{Z} \\
 \implies x = 1 : 1 + \frac{1}{1} &\in \mathbb{Z} \\
 \implies \forall x \in \mathbb{R} \setminus \{0\} : x^0 + \frac{1}{x^0} &\in \mathbb{Z} \\
 \implies \forall x \in \mathbb{R} \setminus \{0\} : n = 0 : x^n + \frac{1}{x^n} &\in \mathbb{Z}
 \end{aligned}$$

**IB:**  $n = 1$

$$\begin{aligned}
 \forall x \in \mathbb{R} \setminus \{0\} : x + \frac{1}{x} &\in \mathbb{Z} \\
 \implies \forall x \in \mathbb{R} \setminus \{0\} : x^1 + \frac{1}{x^1} &\in \mathbb{Z} \\
 \implies \forall x \in \mathbb{R} \setminus \{0\} : n = 1 : x^n + \frac{1}{x^n} &\in \mathbb{Z}
 \end{aligned}$$

**IS:**  $n \rightarrow n + 1$  Okay, how does the induction step for an implication look like?

$$\begin{aligned}
 ((a \rightarrow b) \rightarrow (a \rightarrow d)) &= \neg(\neg a \vee b) \vee (\neg a \vee d) \\
 &= (a \wedge \neg b) \vee \neg a \vee d \\
 &= ((a \vee \neg a) \wedge (\neg a \vee \neg b)) \vee d \\
 &= (\neg a \vee \neg b) \vee d \\
 &= (a \wedge b) \rightarrow d
 \end{aligned}$$

Therefore we can assume

$$\left(x + \frac{1}{x} \in \mathbb{Z}\right) \wedge \left(x^n + \frac{1}{x^n} \in \mathbb{Z}\right)$$

and need to prove that this follows:

$$x^{n+1} + \frac{1}{x^{n+1}} \in \mathbb{Z}$$

$$\begin{aligned}
 \left(x^n + \frac{1}{x^n} \in \mathbb{Z}\right) &= \left(x^n + \frac{1}{x^n}\right) \left(x + \frac{1}{x}\right) \in \mathbb{Z} \\
 &= \left(x^n \cdot x + \frac{1}{x^n} \cdot x + x^n \cdot \frac{1}{x} + \frac{1}{x^n} \cdot \frac{1}{x}\right) \in \mathbb{Z} \\
 &= \left(x^{n+1} + x^{-n+1} + x^{n-1} + x^{-n-1}\right) \in \mathbb{Z} \\
 &= \left(x^{n+1} + x^{-n-1} + x^{n-1} + x^{-n+1}\right) \in \mathbb{Z} \\
 &= \left(x^{n+1} + \frac{1}{x^{n+1}}\right) + \underbrace{\left(x^{n-1} + \frac{1}{x^{n-1}}\right)}_{\substack{\in \mathbb{Z} \text{ because of induction hypothesis} \\ \text{and we have a 2-step induction}}} \in \mathbb{Z} \\
 &= \left(x^{n+1} + \frac{1}{x^{n+1}}\right) \in \mathbb{Z}
 \end{aligned}$$

□

## 23 Exercise 28

**Exercise 21.** Let  $K$  be an ordered field and  $a, b \in K_+$ . Show:

$$a < b \Rightarrow a^2 < b^2$$

Especially the mapping  $f : K_+ \cup \{0\} \rightarrow K_+ \cup \{0\}, a \mapsto a^2$  is injective.

We already know,

$$\mathbf{U1} \quad \forall a, b \in K : a < b \Leftrightarrow b > a$$

$$\mathbf{U2} \quad \forall a \in K : a^2 = a \cdot a$$

$$\mathbf{U3} \quad \forall c \in K_+ : a > b \Rightarrow ac > bc$$

$$\mathbf{M1} \quad \forall a, b \in K : a \cdot b = b \cdot a$$

*Proof.*

$$\begin{aligned} a < b : \quad & \mathbf{U1} \Rightarrow b > a \\ & \mathbf{U1} \Rightarrow b \cdot a > a \cdot a && [\text{yes, } a \text{ originates from } K_+] \\ & \mathbf{U2} \Rightarrow b \cdot a > a^2 \\ b > a : \quad & \mathbf{U1} \Rightarrow b \cdot b > a \cdot b && [\text{yes, } b \text{ originates from } K_+] \\ & \mathbf{U2} \Rightarrow b^2 > a \cdot b \\ & \mathbf{M1} \Rightarrow b^2 > b \cdot a \\ b^2 > b \cdot a \wedge b \cdot a > a^2 : \quad & \mathbf{U3} \Rightarrow b^2 > a^2 \\ & \mathbf{U1} \Rightarrow a^2 < b^2 \\ & \Rightarrow \forall a, b \in K_+ : a < b \Rightarrow a^2 < b^2 \end{aligned}$$

□

Injectivity:

$$\forall a_1, a_2 \in K_+ \cup \{0\} : a_1 \neq a_2 \Rightarrow a_1^2 \neq a_2^2$$

*Proof.* First we consider  $a = 0$ . In this case,  $a = 0$  and  $a^2 = a \cdot a = 0 \cdot 0 = 0$  according to the axiom  $0 \cdot a = 0$  we have proven in the lecture. So for  $a = 0$ , there is only one  $a$  for which the square is zero, which is 0.

We can proceed in  $K_+$ . Proof by contradiction:

$$\exists a_1, a_2 \in K_+ : a_1 \neq a_2 \Rightarrow a_1^2 = a_2^2$$

$$a_1 \neq a_2 \Leftrightarrow a_1 < a_2 \vee a_1 > a_2$$

because  $a_1$  and  $a_2$  are elements of an ordered field.

**Case 1:**  $a_1 < a_2$

$$a_1 < a_2 \Rightarrow a_1^2 < a_2^2$$

**Case 2:**  $a_1 > a_2$

$$a_1 > a_2 \Rightarrow a_1^2 > a_2^2$$

Therefore either  $a_1^2 < a_2^2$  or  $a_1^2 > a_2^2$ . So

$$a_1^2 \neq a_2^2$$

This contradicts and therefore  $\nexists a_1, a_2 \in K_+ : a_1 \neq a_2 \Rightarrow a_1^2 = a_2^2$  or because we covered  $a = 0$ ,

$$\nexists a_1, a_2 \in K_+ \cup \{0\} : a_1 \neq a_2 \Rightarrow a_1^2 = a_2^2$$

□

## 24 Exercise 29

**Exercise 22.** Let  $K$  be an ordered field and  $a, b \in K$ . Show:

$$|a + b| = |a| + |b| \Leftrightarrow ab \geq 0$$

Triangular inequality:

$$\forall a, b \in K : |a + b| \leq |a| + |b|$$

Absolute values are defined with,

$$|a| = \begin{cases} a & a \in K_+ \\ 0 & a = 0 \\ -a & a \in K_- \end{cases}$$

*Proof.* Case distinction:

$$a = 0, b = 0$$

$$\begin{aligned} |a + b| &\leq |a| + |b| \\ |a + 0| &\leq |a| + |0| \\ A3 \Rightarrow |a| &\leq |a| + 0 \\ A3 \Rightarrow |a| &= |a| \end{aligned}$$

$$a > 0, b = 0$$

$$\begin{aligned} |a + b| &\leq |a| + |b| \\ |a + 0| &\leq |a| + |0| \\ A3 \Rightarrow |a| &\leq |a| + 0 \\ A3 \Rightarrow |a| &= |a| \end{aligned}$$

$$a = 0, b > 0$$

$$\begin{aligned} |a + b| &\leq |a| + |b| \\ |0 + b| &\leq |0| + |b| \\ A3 \Rightarrow |b| &\leq 0 + |b| \\ A3 \Rightarrow |b| &= |b| \end{aligned}$$

$$a > 0, b > 0$$

$$\begin{aligned} \underbrace{|a + b|}_{\in K_+} &\leq \underbrace{|a|}_{\in K_+} + \underbrace{|b|}_{\in K_+} \\ (a + b) &\leq (a) + (b) \\ A2 \Rightarrow a + b &\leq a + b \\ a + b &= a + b \end{aligned}$$

□

## 25 Exercise 33

$$\begin{aligned}
 & [a_n, b_n], [c_n, d_n], a_n \leq \alpha \leq b_n, c_n \leq \gamma \leq d_n \\
 & \forall \varepsilon > 0 \exists N(\varepsilon) : |a_n - b_n| < \varepsilon \forall n \geq N(\varepsilon) \\
 & \left[ \frac{1}{b_n}, \frac{1}{a_n} \right] \rightarrow \frac{1}{b_n} \leq \frac{1}{b_{n+1}} \leq \frac{1}{\alpha} \leq \frac{1}{a_{n+1}} \leq \frac{1}{a_n} \\
 & \left| \frac{1}{b_n} - \frac{1}{a_n} \right| = \frac{a_n - b_n}{a_n b_n} = \frac{|a_n - b_n|}{|a_n| |b_n|} \leq \frac{\varepsilon}{|a_1| \alpha} = \varepsilon'
 \end{aligned}$$

Important: our approximation  $a_n \geq a_1 > 0$  and  $b_n \geq \alpha$  is independent of  $n$ !

$$\begin{aligned}
 & \forall \varepsilon' > 0 \exists N(\varepsilon') : \left| \frac{1}{b_n} - \frac{1}{a_n} \right| < \varepsilon' \\
 & |a_n c_n - b_n d_n| = |a_n c_n - \alpha c_n + \alpha c_n - \alpha c_n + \alpha \gamma - b_n \gamma + b_n \gamma - b_n d_n| \\
 & \leq \underbrace{|a_n - \alpha|}_{< \varepsilon} \underbrace{|c_n|}_{\leq \gamma} + \underbrace{|\alpha|}_{< \varepsilon} \underbrace{|c_n - \gamma|}_{< \varepsilon} + \underbrace{|\gamma|}_{1} \underbrace{|\alpha - b_n|}_{< \varepsilon} + \underbrace{|b_n|}_{b_1} \underbrace{|\gamma - d_n|}_{< \varepsilon} < \varepsilon \underbrace{(2\gamma + \alpha + b_1)}_{=c} = \varepsilon'
 \end{aligned}$$

## 26 Exercise 34

**Exercise 23.** Let  $f : X \rightarrow Y$  be a mapping. Prove that:

1. If a mapping  $g : Y \rightarrow X$  with  $g \circ f = \text{id}_X$  exists,  $f$  is injective.
2. If a mapping  $h : Y \rightarrow X$  with  $f \circ h = \text{id}_Y$  exists,  $f$  is surjective.

**Remark.**  $\text{id}_X$  is the identity function over the set  $X$ . The identity function is always defined as  $f : X \rightarrow X$  with  $x \mapsto x$ .

### 26.1 Exercise 34.1

So given that  $g : Y \rightarrow X$  exists with  $g \circ f = \text{id}_X$ , let  $x \in X$ .

$$x \in X \Rightarrow f(x) \in Y \Rightarrow g(f(x)) = x \Leftrightarrow \text{id}_X(x) = x$$

To show injectivity, we need to show for all  $x_1, x_2 \in X$ :

$$f(x_1) = f(x_2) \Rightarrow x_1 = x_2$$

Consider two arbitrary values  $x_1, x_2 \in X$ .

$$\begin{aligned}
 & f(x_1) = f(x_2) \\
 & \Rightarrow g(f(x_1)) = g(f(x_2)) \\
 & \Rightarrow x_1 = x_2
 \end{aligned}$$

As far as  $x_1$  and  $x_2$  are two arbitrary elements of  $X$ , this holds for any pair of elements of  $X$ . We have directly proven injectivity of  $f$ .

## 26.2 Exercise 34.2

Given that  $h : Y \rightarrow X$  exists with  $f \circ h = \text{id}_Y$ , let  $y \in Y$ .

$$y \in Y \Rightarrow h(y) \in X \Rightarrow f(h(y)) \in Y \Leftrightarrow \text{id}_Y(y) = y$$

To show surjectivity, we need to show for all  $y_1, y_2 \in Y$ :

$$\forall y \in Y \exists x \in X : f(x) = y$$

Consider an arbitrary value  $y \in Y$ . Because of the existence of the identity function, it holds that:

$$f(h(y)) = y$$

We define  $h(y)$  as an intermediate value with a different name:

$$x := h(y)$$

$$\Rightarrow \exists x \in X : f(x) = y$$

We have show that for any arbitrary value  $y \in Y$ . So it holds for any value of  $Y$ :

$$\Rightarrow \forall y \in Y \exists x \in X : f(x) = y$$

We have directly proven surjectivity of  $f$ .

## 27 Exercise 35

This exercise was delayed until 26th of November 2015 (later then the other exercises here).

$(a_n)_{n \in \mathbb{N}}$  is sequence with  $\lim_{n \rightarrow \infty} a_n = a \in \mathbb{R}$

$(b_n)_{n \in \mathbb{N}}$  is sequence with  $\lim_{n \rightarrow \infty} b_n = b \in \mathbb{R}$

Furthermore  $b \neq 0$ .

### 27.1 Part 1

$$\lim_{n \rightarrow \infty} a_n = a \wedge \lim_{n \rightarrow \infty} b_n = b \neq 0$$

Let  $\varepsilon > 0$  be arbitrary.

Claim:  $\exists k \in \mathbb{N} \forall n \geq k : |b_n| > \frac{|b|}{2}$ .

Proof: Let  $\varepsilon > 0$ . Consider  $\varepsilon = \frac{|b|}{2}$ .

For  $\varepsilon = \frac{|b|}{2} > 0$ :

$$\exists k \in \mathbb{N} : \forall n \geq k : |b_n - b| < \frac{|b|}{2} = \varepsilon$$

$$\begin{aligned} \forall n \geq k : |b_n| &= |b_n - b + b| \geq \left| |b| - \underbrace{|b - b_n|}_{< \frac{|b|}{2}} \right| \\ &> |b| - |b - b_n| \\ &> |b| - \frac{|b|}{2} \\ &= \frac{|b|}{2} \end{aligned}$$

Claim:

$$\text{sequence } \left( \frac{1}{b_n} \right)_{n \in \mathbb{N}} \wedge \exists \lim \left( \frac{1}{b_n} \right) = \frac{1}{b}$$

Proof: For  $\frac{\varepsilon |b|^2}{2}$  :

$$\exists N \in \mathbb{N} : \forall n \geq N : |b_n - b| < \frac{\varepsilon |b|^2}{2}$$

It holds that  $\forall n \geq N$  :

$$\left| \frac{1}{b_n} - \frac{1}{b} \right| = \left| \frac{b - b_n}{b_n \cdot b} \right| = \frac{|b - b_n|}{|b_n| \cdot |b|} < \frac{\varepsilon \cdot \frac{|b|^2}{2}}{\frac{|b|}{b} \cdot |b|} = \varepsilon$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} a_n \cdot \lim_{n \rightarrow \infty} \frac{1}{b_n} = a \cdot \frac{1}{b} \cdot \frac{a}{b}$$

Or a direct proof:

$$\begin{aligned} \left| \frac{a_n}{b_n} - \frac{a}{b} \right| &= \left| \frac{a_n b - a b_n}{b_n b} \right| \\ &= \frac{|a_n b - a b + a b - a b_n|}{|b_n| |b|} \\ &\leq \frac{|b| |a_n - a| + |a| |b_n - b|}{\frac{|b|}{2} \cdot |b|} \\ &\leq C \cdot \varepsilon \end{aligned}$$

## 27.2 Part 2

## 28 Exercise 36

$$A = \left\{ \frac{1}{2^m} + \frac{1}{n} \mid m, n \in \mathbb{N}_+ \right\}$$

Assumption:  $\min a = 0$

$$0 \notin A$$

$$\frac{1}{2^N} + \frac{1}{N} < 2\varepsilon$$

$$\forall \varepsilon > 0 \exists N \in \mathbb{N}_+ : (m \geq N \Rightarrow \left| \frac{1}{2^m} - 0 \right| < \varepsilon)$$

$$\frac{1}{2^N} < \varepsilon$$

$$n \geq N \Rightarrow \left| \frac{1}{n} - 0 \right| < \varepsilon$$

Assume  $\exists s > 0$  is our lower bound.

$$\exists m : \frac{1}{2^m} < \frac{s}{2} = \varepsilon$$

$$\varepsilon = \frac{s}{2} : \exists N : \frac{1}{N} < \frac{s}{2}$$

$$\rightarrow \underbrace{\frac{1}{sm} + \frac{1}{N}}_{\in A} < s$$

$$\Rightarrow \inf A = 0$$

Remark: When starting this exercise, always estimate whether a maximum/minimum exists. If so, you can save time to prove supremum/infimum.

$$\frac{1}{2^{m+1}} < \frac{1}{2^m} \forall m$$

$$\frac{1}{N+1} < \frac{1}{N} \forall N$$

Therefore  $\max A$  is when  $m, n$  is as small as possible:

$$\frac{1}{2} + \frac{1}{1} = \frac{3}{2}$$

$$\max(A) = \frac{3}{2} = \sup(A)$$

$$B = \left\{ \frac{x}{1+x} \mid x \in \mathbb{R}, x \geq 0 \right\}$$

$\min(B) = 0$  because  $0 \leq \frac{x}{1+x} \forall x \geq 0 \wedge \frac{x}{1+x} \Big|_{x=0} = 0$ .

$$\frac{x}{1+x} < 1 \Leftrightarrow x < 1+x \Leftrightarrow 0 < 1 \forall x \geq 0$$

Is 1 an upper bound and  $1 \notin B$ ?

Assume  $\exists s < 1$ :

$$\frac{x}{1+x} \leq s$$

$$x \leq s(1+x)$$

$$x(1-s) \leq 0$$

$$1-s > 0$$

$$\Rightarrow \sup(B) = 1 \wedge \nexists \max(B)$$

## 29 Exercise 37

$$I = [a, b) = \{x \in \mathbb{R} \mid a \leq x < b\}$$

$$a = \min[a, b) \Rightarrow a \text{ is } \inf([a, b))$$

$$a \in [a, b) : \forall x \in [a, b) : a \leq x \Rightarrow \min(I) = a \Rightarrow \inf(I) = a$$

$b$  is upper bound:

$$b \notin [a, b) \text{ by definition } \forall x \in [a, b) : b > x$$

Claim:  $b$  is the smallest upper bound.

Assume:  $\exists b' < b : b'$  is upper bound.

$$b' \in [a, b)$$

”  
because  $\mathbb{R}$  is complete

## 30 Exercise 38

**Exercise 24.** Let  $A$  and  $B$  two non-empty, bounded by below subseq of  $\mathbb{R}$ . Prove that

$$\inf(A \cup B) = \min \{\inf(A), \inf(B)\}$$

Without loss of generality,  $\inf A \leq \inf B$ :

Let  $a \in A$  and  $b \in B$  arbitrary. This implies that  $a \geq \inf A$  and  $b \geq \inf B \geq \inf A$ .

$$\Rightarrow \forall x \in (A \cup B) : x \geq \inf A$$

$$\Rightarrow \inf(A) \geq \inf(A \cup B)$$

Because extending a set  $A$  with additional elements, the infimum cannot be increased, but only decreased.

$$\Rightarrow \inf(A) \leq \inf(A \cup B)$$

$$x \in A : \inf \{\inf A, \inf B\} \leq \inf(A) \leq x$$

$$x \in B : \inf \{\inf A, \inf B\} \leq \inf(B) \leq x$$

$$\forall x \in A \cup B : \underbrace{\min \{\inf A, \inf B\}}_{\text{lower bound}} \leq x$$

$$\Rightarrow \inf(A \cup B) \leq \min \{\inf(A), \inf(B)\}$$

$$\Rightarrow \inf(A) = \inf(A \cup B)$$

## 31 Exercise 39

### 31.1 Exercise 39a

$$\begin{aligned} \sup_{y \in Y} \inf_{x \in X} f(x, y) &\leq \inf_{x \in X} \sup_{y \in Y} f(x, y) \\ \underbrace{\inf_{x \in X} f(x, y)}_{\sup_{y \in Y}} &\leq f(x, y) \leq \sup_{y \in Y} f(x, y) \\ \sup_{y \in Y} \inf_{x \in X} f(x, y) &\leq \sup_{y \in Y} f(x, y) \\ \sup_{y \in Y} \inf_{x \in X} f(x, y) &= \inf_{x \in X} \sup_{y \in Y} \inf_{x \in X} f(x, y) \leq \inf_{x \in X} \sup_{y \in Y} f(x, y) \quad \checkmark \end{aligned}$$

### 31.2 Exercise 39b

$$\begin{aligned} f : (x, y) &\mapsto 1_{\{x \geq 0, y \geq 0\} \cup \{x < 0, y < 0\}} \\ \sup_{y \in Y} f(x, y) &= 1 \forall x \\ \inf_{x \in X} \sup_{y \in Y} f(x, y) &= 1 \\ \inf_{x \in X} f(x, y) &= 0 \forall y \in [-1, 1] \\ \sup_{y \in Y} \inf_{x \in X} f(x, y) &= 0 < 1 \end{aligned}$$



## 32 Exercise 40

### 32.1 Exercise 40.a.1

$$\frac{5+i}{2+3i} \cdot \frac{2-3i}{2-3i} = \frac{10-15i+2i-3i^2}{-6i+4+6i-9i^2} = \frac{10-13i+3}{4+9} = \frac{13-13i}{13} = 1-i$$

### 32.2 Exercise 40.a.2

$$\begin{aligned} z^2 &= \frac{1+\sqrt{3}i}{2} \\ z^2 &= \pm \sqrt{\frac{1}{2} + \frac{\sqrt{3}i}{2}} \\ z^2 &= \pm \sqrt{\frac{9+6\sqrt{3}i-3}{12}} \\ z^2 &= \pm \sqrt{\frac{(3+\sqrt{3}i)^2}{12}} \\ z^2 &= \pm \frac{3+\sqrt{3}i}{\sqrt{12}} \\ z^2 &= \pm \left( \frac{\sqrt{3}}{2} + i\frac{1}{2} \right) \end{aligned}$$

### 32.3 Exercise 40.b.1

$$\begin{aligned} M_1 &= \left\{ z \in \mathbb{C} \setminus \{0\} \mid \left| \frac{1}{z} \right| < 2 \right\} \\ \left| \frac{1}{z} \right| &= \left| \frac{1}{a+bi} \right| = \frac{|1|}{|a+bi|} = \frac{1}{\sqrt{a^2+b^2}} \\ &\Rightarrow \frac{1}{\sqrt{a^2+b^2}} < 2 \\ &\Rightarrow \frac{1}{2} < \sqrt{a^2+b^2} \\ &\Rightarrow \frac{1}{4} < a^2+b^2 \end{aligned}$$

Illustrated we draw a circle originating in  $(0,0)$  with radius  $\frac{1}{2}$ . The solution set is the whole plane excluding everything what is part of the circle.

### 32.4 Exercise 40.b.2

$$\begin{aligned} M_2 &= \{ z \in \mathbb{C} \mid \Im((1+i)z) = 0 \} \\ &\quad \Im(z+zi) \end{aligned}$$

TODO

### 33 Exercise 41

$$A_n := (-\infty, a_n)_{n \in \mathbb{N}} \quad A := \bigcup_{n \in \mathbb{N}} A_n$$

$$B_n := (b_n, \infty)_{n \in \mathbb{N}} \quad B := \bigcup_{n \in \mathbb{N}} B_n$$

$$\forall n \in \mathbb{N} : x \in I_n$$

Show that  $x = \sup A = \inf B$ .

Because  $I_n$  are nested intervals it holds that

$$a_1 \leq \dots \leq a_n \leq a_{n+1} \leq x$$

Because

$$\forall \varepsilon > 0 \exists N : N \geq n : 0 \leq x - a_n \leq b_n - a_n \leq \varepsilon$$

it holds that

$$x = \sup(a_n)$$

Let  $y \in A$ .

$$\exists n \in \mathbb{N} : y \in A_n \Rightarrow y < a_n \leq x$$

$$\Rightarrow y \in A : y < x$$

Therefore  $x$  is an upper bound. Is it the only upper bound?

Assume another upper bound  $x'$  exists.

$$x' < x = \lim_{n \rightarrow \infty} a_n$$

$$\Rightarrow \exists N \in \mathbb{N} : x' < a_n \quad \forall n \geq N$$

$$\varepsilon = \frac{x - x'}{2}$$

$$\Rightarrow \exists y \in A_{n+1}$$

$$y > x'$$

This is a contradiction and therefore  $x$  is the distinct upper bound.

The proof for the infimum works analogously.

It only remains to show that  $x \notin A$ .

$$\forall a_n \neq x \Rightarrow \exists a_{n+k} : a_n < a_{n+k}$$

### 34 Exercise 42

Give the limits for the following sequences:

#### 34.1 Exercise 42.a

$$a_n = \frac{5n+2}{3n+7}$$

$$\begin{aligned}
\lim_{n \rightarrow \infty} a_n &= \frac{5n+2}{3n+7} \\
&= \frac{\lim_{n \rightarrow \infty} 5n+2}{\lim_{n \rightarrow \infty} 3n+7} \\
&= \frac{\lim_{n \rightarrow \infty} 5n + \lim_{n \rightarrow \infty} 2}{\lim_{n \rightarrow \infty} 3n + \lim_{n \rightarrow \infty} 7} \\
&= \frac{n(5 + \frac{2}{n})}{n(3 + \frac{7}{n})} \\
&= \frac{5 + \overbrace{\frac{2}{n}}^{\rightarrow 0}}{3 + \underbrace{\frac{7}{n}}_{\rightarrow 0}} \\
&= \frac{5}{3}
\end{aligned}$$

This works only if the denominator is non-zero.  $\lim_{n \rightarrow \infty} (3 + \frac{7}{n})$  turns out to be non-zero.

### 34.2 Exercise 42.b

$$b_n = \frac{2n^2 - 4n + 5}{n^3 + 2\sqrt{n}}$$

First, we make a remark, that  $\lim_{n \rightarrow \infty} \frac{1}{n^2} = 0$ . Why, because

$$\lim_{n \rightarrow \infty} \frac{1}{n^2} = \left( \lim_{n \rightarrow \infty} \frac{1}{n} \right) \cdot \left( \lim_{n \rightarrow \infty} \frac{1}{n} \right) = 0$$

This can be generalized for  $\lim_{n \rightarrow \infty} \frac{1}{n^k} = 0$  with  $k \in \mathbb{N}_+$ .

$$\begin{aligned}
\lim_{n \rightarrow \infty} b_n &= \frac{2n^2 - 4n + 5}{n^3 + 2\sqrt{n}} \\
&= \frac{n^3 \cdot \left( \frac{2}{n} - \frac{4}{n^2} + \frac{5}{n^2} \right)}{n^3 \cdot \left( 1 + 2\frac{n^{0.5}}{n^3} \right)} \\
&= \frac{\frac{2}{n} - \frac{4}{n^2} + \frac{5}{n^3}}{1 + 2 \cdot \frac{1}{n^{2.5}}} \\
&= \frac{\frac{2}{n} - \frac{4}{n^2} + \frac{5}{n^3}}{\frac{n^{2.5}}{n^{2.5}}} \\
&= \frac{2n^{1.5} - 4n^{0.5} + 5n^{0.5}}{n^{2.5} + 2} \cdot \frac{1}{\frac{1}{n^{2.5}}} \\
&= \frac{2n^{-1} - 4n^{-2} + 5n^{-3}}{1 + 2n^{-2.5}} \\
&= \frac{0}{1}
\end{aligned}$$

Or generally:

$$2n^2 - 4n + 5 \leq 2n^2 + 4n^2 + 5n^2 \leq 11n^2$$

$$0 \leq b_n \leq \frac{11n^2}{n^3} = \underbrace{\frac{11}{n}}_{\rightarrow 0}$$

### 34.3 Exercise 42.c

$$c_n = \sqrt{4n^2 + 2n + 3}$$

$$\begin{aligned} c_n &= \sqrt{4n^2 + 2n + 3} \cdot \frac{\sqrt{4n^2 + 2n + 3}}{\sqrt{4n^2 + 2n + 3} + 2n} \\ &= \dots \\ &= \frac{2 + \frac{3}{n}}{\sqrt{4 + \frac{2}{n} + \frac{3}{n^2}} + 2} \\ &= \frac{2}{4} \\ &= \frac{1}{2} \end{aligned}$$

### 34.4 Exercise 42.d

$$d_n = \binom{n}{k} n^{-k} \text{ with } n \in \mathbb{N} \text{ for a fixed } k \in \mathbb{N}_+$$

$$\begin{aligned} d_n &= \binom{n}{k} n^{-k} \\ &= \frac{n!}{k!(n-k)!n^k} \\ &= \frac{n \cdot (n-1) \cdot \dots \cdot 1}{k!(n-k)!n^k} \\ &= \frac{n \cdot (n-1) \cdot \dots \cdot 1}{k!(n-k)!n^k} \\ &= \frac{(1 - \frac{1}{n})(1 - \frac{2}{n}) \cdot \dots \cdot (1 - \frac{k-1}{n}) \cdot (n-k) \cdot \dots \cdot 1}{k!(n-k)!} \\ &= \frac{(n-k)!}{k!(n-k)!} \\ &= \frac{1}{k!} \end{aligned}$$

Or better we write:

$$\begin{aligned}
\frac{n!}{k!(n-k)!} &= \frac{\prod_{i=0}^{n-1} (n-i)}{\prod_{j=k}^{n-1} (n-j)} \\
&= \frac{1}{k!} \prod_{j=0}^{k-1} (n-j) n^{-k} \\
&= \frac{1}{k!} \prod_{j=0}^{k-1} \left[ (n-j) \cdot \frac{1}{n} \right] \\
&= \frac{1}{k!} \prod_{j=0}^{k-1} \left( 1 - \frac{j}{n} \right) \\
\lim_{n \rightarrow \infty} \frac{1}{k!} \prod_{j=0}^{k-1} \left( 1 - \frac{j}{n} \right) &= \frac{1}{k!} \lim_{n \rightarrow \infty} \prod_{j=0}^{k-1} \left( 1 - \frac{j}{n} \right) \quad [\text{if limes exist}] \\
&= \frac{1}{k!} \prod_{j=0}^{k-1} \underbrace{\lim_{n \rightarrow \infty} \left( 1 - \frac{j}{n} \right)}_{=1} \\
&= \frac{1}{k!} \forall j = 0, \dots, k-1
\end{aligned}$$

## 35 Exercise 43

**Exercise 25.** Let  $(a_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathbb{R}_+$  with  $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = q$ . Prove that

$$(a_n)_{n \in \mathbb{N}} \begin{cases} \text{converges} & \text{if } q < 1 \\ \text{diverges} & \text{if } q > 1 \end{cases}$$

In case  $q = 1$  no statement about the convergence of  $(a_n)_{n \in \mathbb{N}}$  can be made.

$$\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = q$$

### 35.1 Examples for $q = 1$

$$\begin{aligned}
a_n &= \frac{1}{n+1} & \frac{a_{n+1}}{a_n} \frac{n+1}{n+2} &\rightarrow_{n \rightarrow \infty} 1 & a_n &\searrow 0 \\
a_n &= n+1 & \frac{a_{n+1}}{a_n} &= \frac{n+2}{n+1} \rightarrow_{n \rightarrow \infty} 1 & a_n &\nearrow 0
\end{aligned}$$

### 35.2 Proof for $q < 1$

$$\exists \underbrace{\varepsilon}_{= \frac{q+1}{2} - a} > 0 : q + \varepsilon < 1$$

If  $n$  is sufficiently large:

$$\left| \frac{a_{n+1}}{a_n} - q \right| < \varepsilon \Rightarrow \frac{a_{n+1}}{a_n} \in (q - \varepsilon, q + \varepsilon)$$

$$\begin{aligned}
0 &\leq a_{n+1} \leq (q + \varepsilon)a_n \\
0 &\leq a_{n+2} \leq (q + \varepsilon)^2 a_n \\
&\dots \\
0 &\leq a_{n+k} \leq (q + \varepsilon)^k a_n
\end{aligned}$$

By induction it holds that

$$0 \leq a_{n+k} \leq \underbrace{(q + \varepsilon)^k}_{\tilde{q} < 1} a_1 \xrightarrow{k \rightarrow \infty} 0$$

This follows from the squeeze theorem.

$$\forall q > 1 \exists \varepsilon > 0 : q - \varepsilon > 1$$

$$\begin{aligned}
a_{n+1} &> (q - \varepsilon)a_n \\
a_{n+k} &> \underbrace{(q - \varepsilon)^k}_{\tilde{q} > 1} a_n
\end{aligned}$$

$$\begin{aligned}
\lim_{n \rightarrow \infty} \tilde{q}^k &= +\infty \\
\tilde{q} &> 1
\end{aligned}$$

## 36 Exercise 44

**Exercise 26.** Let  $(a_n)_{n \in \mathbb{N}}$  be a zero sequence in  $\mathbb{R}$  and  $(b_n)_{n \in \mathbb{N}}$  a bounded sequence in  $\mathbb{R}$ . Prove that  $(a_n b_n)_{n \in \mathbb{N}}$  is a zero sequence.

Because  $(b_n)_{n \in \mathbb{N}}$  is bounded some  $d$  exists such that

$$\forall \varepsilon > 0 : \exists N \in \mathbb{N} : n \geq N : |a_n - 0| < \varepsilon$$

Consider  $\lim_{n \rightarrow \infty} (a_n \cdot b_n) = 0$ .

We need to show that

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} : n \geq N : |a_n \cdot b_n - 0| < \varepsilon \cdot d$$

Where  $\varepsilon \cdot d$  is epsilon multiplied with constant  $d$ . This is a hand-crafted value (meaning that we selected it intentionally and will turn out to solve our problem). Now we elaborate on the relation:

$$\begin{aligned}
|a_n \cdot b_n| &< \varepsilon \cdot d \\
|a_n| \cdot |b_n| &< \varepsilon \cdot d \\
|a_n| \cdot d &< \varepsilon \cdot d \\
|a_n| &< \varepsilon
\end{aligned}$$

Because  $a_n < \varepsilon$  it holds that some constant exists for a sufficiently large  $N$  such that  $|a_n \cdot b_n|$  is always smaller than some constant  $\varepsilon$ .

## 37 Exercise 45

**Exercise 27.** Let  $a, b, c \in [0, \infty)$ . Show that,

$$\lim_{n \rightarrow \infty} \sqrt[n]{a^n + b^n + c^n} = \max \{a, b, c\}$$

Without loss of generality, let  $a = \max \{a, b, c\}$ . Because  $a, b, c$  is non-negative,

$$\begin{aligned} a^n &\leq a^n + b^n + c^n \leq 3a^n \\ \sqrt[n]{a^n} &\leq \sqrt[n]{a^n + b^n + c^n} \leq \sqrt[n]{3} \cdot \sqrt[n]{a^n} \\ \lim_{n \rightarrow \infty} \sqrt[n]{a^n} &= a \\ \lim_{n \rightarrow \infty} \sqrt[n]{3} \cdot \sqrt[n]{a^n} &= a \lim_{n \rightarrow \infty} \sqrt[n]{3} = a \cdot 1 \end{aligned}$$

Due to the squeeze theorem, it holds that  $\lim_{n \rightarrow \infty} \sqrt[n]{a^n + b^n + c^n} = a = \max \{a, b, c\}$ .

### 38 Exercise 46

**Exercise 28.** Let  $a_0 \in (0, 1)$  and a sequence  $(a_n)_{n \in \mathbb{N}}$  is recursively defined with

$$a_{n+1} = 1 - \sqrt{1 - a_n} \text{ for } n \geq 0$$

**Induction base**

$$a_1 = 1 - \sqrt{1 - a_0} \Rightarrow 0 < a_1 < 1 \quad a_1 \in (0, 1)$$

**Induction step** Let  $a_n \in (0, 1)$ .

$$\begin{aligned} 0 < a_n < 1 &\Rightarrow -1 < -a_n < 0 \Rightarrow 0 < \sqrt{1 - a_n} < 1 \\ 0 < 1 - \underbrace{\sqrt{1 - a_n}}_{a_{n+1}} &< 1 \end{aligned}$$

So  $a_{n+1} < a_n$ .

$$\begin{aligned} 1 - \sqrt{1 - a_n} < a_n &\Leftrightarrow (1 - a_n)^2 < 1 - a_n \\ &\Rightarrow 1 - a_n < \sqrt{1 - a_n} \\ &\Rightarrow x^2 < x \Leftrightarrow x \in (0, 1) \end{aligned}$$

$$\begin{aligned} a_{n+1} &= 1 - \sqrt{1 - a_n} \\ a &= 1 - \sqrt{1 - a} \end{aligned}$$

Therefore only 0 or 1 are possible limes for this sequence. But monotonically decreasing implies that 0 is the limes (bounded below and monotonically decreasing sequences are convergent).

### 39 Exercise 47

**Exercise 29.** For a sequence  $(a_n)_{n \in \mathbb{N}}$  in  $\mathbb{R}$  we assign the sequence  $(s_n)_{n \in \mathbb{N}}$ , where

$$s_n = \frac{1}{n+1} \sum_{k=0}^n a_k \text{ for } n \geq 0$$

is the mean value of the first  $n+1$  sequence numbers.

- Show that: If  $\lim_{n \rightarrow \infty} a_n = a$  with  $a \in \mathbb{R}$ , then  $\lim_{n \rightarrow \infty} s_n = a$ .

- Give an example for a divergent sequence  $(a_n)_{n \in \mathbb{N}}$  for which the sequence of mean values converges anyways.

### 39.1 Exercise 47.a

We show that  $\exists a \in \mathbb{R} : \lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} s_n = a$ .

$$\begin{aligned}
 \lim_{n \rightarrow \infty} s_n &= \lim_{n \rightarrow \infty} \left( \frac{1}{n+1} \sum_{k=0}^n a_k \right) = \lim_{n \rightarrow \infty} \left( \frac{1}{n+1} \sum_{k=0}^n (a_n - (a_n - a_k)) \right) \\
 &= \lim_{n \rightarrow \infty} \left( \frac{1}{n+1} \sum_{k=0}^n a_n \right) - \lim_{n \rightarrow \infty} \left( \frac{1}{n+1} \sum_{k=0}^n (a_n - a_k) \right) \\
 &= \lim_{n \rightarrow \infty} \frac{1}{n+1} (n+1) a_n - \lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{a_n - a_k}{n+1} \\
 &\quad [\forall \varepsilon > 0 \exists N : n \geq N : |a_n - a| < \varepsilon] \\
 &= a - \lim_{n \rightarrow \infty} \underbrace{\sum_{k=0}^N \frac{a_n - a_k}{n+1}}_{(N+1) \frac{C}{n+1}} + \sum_{N+1}^n \frac{a_n - a_k}{n+1} \\
 &\leq \underbrace{\lim_{n \rightarrow \infty} \frac{(N+1)C}{n+1}}_{\rightarrow 0} + \lim_{n \rightarrow \infty} \sum_{n=N+1}^n \frac{a_n - a_b}{n+1} \\
 &\quad \left[ \lim_{n \rightarrow \infty} \sum_{n=N+1}^n \frac{\varepsilon}{n+1} = \frac{n-N-1}{n+1} \varepsilon \rightarrow 1 \right] \\
 &= \lim_{n \rightarrow \infty} \frac{(N+1)C}{n+1} +
 \end{aligned}$$

### 39.2 Exercise 47.a, radical variant

$$\sum_{n=N-1}^n a - \varepsilon \leq \dots \leq \frac{\sum_{k=0}^N a_k + \sum_{k=N+1}^n (a + \varepsilon)}{n+1}$$

### 39.3 Exercise 47.b

$$\begin{aligned}
 (a_n)_{n \in \mathbb{N}} &= (-1)^n \\
 \Rightarrow \lim_{n \rightarrow \infty} s_n &= \lim_{n \rightarrow \infty} \frac{\sum_{k=0}^n (-1)^k}{n+1} = 0
 \end{aligned}$$

## 40 Exercise 48

**Exercise 30.** Let  $M \subseteq \mathbb{R}$  be a bounded above set and  $s \in \mathbb{R}$ . Prove that:

$$s = \sup(M) \Leftrightarrow \begin{cases} \forall x \in M : s \geq x \\ \exists (x_n)_{n \in \mathbb{N}}, x_n \in M : \lim_{n \rightarrow \infty} x_n = s \end{cases} \quad \text{and}$$

$$\exists (x_n)_{n \in \mathbb{N}}, x_n \in M : \lim_{n \rightarrow \infty} x_n = s \Leftrightarrow \varepsilon > 0 \exists N \in \mathbb{N} : |x_n - s| < \varepsilon$$

We prove the first direction  $\Leftarrow$ .



Let  $s$  be an upper bound of  $M$ . Let  $(x_n)_{n \in \mathbb{N}}$  in  $M$  with  $\lim_{n \rightarrow \infty} x_n = s$ .

$$\begin{aligned} \Rightarrow \forall t < s \exists N \in \mathbb{N} : n \geq N \Rightarrow (s - x_n) \leq |s - x_n| < \frac{\varepsilon}{2} \\ \Rightarrow t < x_n \end{aligned}$$

We prove the second direction  $\Rightarrow$ .

Therefore

$$\begin{aligned} s - \frac{1}{n} \text{ is not an upper bound of } M \\ \Rightarrow \exists x_n \in M : s - \frac{1}{n} < x_n \\ \exists x_n \in M : s - \frac{1}{n} < x_n \\ s - \frac{1}{n} \leq x_n < s \\ s \leq x_n < s \Rightarrow (x_n)_{n \in \mathbb{N}} \rightarrow s \end{aligned}$$

## 41 Exercise 49

**Exercise 31.** Let  $(a_n)_{n \in \mathbb{N}}$  be a convergent sequence of non-negative real numbers with  $\lim_{n \rightarrow \infty} a_n = a$  and  $k \in \mathbb{N}_+$ . Show that

$$\lim_{n \rightarrow \infty} \sqrt[k]{a_n} = \sqrt[k]{a}$$

Hint:  $a_n - a = \sqrt[k]{a_n^k} - \sqrt[k]{a^k} = (\sqrt[k]{a_n} - \sqrt[k]{a})(\dots)$ .

$$\lim_{n \rightarrow \infty} \sqrt[k]{a_n} = \sqrt[k]{a}$$

$$\lim_{n \rightarrow \infty} a_n = a \quad \text{i.e.} \quad \forall \varepsilon \exists N \in \mathbb{N} : (\sqrt[k]{a_n} - \sqrt[k]{a}) \left( \sum_{i=0}^{k-1} \sqrt[k]{a_n^{k-1-j}} \cdot \sqrt[k]{a} \right)$$

**Case 1:**  $a > 0$

$$|a_n - a| < \frac{a}{2}$$

$$|a_n| > \frac{|a|}{2}$$

$\Rightarrow$  the product is always positive:

$$\underbrace{(\sqrt[k]{a_n} - \sqrt[k]{a}) \left( \sum_{i=0}^{k-1} \sqrt[k]{a_n^{k-1-j}} \cdot \sqrt[k]{a} \right)}_{b_n \geq b > 0}$$

Done.

**Case 2**

$$\sqrt[k]{a_n} < \varepsilon \Leftrightarrow a_n < \varepsilon^k = \tilde{\varepsilon}$$

$$\forall \tilde{\varepsilon} > 0 \exists N \in \mathbb{N} : n \geq N : |a_n - 0| < \tilde{\varepsilon}$$

$$\Rightarrow \sqrt[k]{a_n}$$

### 41.1 Shorter valid solution

$$b_n = \frac{a_n - a}{(\sqrt[k]{a_n} - \sqrt[k]{a}) \left( \sum_{i=0}^{k-1} \sqrt[k]{a_n}^{k-1-i} \cdot \sqrt[k]{a} \right)} = \sqrt[k]{a_n} - \sqrt[k]{a}$$

$$\sqrt[k]{a_n} - \sqrt[k]{a} = \frac{a_n - a}{b_n}$$

We already know that

$$\lim_{n \rightarrow \infty} \frac{a_n - a}{b_n} = \frac{\lim_{n \rightarrow \infty} (a_n - a)}{\lim_{n \rightarrow \infty} b_n} = \frac{0}{b} = 0$$

## 42 Exercise 50

**Exercise 32.** Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathbb{C}$  and  $q \in [0, 1)$  such that for all  $n \geq 1$ ,

$$|x_{n+1} - x_n| \leq q \cdot |x_n - x_{n-1}|.$$

Prove that  $(x_n)_{n \in \mathbb{N}}$  is a Cauchy sequence. Does the implication still hold, if only  $|x_{n+1} - x_n| < |x_n - x_{n-1}|$ ?

This is actually the fixed point theorem by Banach. Contraction property.

### 42.1 Special case: $q = 0$ implies a Cauchy sequence

If  $q = 0$ , then the sequence  $(x_n)_{n \in \mathbb{N}}$  is constant. A constant sequence is always a Cauchy sequence.

### 42.2 An observation

Let  $q \in [0, 1)$ .

$$\begin{aligned} \forall n \geq 1 : |x_{n+2} - x_{n+1}| &\leq q |x_{n+1} - x_n| \\ \forall n \geq 1 : q |x_{n+1} - x_n| &\leq q^2 |x_n - x_{n-1}| \\ \Rightarrow \forall n \geq 1 : |x_{n+2} - x_{n+1}| &\leq q^2 |x_n - x_{n-1}| \end{aligned}$$

We make a more general claim:

### 42.3 Complete induction over $k \in \mathbb{N}$

$$|x_{n+k+1} - x_{n+k}| \leq q^{k+1} |x_n - x_{n-1}| \quad \forall n \geq N : N = 1$$

**Induction base:**  $k = 0$

$$|x_{n+1} - x_n| \leq q^1 |x_n - x_{n-1}| \quad \checkmark$$

**Induction step:**  $k \rightarrow k + 1$

$$\begin{aligned} |x_{n+k+2} - x_{n+k+1}| &\leq q^{k+2} \cdot |x_n - x_{n-1}| \\ \text{we know: } |x_{n+k+2} - x_{n+k+1}| &\leq q \cdot |x_{n+k+1} - x_{n+k}| \\ q |x_{n+k+1} - x_{n+k}| &\stackrel{!}{\leq} q^{k+2} \cdot |x_n - x_{n-1}| \\ |x_{n+k+1} - x_{n+k}| &\leq q^{k+1} \cdot |x_n - x_{n-1}| \quad \checkmark \end{aligned}$$

## 42.4 Selection of $\varepsilon$

Let  $k \in \mathbb{N}$ . We have proven that  $\forall N \geq 1$ , it holds that

$$|x_{N+k+1} - x_{N+k}| \leq q^{k+1} |x_N - x_{N-1}|$$

Consider  $N = 1$ .

$$|x_{k+2} - x_{k+1}| \leq q^{k+1} |x_1 - x_0|$$

Now consider  $\varepsilon_0 = q^{k+1} |x_1 - x_0|$ . Let's recognize that we can choose some  $n, m \in \mathbb{N}$  without loss of generality  $n \geq m$ . Then it holds that,

$$|x_n - x_{n-1}| \leq \varepsilon_0 \quad \wedge \quad |x_m - x_{m-1}| \leq \varepsilon_0$$

Let  $\varepsilon = \varepsilon_0(n - m) + 1 = q^{k+1} |x_1 - x_0| (n - m) + 1$ .

TODO:  $\sum_{k=0}^{m-n-1} q^k = \frac{1-q^{n-m}}{1-q}$ .  $\varepsilon = \frac{1}{1-q} |x_{n+1} - x_n|$

## 42.5 Is $\varepsilon$ an appropriate constant?

We need to show the Cauchy criterion:

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} : n \geq N \wedge m \geq N : |a_n - a_m| < \varepsilon$$

$$\begin{aligned} |a_n - a_m| &= |a_n - a_{n-1} + a_{n-1} - a_{n-2} + a_{n-2} - \cdots + a_{m+1} - a_m| \\ &\leq |a_n - a_{n-1}| + |a_{n-1} - a_{n-2}| + \cdots + |a_{m+1} - a_m| && \text{[triangle inequality]} \\ &= \sum_{i=n}^{m+1} |a_i - a_{i-1}| \\ &\leq \sum_{i=n}^{m+1} q^{k+1} |x_1 - x_0| \\ &= (n - m) \cdot q^{k+1} |x_1 - x_0| \\ &< (n - m) \cdot q^{k+1} |x_1 - x_0| + 1 \\ &= \varepsilon \end{aligned}$$

## 42.6 Does it also hold for $|x_{n+1} - x_n| < |x_n - x_{n-1}|$ ?

The proof stays the same (so this statement is true as well).

- Because  $q$  cannot be  $\geq 1$  in  $q \in [0, 1)$ ,  $q$  cancels out with smaller-than.
- $\varepsilon$  can be chosen differently (we can skip  $+1$ ):

$$\varepsilon = q^{k+1} |x_1 - x_0| (n - m)$$

## 43 Exercise 51

**Exercise 33.** A sequence  $(a_n)_{n \in \mathbb{N}}$  in  $\mathbb{R}$  is recursively defined by  $a_0 = 0, a_1 = 1$  and

$$a_n = \frac{a_{n-1} + a_{n-2}}{2} \quad \text{for } n \geq 2$$

Show (using the result from Exercise 50), that the sequence  $(a_n)_{n \in \mathbb{N}}$  converges.

*Bonus:* Can you determine the limes of this sequence?

$$(a_n)_{n \in \mathbb{N}} = \left(0, 1, \frac{1}{2}, \frac{3}{4}, \frac{5}{8}, \frac{11}{16}, \frac{21}{32}, \dots\right)$$

We will also rephrase the equation:

$$\begin{aligned} a_n &= \frac{1}{2}(a_{n-1} + a_{n-2}) \\ a_{n-1} &= 2a_n - a_{n-2} \end{aligned}$$

We revise the previous theorem:

Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathbb{C}$  and  $q \in [0, 1)$  such that for all  $n \geq 1$ ,

$$|x_{n+1} - x_n| \leq q \cdot |x_n - x_{n-1}|.$$

Then  $(x_n)_{n \in \mathbb{N}}$  is a Cauchy sequence.

$$\begin{aligned} |a_{n+1} - a_n| &\leq q |a_n - a_{n-1}| \\ \left| \frac{a_n + a_{n-1}}{2} - \frac{a_{n-1} + a_{n-2}}{2} \right| &\leq q \left| \frac{a_{n-1} + a_{n-2}}{2} - \frac{2a_{n-1}}{2} \right| \\ \frac{1}{2} |a_n + a_{n-1} - a_{n-1} - a_{n-2}| &\leq \frac{q}{2} \cdot |a_{n-1} + a_{n-2} - 2a_{n-1}| \\ |a_n - a_{n-2}| &\leq q \cdot |a_{n-2} - a_{n-1}| \end{aligned}$$

We have observed previously that  $a_{n-1} = 2a_n - a_{n-2}$

$$\begin{aligned} |a_n - a_{n-2}| &\leq q \cdot |2a_{n-2} - 2a_n| \\ |a_n - a_{n-2}| &\leq 2q \cdot |a_{n-2} - a_n| \\ |a_n - a_{n-2}| &\leq 2q \cdot |(-1)(a_n - a_{n-2})| \\ |a_n - a_{n-2}| &\leq 2q \cdot |a_n - a_{n-2}| \\ 1 &\leq 2q \end{aligned}$$

So this can be proven for any  $q \geq \frac{1}{2}$ , so we choose  $q = \frac{1}{2}$ .

## 44 Exercise 52

**Exercise 34.** Let  $(a_n)_{n \in \mathbb{N}}$  be a bounded above sequence in  $\mathbb{R}$ . Show that a number  $s \in \mathbb{R}$  is the limes superior of  $(a_n)_{n \in \mathbb{N}}$  if and only if for every  $\varepsilon > 0$  the following conditions hold:

1. the set  $\{n \in \mathbb{N} \mid a_n > s - \varepsilon\}$  is infinite.
2. the set  $\{n \in \mathbb{N} \mid a_n > s + \varepsilon\}$  is finite.

We shortly point out that a limes superior must not necessarily exist, because bounded sequences in  $\mathbb{C}$  have a limit point, but not sequences which are only bounded above.

### 44.1 Proof direction $\Leftarrow$

Given a limes superior  $s$ . So  $\forall \varepsilon > 0 : |a_n - s| < \varepsilon$  for infinitely many indices.

1. If  $s$  is a limit point, it has infinitely many sequence numbers within its  $\varepsilon$ -environment.

$$\forall \varepsilon > 0 : |a_n - s| < \varepsilon \quad \Rightarrow \quad |\{n \in \mathbb{N} \mid a_n > s - \varepsilon\}| = \infty$$




Figure 2: Illustration of the  $\varepsilon$  environment

2. So  $s$  is the supremum of all limit points. Proof by contradiction:

Assume infinitely many elements are greater  $s + \varepsilon$ . Consider the upper bound of  $(a_n)_{n \in \mathbb{N}}$  and  $s + \varepsilon$  as lower bound. This subsequence is bounded in  $\mathbb{C}$  (actually  $\mathbb{R}$ ). Due to the Weierstrass-Bolzano theorem, a convergent subsequence must exist which necessarily provides a limit point. This limit point must be  $\geq s + \varepsilon$ , because of the lower bound. So we found another limit point which contradicts with our assumption that  $s$  is the supremum of all limit points.

#### 44.2 Proof direction $\Rightarrow$

From the first property it holds that  $|\{n \in \mathbb{N} \mid a_n > s - \varepsilon\}| = \infty \Rightarrow s - a_n < \varepsilon$  for infinitely many sequence numbers.

From the second property it holds that  $|\{n \in \mathbb{N} \mid a_n > s + \varepsilon\}| < \infty \Rightarrow a_n > s + \varepsilon$ . From that it follows that  $s - a_n > \varepsilon \Rightarrow a_n - s < \varepsilon$ . So  $a_n - s > \varepsilon$  for infinitely many sequence numbers. So  $a_n - s < \varepsilon$  for infinitely many sequence numbers.

From that it follows that  $|a_n - s| < \varepsilon$  holds for infinitely many.

### 45 Exercise 53

**Exercise 35.** Determine the limes superior and limes inferior of the sequence  $(a_n)_{n \in \mathbb{N}}$  defined by

$$a_n = (1 + (-1)^n) (-1)^{\frac{n(n-1)}{2}}.$$

$$a_n = (1 + (-1)^n) (-1)^{n(n-1)/2}$$

#### 45.1 Case 1: $n = 2k$ with $k \in \mathbb{N}$

$$\begin{aligned} a_n &= (1 + (-1)^{2k}) (-1)^{2k(2k-1)/2} \\ &= \left(1 + ((-1)^2)^k\right) (-1)^{k(2k-1)} \\ &= 2(-1)^{k(2k-1)} \end{aligned}$$

#### 45.2 Case 1.a: $k = 2l$ with $l \in \mathbb{N}$

$$\begin{aligned}
 &= 2(-1)^{2l(4l-1)} \\
 &= 2 \left( (-1)^2 \right)^{l(4l-1)} \\
 &= 2
 \end{aligned}$$

#### 45.3 Case 1.b: $k = 2l + 1$ with $l \in \mathbb{N}$

$$\begin{aligned}
 &= 2(-1)^{(2l+1)(4l+1)} \\
 &= 2(-1)^{2l(4l+1)} \cdot (-1)^{(4l+1)} \\
 &= 2(-1)^{2l(4l+1)} \cdot \left( (-1)^4 \right)^l \cdot (-1) \\
 &= 2 \cdot 1 \cdot 1 \cdot (-1) \\
 &= -2
 \end{aligned}$$

#### 45.4 Case 2: $n = 2k + 1$ with $k \in \mathbb{N}$

$$\begin{aligned}
 a_n &= \left( 1 + (-1)^{2k+1} \right) (-1)^{(2k+1)(2k)/2} \\
 &= \left( 1 + (-1)^{2k} \cdot (-1) \right)^{(2k+1)k} \\
 &= (1 + 1 \cdot (-1))^{(2k+1)k} \\
 &= 0^{(2k+1)k} \\
 &= 0
 \end{aligned}$$

#### 45.5 Putting it together

We have isolated three subsequences. Every subsequence itself is a constant sequence. Every constant sequence has a limit point. So the set of limit points is given with:

$$L = \{-2, 0, 2\}$$

$$\begin{aligned}
 \sup(L) &= \max(L) = 2 \Rightarrow \limsup_{n \rightarrow \infty} (a_n) = 2 \\
 \inf(L) &= \min(L) = -2 \Rightarrow \liminf_{n \rightarrow \infty} (a_n) = -2
 \end{aligned}$$

#### 45.6 Limes

$$\begin{aligned}
 0 + \sum_{k=0}^n \frac{1}{2^{2k}} - \frac{1}{2} \sum_{k=0}^n \frac{1}{2^{2k}} &= \sum_{k=0}^n \left( \frac{1}{2^{2k}} \right) \left( 1 - \frac{1}{2} \right) \\
 &= \sum_{k=0}^n \left( \frac{1}{4} \right)^k \left( \frac{1}{2} \right) = \frac{1}{2} \frac{\left( \frac{1}{4} \right)^{n+1} - 1}{\frac{1}{4} - 1} \\
 &= \frac{1}{2} \frac{1}{\frac{3}{4}} = \frac{2}{3} \\
 \lim_{n \rightarrow \infty} a_n &= \frac{2}{3}
 \end{aligned}$$

### 46 Exercise 54

**Exercise 36.** Let  $a \in \mathbb{R} \setminus \mathbb{C}$  be an irrational number and set  $(p_n)_{n \in \mathbb{N}}, (q_n)_{n \in \mathbb{N}}$  be sequences in  $\mathbb{Z}$  or  $\mathbb{N}_+$  such that it holds

$$\lim_{n \rightarrow \infty} \frac{p_n}{q_n} = a.$$

Prove that  $\lim_{n \rightarrow \infty} q_n = \infty$

*Hint:* Prove by contradiction. Assume the sequence  $(q_n)_{n \in \mathbb{N}}$  does not converge to  $\infty$ . Consider that there must exist a bounded subsequence and use the Bolzano-Weierstrass theorem.

## 47 Exercise 55

### 47.1 Exercise 55.a

$$\sum_{n=0}^{\infty} \frac{n-5}{n^2+1}$$

Let  $n \geq 5$ . Direct comparison test:

$$\begin{aligned} \frac{1}{2^n} &\leq \left| \frac{n-5}{n^2+1} \right| \\ n^2+1 &\leq 8n^2-10n \\ 0 &\leq n^2-10n-1 \\ n &\geq 5 \pm \sqrt{26} = 11 \end{aligned}$$

So  $n \geq 11$  shows that it converges.

Ratio test:

$$\begin{aligned} &\lim_{n \rightarrow \infty} \left| \frac{n-4}{(n+1)^2+1} \cdot \frac{n^2+1}{n+5} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{(n-4)(n^2+1)}{((n+1)^2+1)(n+5)} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{n^3-4n^2+n-4}{(n^2+2n+1)(n+5)} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{\frac{n^3}{n^3} - \frac{4n^2}{n^3} + \frac{n^3}{n^3} - \frac{4}{n^3}}{\frac{n^3}{n^3} + \frac{2n^2}{n^3} - \frac{9n}{n^3} - \frac{5}{n^3}} \right| \end{aligned}$$

Shows convergence.

But this solution is wrong.

#### 47.1.1 Lathos' solution

$$\begin{aligned} &n \geq 10 \\ &\frac{n-5}{n^2+1} \geq \frac{n-\frac{n}{2}}{n^2+n^2} = \frac{1}{2} \frac{1}{n+n} = \frac{1}{4n} \\ &\sum_{n=1}^{\infty} \frac{1}{4n} \xrightarrow{\text{direct comparison crit.}} \sum_{n=10}^{\infty} \frac{n-5}{n^2+1} \text{ div. } \Rightarrow \sum_{n=0}^{\infty} \frac{n-5}{n^2+1} \text{ diverges} \end{aligned}$$

### 47.2 Exercise 55.b

$$\sum_{n=2}^{\infty} \frac{2n+1}{n(n-1)^2}$$

Direct comparison test:

$$\begin{aligned} \sum_{n=2}^{\infty} \frac{2n+1}{n(n-1)^2} &\leq \sum_{n=2}^{\infty} \left| \frac{4n}{n(\frac{1}{4}n)^2} \right| \cdot \sum_{n=2}^{\infty} \frac{4n}{n \frac{1}{16} n^2} \\ &= \sum_{n=2}^{\infty} \left| \frac{4 \cdot 16n}{n^3} \right| \\ &= 4 \cdot 16 \sum_{n=2}^{\infty} \left| \frac{n}{n^3} \right| \leq 64 \cdot \sum_{n=2}^{\infty} \left| \frac{4 \cdot 16n}{n^3} \right| \end{aligned}$$

And this converges absolutely.

### 47.3 Exercise 55.c

$$\begin{aligned} &\lim_{n \rightarrow \infty} \left| \left( \frac{4}{5} \right)^{n+1} (n+1)^5 \left( \frac{4}{5} \right)^{-n} n^{-5} \right| \\ &= \lim_{n \rightarrow \infty} \left| \frac{4}{5} \left( \frac{n^5}{n^5} + 5 \frac{n^4}{n^5} + 10 \cdot \frac{n^3}{n^5} + 10 \frac{n^2}{n^5} + 5 \frac{n}{n^5} + \frac{1}{n^5} \right) \right| \\ &= \left| \frac{4}{5} \right| < 1 \end{aligned}$$

So  $\sum_{n=5}^{\infty} \left( \frac{4}{5} \right)^n n^5$  converges.

### 47.4 Exercise 55.d

We use the Leibniz criterion.

We need to show that  $|a_{n+1}| < |a_n|$ .

$$\begin{aligned} &\left| \frac{(-1)^{n+1}}{\sqrt{n+1}} \right| < \left| \frac{(-1)^n}{\sqrt{n}} \right| \\ &\frac{1}{\sqrt{n+1}} < \frac{1}{\sqrt{n}} \\ &\sqrt{n} < \sqrt{n+1} \end{aligned}$$

$n < n+1$  so it is monotonically increasing.

Does it converge absolutely?

$$\sum_{n=1}^{\infty} \left| \frac{(-1)^n}{\sqrt{n}} \right| = \sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} \geq \sum_{n=1}^{\infty} \frac{1}{n}$$

No, this does not converge. So it does not.

### 47.5 Exercise 55.e

TODO



## 48 Exercise 56

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)(n+2)} = \sum_{n=1}^{\infty} \left( \frac{A}{n} + \frac{B}{n+1} + \frac{C}{n+2} \right)$$

$$1 = A(n+1)(n+2) + B(n+2)n + C(n+1)n$$

$$1 = An^2 + 3An + 2A + Bn^2 + 2Bn + Cn^2 + Cn$$

For terms of degree 2 it holds that,  $0 = A + B + C$ . For terms of degree 1 it holds that,  $0 = 3A + 2B + C$ . For terms of degree 0 it holds that,  $1 = 2A$ .

We solve this equation system and get  $A = \frac{1}{2}$ ,  $B = -1$  and  $C = \frac{1}{2}$ .

$$\begin{aligned} \sum_{n=1}^{\infty} \left( \frac{1}{2n} - \frac{1}{n+1} + \frac{1}{2(n+2)} \right) &= \frac{1}{2} \sum_{n=1}^{\infty} \left( \frac{1}{n} - \frac{2}{n+1} + \frac{1}{n+2} \right) \\ &= \lim_{N \rightarrow \infty} \frac{1}{2} \left( \left( \sum_{n=1}^N \frac{1}{n} - \frac{1}{n+1} \right) - \left( \sum_{n=1}^N \frac{1}{n+1} - \frac{1}{n+2} \right) \right) \\ &= \lim_{N \rightarrow \infty} \frac{1}{2} \left( \left( \sum_{n=1}^N \frac{1}{n} - \frac{1}{n+1} \right) - \left( \sum_{n=2}^{N+1} \frac{1}{n} - \frac{1}{n+1} \right) \right) \\ &= \lim_{N \rightarrow \infty} \frac{1}{2} \left( \left( \sum_{n=2}^N \frac{1}{n} - \frac{1}{n+1} \right) + \left( \frac{1}{1} - \frac{1}{2} \right) - \left( \sum_{n=2}^N \frac{1}{n} - \frac{1}{n+1} \right) - \frac{1}{N+1} + \frac{1}{N+2} \right) \\ &= \lim_{N \rightarrow \infty} \frac{1}{2} \left( \left( \frac{1}{1} - \frac{1}{2} \right) - \frac{1}{N+1} + \frac{1}{N+2} \right) \\ &= \lim_{N \rightarrow \infty} \frac{1}{2} \left( \frac{1}{2} - \frac{1}{N+1} + \frac{1}{N+2} \right) \\ &= \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4} \end{aligned}$$

## 49 Exercise 57

## 50 Exercise 58

TODO: I think I only considered real numbers. Not complex numbers.

Direct comparison test: The series  $\sum_{n=0}^{\infty} \left| \frac{a_n}{a_n+1} \right|$  converges if

1.  $0 < \left| \frac{a_n}{a_n+1} \right| \leq |a_n|$ , and
2.  $\sum_{n=0}^{\infty} |a_n|$  converges absolutely.

Firstly we cover a special case: Assume  $\exists n \in \mathbb{N}$  such that  $a_n = 0$ . Followingly this means there exists some  $N$  such that all following  $n$  are 0.

$$\exists N \in \mathbb{N} : \forall n \in \mathbb{N} : n \geq N : a_n = 0$$

In this case the partial sum will not change at some sufficiently large  $n$ . So  $\sum_{n=0}^{\infty} \left| \frac{a_n}{a_n+1} \right|$  converges.

In the following we can assume  $\forall n \in \mathbb{N} : a_n > 0$ .

1. We show that  $0 < \left| \frac{a_n}{a_n+1} \right| \leq |a_n|$ .

Here  $0 < \left| \frac{a_n}{a_n+1} \right|$  certainly holds because otherwise  $a_n = 0$  for some  $n$ . This would be the special case we have already covered and we don't need to look at any more.

We show  $\left| \frac{a_n}{a_n+1} \right| \leq |a_n|$ .

$$\begin{aligned} \frac{|a_n|}{|a_n+1|} &\leq |a_n| \\ \Rightarrow |a_n| &\leq |a_n| |a_n+1| \\ \Rightarrow 1 &\leq |a_n+1| \end{aligned}$$

2. This holds by precondition.

## 51 Exercise 59

## 52 Exercise 60

### 52.1 Exercise 60.a

wlog  $|a_n| \leq |b_n| \leq |a_n \cdot b_n| \leq |a_n^2|$ .

$$\begin{aligned} \Rightarrow |a_n^2| &\leq |a_n b_n| \leq |b_n^2| \\ \Rightarrow \sum_{n=0}^{\infty} |a_n b_n| & \end{aligned}$$

is absolute convergent.

### 52.2 Exercise 60.b

$$\sum_{n=0}^{\infty} a_n^2 + \sum_{n=0}^{\infty} (2a_n b_n) + \sum_{n=0}^{\infty} b_n^2$$

## 53 Exercise 61

**Exercise 37.** Determine the convergence radius of the following power series

**53.1 Exercise 61.a:**  $\sum_{n=1}^{\infty} \frac{1}{n^n} z^n$

**53.2 Exercise 61.b:**  $\sum_{n=1}^{\infty} \frac{(n!)^2}{(2n)!} z^n$

**53.3 Exercise 61.c:**  $\sum_{n=0}^{\infty} \frac{2^n}{n!} (z+i)^n$

**53.4 Exercise 61.d:**  $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{z^{2n-1}}{(2n-1)!}$

*Hint: Investigate convergence of this series, for fixed  $z \in \mathbb{C}$ , directly with help of the quotient criterion.*

## 54 Exercise 62

**Exercise 38.** We consider the function  $f : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$ ,  $x \mapsto \frac{1}{x}$ . Determine for appropriate subsets

of  $\mathbb{C} \setminus \{0\}$ , representations of  $f$  as power series of structure

$$\frac{1}{x} = \sum_{n=0}^{\infty} a_n(x-1)^n \quad \text{and} \quad \frac{1}{x} = \sum_{n=0}^{\infty} b_n(x-2)^n \quad (a_n, b_n \in \mathbb{C}).$$

Determine the convergence radius of both power series.

*Hint:* Geometric series.

## 55 Exercise 63

**Exercise 39.** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f(x) = x^3 - 2x^2 + x + 1$ . Show, with the help of its  $\varepsilon$ - $\delta$ -definition of continuity, that  $f$  is continuous at  $-1$ .

## 56 Exercise 64

**Exercise 40.** Let  $D \subseteq \mathbb{R}$ , and  $f, g : D \rightarrow \mathbb{R}$  in  $x_0 \in D$  are continuous functions and it holds that  $f(x_0) = g(x_0)$ . Show: Let  $h : D \rightarrow \mathbb{R}$  be a function, such that for all  $x \in D$

$$f(x) \leq h(x) \leq g(x).$$

Prove that  $h$  is continuous in  $x_0$ .

## 57 Exercise 65

**Exercise 41.** Let  $D \subseteq \mathbb{C}$ , and  $f, g : D \rightarrow \mathbb{C}$  are continuous in  $z_0 \in D$ . Show, with the use of the  $\varepsilon$ - $\delta$ -definition of continuity, that also the function

$$fg : \begin{cases} D & \rightarrow \mathbb{C}, \\ z & \mapsto f(z)g(z) \end{cases}$$

is continuous in  $z_0$ .

## 58 Exercise 66

**Exercise 42.** Determine all  $a \in \mathbb{R}$  such that the function is continuous in  $\mathbb{R}$ :

$$g : \mathbb{R} \rightarrow \mathbb{R}$$

$$g(x) = \begin{cases} x^3 + 2ax^2 + a^2 & \text{if } x \leq 1 \\ ax^2 + \frac{4a^2}{1+x^2} & \text{if } x > 1 \end{cases}$$

Let  $f(x) = x^3 + 2ax^2 + a^2$  and  $h(x) = ax^2 + \frac{4a^2}{1+x^2}$ .

1.  $f$  is a polynomial. A polynomial is continuous.
2.  $g$  is not a polynomial. But  $(ax^2)(1+x^2) + (4a^2)$  is a polynomial. And  $\frac{1}{1+x^2}$  is continuous in  $\mathbb{R}$ .
3. If the limes of  $f$  and  $g$  yields the same value at point  $x_0$ , then the function is continuous at this  $x_0$ .

$$\begin{aligned} \lim_{x \nearrow 1} ax^2 + \frac{4a^2}{1+x^2} &= a + \frac{4a^2}{2} \\ &= a + 2a^2 - a \end{aligned}$$

We use:

$$\lim_{n \rightarrow \infty} (a_n + b_n) = \lim_{n \rightarrow \infty} a_n + \lim_{n \rightarrow \infty} b_n$$

This holds only if  $\lim_{n \rightarrow \infty} a_n$  and  $\lim_{n \rightarrow \infty} b_n$  exist. Furthermore:

$$\lim_{x \searrow 1} ax^2 = a \cdot \lim_{x \searrow 1} x^2$$

$g$  is continuous in

$$a = \frac{1 \pm \sqrt{5}}{2}$$

Fundamental theorem supporting this solution:  $f$  is continuous in  $x_0$  if and only if  $\lim_{x \nearrow x_0} f(x) = \lim_{x \searrow x_0} f(x)$ .

Alternative approach: Consider  $g(x) - d(x)$  where where

$$d(x) = \begin{cases} x^3 + 2ax^2 + a^2 - (1+a)^2 & \text{if } x \leq 1 \\ ax^2 + \frac{4a^2}{1+x^2} - a(1+2a) & \text{if } x > 1 \end{cases}$$

Then

$$d_{x \leq 1}(x) = (x^3 - 1) + 2a(x^2 - 1) = (x - 1)[x^2 + x + 1 + 2a(x + 1)]$$

Consider the sequence  $d_{x \leq 1} \forall x_n \rightarrow 1$ . Then

$$d_{x \leq 1} \rightarrow 0(3 + 4a).$$

Also

$$d_{x > 1}(x) = ax^2 + \frac{4a^2}{1+x^2} - a - 2a^2 = (x - 1)(\text{some bounded value})$$

## 59 Exercise 67

**Exercise 43.** Let  $f, g : D \rightarrow \mathbb{R}$  with  $D \subseteq \mathbb{R}$ . The functions  $|f| : D \rightarrow \mathbb{R}$  and  $\max\{f, g\} : D \rightarrow \mathbb{R}$  is defined by

$$\forall x \in D : |f|(x) = |f(x)| \quad \max\{f, g\}(x) = \max\{f(x), g(x)\}$$

### 59.1 Exercise 67.a: $|f|$ is continuous, then $f$ is continuous

Counterexample:

$$|f| = \begin{cases} -1 & \text{if } x \leq 0 \\ 1 & \text{if } x > 0 \end{cases} \Rightarrow |f|(x) = 1$$

Is continuous. But

$$f(x) = \begin{cases} -1 & \text{if } x \leq 0 \\ 1 & \text{if } x > 0 \end{cases}$$

is not.

### 59.2 Exercise 67.b: $f$ and $g$ are continuous, then $\max\{f, g\}$ is continuous

We use this definition of  $\max\{f, g\}$ .

$$\max_{f, g}(x) = \frac{f(x) + g(x)}{2} + \frac{|f(x) - g(x)|}{2}$$

We know that the addition of two continuous functions is continuous. Also the multiplication of a continuous function with a scalar is continuous. It is easy to see that also the absolute value of the difference of two continuous functions is continuous.

So all involved functions are continuous and therefore  $\max_{f, g}$  is also continuous.

### 59.3 Exercise 67.c: $fg$ is continuous, then $f$ and $g$ are continuous

Counterexample:

$$f(x) = \begin{cases} 1 & \text{if } x = 1 \\ 0 & \text{otherwise} \end{cases} \quad g(x) = 0$$

$fg(x)$  is continuous, but not  $f(x)$ .

## 60 Exercise 68

**Exercise 44.** Show that the function  $h$  is continuous, but not uniformly continuous.

$$h : \mathbb{R} \rightarrow \mathbb{R} \quad x \mapsto x^2$$

$f : D \subseteq \mathbb{C} \rightarrow \mathbb{C}$  is uniformly continuous iff

$$\forall \varepsilon > 0 \exists \delta > 0 : \forall z_0, z_1 \in D \text{ with } |z_1 - z_0| < \delta \Rightarrow |f(z_1) - f(z_0)| < \varepsilon$$

$f : D \subseteq \mathbb{C} \rightarrow \mathbb{C}$  is continuous iff

$$\forall \varepsilon > 0 \exists \delta > 0 : \forall x \in D \wedge |x - x_0| < \delta \Rightarrow |f(x) - f(x_0)| < \varepsilon$$

It is not uniformly continuous because ...

Choose  $\varepsilon = 1$  and  $\delta$  arbitrary. Consider  $x_0 = \sqrt{n}$  and  $x = \sqrt{n+1}$ .

$$|x_0 - x| = |\sqrt{n} - \sqrt{n+1}| = -\sqrt{n} + \sqrt{n+1} \cdot \frac{\sqrt{n} + \sqrt{n+1}}{\sqrt{n} + \sqrt{n+1}} = \frac{n - n + 1}{\sqrt{n} + \sqrt{n+1}} \stackrel{!}{<} \delta$$

This holds with  $n > (\frac{1}{2}\delta)^2$ .

$x^2$  is a polynomial and polynomials are continuous.

## 61 Exercise 72

**Exercise 45.** Show that the equation  $x^7 + x = 1$  has exactly one real solution.

$$x^7 + x = 1$$

$$f(x) = x^7 + x - 1$$

Choose

- $x_1 = 0$ :  $f(x_1) = f(0) = -1 < 0$
- $x_2 = 1$ :  $f(x_2) = 1 > 0$

Because  $f$  is continuous, the intermediate value theorem applies. So,

$$\exists x_0 \in [0, 1] : f(x_0) = 0 \Rightarrow \exists \text{ solution}$$

Assume, there exists some  $x_0, \tilde{x}_0$  with  $x_0 \neq \tilde{x}_0$ . Then it holds that

$$f(x_0) = 0 = f(\tilde{x}_0) \Rightarrow \exists \xi \in (x_0, \tilde{x}_0) : f'(\xi) = 0$$

but

$$f'(x) = 7 \underbrace{x^6}_{\geq 1} + 1 \geq 1$$

This contradicts with our assumption that a solution to  $f'(\xi) = 0$  exists (Rolle's Theorem).

## 62 Exercise 73

**Exercise 46.**  $f : [a, b] \rightarrow \mathbb{R}$  is a continuous function and let  $x_1, \dots, x_n \in [a, b]$ . Show that:

$$\exists y \in [a, b] : f(y) = \frac{1}{n} \sum_{i=1}^n f(x_i)$$

We look at the minimum and maximum:

$$\begin{aligned} \min_{x \in [a, b]} f(x) &\leq f(x) \leq \max_{x \in [a, b]} f(x) \\ n \cdot \min_{x \in [a, b]} f(x) &\leq \sum_{i=1}^n f(x_i) \leq n \cdot \max_{x \in [a, b]} f(x) \\ \min_{x \in [a, b]} f(x) &\leq \frac{1}{n} \sum_{i=1}^n f(x_i) \leq \max_{x \in [a, b]} f(x) \end{aligned}$$

by the Intermediate Value Theorem it holds that

$$\exists y \in [a, b] : f(y) = \frac{1}{n} \sum_{i=1}^n f(x_i)$$

## 63 Exercise 74

**Exercise 47.**  $f : \mathbb{Q} \rightarrow \mathbb{Q}$  is defined as  $f(x) = x^2 - 2$ . Show that  $f$  is continuous and furthermore that  $f(0) = -1 < 2 = f(2)$  but  $f(x) \neq 0$  for all  $x \in [0, 2] \cap \mathbb{Q}$ . (This means that the Intermediate Value Theorem does not hold for continuous  $f : \mathbb{Q} \rightarrow \mathbb{Q}$ )

You can use  $\sqrt{2} \notin \mathbb{Q}$  without proof.

### 63.1 $f$ is continuous

$$\forall \varepsilon > 0 \exists \delta > 0 \forall x \in \mathbb{Q} : |x - x_0| < \delta \Rightarrow |f(x) - f(x_0)| < \varepsilon$$

Let  $\varepsilon > 0$ . Choose  $\delta := \min \left\{ 1, \frac{\varepsilon}{1+2x_0} \right\}$ .

Consider

$$\begin{aligned} |f(x) - f(x_0)| &= |x^2 - 2 - x_0^2 + 2| \\ &= |(x - x_0)| |(x + x_0)| \\ &< \delta |x - x_0 + 2x_0| \\ &\leq \delta (|x - x_0| + |2x_0|) \\ &\leq \delta (\delta + 2x_0) \\ &\leq \delta (1 + 2x_0) \\ &\leq \frac{\varepsilon}{1 + 2x_0} \cdot (1 + 2x_0) \\ &= \varepsilon \end{aligned}$$

### 63.2 Find $f(x) = 0 = x^2 - 2$

$$\begin{aligned} x^2 &= 2 \\ x &= \pm \sqrt{2} \notin \mathbb{Q} \end{aligned}$$

$$\Rightarrow f(x) \neq 0 \quad \forall x \in \mathbb{Q}$$

Hence, the intermediate value theorem only holds for compact fields.

### 63.3 Using the sequence theorem

Actually, you can also apply the sequence theorem to show this exercise.

$$\left( \lim_{x_n \rightarrow x \in \mathbb{Q}} x_n \right)^2 - 2 = x^2 - 2 = f(x)$$

## 64 Exercise 75

**Exercise 48.** Determine, if possible, the following limits (for  $z \in \mathbb{C}, x \in \mathbb{R}$ ):

**64.1 Exercise 75.a:**  $\lim_{z \rightarrow 1} \frac{z^m - 1}{z^n - 1}$  **with**  $m, n \in \mathbb{N}_+$

$$\begin{aligned} \frac{z^m - 1}{z^n - 1} &= \frac{(z - 1)(z^{m-1} + \dots + 1)}{(z - 1)(z^{n-1} + \dots + 1)} = \frac{\sum_{k=0}^{m-1} z^k}{\sum_{l=0}^{n-1} z^l} \\ \lim_{z \rightarrow 1} \frac{\sum_{k=0}^{m-1} z^k}{\sum_{l=0}^{n-1} z^l} &= \frac{\lim_{z \rightarrow 1} \sum_{k=0}^{m-1} z^k}{\lim_{z \rightarrow 1} \sum_{l=0}^{n-1} z^l} \\ &= \frac{\sum_{k=0}^{m-1} \lim_{z \rightarrow 1} z^k}{\sum_{l=0}^{n-1} \lim_{z \rightarrow 1} z^l} = \frac{1 + (m-1) \cdot 1}{1 + (n-1) \cdot 1} = \frac{m}{n} \end{aligned}$$

**64.2 Exercise 75.b:**  $\lim_{x \rightarrow \infty} \sqrt{x - \sqrt{x}} - \sqrt{x}$

$$\begin{aligned} \sqrt{x - \sqrt{x}} - \sqrt{x} &= \frac{(\sqrt{x - \sqrt{x}} - \sqrt{x}) \cdot (\sqrt{x - \sqrt{x}} + \sqrt{x})}{\sqrt{x - \sqrt{x}} + \sqrt{x}} = \frac{(\sqrt{x - \sqrt{x}})^2 - \sqrt{x}^2}{\sqrt{x - \sqrt{x}} + \sqrt{x}} \\ &= \frac{x - \sqrt{x} - x}{\sqrt{x - \sqrt{x}} + \sqrt{x}} = \frac{-1}{\sqrt{1 - \frac{\sqrt{x}}{x}} + 1} = \frac{-1}{\sqrt{1 - \frac{1}{\sqrt{x}}} + 1} \end{aligned}$$

This is a composition of continuous functions.

$$\lim_{x \rightarrow \infty} \sqrt{x - \sqrt{x}} - \sqrt{x} = \frac{-1}{\sqrt{1 - \lim_{x \rightarrow \infty} \frac{1}{\sqrt{x}}} + 1} = \frac{-1}{2}$$

## 65 Supporting sessions

Fr, 29.1 12:00–13:15, Kniely, HS 11.02

Mo, 1.2 13:00–15:00, Peichl, HS 11.02

Exam: Sheet 7 and following

Convergence, sequences, recursive sequences, partial sums, series, Cauchy convergence, absolute convergence, convergence criteria, convergence radius, continuity,  $\varepsilon$ - $\delta$ -criterion, intermediate value theorem, Rolle's theorem, convergence of functions, differentiability

## 66 Exercise 79

$$f(x) = \frac{(x+1)^2}{x^2+1} \geq 0$$

$$f'(x) = \frac{-2(x^2-1)}{(x^2+1)^2}$$

$$f''(x) = \frac{4x(x^2-3)}{(x^2+1)^3}$$

Roots:  $x = 1$  and  $x = -1$ .