

# Analysis 2

Lecture notes, University (of Technology) Graz  
based on the lecture by Wolfgang Ring

Lukas Prokop

July 26, 2020

## Contents

<b>1</b>	<b>Mathematical Redux and topological fundamentals</b>	<b>3</b>
1.1	Metric . . . . .	3
1.2	Cauchy-Schwarz inequality . . . . .	4
1.3	Euclidean norm . . . . .	5
1.4	Metric space . . . . .	5
1.5	Open sets, convergence and accumulation points . . . . .	6
1.6	Norm . . . . .	7
1.7	Contact point . . . . .	9
1.8	Topology . . . . .	13
1.9	Variations of continuity notions . . . . .	15
1.9.1	Uniform continuity . . . . .	15
1.9.2	Lipschitz continuity . . . . .	15
1.10	Banach Fixed Point Theorem . . . . .	15
<b>2</b>	<b>Compactness</b>	<b>18</b>
2.1	Definition . . . . .	18
2.2	Boundedness . . . . .	19
2.3	Total boundedness . . . . .	20
2.4	Compactness, continuity and openness . . . . .	20
2.5	Normed vector spaces . . . . .	24

<b>3</b>	<b>Integration calculus</b>	<b>28</b>
3.1	Partitions and refinements . . . . .	28
3.2	Characteristic functions . . . . .	33
3.3	Limit points . . . . .	35
3.4	Regulated functions . . . . .	38
3.5	Bounded functions on bounded sets . . . . .	39
<b>4</b>	<b>Integration of regulated functions</b>	<b>43</b>
4.1	Elementary properties of an integral . . . . .	45
4.2	Mean value theorem of integration calculus . . . . .	48
4.3	One-sided derivatives . . . . .	52
4.4	Fundamental theorem of differential/integration calculus . . . .	53
4.5	Integration methods . . . . .	59
4.5.1	Partial integration . . . . .	59
4.5.2	Integration by substitution . . . . .	62
4.6	Improper integrals . . . . .	66
4.7	Young's inequality . . . . .	71
4.8	Hölder's inequality . . . . .	75
<b>5</b>	<b>Elaboration on differential calculus</b>	<b>76</b>
5.1	Multiple times differentiable . . . . .	80
5.2	Best polynomial approximation of a function . . . . .	80
5.3	Taylor polynomial . . . . .	82
5.4	Integral form of the Taylor remainder term . . . . .	84
5.5	Lagrange form of the Taylor remainder term . . . . .	85
5.6	Sufficient criteria for extreme values . . . . .	87
5.7	Qualitative Taylor equation . . . . .	88
5.8	Taylor series . . . . .	89
<b>6</b>	<b>Multidimensional differential calculus</b>	<b>92</b>
6.1	Frobenius and matrix norm . . . . .	92
6.2	Operator norm and bounded linear operators . . . . .	93
6.3	Landau notation . . . . .	96
6.4	Multidimensional derivative of a function . . . . .	97

6.5	Fréchet derivative . . . . .	98
6.6	Chain rule . . . . .	100
6.7	Differentiability on $D$ . . . . .	103
6.7.1	Determination of $Df(x_0) \in \mathbb{R}^{n \times m}$ . . . . .	103
6.8	Partial derivative and Jacobi matrix . . . . .	105
6.9	Total differentiability, nabla operator and gradients . . . . .	108
6.10	Optimality criteria for multi-dimensional functions . . . . .	113
6.11	Diffeomorphism . . . . .	114
6.12	Local inversion theorem . . . . .	114
6.13	Implicit functions . . . . .	118
6.14	Partial derivatives and Hessian Matrix . . . . .	122
6.15	Schwarz' Theorem . . . . .	123
6.16	Multiindex notation . . . . .	126
6.17	Qualitative Taylor Theorem . . . . .	130
6.18	Sufficient optimality criteria . . . . .	132
<b>7</b>	<b>Differential geometry of curves</b>	<b>134</b>
7.1	Curves, regular curves and tangential vectors . . . . .	134
7.2	Reparameterization of parameterized curves . . . . .	136
7.3	Arc length of parameterized curves . . . . .	142
7.4	Curvature, Torsion, Frenet's formulas . . . . .	150
7.5	Curvatures of plane curves . . . . .	159

↓ This lecture took place on 2018/03/06.

# 1 Mathematical Redux and topological fundamentals

## 1.1 Metric

**Definition 1.1.** Let  $X \neq \emptyset$  be a set. We define a map  $d : X \times X \rightarrow [0, \infty)$ .  $d$  should behave like a geometrical distance. We require  $\forall x, y, z \in X$ :

- $d(x, y) = d(y, x)$  [called symmetry]

- $d(x, y) = 0 \iff x = y$  [called positive definiteness]
- $\forall x, y, z \in X : d(x, z) \leq d(x, y) + d(y, z)$  [called triangle inequality]

Then  $d$  is called metric or distance function on  $X$ .  $(X, d)$  is called metric space.

**Example 1.1.**

- $X \subseteq \mathbb{C}$ ,  $d(x, y) = |x - y|$ . It satisfies  $|x - z| \leq |x - y| + |y - z|$
- $X \subseteq \mathbb{R}^n$ ,  $\|x - y\| = \langle x - y, x - y \rangle^{\frac{1}{2}}$

**Claim.**

$$\langle x, y \rangle = \sum_{i=1}^n x_i y_i$$

$$\|x\| = \langle x, x \rangle^{\frac{1}{2}} = \sqrt{\sum_{i=1}^n x_i^2}$$

$$\|x\| = \sqrt{x_1^2 + x_2^2}$$

The triangle inequality holds:  $\|x + y\| \leq \|x\| + \|y\|$ .

*Proof.*

$$\begin{aligned} \|x + y\|^2 &= \langle x + y, x + y \rangle \\ &= \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle \\ &= \|x\|^2 + 2 \langle x, y \rangle + \|y\|^2 \\ &\leq \|x\|^2 + 2 \|x\| \|y\| + \|y\|^2 && \text{[by Cauchy-Schwarz ineq.]} \\ &= (\|x\| + \|y\|)^2 \\ \|x - y\|^2 &= \langle x - y, x - y \rangle \\ &= \|x\|^2 - 2 \langle x, y \rangle + \|y\|^2 \\ \|x + y\|^2 + \|x - y\|^2 &= 2(\|x\|^2 + \|y\|^2) \end{aligned}$$

□

## 1.2 Cauchy-Schwarz inequality

**Theorem 1.1** (Cauchy-Schwarz inequality).

$$|\langle x, y \rangle| \leq \|x\| \|y\|$$

*Proof.*

$$0 \leq \|x - \lambda y\|^2 = \langle x - \lambda y, x - \lambda y \rangle = \|x\|^2 - 2\lambda \langle x, y \rangle + \lambda^2 \|y\|^2 \quad \forall \lambda \in \mathbb{R}$$

$$\lambda := \frac{\langle x, y \rangle}{\|y\|^2} \implies 0 \leq \|x\|^2 - 2 \frac{|\langle x, y \rangle|^2}{\|y\|^2} + \frac{|\langle x, y \rangle|^2}{\|y\|^4} \cdot \|y\|^2$$

$$\iff 0 \leq \|x\|^2 - \frac{|\langle x, y \rangle|^2}{\|y\|^2} \iff \frac{|\langle x, y \rangle|^2}{\|y\|^2} \leq \|x\|^2 \iff |\langle x, y \rangle|^2 \leq \|x\|^2 \cdot \|y\|^2$$

□

### 1.3 Euclidean norm

**Definition 1.2.**  $\|x\| = \sqrt{\sum_{i=1}^n x_i^2}$  is called Euclidean norm (length) of vector  $x \in \mathbb{R}^n$ .

$\|x\| = \langle x, x \rangle^{\frac{1}{2}}$ . It satisfies:

1.  $\|\lambda x\| = |\lambda| \|x\| \quad \forall x \in \mathbb{R}^n, \lambda \in \mathbb{R}$
2.  $\|x\| = 0 \iff x = 0 \text{ in } \mathbb{R}^n$
3.  $\|x + y\| \leq \|x\| + \|y\|$

In general: Let  $V$  be a vector space over  $\mathbb{R}$ . A map  $\|\cdot\|$ , which assigns every vector  $x$  a non-negative real number satisfying the properties above, is called **norm** on  $V$ . Then  $(V, \|\cdot\|)$  is called a **normed vector space**.

Let  $X \subseteq \mathbb{R}^n$  ( $V$  is a normed vector space), then  $d(x, y) = \|x - y\|$  is a metric on  $X$ .

$$\|y - x\| = \|(-1)(x - y)\| = |-1| \cdot \|x - y\| = \|x - y\|$$

$$d(x, y) = 0 \iff \|x - y\| = 0 \iff x - y = 0 \iff x = y$$

$$d(x, z) = \|z - x\| = \|z - y + y - x\| \leq \|z - y\| + \|y - x\| = d(z, y) + d(y, x)$$

### 1.4 Metric space

**Example 1.2** (metric space). Distance is not a norm. Consider an area in  $\mathbb{R}^3$ .

$d(x, y)$  is the shortest path, connecting  $x$  and  $y$  in  $X$ . See Figure 1.

**Example 1.3** (French railway metric). All connections between two cities pass through Paris except one of the cities is Paris.

**Example 1.4.**  $X = \mathbb{R}^2$ . Let  $p \in \mathbb{R}^2$  be fixed.

$$d(x, y) = \begin{cases} |x - y| & \text{if } x, y, p \text{ are on one line} \\ |x - p| + |p - y| & \text{if } x, y, p \text{ are not on one line} \end{cases}$$




Figure 1: Example in  $\mathbb{R}^3$ . The red line illustrates the shortest path, which is not necessarily "straight".

## 1.5 Open sets, convergence and accumulation points

Now we put some terminology into the context of a metric space.  $(X, d)$  is a metric space.

**Definition 1.3.** Let  $x \in X, r \geq 0$ .  $K_r(x)$  is an open sphere with radius  $r$  at center  $x$ .

$$K_r(x) := \{z \in X \mid d(x, z) < r\}$$

**Definition 1.4.**  $\overline{K_r(x)}$  is a closed sphere with center  $x$  and radius  $r$ .

$$\overline{K_r(x)} := \{z \in X \mid d(x, z) \leq r\}$$

**Definition 1.5** (Sequences in  $X$ ). Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence in  $X$  (hence,  $\forall n \in \mathbb{N} : x_n \in X$ )

1.  $(x_n)_{n \in \mathbb{N}}$  is called convergent with limit  $x \in X$  if

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} : n \geq N \implies d(x_n, x) < \varepsilon$$

Denoted as  $\lim_{n \rightarrow \infty} x_n = x$ .

2.  $(x_n)_{n \in \mathbb{N}}$  is a Cauchy sequence if

$$\forall \varepsilon > 0 \exists N \in \mathbb{N} : n, m \geq N \implies d(x_n, x_m) < \varepsilon$$

**Claim.** Every convergent sequence is also a Cauchy sequence.

*Proof.* Let  $(x_n)_{n \in \mathbb{N}}$  be convergent with limit  $x$ . Let  $\varepsilon > 0$  be arbitrary. Because  $(x_n)_{n \in \mathbb{N}}$  is convergent, there exists  $N \in \mathbb{N}$  such that  $n \geq N \implies d(x_n, x) < \frac{\varepsilon}{2}$ . Now let  $n, m \geq N$ . Then,

$$d(x_n, x_m) \leq \underbrace{d(x_n, x)}_{< \frac{\varepsilon}{2}} + \underbrace{d(x, x_m)}_{< \frac{\varepsilon}{2}} < \varepsilon$$

□

**Definition 1.6.**  $(X, d)$  is called complete metric space if every Cauchy sequence in  $X$  is also convergent (has a limit) in  $X$ .

$\mathbb{R}$  is complete.  $\mathbb{R}^n$  is also complete.  $\mathbb{Q} \subseteq \mathbb{R}$  is incomplete.

**Definition 1.7.** Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence of  $X$ .  $x$  is called “accumulation point” (dt. Häufungspunkt) of the sequence if  $\forall \varepsilon > 0 : K_\varepsilon(x)$  contains infinitely many sequence elements.

↓ This lecture took place on 2018/03/08.

$(X, d)$  is called metric space.

$$\begin{aligned} d(x, y) = 0 &\iff x = y \\ \forall x, y \in X : d(x, y) &= d(y, x) \\ d(x, z) &\leq d(x, y) + d(y, z) \forall x, y, z \in X \end{aligned}$$

## 1.6 Norm

Let  $V$  be a vector space.  $\|\cdot\|$  is called norm on  $V$ .

$$\begin{aligned} \|x\| = 0 &\iff x = 0 \\ \forall \lambda \in \mathbb{R}, \mathbb{C} : \forall x \in V : \|\lambda x\| &= |\lambda| \|x\| \\ \forall x, y, z \in V : \|x + y\| &\leq \|x\| + \|y\| \end{aligned}$$

Let  $X \subseteq V$  be a subset of normed vector space  $V$ . Then  $X$  is a metric space with induced metric  $d(x, y) = \|x - y\|$ .

Consider  $V = \mathbb{R}^n$ . Then  $\|x\|_2 = \left(\sum_{i=1}^n x_i^2\right)^{\frac{1}{2}}$  is a norm on  $\mathbb{R}^n$ .  $\|x\|_2$  is called Euclidean norm on  $\mathbb{R}^n$ .

Other norms in  $\mathbb{R}^n$ :

$$\|x\|_\infty = \max \{|x_i| \mid i = 1, \dots, n\}$$

$$\|x\|_1 = \sum_{i=1}^n |x_i|$$

for  $1 \leq p < \infty$ .

$$\|x\|_p = \left( \sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}}$$

e.g.  $\|x\|_1$  in  $\mathbb{R}^2$




Figure 2: Visualizing  $\|x\|_1$ , the Manhattan norm. You can think of walking the distance along the x-axis combined with the distance along the y-axis.

$$\|x - y\| = |x_1 - y_1| + |x_2 - y_2|$$

is the so-called *Manhattan metric*.

The concepts “subsequence”, “final element of a sequence”, “reordering of a sequence” correspond one-by-one to metric spaces.

**Definition 1.8** (Accumulation point). Let  $(X_n)_{n \in \mathbb{N}}$  be a sequence in  $X$ .  $x \in X$  is called accumulation point of sequence  $X$  if  $\forall \varepsilon > 0$  the sphere  $K_\varepsilon(x)$  contains infinitely many elements.

**Lemma 1.1.**  $x \in X$  is accumulation point of sequence  $(x_n)_{n \in \mathbb{N}}$  iff there exists a subsequence  $(x_{n_k})_{k \in \mathbb{N}}$  such that  $x = \lim_{k \rightarrow \infty} x_{n_k}$ .

*Proof.* See Analysis 1 course

□





Figure 3: Contact points in set  $B$

### 1.7 Contact point

Let  $B \subseteq X$  and  $X$  is a metric space. Then  $B$  with  $d$  is a metric space itself.

**Definition 1.9.** Let  $B \subseteq X$  and  $x \in X$ . We say,  $x$  is a contact point of  $B$  if  $\forall \varepsilon > 0 : K_\varepsilon(x) \cap B \neq \emptyset$ .

Equivalently,  $y \in X$  is not a contact point of  $B \iff \exists \varepsilon > 0 : K_\varepsilon(y) \cap B = \emptyset$ .

See Figure 3.

We let  $\bar{B} = \{x \in X \mid x \text{ is contact point of } B\}$ .

$\bar{B}$  is called closed hull of  $B$ .

$B$  is called closed if  $B = \bar{B}$ , hence, every contact point is also element of  $B$ .

**Remark 1.1.** Because  $\forall x \in B$  holds  $K_r(x) \cap B \supseteq \{x\} \forall r > 0$  is  $x$  always contact point of  $B$ . Also  $B \subseteq \bar{B}$  (always)

**Lemma 1.2.**  $x$  is contact point of  $B \iff \exists (x_n)_{n \in \mathbb{N}}$  with  $x_n \in B$  and  $\lim_{n \rightarrow \infty} x_n = x$ .

*Proof.* Let  $x$  be a contact point of  $B$ .

**Direction  $\Rightarrow$**  Because  $K_{\frac{1}{n}}(x) \cap B \neq \emptyset$ , choose  $x_n \in K_{\frac{1}{n}}(x) \cap B$ . The sequence  $(x_n)_{n \in \mathbb{N}}$  has property  $d(x_n, x) < \frac{1}{n}$ . Let  $\varepsilon > 0$  be arbitrary. Choose  $N \in \mathbb{N}$  such that  $N > \frac{1}{\varepsilon}$  (consider the Archimedean axiom). Then for  $n \geq N$ ,  $d(x_n, x) < \frac{1}{n} \leq \frac{1}{N} < \varepsilon$ , hence  $\lim_{n \rightarrow \infty} x_n = x$ .

**Direction  $\Leftarrow$**  Let  $x = \lim_{n \rightarrow \infty} x_n$  and  $x_n \in B$ . Let  $\varepsilon > 0$  be arbitrary and  $N \in \mathbb{N}$  such that  $d(x_n, x) < \varepsilon \forall n \geq N$ . Then  $d(x_N, x) < \varepsilon$ , hence

$$x_N \in \underbrace{K_\varepsilon(x) \cap B}_{\neq \emptyset}$$

So  $x$  is contact point of  $B$ .

□

**Lemma 1.3** (“ $\overline{B}$  is closed in itself”).  $\forall B \subseteq X : \overline{B} = \overline{\overline{B}}$ .

*Proof.*  $\overline{B} \subseteq \overline{\overline{B}}$  is trivial, since this holds for all closures. It remains to show that  $\overline{\overline{B}} \subseteq \overline{B}$ . Let  $x \in \overline{\overline{B}}$ , show that  $x \in \overline{B}$ .

Since  $x \in \overline{\overline{B}}$ ,  $x$  is a contact point. Thus,  $\forall \gamma > 0 : K_\gamma(x) \cap \overline{B} \neq \emptyset$ . Hence, for arbitrary  $\gamma > 0$ , there exists  $y : y \in K_\gamma(x) \wedge y \in \overline{B}$ .

Since  $y \in \overline{B}$ ,  $y$  is a contact point. Thus  $\forall \delta > 0 : B_\delta(y) \cap B \neq \emptyset$ . Hence, for arbitrary  $\delta > 0$ , there exists  $z : z \in K_\delta(y) \wedge z \in B$ .

Let  $\varepsilon > 0$  and  $\gamma := \frac{\varepsilon}{2}$  and  $\delta := \frac{\varepsilon}{2}$ . Then,

$$d(z, x) \leq \underbrace{d(z, y)}_{< \frac{\varepsilon}{2}} + \underbrace{d(y, x)}_{< \frac{\varepsilon}{2}} < \varepsilon$$

Thus,  $z \in K_\varepsilon(x)$ . We already know that  $z \in B$ . Since  $z$  exists, we showed that  $K_\varepsilon(x) \cap B \neq \emptyset$ . Since  $\varepsilon$  was chosen arbitrarily, this is true for any  $\varepsilon > 0$ .

$$\implies \forall \varepsilon > 0 : B_\varepsilon(x) \cap B \neq \emptyset \implies x \in \overline{B}$$

□

**Lemma 1.4.** Let  $X$  be a metric space.

- $A_i \subseteq X$  be closed  $\forall i \in I$ . Then  $A = \bigcap_{i \in I} A_i = \{x \in X \mid x \in A_i \forall i \in I\}$  is closed itself.
- $A_1, \dots, A_n \subseteq X$  are closed. Then  $\bigcup_{k=1}^n A_k$  is closed in  $X$ .
- $\varphi$  is closed,  $X$  is closed.

*Proof.* See Analysis 1 course. □

**Definition 1.10.** Let  $x \in X$  be called accumulation point of set  $B \subseteq X$  if  $\forall \varepsilon > 0 : (K_\varepsilon(x) \setminus \{x\}) \cap B \neq \emptyset$ .

**Remark 1.2.** Accumulation points only exist in the context of sets. Accumulation values only exist in the context of sequences.

For example  $(+1, -1, +1, -1, +1, \dots)$  has accumulation values  $+1$  and  $-1$ .

**Lemma 1.5.** Let  $x \in X$  be accumulation point on  $B \iff$  every sphere  $K_\varepsilon(x)$  contains infinitely many points of  $B$ .




Figure 4:  $x$  is an inner point of  $U$  if  $\exists r > 0 : K_r(x) \subseteq U$

*Proof.* Direction  $\Leftarrow$  is trivial.

Direction  $\Rightarrow$ : Choose  $x_1 \in (K_1(x) \setminus \{x\}) \cap B$ , hence  $x_1 \neq x$ ,  $x_1 \in B$  and  $d(x_1, x) < 1$ . Let  $r_1 = 1$ .

Inductive: choose  $r_n = \min(\frac{1}{n}, d(x_{n-1}, x))$  and  $x_n \in (K_{r_n}(x) \setminus \{x\}) \cap B$ . Then  $d(x_n, x) > 0$  (because  $x_n \neq x$ ) where  $d(x_n, x) < r_n < \frac{1}{n}$ .

$$0 < d(x_n, x) < \frac{1}{n}$$

Furthermore,  $d(x_n, x) < r_n \leq d(x_{n-1}, x)$ . So  $x_n \neq x_{n-1}$ .

Inductive:  $x_n \neq x_{n-1} \neq x_{n-2} \neq \dots \neq x_1$ . Now consider arbitrary  $\varepsilon > 0$  and  $N$  large enough such that  $\frac{1}{N} < \varepsilon$ .

Then  $\forall n \geq N : 0 < d(x_n, x) < \frac{1}{n} \leq \frac{1}{N} < \varepsilon$ . So  $K_\varepsilon(x) \cap B$  contains infinitely many points  $x_N, x_{N+1}, x_{N+2}, \dots$ .  $\square$

**Definition 1.11.** Let  $U \subseteq X$  and  $x \in U$ . We say  $x$  is an inner point of  $U$  if  $\exists r > 0 : K_r(x) \subseteq U$ . We let  $\mathring{U} = \{x \in U \mid x \text{ is inner point of } U\}$  and call it interior of  $U$  (dt. offenen Kern von  $U$  or das Innere von  $U$ ).  $O \subseteq X$  is called open (open set), if every point  $x \in O$  is also an inner point of  $O$ . Hence  $\mathring{O} = O$ . Compare with Figure 4.

**Example 1.5.** Let  $K_r(x)$  with  $r > 0$  be an open sphere in  $X$ . Then  $K_r(x)$  is an open set in  $X$ . Compare with Figure 5.

*Proof.* Why? Let  $y \in K_r(x)$ . Show that  $y$  is an inner point of the sphere.  $d(y, x) = s < r$ . Define  $r' = r - s > 0$ .

**Claim.**  $K_{r'}(y) \subseteq K_r(x)$ .

Let  $z \in K_{r'}(y)$ , hence  $d(z, y) < r'$ . Then,

$$d(z, x) \leq \underbrace{d(x, y)}_{< r'} + \underbrace{d(y, z)}_{= s} < r' + s = r$$

So  $z \in K_r(x)$  and therefore  $K_{r'}(y) \subseteq K_r(x)$ .  $\square$




Figure 5: Let  $K_r(x)$  with  $r > 0$  be an open sphere in  $X$ . Then  $K_r(x)$  is an open set in  $X$ .

**Lemma 1.6.** *Let  $U \subseteq X$  be arbitrary. Then  $\mathring{U} \subseteq X$  is an open set in  $X$ .*

*Proof.* Let  $x \in \mathring{U}$ . Hence  $x$  is an inner point of  $U$ . Show that  $x$  is an inner point of  $\mathring{U}$ , also  $\exists r > 0 : K_r(x) \subseteq \mathring{U}$ .

Because  $x \in \mathring{U}$ ,  $r > 0$  exists:  $K_r(x) \subseteq U$ . Claim: Every point  $y \in K_r(x)$  is also an inner point of  $U$ . Obvious (previous example), because  $r' > 0$  exists such that  $K_{r'}(y) \subseteq K_r(x) \subseteq U$  so  $y \in \mathring{U}$  and  $K_r(x) \subseteq \mathring{U}$ .  $\square$

**Theorem 1.2.** *Let  $X$  be a metric space.*

$$A \subseteq X \text{ is closed in } X \iff O = X \setminus A = A^C \text{ is open}$$

*Indirect proof.* Let  $A$  be closed and  $O := A^C$ . For any  $x \in O$ , we show  $x \in \mathring{O}$ .

Assume the opposite. With  $O^C = A$ ,

$$\begin{aligned} \forall \varepsilon > 0 : \neg (K_\varepsilon(x) \subseteq O) \\ \iff K_\varepsilon(x) \cap O^C \neq \emptyset \end{aligned}$$

**Direction  $\Leftarrow$ :** So  $x$  is contact point of  $A$ . Because  $A$  is closed,  $x \in A$ . This contradicts with  $x \in O = A^C$ . Thus  $O$  is open.

**Direction  $\Rightarrow$ :** Let  $O = A^C$  be open and let  $x$  be a contact point of  $A$ . Show that  $x \in A$ .

Assume the opposite, hence  $x \in A^C = O$  and  $O$  is open. So  $\exists r > 0 : K_r(x) \subseteq O$ , so  $K_r(x) \cap A = \emptyset$  where  $A = O^C$ . Hence  $x$  is not a contact point of  $A$ .

So every contact point of  $A$  is also an element of  $A$  and  $A$  is closed.

□

**Theorem 1.3.** *Let  $X$  be a metric space. Then,*

- *If  $O_i \subseteq X$  is open in  $X \forall i \in I$ , then also  $O = \bigcup_{i \in I} O_i$  is open in  $X$ .*
- *If  $O_1, O_2, \dots, O_n$  is open in  $X$ , then  $\bigcap_{k=1}^n O_k$  is open in  $X$ .*
- *$X$  is open and  $\emptyset$  is open.*

*Proof.* By Lemma 1.4, Theorem 1.2 and De Morgan's Laws:

$$\left( \bigcup_{i \in I} A_i \right)^C = \bigcap_{i \in I} A_i^C$$

□

## 1.8 Topology

**Definition 1.12.** *Given a set  $X$ . If a subset  $T \subseteq \mathcal{P}(X)$  is defined such that the elements  $O \in T$  (hence  $O \subseteq X$ ) satisfy the conditions of Theorem 1.3, then  $T$  is called topology on  $X$ .  $(X, T)$  is called topological space.*

*The sets  $O \in T$  are called open sets in terms of  $T$ . The complements  $A = O^C$  for  $O \in T$  are called closed sets.*

**Definition 1.13.** *Let  $x \in U \subseteq X$ . We claim that  $U$  is a neighborhood of  $x$ , if  $r > 0$  exists such that  $x \in K_r(X) \subseteq U$*

*See Figure 6*

**Remark 1.3.**  *$O \subseteq X$  is open iff  $O$  is neighborhood of every point  $x \in O$ .*

**Definition 1.14.** *Let  $X$  and  $Y$  be metric spaces and  $x_0 \in X$ . Let  $f : X \rightarrow Y$  be given. We say  $f$  is continuous in  $x_0$  if*

$$\forall \varepsilon > 0 \exists \delta > 0 : \forall x \in X : d_X(x, x_0) < \delta \implies d_Y(f(x), f(x_0)) < \varepsilon$$

*Here,  $d_X$  is a metric on  $X$  and  $d_Y$  is a metric on  $Y$ .*

↓ This lecture took place on 2018/03/13.




Figure 6: Neighborhood of  $x$

---

**Theorem 1.4.** Let  $X$  and  $Y$  be metric spaces.  $f : X \rightarrow Y$ . Let  $x_0 \in X$  be given. Then the following statements are equivalent:

1.  $f$  is continuous in  $x_0$
2. For every neighborhood  $V$  of  $y_0 = f(x_0)$  it is given that  $f^{-1}(V)$  is a neighborhood of  $x_0$
3. For every sequence  $(x_n)_{n \in \mathbb{N}}$  with  $\lim_{n \rightarrow \infty} x_n = x_0$  it holds that  $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$

*Proof.* See Analysis 1. □

**Definition 1.15.** Let  $f : X \rightarrow Y$  be called continuous on  $X$ , if  $f$  is continuous in every point  $x_0 \in X$ .

**Theorem 1.5.** Let  $f : X \rightarrow Y$  be given. Then  $f$  is continuous on  $X \iff \forall$  open  $O \subseteq Y : U = f^{-1}(O)$  open in  $X$ .

**Remark 1.4.** This characterization of continuity also works in topological spaces.

*Proof.* Direction  $\Rightarrow$ .

Let  $f$  be continuous in  $X$  and let  $O \subseteq Y$  be open. Let  $U = f^{-1}(O)$  and choose  $x_0 \in U$ . Then  $f(x_0) \in O$ , hence  $O$  is a neighborhood of  $f(x_0)$ . By Theorem 1.4 (b), it follows that  $U = f^{-1}(O)$  is a neighborhood of  $x_0$ .

Hence,  $U$  is neighborhood of every of its points, hence open in  $X$ .

Direction  $\Leftarrow$ .

Let the preimages of open sets be open and  $x_0 \in X$  and  $y_0 = f(x_0)$ . Let  $V$  be a neighborhood of  $y_0 = f(x_0)$ , hence  $\exists \varepsilon > 0 : K_\varepsilon(f(x_0)) \subseteq V$ . Because  $K_\varepsilon(f(x_0))$  is an open set,  $f^{-1}(K_\varepsilon(f(x_0))) \in x_0$  is open in  $X$ .

Therefore, there exists  $\delta > 0$  such that  $K_\delta(x_0) \subseteq f^{-1}(K_\varepsilon(f(x_0))) \subseteq f^{-1}(V)$ . Hence,  $f^{-1}(V)$  is a neighborhood of  $x_0$ . Then by Theorem 1.4 (b), it follows that  $f$  is continuous in  $x_0$  (chosen arbitrarily). Hence  $f$  is continuous on  $X$ .  $\square$

## 1.9 Variations of continuity notions

### 1.9.1 Uniform continuity

**Definition 1.16.** Let  $f : X \rightarrow Y$  be given. We call “ $f$  uniformly continuous on  $X$ ” if

$$\forall \varepsilon > 0 \exists \delta > 0 \forall x, y \in X : d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \varepsilon$$

**Remark 1.5.** Compare it with the definition of “continuous in  $X$ ”:

$$\forall x \in X \forall \varepsilon > 0 \exists \delta > 0 : \forall y \in X : d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \varepsilon$$

The difference is the location of the  $\forall x \in X$  quantifier.

Every uniformly continuous map is continuous.

Example:  $f : (0, \infty) \rightarrow (0, \infty)$  with  $f(x) = \frac{1}{x}$  is continuous, but not continuously continuous.

### 1.9.2 Lipschitz continuity

**Remark 1.6.** Rudolf Lipschitz [1832–1903], University of Bonn

**Definition 1.17.**  $f : X \rightarrow Y$  is called Lipschitz continuous with Lipschitz constant  $L \geq 0$  if  $\forall x, y \in X : d_Y(f(x), f(y)) \leq L \cdot d_X(x, y)$ .

**Theorem 1.6.** Every Lipschitz continuous function is uniformly continuous.

*Proof.* For  $\varepsilon > 0$ , choose  $\delta = \frac{\varepsilon}{L+1}$ . Then  $d_X(x, y) < \delta = \frac{\varepsilon}{L+1} \implies d_Y(f(x), f(y)) \leq L \cdot d_X(x, y) < \frac{L}{L+1} \cdot \varepsilon < \varepsilon$ .  $\square$

Most often  $X \subseteq V$ ,  $Y \subseteq W$ .  $V$  and  $W$  are normed vector spaces and  $d(x, y) = \|x - y\|$ .

## 1.10 Banach Fixed Point Theorem

**Definition 1.18.** A Lipschitz continuous map  $f : X \rightarrow X$  with Lipschitz constant  $L < 1$  is called contraction on  $X$ . Compare with Figure 7.




Figure 7: A contraction maps to points closer to each other




Figure 8: Banach Fixed Point Theorem states that applying  $f$  iteratively gives a point coming closer and closer to the previous one

**Theorem 1.7** (Banach fixed-point theorem). *Let  $f : X \rightarrow X$  be a contraction and  $X$  be complete. Then there exists a uniquely defined  $\hat{x} \in X$  such that  $\hat{x} = f(\hat{x})$ .  $\hat{x}$  is called fixed point on  $f$ . Furthermore  $x_0 \in X$  is arbitrary and  $x_n = f(x_{n-1})$  for all  $n \geq 1$ . Compare with Figure 8.*

$$\lim_{n \rightarrow \infty} x_n = \hat{x}$$

**Remark 1.7.** *The following proof is a very common exam question.*

*Proof.* Let  $x_0 \in X$  be arbitrary.  $x_n$  is constructed inductively by  $x_n = f(x_{n-1})$  for all  $n \geq 1$ .



**Prove that  $(x_n)_{n \in \mathbb{N}}$  is a Cauchy sequence in  $X$**

$$\begin{aligned}
 d(x_n, x_{n+k}) &\leq d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + \cdots + d(x_{n+k-1}, x_{n+k}) \quad \text{by triangle ineq.} \\
 &= d(x_n, x_{n+1}) + d(f(x_n), f(x_{n+1})) + \cdots + d(f(x_{n+k-2}), f(x_{n+k-1})) \\
 &\leq d(x_n, x_{n+1}) + L(d(x_n, x_{n+1}) + d(x_{n+1}, x_{n+2}) + \cdots + d(x_{n+k-2}, x_{n+k-1}))
 \end{aligned}$$

this inequality is given by contraction

$$\begin{aligned}
 &= d(x_n, x_{n+1})(1 + L) + L(d(f(x_n), f(x_{n+1})) + \cdots + d(f(x_{n+k-3}), f(x_{n+k-2}))) \\
 &\leq d(x_n, x_{n+1})(1 + L) + L^2[d(x_n, x_{n+1}) + \cdots + d(x_{n+k-3}, x_{n+k-2})] \\
 &\leq \cdots \leq d(x_n, x_{n+1})(1 + L + L^2 + \cdots + L^{k-1}) \\
 &= d(f(x_{n-1}), f(x_n)) \left( \sum_{j=0}^{k-1} L^j \right) \leq L d(x_{n-1}, x_n) \cdot \left( \sum_{j=0}^{k-1} L^j \right) \\
 &\leq L^n d(x_0, x_1) \cdot \underbrace{\left( \sum_{j=1}^{k-1} L^j \right)}_{\leq \sum_{j=0}^{\infty} L^j = \frac{1}{1-L}} \\
 &\leq \frac{L^n}{1-L} d(x_0, x_1) \\
 d(x_n, x_{n+k}) &\leq \frac{L^n}{1-L} d(x_0, x_1) \quad \forall n \in \mathbb{N} \forall k \in \mathbb{N}_0 \text{ with } 0 \leq L < 1
 \end{aligned}$$

$$\begin{aligned}
 &\underbrace{\frac{L^n}{1-L} d(x_0, x_1)}_{>0} < \varepsilon \iff \\
 L^n &< \frac{\varepsilon}{d(x_0, x_1) + 1} (1 - L) \quad (L > 0) \\
 \iff n \underbrace{\ln L}_{<0} &< \ln \frac{\varepsilon}{d(x_0, x_1) + 1} (1 - L) \\
 \iff n &> \frac{1}{\ln L} \ln \frac{\varepsilon}{d(x_0, x_1) + 1} (1 - L)
 \end{aligned}$$

Hence  $(x_n)_{n \in \mathbb{N}}$  is a Cauchy sequence in  $X$ .  $X$  is complete, hence  $\exists \hat{x} \in X$  :  $\hat{x} = \lim_{n \rightarrow \infty} x_n$ . Because  $\hat{x} = \lim_{n \rightarrow \infty} x_{n+1} = \lim_{n \rightarrow \infty} f(x_n) = f(\hat{x})$  where the last equality is given by continuity of  $f$ . Therefore  $\hat{x} = f(\hat{x})$  is a fixed point on  $f$ .

**Prove uniqueness of point  $\hat{x}$**  Let  $\tilde{x} = f(\tilde{x})$ . Then  $d(\hat{x}, \tilde{x}) = d(f(\hat{x}), f(\tilde{x})) \leq L d(\hat{x}, \tilde{x})$  with  $L < 1$ . If  $d(\hat{x}, \tilde{x}) > 0$ , then  $1 \leq L$ . This is a contradiction. Hence  $d(\hat{x}, \tilde{x}) = 0$  must hold, hence  $\hat{x} = \tilde{x}$ .

□

**Remark 1.8.** • The Fixed Point Theorem provides an algorithm for numeric computation of  $\hat{x}$ .

- It can reformulate problems  $f(x) = 0$  (in  $\mathbb{R}^n$ ) to

$$f(x) + x = g(x) = x$$

- Attention: The conditions of the Fixed Point Theorem cannot be changed to the structure

$$d(f(x), f(y)) < L \cdot d(x, y) \wedge L \leq 1$$

or

$$d(f(x), f(y)) \leq L \cdot d(x, y) \wedge L < 1$$

This will be discussed in the practicals.

**Lemma 1.7.** Let  $X$  be a complete metric space. Let  $A \subseteq X$  be closed. Then  $(A, d)$  is itself a complete, metric space.

*Proof.* Let  $(x_n)_{n \in \mathbb{N}}$  be a Cauchy sequence in  $A$  ( $x_n \in A$ ). Then  $(x_n)_{n \in \mathbb{N}}$  is also a Cauchy sequence in  $X$ . Because  $X$  is complete, there exists  $\hat{x} = \lim_{n \rightarrow \infty} x_n$ . Therefore  $\hat{x}$  is a contact point of  $A$ . Because  $A$  is closed,  $\hat{x} \in A$ .

Therefore every Cauchy sequence in  $A$  has a limit point in  $A$ , hence  $A$  is complete. □

## 2 Compactness

### 2.1 Definition

**Definition 2.1.** A metric space  $(X, d)$  is called compact if every sequence  $(x_n)_{n \in \mathbb{N}}$  has a convergent subsequence.

Specifically, this definition is called sequence compactness. The other definition defines compactness as closed and bounded subset of an Euclidean space. The latter definition only works for spaces with the Heine-Borel property (counterexample: any incomplete metric space like the space of rational numbers is closed and bounded but not compact). Therefore the generalization is recommended to be remembered.

**Lemma 2.1.** Let  $X$  be a compact, metric space. Then  $X$  is complete.

*Proof.* Let  $(x_n)_{n \in \mathbb{N}}$  be a Cauchy sequence in  $X$ . By compactness, it follows that  $\exists (x_{n_k})_{k \in \mathbb{N}}$  with  $\lim_{k \rightarrow \infty} x_{n_k} = \hat{x}$ . Choose  $\varepsilon > 0$  arbitrary and  $L$  large enough such that  $k \geq L \implies d(x_{n_k}, \hat{x}) < \frac{\varepsilon}{2}$ . Furthermore choose  $N \in \mathbb{N}$  large enough such that  $n, m \geq N \implies d(x_n, x_m) < \frac{\varepsilon}{2}$  (satisfied, because  $(x_n)_{n \in \mathbb{N}}$  is a Cauchy sequence). Choose  $k \geq L$  and  $n_k \geq N$ . Let  $n_k$  be fixed this way. Then it holds

$\forall n \geq N : d(x_n, \hat{x}) \leq d(x_n, x_{n_k}) + d(x_{n_k}, \hat{x}) < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$ . The first summand  $\frac{\varepsilon}{2}$  results from the Cauchy sequence property, the second summand  $\frac{\varepsilon}{2}$  results by convergence of  $(x_{n_k})$ . Hence  $(x_n)_{n \in \mathbb{N}}$  is convergent with limit  $\hat{x}$ .  $\square$

## 2.2 Boundedness

**Definition 2.2.** A metric space  $X$  is called bounded if there exists  $M \geq 0$ , such that  $d(x, y) \leq M \forall x, y \in X$ .

It holds for arbitrary  $x \in X$  that  $\forall y \in X : y \in K_M(x)$ . So,  $X \subseteq K_M(x)$ . On the contrary, let  $X \subseteq \overline{K_M(x)}$  and let  $y \in X$  and  $z \in X$  be arbitrary. Then  $d(y, z) \leq d(y, x) + d(x, z) \leq M + M = 2M$ . Hence,  $X$  is bounded.

So,  $X$  is bounded  $\iff \exists x \in X \wedge M \geq 0 : X \subseteq \overline{K_M(x)}$ .

**Lemma 2.2.** Every compact, metric space is also bounded.

*Proof.* Assume  $X$  is unbounded.

We construct a sequence of points  $(x_n)_{n \in \mathbb{N}}$  with  $d(x_n, x_m) \geq 1 \forall n, m \in \mathbb{N}$  with  $n \neq m$ .

We use the following auxiliary result: Let  $B = \bigcup_{j=1}^n K_1(z_j)$  for arbitrary  $n \in \mathbb{N}$  and arbitrary  $z_j \in X$ . Then  $B$  is bounded. This result will be part of the practicals.

We construct  $(x_n)_{n \in \mathbb{N}}$  inductively. Choose arbitrary  $x_0 \in X$ . Assume  $(x_1, \dots, x_{n-1})$  are already found. Then,

$$\underbrace{X}_{\text{unbounded}} \not\subseteq \underbrace{\bigcup_{j=1}^{n-1} K_1(x_j)}_{\text{bounded}}$$

hence  $\exists x_n \in X \setminus \bigcup_{j=1}^{n-1} K_1(x_j)$ . Because  $x_n \notin K_1(x_j)$  for  $j = 0, \dots, n-1$  it is given that  $d(x_n, x_j) \geq 1 \forall j < n$ . We get  $(x_n)_{n \in \mathbb{N}}$  with  $d(x_n, x_m) \geq 1 \forall n \in \mathbb{N} \forall m < n$ , hence  $m \neq n$ . Because  $d(x_n, x_m) \geq 1$ , i.e.  $(x_n)_{n \in \mathbb{N}}$  does not contain any Cauchy sequence as subsequence,  $(x_n)_{n \in \mathbb{N}}$  does not have a convergent subsequence. Therefore  $X$  is not compact.  $\square$

↓ This lecture took place on 2018/03/15.

Every compact metric space is bounded. Every compact metric space is complete. In  $\mathbb{C}(\mathbb{R}^n)$ , any  $A \subseteq \mathbb{C}$  is closed. Then  $A$  with metric  $d(x, y) = |x - y|$  is complete as metric space.

If  $A$  is additionally bounded, then  $A$  is compact (see course Analysis 1, Bolzano-Weierstrass).

*Attention!* Let  $V$  be an infinite-dimensional, complete, normed vector space.

- For example,  $V = C([a, b], \mathbb{R}) = \{f : [a, b] \rightarrow \mathbb{R} \mid f \text{ is continuous in } [a, b]\}$
- with norm  $\|f\|_\infty = \max\{|f(x)| : x \in [a, b]\}$
- and metric  $\|f - g\|_\infty = \max\{|f(x) - g(x)| : x \in [a, b]\}$

$C([a, b], \mathbb{R})$  is a complete, normed vector space. So,  $\overline{K_1(0)}$  is not compact in  $C([a, b], \mathbb{R})$  (i.e.  $V$ , for every infinite-dimensional vector space).

*Again:* do not remember “compactness” as closed and bounded, because this definition only holds in the finite-dimensional case.

In the last proof, we have shown: If a sequence  $(x_n)_{n \in \mathbb{N}}$  with  $x_n \in X$  and  $d(x_n, x_m) \geq 1$  (or  $\geq \varepsilon$ )  $\forall n \neq m \implies X$  is not compact.

## 2.3 Total boundedness

**Definition 2.3.**  $X$  is called totally bounded, if for every  $\varepsilon > 0$ , finitely many points  $X_1^\varepsilon, X_2^\varepsilon, \dots, X_{N(\varepsilon)}^\varepsilon$  such that  $X \subseteq \bigcup_{i=1}^{N(\varepsilon)} K_\varepsilon(X_i^\varepsilon)$ .

Hence, for every  $x \in X$ , there exists some  $X_j^\varepsilon$  such that  $d(x, X_j^\varepsilon) < \varepsilon$ .

**Remark 2.1** (For the practicals). Let  $X$  be totally bounded, then there does not exist some sequence  $(x_n)_{n \in \mathbb{N}}$  with  $d(x_n, x_m) \geq \varepsilon \forall n \neq m$ . It holds, that  $X$  is compact if and only if  $X$  is totally bounded and complete.

## 2.4 Compactness, continuity and openness

**Theorem 2.1.** Let  $f : X \rightarrow Y$  be continuous. Let  $X$  be compact. Then image  $f(X) \subseteq Y$  is also compact.

Be aware, that this proof is a common exam question and students often begin with the wrong order.

*Proof.* Let  $(y_n)_{n \in \mathbb{N}}$  be an arbitrary sequence in  $f(X)$ . Show that  $(y_n)_{n \in \mathbb{N}}$  has a convergent subsequence. Because  $y_n \in f(X)$ , there exists at least one  $x_n$  with  $y_n = f(x_n)$ . Then  $(x_n)_{n \in \mathbb{N}}$  is a sequence in  $X$ .  $X$  is compact, hence there exists a subsequence  $(x_{n_k})_{k \in \mathbb{N}}$  with  $\lim_{k \rightarrow \infty} x_{n_k} = \hat{x} \in X$ . Because  $f$  is continuous,  $\lim_{k \rightarrow \infty} f(x_{n_k}) = \lim_{k \rightarrow \infty} y_{n_k} = f(\hat{x}) =: \hat{y}$ . So  $(y_n)_{n \in \mathbb{N}}$  has a convergent subsequence. Hence  $f(X) \subseteq Y$  is compact.  $\square$

**Theorem 2.2** (Conclusion). *Let  $X$  be compact,  $f : X \rightarrow \mathbb{R}$  continuous on  $X$ . Then there exists  $\underline{x}$  and  $\bar{x} \in X$ , such that*

$$f(\underline{x}) \leq f(x) \leq f(\bar{x}) \quad \forall x \in X$$

*Hence,  $f$  has a maximum and a minimum.*

*Proof.*  $f(X) \subseteq \mathbb{R}$  is compact (Theorem 2.1), hence  $f(X)$  is bounded and complete, hence closed in  $\mathbb{R}$ . There exists  $\xi \in \mathbb{R}$  with  $\xi = \sup f(X)$ , because  $f(X)$  is complete and  $\xi$  is a contact point of  $f(X)$ , so  $\xi \in f(X)$ , hence  $\exists \bar{x} \in X : \xi = f(\bar{x})$ . Furthermore,  $\xi$  is an upper bound of  $f(X) \rightarrow f(x) \leq \xi = f(\bar{x}) \forall x \in X$ .

For  $\underline{x}$ , it works the same way.  $\square$

**Theorem 2.3.** *Let  $f : X \rightarrow Y$  be continuous on  $X$  and  $X$  is compact. Then  $f$  is uniformly continuous on  $X$ .*

*Indirect proof.* Assume  $X$  is compact,  $f : X \rightarrow Y$  is continuous, but not uniformly continuous. Uniform continuity:

$$\forall \varepsilon > 0 \exists \delta > 0 : \forall x, y \in X : d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \varepsilon$$

Not uniformly continuous:

$$\exists \varepsilon > 0 \forall \delta_n = \frac{1}{n} (n \in \mathbb{N}) \exists x_n, y_n \in X : d_X(x_n, y_n) < \frac{1}{n} \wedge d_Y(f(x_n), f(y_n)) \geq \varepsilon$$

Now choose some  $(x_n)$  and  $(y_n)$ . We will use a specific  $\varepsilon$  later. Because  $X$  is compact, there exists a convergent subsequence of  $(x_n)_{n \in \mathbb{N}}$ , hence  $\lim_{k \rightarrow \infty} x_{n_k} = \hat{x}$ . The sequence  $(y_{n_k})_{k \in \mathbb{N}}$  has a convergent subsequence itself:

$$\lim_{l \rightarrow \infty} y_{(n_k)_l} = \hat{y}$$

Because  $(x_{n_k})_{k \in \mathbb{N}}$  is convergent, the subsequence  $(x_{(n_k)_l})_{l \in \mathbb{N}}$  converges towards the same limit  $\hat{x}$ .

$$\tilde{x}_l := x_{n_{k_l}} \quad \tilde{y}_l := y_{n_{k_l}}$$

because  $l \leq n_{k_l}$  and

$$d_X(\tilde{x}_l, \tilde{y}_l) = d_X(x_{n_{k_l}}, y_{n_{k_l}}) \underbrace{\leq}_{\text{by assumption}} \frac{1}{n_{k_l}} \leq \frac{1}{l}$$

**Claim.** *For  $\hat{x} = \lim_{l \rightarrow \infty} \tilde{x}_l$  and  $\hat{y} = \lim_{l \rightarrow \infty} \tilde{y}_l$ , it holds that  $\hat{x} = \hat{y}$ .*

*Proof.* Let  $\varepsilon' > 0$  be arbitrary,  $l$  large enough such that

- $\frac{1}{l} < \frac{\varepsilon'}{3}$
- $d_X(\tilde{x}_l, \hat{x}) < \frac{\varepsilon'}{3}$

- $d_X(\tilde{y}_l, \hat{y}) < \frac{\varepsilon'}{3}$

Therefore

$$d_X(\hat{x}, \hat{y}) \leq d_X(\hat{x}, \tilde{x}_l) + d_X(\tilde{x}_l, \tilde{y}_l) + d_X(\tilde{y}_l, \hat{y}) < \frac{\varepsilon'}{3} + \frac{1}{l} + \frac{\varepsilon'}{3} < \varepsilon'$$

Therefore  $d_X(\hat{x}, \hat{y}) = 0$ , hence  $\hat{x} = \hat{y}$ .  $\square$

Because  $f$  is continuous and  $\tilde{x}_l \rightarrow \hat{x}$  and  $\tilde{y}_l \rightarrow \hat{x}$ , there exists  $l \in \mathbb{N}$  such that

$$d_Y(f(\tilde{x}_l), f(\hat{x})) < \frac{\varepsilon}{2}$$

and also

$$d_Y(f(\tilde{y}_l), f(\hat{x})) < \frac{\varepsilon}{2}$$

where  $\varepsilon$  is the epsilon from the very beginning of the proof.

$$\implies d_Y(f(\tilde{x}_l), f(\hat{x})) + d_Y(f(\tilde{y}_l), f(\hat{x})) < \varepsilon$$

This contradicts to

$$d_Y(f(\tilde{x}_l), f(\tilde{y}_l)) = d_Y(f(x_{n_{k_l}}), f(y_{n_{k_l}})) \geq \varepsilon$$

Hence,  $f$  is uniformly continuous.  $\square$

Subsets of  $(\mathbb{R}^n, \|\cdot\|)$  (or  $(V, \|\cdot\|)$ ) as metric spaces.

We consider  $\Omega \subseteq V$  where  $V$  is a normed vector space.  $(\Omega, d)$  is  $d(x, y) = \|x - y\|$  is a metric space.

$$K_r^\Omega(x) = \{y \in \Omega \mid \|y - x\| < r\}$$

is a sphere with center  $x$  and radius  $r$  in  $\Omega$ .

$$K_r^V(x) = \{y \in V \mid \|y - x\| < r\}$$

obvious:  $K_r^\Omega(x) = \Omega \cap K_r^V(x)$ .

**Lemma 2.3.** Let  $O' \subseteq \Omega \subseteq V$ .

Then  $O'$  is open in  $\Omega \iff$  there exists  $O \subseteq V$  is open in  $V$  such that  $O' = O \cap \Omega$ .

*Proof.*  $\Rightarrow$  Let  $O' \subseteq \Omega$  be open in  $\Omega$  and  $x \in O'$  be arbitrary. Then there exists  $r(x) > 0 : x \in K_{r(x)}^\Omega(x) = K_{r(x)}^V(x) \cap \Omega \subseteq O'$ . Then

$$O' = \bigcup_{x \in O'} \{x\} \subseteq \bigcup_{x \in O'} K_{r(x)}^\Omega(x) = \left( \bigcup_{x \in O'} (K_{r(x)}^V(x) \cap \Omega) \right) = \underbrace{\left( \bigcup_{x \in O'} K_{r(x)}^V(x) \right)}_{=O \subseteq V \text{ is open in } V} \cap \Omega \subseteq O'$$

So every  $\subseteq$  in this inclusion chain is actually an equality. So  $O' = O \cap \Omega$ .




Figure 9: Subsets of  $(\mathbb{R}^n, \|\cdot\|)$  as metric spaces

$\Leftarrow$  Let  $O' = O \cap \Omega$  and  $x \in O'$  be chosen arbitrarily. Because  $x \in O$  and  $O$  is open in  $V$ .

$$\exists r > 0 : K_r^V(x) \subseteq O \implies \underbrace{K_r^V(x) \cap \Omega}_{=K_r^\Omega(x)} \subseteq O \cap \Omega = O'$$

So  $O'$  is open in  $\Omega$ .

□

**Remark 2.2.**  $A' \subseteq \Omega$  is closed in  $\Omega \iff \exists A \subseteq V$  closed in  $V$  with  $A' = A \cap \Omega$ .

**Remark 2.3.** Let  $T$  be an arbitrary topological space with topology  $\tau$  on  $T$  (a system of open sets). Furthermore let  $\Omega \subseteq T$ .

Then  $\Omega$  itself is a topological space with  $O' \subseteq \Omega$  is open  $\iff \exists O \subset T$  open in  $T$  with  $O' = O \cap \Omega$ .

Also called “subspace topology”, “trace topology” or “relative topology”.

Attention!

$$O' \subseteq \Omega \text{ open in } \Omega \implies O' \text{ open in } V$$

does *not* hold in general.

**Example 2.1.**

$$\Omega = [0, 1] \cap [0, 1)$$

$K_{\frac{1}{2}}(p) \cap \Omega$  is open in  $\Omega$  but not open in  $\mathbb{R}^2$ .

Analogously,

$$A' \subseteq \Omega \text{ is closed} \implies A' \text{ closed in } V$$

does *not* hold in general.

**Remark 2.4.**  $K \text{ is compact in } \Omega \implies K \text{ is compact in } V$

Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence in  $K$ . Compactness  $\implies \exists (x_{n_k})_{k \in \mathbb{N}} : x_{n_k} \rightarrow \hat{x}$  for  $k \rightarrow \infty$  and  $K \subseteq \Omega \subseteq V$ .

Then  $(x_n)_{n \in \mathbb{N}}$  also has a convergent subsequence in  $V$ .

## 2.5 Normed vector spaces

**Definition 2.4.** Let  $V$  be a vector space and  $\|\cdot\|_1$  and  $\|\cdot\|_2$  are normed on  $V$ . We say,  $\|\cdot\|_1$  is equivalent to norm  $\|\cdot\|_2$ , if  $0 < m \leq M$  exist such that

$$m \|v\|_1 \leq \|v\|_2 \leq M \|v\|_1 \quad \forall v \in V$$

**Remark 2.5.** Equivalence of norms is an equivalence relation.

**reflexivity**  $\|\cdot\|_1$  is equivalent to  $\|\cdot\|_1$  with  $m = M = 1$ .

**symmetry**

$$\begin{aligned} m \|v\|_1 \leq \|v\|_2 &\implies \|v\|_1 \leq \frac{1}{m} \|v\|_2 \wedge \|v\|_2 \leq M \cdot \|v\|_1 \implies \frac{1}{M} \|v\|_2 \leq \|v\|_1 \\ &\implies \underbrace{\frac{1}{M}}_{m'} \|v\|_2 \leq \|v\|_1 \leq \underbrace{\frac{1}{m}}_{M'} \|v\|_2 \end{aligned}$$

hence the equivalence relations of norms are symmetrical.

**transitivity** Let  $\|\cdot\|_1$  and  $\|\cdot\|_2$  be equivalent. Let  $\|\cdot\|_2$  and  $\|\cdot\|_3$  be equivalent.

$$\begin{aligned} m \cdot \|v\|_1 &\leq \|v\|_2 \leq M \|v\|_1 \quad \forall v \in V \\ m' \cdot \|v\|_2 &\leq \|v\|_3 \leq M' \|v\|_2 \quad \forall v \in V \\ \implies m \cdot m' \|v\|_1 &\leq m' \|v\|_2 \leq \|v\|_3 \leq M' \|v\|_2 \leq M \cdot M' \|v\|_1 \end{aligned}$$

↓ This lecture took place on 2018/03/20.

Addendum:



- Let  $(x_n)_{n \in \mathbb{N}}$  be in  $(X, d)$ , then

$$\underbrace{x = \lim_{n \rightarrow \infty} x_n}_{\text{in } X} \iff \underbrace{\lim_{n \rightarrow \infty} d(x_n, x) = 0}_{\text{in } \mathbb{R}}$$

$$(\iff \lim_{n \rightarrow \infty} \|x_n - x\| = 0 \text{ in normed vector spaces } V)$$

- Reversed triangle inequality: Let  $V$  be a normed vector space. Let  $x, y \in V$ .

$$\|x\| = \|x - y + y\| \leq \|x - y\| + \|y\|$$

Hence,

$$\|x\| - \|y\| \leq \|x - y\|$$

By exchanging  $x$  and  $y$ ,

$$\|y\| - \|x\| \leq \|x - y\|$$

$$\implies \left| \|x\| - \|y\| \right| \leq \|x - y\|$$

- Define the map  $n : V \rightarrow [0, \infty)$  on  $(V, \|\cdot\|)$  with  $n(x) = \|x\|$ . Then  $n$  is continuous on  $V$  because

$$|n(x_1) - n(x_2)| = \left| \|x_1\| - \|x_2\| \right| \leq \|x_1 - x_2\|$$

Hence,  $n$  is Lipschitz continuous with constant 1.

Regarding the equivalence of norms:

**Lemma 2.4.** Let  $\|\cdot\|_1$  and  $\|\cdot\|_2$  be equivalent norms on  $V$ . Then,

1.  $\lim_{n \rightarrow \infty} \|x_n - x\|_1 = 0 \iff \lim_{n \rightarrow \infty} \|x_n - x\|_2 = 0$ , hence  $(x_n)_{n \in \mathbb{N}}$  is convergent with limit  $x$  with respect to  $\|\cdot\|_1 \iff (x_n)_{n \in \mathbb{N}}$  is convergent with limit  $x$  with respect to  $\|\cdot\|_2$ .
2.  $O \subseteq V$  is open with respect to  $\|\cdot\|_1 \iff O$  is open with respect to  $\|\cdot\|_2$ , hence  $\tau_1 = \tau_2$  (topologies are equivalent).
3.  $K \subseteq V$  is compact with respect to  $\|\cdot\|_1 \iff K$  is compact with respect to  $\|\cdot\|_2$ .

*Proof.* Let  $\|\cdot\|_1$  and  $\|\cdot\|_2$  are equivalent, hence  $\exists m, M > 0 : m\|x\|_1 \leq \|x\|_2 \leq M\|x\|_1 \forall x \in V$ .

1. Let  $\varepsilon > 0$  and  $\lim_{n \rightarrow \infty} \|x_n - x\|_1 = 0$ . Choose  $N \in \mathbb{N}$  such that  $n \geq N \implies \|x_n - x\|_1 < \frac{\varepsilon}{M}$ . For those  $n$ ,

$$\|x_n - x\|_2 \leq M\|x_n - x\|_1 < \frac{\varepsilon}{M} \cdot M = \varepsilon$$

Hence,  $\lim_{n \rightarrow \infty} \|x_n - x\|_2 = 0$ .

2.  $K_r^2(x) = \{y \in V \mid \|y - x\|_2 < r\}$ . For  $y \in K_r^2(x)$ ,

$$m \|y - x\|_1 \leq \|y - x\|_2 < r$$

hence,

$$\|y - x\|_1 < \frac{r}{m} \implies y \in K_{\frac{r}{m}}^1(x)$$

hence  $K_r^2(x) \subseteq K_{\frac{r}{m}}^1(x)$ . Let  $y \in K_{\frac{r}{m}}^1(x)$ . Then,

$$\|y - x\|_2 \leq M \|y - x\|_1 < M \cdot \frac{r}{M} = r$$

hence  $y \in K_r^2(x) \implies K_{\frac{r}{M}}^1(x) \subseteq K_r^2(x)$ . Now let  $O$  be open with respect to  $\|\cdot\|_2$ , hence

$$\forall x \in O \exists r > 0 : K_r^2(x) \subseteq O \implies K_{\frac{r}{M}}^1(x) \subseteq K_r^2(x) \subseteq O$$

so  $O$  is open with respect to  $\|\cdot\|_1 \implies O$  is open with respect to  $\|\cdot\|_2$  analogously.

3. Let  $K$  be compact with respect to  $\|\cdot\|_1$  and  $(x_n)_{n \in \mathbb{N}}$  be a sequence in  $K$ . Then there exists a subsequence  $(x_{n_k})_{k \in \mathbb{N}}$  with  $\|x_{n_k} - x\|_1 \rightarrow 0$  for  $k \rightarrow \infty$   $\xrightarrow{\text{by the first property}}$   $\|x_{n_k} - x\|_2 \rightarrow 0$ . Hence  $(x_{n_k})_{k \in \mathbb{N}}$  is also a convergent subsequence with respect to  $\|\cdot\|_2$ .

□

**Remark 2.6** (Proven in the practicals). Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathbb{R}^k$

$$\|x\|_\infty = \max \{ |x^i| \mid i = 1, \dots, k \}$$

$$x = \begin{bmatrix} x^1 \\ x^2 \\ \vdots \\ x^k \end{bmatrix}$$

Then  $\lim_{n \rightarrow \infty} \|x_n - x\|_\infty = 0 \iff \lim_{n \rightarrow \infty} |x_n^i - x^i| = 0$  for all  $i \in \{1, \dots, k\}$ .

**Theorem 2.4** (Bolzano-Weierstrass theorem in  $\mathbb{R}^k$ ). Let  $K \subseteq \mathbb{R}^k$  be closed and bounded. Then  $K$  is compact in  $(\mathbb{R}^k, \|\cdot\|_\infty)$ .

*Proof.* Let  $\|x\|_\infty \leq M \forall x \in K \iff |x^i| \leq M \forall x \in K$  and  $i \in \{1, \dots, k\}$ . Choose  $(x_n)_{n \in \mathbb{N}}$  an arbitrary sequence in  $K$ .  $(x_n^i)_{n \in \mathbb{N}}$  is a bounded sequence in  $\mathbb{R}$ . Because  $(x_n^1)_{n \in \mathbb{N}}$  is bounded, there exists a convergent subsequence  $(x_{n_{l_1}}^1)_{l_1 \in \mathbb{N}}$

$$\lim_{l_1 \rightarrow \infty} x_{n_{l_1}}^1 = x^1$$

Consider  $(x_{n_{l_1}}^2)_{l_1 \in \mathbb{N}}$ , a subsequence of a bounded sequence, hence bounded itself. By the Bolzano-Weierstrass theorem in  $\mathbb{R}$ , there exists a convergent subsequence  $(x_{n_{l_1 l_2}}^2)_{l_2 \in \mathbb{N}}$  with  $\lim_{l_2 \rightarrow \infty} x_{n_{l_1 l_2}}^2 = x^2$ . Consider  $x_{n_{l_1 l_2}}^1$  as subsequence of  $x_{n_{l_1}}^1$  is already convergent, hence  $\lim_{l_2 \rightarrow \infty} x_{n_{l_1 l_2}}^1 = x^1$ . Furthermore, for indices up to  $i$ :

$$\lim_{l_k \rightarrow \infty} x_{n_{l_1 l_2 \dots l_k}} = x^i \quad \text{for } i = 1, \dots, k$$

Hence, with  $\tilde{x}_{l_k} = x_{n_{l_1 l_2 \dots l_k}}$  gives a subsequence of  $x_n$ , converging by each coordinate. Thus,

$$\lim_{l_k \rightarrow \infty} \|\tilde{x}_{l_k} - x\|_\infty = 0$$

Because  $\tilde{x}_{l_k} \in K$  and  $K$  is closed,  $x \in K$ . Hence  $K$  is compact.  $\square$

**Theorem 2.5** (Norm equivalence in  $\mathbb{R}^k$ ). *In  $\mathbb{R}^k$ , all norms are equivalent.*

*Proof.* We show: Let  $\|\cdot\|$  be an arbitrary norm on  $\mathbb{R}^n$ . Then  $\|\cdot\|$  is equivalent to  $\|\cdot\|_\infty$ . By transitivity of norm equivalence, two arbitrary norms are equivalent to each other.

1. Let  $(e_1, e_2, \dots, e_k)$  be the canonical basis in  $\mathbb{R}^k$ .

$$x = \begin{bmatrix} x^1 \\ \vdots \\ x^k \end{bmatrix} = \sum_{j=1}^k x^j e_j$$

Furthermore let  $M' = \max \{\|e_j\| : j = 1, \dots, k\}$  with  $\|e_j\| \neq 0$  and  $M' > 0$ . Then

$$\begin{aligned} \|x\| &= \left\| \sum_{j=1}^k x^j e_j \right\| \leq \sum_{j=1}^k \|x^j e_j\| = \sum_{j=1}^k |x^j| \|e_j\| \\ &\leq M' \sum_{j=1}^k \underbrace{|x^j|}_{\leq \|x\|_\infty} \leq \underbrace{M' \cdot k}_M \|x\|_\infty = M \|x\|_\infty \end{aligned}$$

2. We consider  $\gamma : \mathbb{R}^k \rightarrow [0, \infty)$ .  $\gamma(x) = \|x\|$  as map on  $(\mathbb{R}^k, \|\cdot\|_\infty)$ .

**Claim.**  $\gamma$  is continuous on  $(\mathbb{R}^k, \|\cdot\|_\infty)$ .

*Proof.* Show that,

$$|\gamma(x) - \gamma(y)| = \left| \|x\| - \|y\| \right| \underbrace{\leq}_{\text{reversed triangle ineq.}} \|x - y\| \underbrace{\leq}_{\text{because of (1)}} M \|x - y\|$$

Hence  $\gamma$  is Lipschitz continuous.  $\square$



Figure 10: Illustration of a partition

We consider  $S_{\infty}^{k-1} = \{x \in \mathbb{R}^k \mid \|x\|_{\infty} = 1\} = \text{boundary}(K_1^{\infty}(0))$ .  $S_{\infty}^{k-1}$  is bounded.

Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence in  $S_{\infty}^{k-1}$  with  $x = \lim_{n \rightarrow \infty} x_n$ . Because  $n(x) = \|x\|_{\infty}$  is continuous,

$$\lim_{n \rightarrow \infty} \underbrace{\|x_n\|_{\infty}}_{=1} = \underbrace{\|x\|}_{=1}$$

Hence  $x \in S_{\infty}^{k-1}$ . Hence,  $S_{\infty}^{k-1}$  is closed in  $(\mathbb{R}^k, \|\cdot\|_{\infty})$ . Hence  $S_{\infty}^{k-1}$  is compact in  $(\mathbb{R}^k, \|\cdot\|_{\infty})$ ,  $\gamma : S_{\infty}^{k-1} \rightarrow [0, \infty)$ , with  $S_{\infty}^{k-1}$  compact, is continuous. Has

a minimum  $m$  on  $S_{\infty}^{k-1}$ . Thus there exists  $\bar{x} \in S_{\infty}^{k-1} : \underbrace{m}_{>0} = \underbrace{\left\| \bar{x} \right\|}_{\neq 0} \leq \|x\| \forall x \in S_{\infty}^{k-1}$ . Let  $x \in \mathbb{R}^k$  be arbitrary with  $x \neq 0$ . Then,  $\frac{x}{\|x\|_{\infty}} \in S_{\infty}^{k-1}$  and

$$m \leq \left\| \frac{x}{\|x\|_{\infty}} \right\| = \frac{1}{\|x\|_{\infty}} \|x\| \implies m \|x\|_{\infty} \leq \|x\|$$

Inequality also holds true for  $x = 0$ .

□

### 3 Integration calculus

#### 3.1 Partitions and refinements

**Definition 3.1.** Let  $a < b$  with  $a, b \in \mathbb{R}$ . We consider functions of  $[a, b]$ . We call  $(x_j)_{j=0}^n$  a partition of  $[a, b]$  if  $a = x_0 < x_1 < x_2 < \dots < x_n = b$ .  $x_j$  decomposes  $[a, b]$  in subintervals  $(x_{j-1}, x_j)$ .  $\varphi : [a, b] \rightarrow \mathbb{R}$  is called step function in  $[a, b]$  with respect to partition  $(x_j)_{j=0}^n$  if  $\varphi|_{(x_{j-1}, x_j)} = c_j$ , so constant for  $j = 1, \dots, n$ .

$\varphi$  is called step function in  $[a, b]$  if there exists a partition such that  $\varphi$  is a subsequence.

$$\tau[a, b] := \left\{ \varphi : [a, b] \rightarrow \mathbb{R} \mid \varphi \text{ is subsequence} \right\}$$

Let  $(\xi_i)_{i=0}^m$  be a partition of  $[a, b]$  and  $(x_j)_{j=0}^n$  as well. Then we call  $(\xi_i)_{i=0}^m$  a refinement of  $(x_j)_{j=0}^n$ . Thus  $(\xi_i)_{i=0}^m$  is a refinement of  $(x_j)_{j=0}^n$  if  $\{x_0, x_1, \dots, x_n\} \subseteq \{\xi_0, \xi_1, \dots, \xi_m\}$ .




Figure 11: (top) Refinement (middle) Function values in points  $x_j$  are unrestricted (bottom) Step functions on a refinement

Compare with Figure 11. Functions values in boundaries  $x_{j-1}$  and  $x_j$  do not have any constraints and this will be relevant for integrals. Some  $\phi$  can be a step function in terms of many, various partitions.

**Lemma 3.1.** Let  $\phi \in \tau[a, b]$  be a step function in terms of partition  $(x_j)_{j=0}^n$  and let  $(\xi_i)_{i=0}^m$  be a refinement of  $(x_j)_{j=0}^n$ . Then  $\phi$  is also a step function in terms of  $(\xi_i)_{i=0}^m$ .

*Proof.* Refinement: For every  $j \in \{0, \dots, n\}$  there exists  $i_j \in \{0, \dots, m\}$  such that  $x_j = \xi_{i_j}$ .  $i_0 = 0, i_n = m$ .  $i_{j-1} < i_j$ .

Let  $i \in \{1, \dots, m\}$ . Then there exists a uniquely determined  $j \in \{1, \dots, n\}$  such that  $\xi_{i-1} < \xi_i \leq \xi_j$  Compare with Figure 12.



Figure 12:  $\xi$  on a refinement  $x_{i_j}$

Then  $(\xi_{i-1}, \xi_i) \subseteq \underbrace{(\xi_{i_{j-1}}, \xi_{i_j})}_{=(x_{j-1}, x_j)}$  and  $\varphi|_{(\xi_{i-1}, \xi_i)} = c_j$  constant. So  $\varphi$  is a subsequence with respect to  $(\xi_i)_{i=0}^m$ .  $\square$

**Definition 3.2.** Let  $\varphi \in \tau[a, b]$  in terms of partition  $(x_j)_{j=0}^n$  with  $\varphi|_{(x_{j-1}, x_j)} = c_j$  and  $\Delta x_j = x_j - x_{j-1} > 0$  for  $j = 1, \dots, n$ . Then we define ...

$$\int_a^b \varphi dx = \sum_{j=1}^n c_j \Delta x_j$$

is called integral of  $\varphi$  in terms of partition  $(x_j)_{j=0}^n$

↓ This lecture took place on 2018/03/22.

Step function  $\varphi$ .  $\varphi|_{(x_{j-1}, x_j)} = c_j$

$$\Delta x_j := x_j - x_{j-1}$$

$$\int_a^b \varphi dx = \sum_{j=1}^n c_j \cdot \Delta x_j$$

**Lemma 3.2.** Let  $(x_i)_{i=0}^n$  be a partition of  $[a, b]$  and  $(\xi_i)_{i=0}^m$  be a refinement of  $(x_j)_{j=0}^n$ . Furthermore let  $\varphi$  be a subsequence with respect to  $(x_j)_{j=0}^n$  (so also with respect to  $(\xi_j)_{j=0}^m$ ). Then the integrals of  $\varphi$  with respect to  $(x_j)_{j=0}^n$  and  $(\xi_i)_{i=0}^m$  are equal.

*Proof.* There exist indices  $i_j$  for  $j = 0, n$  such that  $x_j = \xi_{i_j}$ .

$$i_0 = 0 \quad i_n = m \quad i_{j-1} < i_j$$




Figure 13: Integral of a step function as sum of areas of rectangles

$$\begin{aligned}
 \Delta x_j &= x_j - x_{j-1} = \xi_{i_j} - \xi_{i_{j-1}} = \xi_{i_j} - \xi_{i_{j-1}} = \underbrace{\sum_{i=i_{j-1}+1}^{i_j} (\xi_i - \xi_{i-1})}_{\text{telescoping sum}} = \sum_{i=i_{j-1}+1}^{i_j} \Delta \xi_i \\
 \varphi|_{x_{j-1}, x_j} &= c_j \implies \varphi|_{(\xi_{i-1}, \xi_i)} = c_j \text{ for } i = i_{j-1} + 1, \dots, i_j \\
 \tilde{c}_i &= \varphi|_{(\xi_{i-1}, \xi_i)} \\
 \underbrace{\sum_{i=1}^m \tilde{c}_i \Delta \xi_i}_{\text{integral of } \varphi \text{ w.r.t } (\xi_i)_{i=0}^m} &= \sum_{j=1}^n \sum_{i=i_{j-1}+1}^{i_j} \tilde{c}_i \Delta \xi_i = \sum_{j=1}^n c_j \underbrace{\sum_{i=i_{j-1}+1}^{i_j} \Delta \xi_i}_{=\Delta x_j} = \sum_{j=1}^n c_j \Delta x_j
 \end{aligned}$$

This is the integral of  $\varphi$  with respect to  $(x_j)_{j=0}^n$ . □

**Lemma 3.3.** Let  $\varphi$  be a step function with respect to  $(x_j)_{j=0}^n$  and  $(w_l)_{l=0}^L$ . Then the integrals of  $\varphi$  with respect to  $(x_j)_{j=0}^n$  and with respect to  $(w_l)_{l=0}^L$  equal.

*Proof.* Let  $\{\xi_i | i = 1, \dots, m\} = \{x_j | j = 0, \dots, n\} \cup \{w_l | l = 0, \dots, L\}$  with  $\xi_0 = a$ ,  $\xi_m = x_n = w_L = b$  and  $\xi_{i-1} < \xi_i$  for  $i = 1, \dots, m$ . Then  $(\xi_i)_{i=0}^m$  is a refinement of

$(x_j)_{j=0}^n$  as well as  $(w_l)_{l=0}^L$ . By Lemma 3.2, the integral of  $\varphi$  with respect to  $(x_j)_{j=0}^n =$  integral of  $\varphi$  with respect to  $(\xi_i)_{i=1}^m =$  integral of  $\varphi$  with respect to  $(w_l)_{l=0}^L$ . Here we discard the statement “with respect to  $(x_j)_{j=0}^n$ ”.  $\square$

**Lemma 3.4.** *Let  $f, g$  be step functions on  $[a, b]$ , i.e.  $f, g \in \tau[a, b]$ .*

- for  $\alpha, \beta \in \mathbb{R}$ , let  $\alpha f + \beta g \in \tau[a, b]$  and

$$\int_a^b (\alpha f + \beta g) dx = \alpha \int_a^b f dx + \beta \int_a^b g dx$$

Hence, the integral is linear on  $[a, b]$ .  $\tau[a, b]$  is a vector space.

- $f \leq g$  in  $[a, b]$ , then  $\int_a^b f dx \leq \int_a^b g dx$  (monotonicity).
- $\left| \int_a^b f dx \right| \leq \int_a^b |f| dx$  ( $|f(x)|$  is also a step function)

*Proof.* 1. Let  $f, g \in \tau[a, b]$ . Let  $(\xi_i)_{i=0}^m$  be a partition such that  $f|_{(\xi_{i-1}, \xi_i)} = c_i$  and  $g|_{(\xi_{i-1}, \xi_i)} = d_i$ . Then

$$\begin{aligned} \int_a^b (\alpha f + \beta g) dx &= \sum_{i=1}^m (\alpha c_i + \beta d_i) \Delta \xi_i \\ &= \alpha \sum_{i=1}^m c_i \Delta \xi_i + \beta \sum_{i=1}^m d_i \Delta \xi_i = \alpha \int_a^b f dx + \beta \int_a^b g dx \end{aligned}$$

Furthermore,

$$(\alpha f + \beta g)|_{(\xi_{i-1}, \xi_i)} = \alpha c_i + \beta d_i = \text{constant}$$

Thus,

$$\alpha f + \beta g \in \tau[a, b]$$

2. Let  $h \in \tau[a, b]$  and  $h(x) \geq 0 \forall x \in [a, b]$ , then  $v_i = h|_{(\xi_{i-1}, \xi_i)} \geq 0$  and

$$\int_a^b h dx = \sum_{i=1}^m \underbrace{h_i}_{\geq 0} \underbrace{\Delta \xi_i}_{\geq 0} \geq 0$$

Now let  $f, g \in \tau[a, b]$ ;  $f \leq g$ . Then  $h = g - f \in \tau[a, b]$  (membership because of (1.)) and  $h \geq 0$ . Therefore,

$$0 \leq \int_a^b h dx = \int_a^b (g - f) dx = \int_a^b g dx - \int_a^b f dx$$



3. In general  $f \leq |f|$ , hence  $\int_a^b f \, dx \leq \int_a^b |f| \, dx$  and also  $-f \leq |f|$ , so

$$\int_a^b (-f) \, dx = - \int_a^b f \, dx \leq \int_a^b |f| \, dx \implies \left| \int_a^b f \, dx \right| \leq \int_a^b |f| \, dx$$

It is left to prove:  $|f| \in \tau[a, b]$  (i.e.  $|f|$  is a step function)

**Remark 3.1** (Elaboration).

$$\begin{aligned} f \leq |f| &\implies \int f \, dx \leq \int |f| \, dx \\ -|f| \leq f &\implies \int -|f| \, dx \leq \int f \, dx \iff - \int |f| \, dx \leq \int f \, dx \\ &\implies - \int |f| \, dx \leq \underbrace{\int f \, dx}_{=:J} \leq \underbrace{\int |f| \, dx}_{=:I} \\ &\iff -I \leq J \leq I \iff |J| \leq I \\ &\implies \left| \int f \, dx \right| \leq \int |f| \, dx \end{aligned}$$

Let  $f|_{(\xi_{i-1}, \xi_i)} = c_i \implies |f|_{(\xi_{i-1}, \xi_i)} = |c_i| = \text{constant}$ . Hence  $|f| \in \tau[a, b]$ .

□

## 3.2 Characteristic functions

**Definition 3.3.** Let  $A \subseteq \mathbb{R}^k$ . We call  $\chi_A : \mathbb{R}^n \rightarrow \mathbb{R}$  with

$$\chi_A(x) = \begin{cases} 1 & \text{if } x \in A \\ 0 & \text{else} \end{cases}$$

a characteristic function (indicator function) of set  $A$ . Often denoted as  $\chi_A = \mathbb{1}$ . Compare with Figure 14.

**Remark 3.2.** Let  $A = (a', b')$  with  $a \leq a' < b' \leq b$ . Then  $\chi_{(a', b')} \in \tau[a, b]$ . Also for  $x \in [a, b]$ , it holds that  $\chi_{\{x\}} \in \tau[a, b]$ . Therefore

- every linear combination of characteristic functions of open subintervals  $(a', b')$  of  $[a, b]$  or
- characteristic functions of one-point sets  $\chi_{\{x\}}, x \in [a, b]$




Figure 14: A characteristic function takes value 1 inside a set  $A$  which can be an interval (left) or a single point (right)




Figure 15: Linear combinations of characteristic functions are step functions

are step functions on  $[a, b]$ . Compare with Figure 15.

$$\sum_{j=1}^n \alpha_j \chi_{(a_j, b_j]} + \sum_{k=1}^m \beta_k \chi_{\{x_k\}} \in \tau[a, b]$$

On the opposite,  $f \in \tau[a, b]$ , hence

$$f|_{(x_{j-1}, x_j)} = c_j \text{ and } f(x_j) = d_j$$

$j=1, \dots, n$   $j=0, \dots, n$

$$f = \sum_{j=1}^n c_j \chi_{(x_{j-1}, x_j)} + \sum_{j=0}^n d_j \chi_{\{x_j\}} = (*)$$

for  $x \in (x_{j-1}, x_j)$ ,  $\xi_{(x_{j-1}, x_j)}(x) = 1$ .

$$\chi_{(x_{l-1}, x_l)}(x) = 0 \text{ for } l \neq j$$




Figure 16: The limit value must not be necessarily the function value

$$\chi_{\{x_l\}}(x) = 0 \text{ for } l = 0, \dots, n$$

i.e.  $\sum_{j=1}^n c_l \chi_{(x_{l-1}, x_l)}(x) + \sum_{l=0}^n d_j \chi_{\{x_l\}}(x) = c_j \cdot 1 + 0 = c_j$  hence  $(*) = c_j$  on  $(x_{j-1}, x_j)$ .  
Therefore  $f \in \tau[a, b] \iff f$  is linear combination of characteristic functions of open intervals or one-pointed sets.

### 3.3 Limit points

**Definition 3.4.** Let  $X$  be a metric space  $A \subseteq X$  and  $x \in X$  is an accumulating point<sup>1</sup> of  $A$ . Let  $f : A \rightarrow \mathbb{R}$ . Let  $d$  be the metric of  $X$ . We say,  $f$  has limit  $c \in \mathbb{R}$  in  $x$  ( $\lim_{\xi \rightarrow x} f(\xi) = c$ ) if

$$\forall \varepsilon > 0 \exists \delta > 0 \forall \xi \in A \forall \xi \neq x : d(\xi, x) < \delta \implies |f(\xi) - c| < \varepsilon$$

**Remark 3.3.** Recognize that  $\xi \neq x$  relaxes the precondition. The preconditions look similar to continuity, but fewer  $\xi$  need to satisfy a certain criterion.

**Remark 3.4.**  $x \in A$  and  $c = f(x) \implies f$  is continuous in  $x$ .

We usually consider  $A = [a, b] \subseteq \mathbb{R}$ ,  $x \in [a, b]$ .

It is possible, that  $f$  in  $x$  has a limit,  $x \in A$  and  $c = \lim_{\xi \rightarrow x} f(\xi) \neq f(x)$ . Compare with Figure 16.

**Definition 3.5.** Now let  $A \subseteq \mathbb{R}$  and  $x$  is a accumulation point of  $A$ . Let  $f : A \rightarrow \mathbb{R}$  be given. We say  $f$  has a right-sided limit  $c$  in  $x$  with  $c = \lim_{\xi \rightarrow x^+} f(\xi) = c$  if

$$\forall \varepsilon > 0 \exists \delta > 0 : \forall \xi \in A, \xi > x : |\xi - x| = \xi - x < \delta \implies |f(\xi) - c| < \varepsilon$$

Compare with Figure 17. The left-sided limit follows analogously (with  $\xi < x$ ).

$$c = \lim_{\xi \rightarrow x^+} f(\xi) \quad d = \lim_{\xi \rightarrow x^-} f(\xi)$$

<sup>1</sup>An accumulation point has 3 equivalent definitions (sequence, intersection, infinitely many elements in sphere).




Figure 17: Left-sided and right-sided limit

**Lemma 3.5** (Sequence criterion for limits of functions). *Let  $f : A \subseteq X \rightarrow \mathbb{R}$  be given.  $x$  is an accumulation point of  $A$ . Then*

$$\lim_{\xi \rightarrow x} f(\xi) = c \iff \forall (\xi_n)_{n \in \mathbb{N}} : \xi_n \in A, \xi_n \neq x \text{ and } \lim_{n \rightarrow \infty} \xi_n = x \text{ it holds that } \lim_{n \rightarrow \infty} f(\xi_n) = c$$

For one-sided limits  $A \subseteq \mathbb{R}$ ,

$$c = \lim_{\xi \rightarrow x^+} f(\xi) \iff \forall (\xi_n)_{n \in \mathbb{N}} : \xi_n \in A : \xi_n > x \text{ with } \lim_{n \rightarrow \infty} \xi_n = x \text{ it holds that } \lim_{n \rightarrow \infty} f(\xi_n) = c$$

*Proof.* See Analysis 1 lecture notes. □

**Remark 3.5.** Due to condition  $\xi_n \neq x$ , the function value must not necessarily match the limit point.

**Remark 3.6.** Attention! We, therefore, use two different definitions of limits.

**Lemma 3.6** (Cauchy criterion for limits of functions). *Let  $f : A \subseteq X \rightarrow \mathbb{R}$ . Let  $x$  be an accumulation point of  $A$ . Let  $X$  be a metric space.*

*$f$  has a limit in  $x$  if and only if  $\forall \varepsilon > 0 \exists \delta > 0 \forall \xi, \eta \in A$*

$$(\xi \neq x \wedge \eta \neq x \wedge d(\xi, x) < \delta \wedge d(\eta, x) < \delta) \implies |f(\xi) - f(\eta)| < \varepsilon$$

*Analogously for one-sided limits with  $A \subseteq \mathbb{R}$ . Additionally, we need the constraint that  $\xi > x$  and  $\eta > x$  for  $\lim_{\xi \rightarrow x^+} f(\xi)$ . And accordingly,  $\xi < x$  and  $\eta < x$  for  $\lim_{\xi \rightarrow x^-} f(\xi)$ .*

*Proof.*  $\implies$  Let  $c = \lim_{\xi \rightarrow x} f(\xi)$  and let  $\varepsilon > 0$  be chosen arbitrarily. Then there exists  $\delta > 0$  such that  $d(\xi, x) < \delta$  and  $\xi \neq x$

$$\implies |f(\xi) - c| < \frac{\varepsilon}{2}$$

For  $\xi, \eta$ :  $d(\xi, x) < \delta$  and  $d(\eta, x) < \delta$  with  $\xi, \eta \neq x$  is therefore

$$|f(\xi) - f(\eta)| = |f(\xi) - c + c - f(\eta)| \leq |f(\xi) - c| + |f(\eta) - c| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

$\Leftarrow$  Assume the Cauchy criterion holds. We show that

1. for every sequence  $(\xi_n)_{n \in \mathbb{N}}$ ,  $\xi_n \in A \setminus \{x\}$  with  $\lim_{n \rightarrow \infty} \xi_n = x$  it holds that  $(f(\xi_n))_{n \in \mathbb{N}}$  is a Cauchy sequence in  $\mathbb{R}$  and therefore convergent in  $\mathbb{R}$ .
2. all Cauchy sequences have the *same* limit  $c$ .

We prove (1.)

Let  $(\xi_n)_{n \in \mathbb{N}}$  be as above. Let  $\varepsilon > 0$  be arbitrary. and  $N_\varepsilon$  large enough such that  $\forall n \geq N_\varepsilon : d(\xi_n, x) < \delta$  ( $\delta$  chosen appropriately to  $\varepsilon$  according to the Cauchy criterion).

By the Cauchy criterion,  $|f(\xi_n) - f(\xi_m)| < \varepsilon$  for all  $m, n \geq N_\varepsilon$ . Therefore  $(f(\xi_n))_{n \in \mathbb{N}}$  is a Cauchy sequence in  $\mathbb{R}$ . If  $\mathbb{R}$  is complete, then there exists  $c = \lim_{n \rightarrow \infty} f(\xi_n)$ . QED.

We prove (2.)

Let  $\xi_n \rightarrow x$  as above and  $\xi'_n \rightarrow x$  as above and  $c = \lim_{n \rightarrow \infty} f(\xi_n)$  as well as  $c' = \lim_{n \rightarrow \infty} f(\xi'_n)$ . Let  $\varepsilon > 0$  be arbitrary,  $N_\varepsilon$  such that  $n \geq N_\varepsilon \implies |f(\xi_n) - c| < \frac{\varepsilon}{3}$  and  $N'_\varepsilon \in \mathbb{N}$  such that  $n \geq N'_\varepsilon \implies |f(\xi'_n) - c'| < \frac{\varepsilon}{3}$ .

Furthermore choose  $\delta > 0$  such that

$$d(\xi, x) < \delta \wedge d(\eta, x) < \delta \implies |f(\xi) - f(\eta)| < \frac{\varepsilon}{3}$$

(because of the Cauchy criterion).  $M_\varepsilon$  such that

$$n \geq M_\varepsilon \implies d(\xi_n, x) < \delta \wedge M'_\varepsilon : n \geq M'_\varepsilon \implies d(\xi'_n, x) < \delta$$

Let  $n \geq \max\{N_\varepsilon, N'_\varepsilon, M_\varepsilon, M'_\varepsilon\}$ .

$\downarrow$  This lecture took place on 2018/04/10.

Then

$$|c - c'| \leq \underbrace{|c - f(\xi_n)|}_{< \frac{\varepsilon}{3}} + \underbrace{|f(\xi_n) - f(\xi'_n)|}_{< \frac{\varepsilon}{3}} + \underbrace{|f(\xi'_n) - c'|}_{< \frac{\varepsilon}{3}} \quad \forall \varepsilon > 0$$

Hence,  $c = c'$ . We have shown that  $\exists c \in \mathbb{R} : \forall (\xi_n)_{n \in \mathbb{N}}$  with  $\lim_{n \rightarrow \infty} \xi_n = x$  it holds that  $\lim_{n \rightarrow \infty} f(\xi_n) = c$ . So  $\lim_{\xi \rightarrow \infty} f(\xi) = c$  because of Lemma 3.5. QED.

□

### 3.4 Regulated functions

**Definition 3.6** (Regulated function). Let  $a < b$ ,  $f : [a, b] \rightarrow \mathbb{R}$ . We call  $f$  a regulated function on  $[a, b]$  if

1.  $\forall x \in (a, b)$ ,  $f$  in  $x$  has a right-sided and a left-sided limit.
2. in  $x = a$ ,  $f$  has a right-sided limit.
3. in  $x = b$ ,  $f$  has a left-sided limit.

$$\mathcal{R}[a, b] = \{ f : [a, b] \rightarrow \mathbb{R} \mid f \text{ is a regulated function} \}$$

**Definition 3.7** (Equivalent definition).

1.  $\forall x \in [a, b)$ ,  $f$  has a right-sided limit in  $x$
2.  $\forall x \in (a, b]$ ,  $f$  has a left-sided limit in  $x$

**Example 3.1.** 1. Let  $f$  be continuous in  $[a, b]$ . Then  $f \in \mathcal{R}[a, b]$ .

2. Let  $\varphi \in \tau[a, b]$  be a step function. Then  $\varphi \in \mathcal{R}[a, b]$ .

*Rationale:*

Let  $x_0 = a < x_1 < \dots < x_n = b$  and  $\varphi|_{(x_{j-1}, x_j)} = c_j$ .

Let  $x \in [a, b]$  be chosen arbitrarily.

**Case 1** Let  $x \in (x_{j-1}, x_j)$  for some  $j \in \{1, \dots, n\}$

$$\implies \lim_{\xi \rightarrow x} \varphi(\xi) = c_j$$

Choose  $\delta$  small enough such that  $(x - \delta, x + \delta) \subseteq (x_{j-1}, x_j)$ .  $\forall \xi$  with  $\xi \in (x - \delta, x + \delta)$  it holds that

$$|\varphi(\xi) - c_j| = 0$$

**Case 2** Let  $x = x_j$  for  $j = 1, \dots, n - 1$ .

$$\implies \lim_{\xi \rightarrow x_j^+} \varphi(\xi) = c_{j+1} \quad \lim_{\xi \rightarrow x_j^-} \varphi(\xi) = c_j$$

Compare with Figure 18.




Figure 18: Regulated function

**Case 3** Let  $x = x_0 = a \implies \lim_{\xi \rightarrow a^+} \varphi(\xi) = c_1$ .

$$x = x_n = b \implies \lim_{\xi \rightarrow b^-} \varphi(\xi) = c_n$$

Let  $f : [a, b] \rightarrow \mathbb{R}$  be monotonically increasing or monotonically decreasing. Then  $f \in \mathcal{R}[a, b]$ . The proof will be done in the practicals.

### 3.5 Bounded functions on bounded sets

**Definition 3.8** (Boundedness). Let  $X \neq \emptyset$  be a set.  $f : X \rightarrow \mathbb{K}$  with  $\mathbb{K} = \mathbb{R}$  or  $\mathbb{K} = \mathbb{C}$ . We say:  $f$  is bounded on  $X$ , if  $f(X) \subseteq \mathbb{K}$  is a bounded set in  $\mathbb{K}$ . Hence,  $\exists m \geq 0 : |f(x)| \leq m \forall x \in X$ . We let,

$$\mathcal{B}(X) = \{f : X \rightarrow \mathbb{K} \mid f \text{ is bounded}\}$$

$\mathcal{B}(X)$  has vector space structure.  $f, g \in \mathcal{B}(X), \lambda \in \mathbb{K}$ .

$$(f + g)(x) = f(x) + g(x)$$

$$(\lambda \cdot f)(x) = \lambda \cdot f(x)$$

$f + g \in \mathcal{B}(X)$  and  $\lambda f \in \mathcal{B}(X)$ . Let  $|f(x)| \leq m \forall x \in X$  and  $|g(x)| \leq m' \forall x \in X$ . Then

$$|(f + g)(x)| = |f(x) + g(x)| \leq |f(x)| + |g(x)| \leq m + m'$$

**Remark 3.7.** It is very interesting, that  $X$  does not require any kind of algebraic structure.

We let

$$\|f\|_\infty = \sup \underbrace{\{|f(x)| : x \in X\}}_{\text{bounded in } \mathbb{R}} = \min \{m \geq 0 : |f(x)| \leq m \forall x \in X\}$$

Some work is required to show that  $\|\cdot\|_\infty$  is a norm on  $\mathcal{B}(X)$ .

Hence,  $(\mathcal{B}(X), \|\cdot\|_\infty)$  is a normed vector space. Convergence in  $\mathcal{B}(X)$ : It holds that  $f_n \rightarrow f$  in  $(\mathcal{B}(X), \|\cdot\|_\infty)$  if and only if  $\forall \varepsilon > 0 \exists N \in \mathbb{N} : n \geq N \implies \|f_n - f\|_\infty < \varepsilon$ .

$$\begin{aligned} \|f_n - f\|_\infty < \varepsilon &\iff \sup \{|f_n(x) - f(x)| : x \in X\} < \varepsilon \\ &\iff |f_n(x) - f(x)| \leq \varepsilon \forall x \in X \end{aligned}$$

Hence,  $f_n \rightarrow f$  in  $(\mathcal{B}(X), \|\cdot\|_\infty) \iff \forall \varepsilon > 0 \exists N \in \mathbb{N} : n \geq N \implies |f_n(x) - f(x)| \leq \varepsilon \forall x \in X$ . We say " $f_n$  converges uniformly to  $f$  on  $X$ ".

**Theorem 3.1** (Approximation theorem for regulated function). Let  $f : [a, b] \rightarrow \mathbb{R}$ . Then  $f \in \mathcal{R}[a, b] \iff \forall \varepsilon > 0$  there exists some step function  $\varphi \in \tau[a, b]$  such that  $|\varphi(x) - f(x)| < \varepsilon \forall x \in [a, b]$  ( $\|\varphi - f\|_\infty < \varepsilon$ ).

Especially  $\varepsilon_n = \frac{1}{n}$  and  $\varphi_n$  as above. Then  $\|\varphi_n - f\|_\infty < \frac{1}{n}$ , hence  $f = \lim_{n \rightarrow \infty} \varphi_n$  uniformly on  $[a, b]$ .

*Proof.*

**Direction  $\implies$ .** Let  $f \in \mathcal{R}[a, b]$ .

Proof by contradiction. We negate our hypothesis:

$$\exists \varepsilon > 0 : \forall \varphi \in \tau[a, b] \exists x \in [a, b] : |\varphi(x) - f(x)| \geq \varepsilon \quad (1)$$

Assume (1) holds for  $f \in [a, b]$ . We construct nested intervals  $[a_n, b_n]$  with  $[a_{n+1}, b_{n+1}] \subseteq [a_n, b_n]$  and  $b_{n+1} - a_{n+1} = \frac{1}{2}(b_n - a_n)$  and (1) holds on  $[a_n, b_n] \forall n \in \mathbb{N}$ . Hence  $\forall \varphi \in \tau[a_n, b_n] \exists x \in [a_n, b_n]$  such that  $|\varphi(x) - f(x)| \geq \varepsilon$ . This is what we want to show.

Let  $a_0 = a$  and  $b_0 = b$ . Then (1) holds on  $[a_0, b_0]$  by assumption.  $n \rightarrow n + 1$ : Construction of  $[a_{n+1}, b_{n+1}]$ . Let  $m_n = \frac{1}{2}(a_n + b_n)$ .

**Claim.** (1) holds either on  $[a_n, m_n]$  or on  $[m_n, b_n]$ .



*Proof.* Because if the opposite of (1) holds on  $[a_n, m_n]$  as well as  $[m_n, b_n]$ , then there exists  $\varphi_n^1 \in \tau[a_n, m_n]$  with  $|\varphi_n^1(x) - f(x)| < \varepsilon \forall x \in [a_n, m_n]$  and if the opposite of (1) holds on  $[m_n, b_n]$ :

$$\exists \varphi_n^2 \in \tau[m_n, b_n] : |\varphi_n^2(x) - f(x)| < \varepsilon \forall x \in [m_n, b_n]$$

$$\varphi^n(x) := \begin{cases} \varphi_n^1(x) & \text{if } x \in [a_n, m_n] \\ \varphi_n^2(x) & \text{if } x \in (m_n, b_n] \end{cases}$$

Then  $\varphi^n$  is piecewise constant, hence  $\varphi^n \in \tau[a_n, b_n]$  and

$$|\varphi^n(x) - f(x)| = \begin{cases} \underbrace{|\varphi_n^1(x) - f(x)|}_{< \varepsilon} & \text{for } x \in [a_n, m_n] \\ \underbrace{|\varphi_n^2(x) - f(x)|}_{< \varepsilon} & \text{for } x \in [m_n, b_n] \end{cases}$$

Therefore,  $|\varphi^n(x) - f(x)| < \varepsilon$ , which contradicts with (1) on  $[a_n, b_n]$ . We conclude: (1) holds on  $[a_n, m_n]$  or on  $[m_n, b_n]$ .  $\square$

Now, choose  $[a_{n+1}, b_{n+1}]$  as one of the subintervals in which (1) holds. Let  $X \in \bigcap_{n \in \mathbb{N}} [a_n, b_n]$  (by completeness of  $\mathbb{R}$ ).

**Case  $x \in (a, b)$**  Let  $x \in (a, b)$ . Let  $\varepsilon$  as above such that (1) holds on every interval  $[a_n, b_n]$ . Let  $c_+ = \lim_{\xi \rightarrow x^+} f(\xi)$  and  $c_- = \lim_{\xi \rightarrow x^-} f(\xi)$  (feasible, because  $f \in \mathcal{R}[a, b]$ ).

By the limit property,  $\exists \delta > 0 : |\xi - x| < \delta$  and  $\xi < x$ , then  $|f(\xi) - c_-| < \varepsilon$  and  $|\xi - x| < \delta$  and  $x < \delta$  then  $|f(\xi) - c_+| < \varepsilon$ .

Additionally, choose  $\delta$  sufficiently small enough such that  $(x - \delta, x + \delta) \subseteq [a, b]$ .

$$\hat{\varphi}(\xi) := \begin{cases} 0 & \text{for } \xi \in [a, b] \setminus (x - \delta, x + \delta) \\ c_- & \text{for } \xi \in (x - \delta, x) \\ c_+ & \text{for } \xi \in (x, x + \delta) \\ f(x) & \text{for } \xi = x \end{cases}$$

Compare with Figure 19.  $\hat{\varphi} \in \tau[a, b]$  and

$$\forall \xi \in (x - \delta, x + \delta) : |\hat{\varphi}(\xi) - f(\xi)| = \begin{cases} \underbrace{|c_- - f(\xi)|}_{< \varepsilon} & \text{for } \xi \in (x - \delta, x) \\ \underbrace{|f(x) - f(x)|}_{=0} & \text{for } \xi = x \\ \underbrace{|c_+ - f(\xi)|}_{< \varepsilon} & \text{for } \xi \in (x, x + \delta) \end{cases}$$




Figure 19: Construction of  $\hat{\phi}(\xi)$

Hence,  $|\hat{\phi}(\xi) - f(\xi)| < \varepsilon$ . Now let  $N$  be sufficiently large enough such that  $[a_N, b_N] \subseteq (x - \delta, x + \delta)$  (possible because  $([a_n, b_n])_{n \in \mathbb{N}}$  gives nested intervals tightening on  $x$ ). Then for  $[a_N, b_N]$ :

$$\hat{\phi}|_{[a_N, b_N]} \in \tau[a_N, b_N]$$

and  $\forall \xi \in [a_N, b_N] \subseteq (x - \delta, x + \delta) : |\hat{\phi}(\xi) - f(\xi)| < \varepsilon$ . This contradicts with (1) on  $[a_N, b_N]$ .

**Case  $x = a$  and  $x = b$**  Is analogous to one-sided limits.

**Direction  $\Leftarrow$ .** Let  $f = \lim_{n \rightarrow \infty} \varphi_n$  be uniform on  $[a, b]$ .

**Claim.**  $\forall x \in (a, b)$  there exists a right-sided limit of  $f$  in  $x$ .

*Proof.* Let  $\varepsilon > 0$  be arbitrary.  $N \in \mathbb{N}$  sufficiently large such that  $|f(\xi) - \varphi_N(\xi)| < \frac{\varepsilon}{2} \forall \xi \in [a, b]$ .  $\varphi_N$  is piecewise constant. Choose  $\delta > 0$  such that  $\varphi_N|_{(x, x+\delta)} = c$ . Now let  $\xi, \eta \in (x, x + \delta)$  be chosen arbitrarily. Then

$$\begin{aligned} |f(\xi) - f(\eta)| &\leq |f(\xi) - \underbrace{c}_{=\varphi_N(\xi)}| + |\underbrace{c}_{=\varphi_N(\eta)} - f(\eta)| \\ &= \underbrace{|f(\xi) - \varphi_N(\xi)|}_{< \frac{\varepsilon}{2}} + \underbrace{|\varphi_N(\eta) - f(\eta)|}_{< \frac{\varepsilon}{2}} < \varepsilon \end{aligned}$$

Therefore  $f$  has a right-sided limit in  $x$  by the Cauchy criterion.  $f$  has right-sided limit in every point.  $\square$

$\forall x \in (a, b)$  for left-sided limits follows analogously.  $\square$

**Corollary.** Every regulated function  $f \in \mathcal{R}[a, b]$  is bounded.

*Rationale.* Let  $\varphi \in \tau[a, b]$  with  $\|f - \varphi\|_\infty < 1$ .  $\varphi$  is bounded, hence  $\exists m \in [0, \infty)$ :  $|\varphi(x)| \leq m \forall x \in [a, b]$ . Then  $|f(x)| \leq |f(x) - \varphi(x)| + |\varphi(x)| < 1 + m \forall x \in [a, b]$ , hence  $f \in \mathcal{B}[a, b]$ .

$$\mathcal{R}[a, b] \subseteq \mathcal{B}[a, b]$$

□

**Corollary.** Let  $f \in \mathcal{R}[a, b] \iff f = \sum_{j=0}^{\infty} \psi_j$  with  $\psi_j \in \tau[a, b]$  and the series converges uniformly on  $[a, b]$ .

*Proof.*

**Direction  $\Leftarrow$ .** Let  $f = \sum_{j=0}^{\infty} \psi_j$  with uniform convergence. Let  $\varphi_n := \sum_{j=0}^n \psi_j \in \tau[a, b]$  and  $f = \lim_{n \rightarrow \infty} \varphi_n$  uniform on  $[a, b] \xrightarrow{\text{Theorem 3.1}} f \in \mathcal{R}[a, b]$ .

**Direction  $\Rightarrow$ .** Let  $f \in \mathcal{R}[a, b]$  and  $f = \lim_{n \rightarrow \infty} \varphi_n$  with  $\varphi_n \in \tau[a, b]$  (by Theorem 3.1).

$$\psi_0 = \varphi_0$$

$$\psi_j = \varphi_j - \varphi_{j-1} \quad \text{for } j \geq 1$$

$$\sum_{j=0}^n \psi_j = \varphi_0 + \sum_{j=1}^n (\varphi_j - \varphi_{j-1}) = \varphi_0 + \sum_{j=1}^n \varphi_j - \sum_{j=0}^{n-1} \varphi_j = \varphi_n$$

converges uniformly to  $f$ .

□

## 4 Integration of regulated functions

**Definition 4.1** (Definition with a theorem). Let  $f \in \mathcal{R}[a, b]$  and  $\varphi_n \in \tau[a, b]$  with  $f = \lim_{n \rightarrow \infty} \varphi_n$  is uniform on  $[a, b]$ . We let

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \int_a^b \varphi_n(x) dx$$

for the integral of  $f$  on  $[a, b]$ .

*Theorem:* This limit (on the right-hand side) always exists and is independent of the particular choice of the approximating sequence.

*Proof.*  $\varphi_n$  is chosen as above.

$$i_n = \int_a^b \varphi_n dx$$

Show:  $i_n$  is Cauchy sequence in  $\mathbb{R}$ .

↓ This lecture took place on 2018/04/12.

Let  $\varepsilon > 0$  be chosen arbitrary. Choose  $N \in \mathbb{N}$  such that

$$n \geq N \implies \|f - \varphi_n\|_\infty < \frac{\varepsilon}{2(b-a)}$$

For  $n, m \geq N$  it holds for  $x \in [a, b]$  that

$$\begin{aligned} |\varphi_n(x) - \varphi_m(x)| &\leq |\varphi_n(x) - f(x)| + |f(x) - \varphi_m(x)| \\ &\leq \| \varphi_n - f \|_\infty + \| f - \varphi_m \|_\infty < \frac{\varepsilon}{2(b-a)} + \frac{\varepsilon}{2(b-a)} = \frac{\varepsilon}{b-a} \end{aligned}$$

$|\varphi_n - \varphi_m|$  is a step function.

$$|\varphi_n - \varphi_m| \leq \frac{\varepsilon}{b-a} \cdot \underbrace{\chi_{[a,b]}}_{\in \tau[a,b]}$$

Integral for subsequence is monotonous:

$$\begin{aligned} |i_n - i_m| &= \left| \int_a^b \varphi_n dx - \int_a^b \varphi_m dx \right| = \left| \int_a^b (\varphi_n - \varphi_m) dx \right| \underbrace{\leq}_{\text{Lemma 3.4}} \int_a^b |\varphi_n - \varphi_m| dx \\ &\underbrace{\leq}_{\text{by monotonicity}} \int_a^b \frac{\varepsilon}{b-a} \cdot \chi_{[a,b]} dx = \frac{\varepsilon}{b-a} \underbrace{\int_a^b \chi_{[a,b]} dx}_{1 \cdot (b-a)} = \varepsilon \end{aligned}$$

So  $(i_n)_{n \in \mathbb{N}}$  is a Cauchy sequence.  $\mathbb{R}$  is complete, hence  $i = \lim_{n \rightarrow \infty} i_n$  exists.

Uniqueness: (dt. mithilfe des Reissverschlussprinzips)

Let  $(\varphi_n)_{n \in \mathbb{N}}, (\Phi_n)_{n \in \mathbb{N}}$  be two sequences of step functions, converging uniformly towards  $f$ .

$$\begin{aligned} i_n &= \int_a^b \varphi_n dx \quad \text{and} \quad j_n = \int_a^b \Phi_n dx \\ i &= \lim_{n \rightarrow \infty} i_n \quad \quad j = \lim_{n \rightarrow \infty} j_n \end{aligned}$$

Show that  $i = j$ .

Now we construct a sequence  $(\mu_n)_{n \in \mathbb{N}}$  of step functions.

$$\underbrace{(\varphi_1, \Phi_1, \varphi_2, \Phi_2, \dots)}_{(\mu_n)_{n \in \mathbb{N}}}$$

$\mu_n$  is a sequence of step functions converging uniformly towards  $f$  (the proof is left as an exercise to the reader).

Because of part 1 of the proof:

$$m_n = \int_a^b \mu_n dx \text{ converges with limit } m$$

$(i_n)_{n \in \mathbb{N}}$  as well as  $(j_n)_{n \in \mathbb{N}}$  are subsequences of  $(m_n)_{n \in \mathbb{N}}$ . Hence  $i = \lim_{n \rightarrow \infty} i_n = m = \lim_{n \rightarrow \infty} j_n = j$ .  $\square$

## 4.1 Elementary properties of an integral

**Theorem 4.1** (Elementary properties of an integral). *Let  $f, g \in \mathcal{R}[a, b]$ ,  $\lambda, \mu \in \mathbb{R}$ . Then*

**Linearity**

$$\lambda f + \mu g \in \mathcal{R}[a, b] \text{ and } \int_a^b (\lambda f + \mu g) dx = \lambda \int_a^b f dx + \mu \int_a^b g dx$$

**Monotonicity** *If  $f(x) \leq g(x) \forall x \in [a, b]$  ( $f \leq g$ ) it holds that*

$$\int_a^b f dx \leq \int_a^b g dx$$

**Boundedness**  $|f| \in \mathcal{R}[a, b]$  and

$$\left| \int_a^b f dx \right| \leq \int_a^b |f| dx$$

*Proof.*

**Linearity.** Let  $x \in [a, b]$  and  $c_+ = \lim_{\xi \rightarrow x^+} f(\xi)$  as well as  $d_+ = \lim_{\xi \rightarrow x^+} g(\xi)$  ( $f, g \in \mathcal{R}[a, b]$ ). Then

$$\lim_{\xi \rightarrow x^+} (\lambda f(\xi) + \mu g(\xi)) = \lambda \lim_{\xi \rightarrow x^+} f(\xi) + \mu \lim_{\xi \rightarrow x^+} g(\xi) = \lambda c_+ + \mu d_+$$

exists. Analogously for the left side, hence  $\lambda f + \mu g \in \mathcal{R}[a, b]$ .

**Claim.** Let  $\varphi_n, \Phi_n \in \tau[a, b]$  with  $\varphi_n \rightarrow f$  and  $\Phi_n \rightarrow g$  is uniform on  $[a, b]$ . Hence  $\lambda \varphi_n + \mu \Phi_n \rightarrow \lambda f + \mu g$  is continuous on  $[a, b]$ .

*Proof.* Let  $\varepsilon > 0$  be arbitrary,  $N$  such that  $n \geq N \implies \|\varphi_n - f\|_\infty < \frac{\varepsilon}{2(|\lambda|+1)}$  and  $M$  such that  $n \geq M \implies \|\Phi_n - g\|_\infty < \frac{\varepsilon}{2(|\mu|+1)}$ .

Then

$$\begin{aligned} \|\lambda\varphi_n + \mu\Phi_n - \lambda f - \mu g\|_\infty &\leq |\lambda| \|\varphi_n - f\|_\infty + |\mu| \|\Phi_n - g\|_\infty \\ &< \frac{|\lambda|}{2(|\lambda| + 1)} \cdot \varepsilon + \frac{|\mu|}{2(|\mu| + 1)} \cdot \varepsilon < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon \end{aligned}$$

□

We continue:

$$\begin{aligned} \int_a^b (\lambda f + \mu g) dx &= \lim_{n \rightarrow \infty} \int_a^b (\lambda \varphi_n + \mu \Phi_n) dx = \lim_{n \rightarrow \infty} \left( \lambda \int_a^b \varphi_n dx + \mu \int_a^b \Phi_n dx \right) \\ &= \lambda \underbrace{\lim_{n \rightarrow \infty} \int_a^b \varphi_n dx}_{\text{exists}} + \mu \underbrace{\lim_{n \rightarrow \infty} \int_a^b \Phi_n dx}_{\text{exists}} = \lambda \int_a^b f dx + \mu \int_a^b g dx \end{aligned}$$

**Monotonicity.** Show: Let  $h \in \mathcal{R}[a, b]$  with  $h \geq 0$  in  $[a, b]$ . Then  $\int_a^b h dx \geq 0$ .

**Claim.** There exists  $(\tilde{\varphi}_n)_{n \in \mathbb{N}}$  with  $\tilde{\varphi}_n \rightarrow h$  uniform on  $[a, b]$  and  $\tilde{\varphi}_n \geq 0$ .

*Proof.* Let  $(\varphi_n)_{n \in \mathbb{N}}$ ,  $\varphi_n \in \tau[a, b]$  with  $\varphi_n \rightarrow h$  uniform on  $[a, b]$ . We define  $\tilde{\varphi}_n$  such that

$$\begin{aligned} \varphi_n &= \sum_{j=1}^{m_n} c_j \chi_{(x_{j-1}, x_j)} + \sum_{j=0}^{m_n} d_j \chi_{\{x_j\}} \\ \tilde{\varphi}_n &:= \sum_{j=1}^{m_n} \underbrace{\tilde{c}_j}_{\geq 0} \chi_{(x_{j-1}, x_j)} + \sum_{j=0}^{m_n} \underbrace{h(x_j)}_{\geq 0} \chi_{\{x_j\}} \end{aligned}$$

and  $\tilde{c}_j := \max c_j, 0 \geq 0$ . So  $\tilde{\varphi}_n \geq 0$ .

For  $x = x_l$  ( $l \in \{0, \dots, m_n\}$ )

$$\begin{aligned} |\tilde{\varphi}_n(x_l) - h(x_l)| &= \left| \sum_{j=1}^{m_n} \tilde{c}_j \underbrace{\chi_{(x_{j-1}, x_j)}(x_l)}_{=0 \text{ bc. } x_l \notin (x_{j-1}, x_j)} + \sum_{j=0}^{m_n} h(x_j) \underbrace{\chi_{\{x_j\}}(x_l)}_{=\delta_{j,l}} - h(x_l) \right| \\ &= |h(x_l) - h(x_l)| = 0 \leq |\varphi_n(x_l) - h(x_l)| \end{aligned}$$

For  $x \in (x_{j-1}, x_j)$

$$|\tilde{\varphi}_n(x) - h(x)| = \left| \sum_{j=1}^{m_n} \tilde{c}_j \underbrace{\chi_{(x_{j-1}, x_j)}(x)}_{\delta_{l,j}} + \sum_{j=0}^{m_n} h(x) \cdot \underbrace{\chi_{\{x_j\}}(x)}_{=0 \text{ bc. } x \neq x_j} - h(x) \right|$$

$$\begin{aligned}
= |\tilde{c}_l - h(x)| &= \begin{cases} |c_l - h(x)| & \text{if } c_l \geq 0 \\ |h(x)| = h(x) & \text{if } c_l < 0 \end{cases} \\
&\leq \begin{cases} |c_l - h(x)| & \text{if } c_l \geq 0 \\ h(x) - c_l & \text{if } c_l < 0 \end{cases} \\
&= \begin{cases} |\varphi_n(x) - h(x)| & \text{if } c_l = \varphi_n(x) \geq 0 \\ |h(x) - \varphi_n(x)| & \text{if } c_l = \varphi_n(x) < 0 \end{cases} \\
&= |\varphi_n(x) - h(x)|
\end{aligned}$$

hence,  $|\tilde{\varphi}_n(x) - h(x)| \leq |\varphi_n(x) - h(x)|$  for  $x \in (x_{l-1}, x_l)$  as well as  $x = x_l$ , hence

$$\|\tilde{\varphi}_n - h\|_\infty \leq \underbrace{\|\varphi_n - h\|_\infty}_{\rightarrow 0 \text{ for } n \rightarrow \infty}$$

Hence  $\|\tilde{\varphi}_n - h\|_\infty \rightarrow 0$  for  $n \rightarrow \infty$ , hence  $\tilde{\varphi}_n$  converges uniformly to  $h$ .  
There exists

$$\int_a^b h \, dx = \lim_{n \rightarrow \infty} \underbrace{\int_a^b \tilde{\varphi}_n \, dx}_{\geq 0} \geq 0$$

□

To show monotonicity, we let  $f \leq g$  in  $[a, b]$ , hence  $h = g - f \geq 0$  in  $[a, b]$

$$\begin{aligned}
\Rightarrow 0 &\leq \int_a^b h \, dx = \int_a^b g \, dx - \int_a^b f \, dx \\
&\Rightarrow \int_a^b f \, dx \leq \int_a^b g \, dx
\end{aligned}$$

**Boundedness.** Consider  $|f|$ . Proving  $|f| \in \mathbb{R}[a, b]$  is left as an exercise to the reader.

$$\begin{aligned}
f \leq |f| \text{ on } [a, b] &\xrightarrow{\text{monotonicity}} \int_a^b f \, dx \leq \int_a^b |f| \, dx \\
-f \leq |f| \text{ on } [a, b] &\xrightarrow{\text{monotonicity}} \int_a^b (-f) \, dx = - \int_a^b f \, dx \leq \int_a^b |f| \, dx \\
&\Rightarrow \left| \int_a^b f \, dx \right| \leq \int_a^b |f| \, dx
\end{aligned}$$

□

**Remark 4.1.**  $\mathcal{R}[a, b]$  is a vector space.

1.  $f, g \in \mathcal{R}[a, b] \implies \lambda f + \mu g \in \mathcal{R}[a, b]$ .  $\|\cdot\|_\infty$  is a norm on  $\mathcal{R}[a, b]$ .  $(\mathcal{R}[a, b], \|\cdot\|_\infty)$  is a normed vector space. Subspace of  $(\mathcal{B}[a, b], \|\cdot\|_\infty)$ . We will show in the practicals that  $(\mathcal{R}[a, b], \|\cdot\|_\infty)$  is complete.

## 4.2 Mean value theorem of integration calculus

**Theorem 4.2** (Mean value theorem of integration calculus). *Let  $f$  be continuous on  $[a, b]$  and  $p \in \mathcal{R}[a, b]$  and  $p \geq 0$  in  $[a, b]$ . Then  $f \cdot p \in \mathcal{R}[a, b]$  and there exists  $\xi \in [a, b]$  such that*

$$\int_a^b f \cdot p \, dx = f(\xi) \cdot \int_a^b p \, dx$$

*Proof.* Let  $m = \min \{f(z) : z \in [a, b]\}$  (exists because  $f$  is continuous and  $[a, b]$  is compact).

$$M = \max \{f(z) : z \in [a, b]\}$$

$$f([a, b]) = [m, M] \text{ (by the mean value theorem)}$$

$$\underbrace{m \cdot p(x)}_{\geq 0} \leq f(x) \cdot p(x) \leq M \cdot p(x)$$

By monotonicity,

$$m \int_a^b p(x) \, dx \leq \int_a^b f p \, dx \leq M \int_a^b p \, dx$$

Therefore, there exists  $\eta \in [m, M]$ .

$$\eta \cdot \int_a^b p(x) \, dx = \int_a^b f p \, dx$$

Mean value theorem: For  $\eta \in [m, M]$  there exists  $\xi \in [a, b]$  such that

$$\eta = f(\xi) \text{ (f is continuous!)}$$

Hence,

$$f(\xi) \int_a^b p \, dx = \int_a^b f \cdot p \, dx$$

$f \cdot p$  is regulated function (over one-sided limits). □

**Lemma 4.1.** *Let  $f \in \mathcal{R}[a, b]$  and  $a \leq \alpha < \beta < \gamma \leq b$ . Then*

$$f|_{[\alpha, \beta]} \in \mathcal{R}[\alpha, \beta], f|_{[\beta, \gamma]} \in \mathcal{R}[\beta, \gamma]$$

$$f|_{[\alpha, \gamma]} \in \mathcal{R}[\alpha, \gamma] \text{ (immediate over onesided limit)}$$






Figure 20: Positive and negative area covered by the integral

$$\text{and } \int_{\alpha}^{\gamma} f dx = \int_{\alpha}^{\beta} f dx + \int_{\beta}^{\gamma} f dx$$

Compare with Figure 20.

*Proof.* Show that this statement holds for  $\varphi \in \tau[a, b]$ . Without loss of generality,  $\alpha = a, \gamma = b$ .

$$\gamma = \sum_{j=1}^m c_j \chi_{(x_{j-1}, x_j)} + \sum_{j=0}^m 0 \cdot \chi_{x_j}$$

The zero represents that this term is not relevant for the integral in any way.

**Case 1**  $\beta = x_l$  for some  $l \in \{1, \dots, m-1\}$

$$\int_{\alpha}^{\gamma} \varphi dx = \sum_{j=1}^m c_j (x_j - x_{j-1})$$

$$\int_{\alpha}^{\beta} \varphi dx = \int_{\alpha}^{x_l} \varphi dx = \sum_{j=1}^l c_j (x_j - x_{j-1})$$

$$\int_{\beta}^{\gamma} \varphi dx = \int_{x_l}^{\gamma} \varphi dx = \sum_{j=l+1}^m c_j (x_j - x_{j-1})$$

And now,

$$\sum_{j=1}^l c_j (x_j - x_{j-1}) + \sum_{j=l+1}^m c_j (x_j - x_{j-1}) = \sum_{j=1}^m c_j (x_j - x_{j-1})$$




Figure 21: Setting for  $\beta \in (x_{l-1}, x_l)$

**Case 2**  $\beta \in (x_{l-1}, x_l)$  for some  $l \in \{1, \dots, m\}$ . Compare with Figure 21.

$$\int_{\alpha}^{\beta} \varphi dx = \sum_{j=1}^{l-1} c_j(x_j - x_{j-1}) + c_l \cdot (\beta - x_{l-1})$$

$$\int_{\beta}^{\gamma} \varphi dx = c_l(x_l - \beta) + \sum_{j=l+1}^m c_j(x_j - x_{j-1})$$

Now we consider the addition of these two expressions:

$$\begin{aligned} & \int_{\alpha}^{\beta} \varphi dx + \int_{\beta}^{\gamma} \varphi dx \\ &= \sum_{j=1}^{l-1} c_j(x_j - x_{j-1}) + \underbrace{c_l(\beta - x_{l-1}) + c_l(x_l - \beta)}_{=c_l(x_l - x_{l-1})} + \sum_{j=l+1}^m c_j(x_j - x_{j-1}) \\ &= \sum_{j=1}^m c_j(x_j - x_{j-1}) = \int_{\alpha}^{\gamma} \varphi dx \end{aligned}$$

Let  $\varphi_n \in \tau[\alpha, \beta]$  with  $\varphi_n \rightarrow f$  uniform on  $[\alpha, \beta] \implies \varphi_n|_{[\alpha, \beta]} \rightarrow f|_{[\alpha, \beta]}$  uniform on  $[\alpha, \beta]$  and also  $\varphi_n|_{[\beta, \gamma]} \rightarrow f|_{[\beta, \gamma]}$  uniform on  $[\beta, \gamma]$ .

$$\begin{aligned} \int_{\alpha}^{\gamma} f dx &= \lim_{n \rightarrow \infty} \int_{\alpha}^{\gamma} \varphi_n dx = \lim_{n \rightarrow \infty} \left( \int_{\alpha}^{\beta} \varphi_n dx + \int_{\beta}^{\gamma} \varphi_n dx \right) \\ &= \underbrace{\lim_{n \rightarrow \infty} \int_{\alpha}^{\beta} \varphi_n dx}_{\substack{\text{exists because} \\ \varphi_n|_{[\alpha, \beta]} \rightarrow f|_{[\alpha, \beta]} \text{ uniform}}} + \lim_{n \rightarrow \infty} \int_{\beta}^{\gamma} \varphi_n dx = \int_{\alpha}^{\beta} f dx + \int_{\beta}^{\gamma} f dx \end{aligned}$$

□

**Remark 4.2** (Notation). Let  $\alpha < \beta$ ,  $\alpha, \beta \in [a, b]$  and  $f \in \mathcal{R}[a, b]$ . We let

$$\int_{\beta}^{\alpha} f \, dx := - \int_{\alpha}^{\beta} f \, dx$$

By this convention,

$$\int_{\alpha}^{\alpha} f \, dx = - \int_{\alpha}^{\alpha} f \, dx \implies \int_{\alpha}^{\alpha} f \, dx = 0$$

**Lemma 4.2.** Let  $f \in \mathcal{R}[a, b]$  and  $\alpha, \beta, \gamma \in [a, b]$  (without particular order). Then

$$\int_{\alpha}^{\gamma} f \, dx = \int_{\alpha}^{\beta} f \, dx + \int_{\beta}^{\gamma} f \, dx$$

*Proof.* Special case: 2 points are equal

$$\alpha = \gamma \implies \int_{\alpha}^{\alpha} f \, dx = 0$$

$$\begin{aligned} \int_{\alpha}^{\beta} f \, dx + \int_{\beta}^{\alpha} f \, dx &= \int_{\alpha}^{\beta} f \, dx - \int_{\alpha}^{\beta} f \, dx = 0 \\ \beta &= \gamma \quad \beta = \alpha \end{aligned}$$

Case:  $\alpha < \beta < \gamma$  follows immediately

And just as a representative other case:  $\alpha < \gamma < \beta$

$$\int_{\alpha}^{\beta} f \, dx \quad \underbrace{=}_{\text{by Lemma 1.7}} \quad \int_{\alpha}^{\gamma} f \, dx + \underbrace{\int_{\gamma}^{\beta} f \, dx}_{- \int_{\beta}^{\gamma} f \, dx}$$

$$\int_{\alpha}^{\beta} f \, dx + \int_{\beta}^{\gamma} f \, dx = \int_{\alpha}^{\gamma} f \, dx$$

□

↓ This lecture took place on 2018/04/17.

**Lemma 4.3.** Let  $f \in \mathcal{R}[a, b]$ . Then there exists an at most countable set  $A \subseteq [a, b]$  such that  $f$  is continuous in every point  $x \in [a, b] \setminus A$ .

*Proof.* Let  $f \in \mathcal{R}[a, b]$  and  $(\varphi_n)_{n \in \mathbb{N}}$  with  $\varphi_n \in \tau[a, b]$  and  $\varphi \rightarrow f$  converging uniformly on  $[a, b]$ .

$$\varphi_n = \sum_{j=1}^{m_n} c_j^n \chi_{(X_{j-1}^n, X_j^n)} + \sum_{j=0}^{m_n} d_j^n \chi_{\{x_j^n\}}$$

$$x_0^n = a < x_1^n < \dots < x_{m_n}^n = b$$

are separating points for  $\varphi_n$

$$A = \{X_j^n : n \in \mathbb{N}, j \in \{0, \dots, m_n\}\}$$

$A$  is a countable union of finite sets  $A_n = \{x_0^n, x_{m_n}^n\}$ .  $A$  is countable (as unions of finite sets are).

Now we show:  $f$  is continuous in every point  $x \in [a, b] : x \notin A$ . Let  $\varepsilon > 0$  be arbitrary. Choose  $N \in \mathbb{N}$  sufficiently large such that  $\|\varphi_N - f\|_\infty < \frac{\varepsilon}{2}$ . Because  $x \in A$ , there exists  $j \in \{1, \dots, m_N\}$  such that  $x \in (x_{j-1}^N, x_j^N)$  is open. Choose  $\delta > 0$  such that  $(x - \delta, x + \delta) \subset (x_{j-1}^N, x_j^N)$ , hence  $\forall \xi \in (x - \delta, x + \delta)$  it holds that  $\varphi_N(\xi) = c_j^N$ . Now consider  $\xi \in (x - \delta, x + \delta)$ , hence  $|\xi - x| < \delta$ . Then

$$\begin{aligned} |f(\xi) - f(x)| &= \left| f(\xi) - \underbrace{\varphi_N(x)}_{c_j^N = \varphi_N(\xi)} + \varphi_N(x) - f(x) \right| \\ &\leq \underbrace{|f(\xi) - \varphi_N(\xi)|}_{\leq \|f - \varphi_N\|_\infty} + \underbrace{|\varphi_N(x) - f(x)|}_{\leq \|\varphi_N - f\|_\infty} < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon \end{aligned}$$

Hence  $f$  is continuous in  $x$ . □

**Remark 4.3** (Notation). Let  $f \in \mathcal{R}[a, b]$ . For  $x \in [a, b)$ , there exists  $f_+(x) := \lim_{\xi \rightarrow x_+} f(\xi)$ . For  $x \in (a, b]$ , there exists  $f_-(x) := \lim_{\xi \rightarrow x_-} f(\xi)$ . Because of Lemma 4.3, it holds that  $f_+(x) = f_-(x) = f(x)$  for all  $x \in [a, b] \setminus A$  and  $A$  is at most countable.

### 4.3 One-sided derivatives

**Definition 4.2** (One-sided derivatives). Let  $g : [a, b] \rightarrow \mathbb{R}$  and  $x \in [a, b)$ . We say  $g$  has the right-sided derivative  $g'_+(x)$  if

$$\lim_{\xi \rightarrow x_+} \frac{g(\xi) - g(x)}{\xi - x} =: g'_+(x)$$

exists. Analogously we define the left-sided derivative

$$g'_-(x) = \lim_{\xi \rightarrow x_-} \frac{g(\xi) - g(x)}{\xi - x}$$

for  $x \in (a, b]$ . Compare with Figure 22.




Figure 22: In this example, the left- and right-sided derivatives are not equal.  
 $f'_+(x) \neq f'_-(x)$

**Remark 4.4.** If  $g$  in  $x$  has a one-sided derivative, then

$$\lim_{\xi \rightarrow x_{\pm}} (g(\xi) - g(x)) = 0$$

Hence  $g$  is continuous in  $x$ .

**Remark 4.5.**  $g : [a, b] \rightarrow \mathbb{R}$  is differentiable in point  $x \in (a, b)$  with derivative  $g'(x)$   
 $\iff g$  has a left- and right-sided derivative in  $x$  and  $g'_-(x) = g'_+(x) (= g'(x))$ .

#### 4.4 Fundamental theorem of differential/integration calculus

**Theorem 4.3** (Fundamental theorem of differential/integration calculus, variant 1). *Isaac Barrow (1630–1677), Isaac Newton (1642–1726), Gottfried Wilhelm von Leibniz (1646–1716).*

Let  $f \in \mathcal{R}[a, b]$ ,  $\alpha \in [a, b]$  and we define

$$F(x) := \int_{\alpha}^x f(\xi) d\xi$$

Then  $F$  is right-sided differentiable in every point  $x \in [a, b)$  and left-sided differentiable

in every  $x \in (a, b]$ . Furthermore

$$F'_+(x) = f_+(x) \forall x \in [a, b) \quad (2)$$

$$F'_-(x) = f_-(x) \forall x \in (a, b] \quad (3)$$

**Remark 4.6.**

$$\frac{d}{dx} \left( \int_a^x f d\xi \right) = f(x)$$

for all  $x$  such that  $f$  is continuous in  $x$ . For those  $x$ ,  $F'(x)$  is differentiable in  $x$  with  $F'(x) = f(x)$ .

**Definition 4.3.** Let  $f \in \mathcal{R}[a, b]$  and  $\Phi : [a, b] \rightarrow \mathbb{R}$  such that  $\Phi$  is one-sided differentiable on  $[a, b]$ . If  $\Phi'_+(x) = f_+(x) \forall x \in [a, b)$  and  $\Phi'_-(x) = f_-(x) \forall x \in (a, b]$  then we call  $\Phi$  an antiderivative of regulated function  $f$ .

*Proof of the Theorem 4.3.* Let  $x_1, x_2 \in [a, b]$  be arbitrary. Let  $F$  be defined as above. Then

$$\begin{aligned} |F(x_2) - F(x_1)| &= \left| \int_a^{x_2} f d\xi - \int_a^{x_1} f d\xi \right| = \left| \int_a^{x_2} f d\xi + \int_{x_1}^a f d\xi \right| = \left| \int_{x_1}^{x_2} f d\xi \right| \\ &\leq \int_{x_1}^{x_2} |f| d\xi \leq \int_{x_1}^{x_2} \underbrace{\|f\|_\infty}_{\text{const independent of } \xi} d\xi = \|f\|_\infty \cdot |x_2 - x_1| \end{aligned}$$

Hence  $F$  is Lipschitz continuous with Lipschitz constant  $\|f\|_\infty$ . So  $F$  is continuous in  $[a, b]$ .

One-sided derivatives: Let  $x \in [a, b)$  and  $\varepsilon > 0$  be arbitrary. Choose  $\delta > 0$  such that  $\forall \xi \in [x, x + \delta)$  it holds that  $|f(\xi) - f_+(x)| < \varepsilon$ . For  $\xi \in (x, x + \delta)$ ,

$$\begin{aligned} \left| \frac{F(\xi) - F(x)}{\xi - x} - f_+(x) \right| &= \frac{1}{|\xi - x|} \left| \underbrace{\int_x^\xi f dy}_{F(\xi) - F(x)} - \underbrace{f_+(x)(\xi - x)}_{\int_x^\xi \underbrace{f_+(x)}_{\text{const.}} dy} \right| \\ &= \frac{1}{|\xi - x|} \left| \int_x^\xi (f - f_+(x)) dy \right| \leq \frac{1}{|\xi - x|} \int_x^\xi \underbrace{|f(y) - f_+(x)|}_{< \varepsilon} dy \\ &\quad y \in (x, \xi) \subseteq (x, x + \delta) \end{aligned}$$

$$< \frac{1}{\xi - x} \varepsilon \cdot \underbrace{\int_x^\xi 1 \, dy}_{|\xi - x|} = \varepsilon$$

Hence,  $F'_+(x) = f_+(x)$ . Analogously,  $F'_-(x) = f_-(x)$  for  $x \in (a, b]$ .  $\square$

**Theorem 4.4** (Fundamental theorem of differential/integration calculus, variant 2). *Let  $f \in \mathcal{R}[a, b]$  and  $\phi$  is an arbitrary antiderivative of  $f$  according to Definition 4.3. For  $\alpha, \beta \in [a, b]$  arbitrary,*

$$\int_\alpha^\beta f \, dx = \phi(\beta) - \phi(\alpha)$$

*Proof.* The proof is given in the next lecture.  $\square$

**Remark 4.7.** *Let  $f$  be continuous and  $\phi$  be an antiderivative of  $f$ . Hence,  $\Phi'(x) = f(x) \forall x \in [a, b]$ . Then*

$$\int_\alpha^\beta \Phi' \, dx = \Phi(\beta) - \Phi(\alpha)$$

*“Integral of a derivative of  $\Phi$  gives  $\Phi(\beta) - \Phi(\alpha)$ ”.*

**Lemma 4.4.** *Let  $A \subseteq [a, b]$  countable.  $f : [a, b] \rightarrow \mathbb{R}$  is continuous and  $f$  is differentiable in every point  $x \in [a, b] \setminus A$ . Furthermore let  $|f'(x)| \leq L$  ( $L \geq 0$ ) for all  $x \in [a, b] \setminus A$ . Then  $f$  is Lipschitz continuous on  $[a, b]$  with constant  $L$ , hence*

$$|f(x_2) - f(x_1)| \leq L |x_2 - x_1| \quad \forall x_1, x_2 \in [a, b]$$

**Remark 4.8.** *Some people call it differentiable almost everywhere, but this expression collides with a different definition pronounced the same way from measure theory.*

*Proof.* Let  $x_1, x_2 \in [a, b]$ , without loss of generality:  $x_1 < x_2$ . Let  $\varepsilon > 0$  be arbitrary. We define

$$F_\varepsilon(x) := |f(x) - f(x_1)| - (L + \varepsilon)(x - x_1)$$

for  $x \in [x_1, b]$ .

Let  $\varepsilon > 0$  be arbitrary. We prove:  $F_\varepsilon(x) \leq 0 \forall x \in [x_1, b]$ . In particular:  $F_\varepsilon(x_2) \leq 0$ . Hence,

$$|f(x_2) - f(x_1)| \leq (L + \varepsilon) \underbrace{(x_2 - x_1)}_{|x_2 - x_1|}$$

We prove by contradiction: Assume there exists  $\varepsilon > 0$  and  $x_\varepsilon > x_1$  such that

$$F_\varepsilon(x_\varepsilon) = \eta > 0 \tag{4}$$

We recognize: Let  $A' = [x_1, b] \cap A$  be countable.

1. hence  $F_\varepsilon(A') \subseteq \mathbb{R}$  is countable
2.  $F_\varepsilon(x_1) = 0, F_\varepsilon(x_\varepsilon) > 0 \implies x_\varepsilon > x_1$
3.  $F_\varepsilon$  is continuous on  $[x_1, b]$ . It holds that  $0 \in F_\varepsilon([x_1, x_\varepsilon])$  because  $0 = F_\varepsilon(x_1)$  and  $\eta \in F_\varepsilon([x_1, x_\varepsilon])$  because  $\eta = F_\varepsilon(x_\varepsilon)$ .

By the Intermediate Value Theorem, it follows that  $[0, \eta] \subseteq F_\varepsilon([x_1, x_\varepsilon])$  where  $[0, \eta]$  is uncountable.  $F_\varepsilon(A')$  is countable, hence there exists  $\gamma \in (0, \eta]$  such that  $\gamma = F_\varepsilon(y)$  and  $\gamma \notin A'$  ( $\gamma > 0$ ); compare with inequality (4). Hence,  $y \notin A'$ . So  $f$  in  $y$  is differentiable. Let  $B := F_\varepsilon^{-1}(\{\gamma\}) \cap ([x_1, x_\varepsilon] \setminus A')$ , where we can skip  $A'$ . Then  $B \neq \emptyset$ .

$B \subseteq [x_1, x_\varepsilon]$  is therefore bounded,  $B \neq \emptyset$ . Hence,  $B$  has a supremum. Let  $x = \sup B$ . Choose  $(y_n)_{n \in \mathbb{N}}$  with  $y_n \in B$  and  $y_n \rightarrow x$  for  $n \rightarrow \infty$ . Because  $F_\varepsilon$  is continuous,

$$\lim_{n \rightarrow \infty} \underbrace{F_\varepsilon(y_n)}_\gamma = F_\varepsilon(x)$$

hence  $F_\varepsilon(x) = \gamma$ . This implies  $x \notin A$ .

Furthermore it holds for  $w \in (x, x_\varepsilon]$  that  $F_\varepsilon(w) > \gamma$ . Because assume the opposite ( $F_\varepsilon(w) \leq \gamma$  for  $w > x$ ). Furthermore,  $F_\varepsilon(x_\varepsilon) = \eta \geq \gamma$ . Because of the Intermediate Value Theorem,  $\exists y \geq w$  with  $F_\varepsilon(y) = \gamma$ . This contradicts with the supremum property of  $x$ .

Now let  $y \in (x, x_\varepsilon]$ .

$$\begin{aligned} \varphi(y) &= \frac{F_\varepsilon(y) - F_\varepsilon(x)}{y - x} \\ &= \underbrace{\frac{|f(y) - f(x_1)| - |f(x) - f(x_1)|}{y - x}}_{\substack{\text{definition of} \\ F_\varepsilon}} - \frac{(L + \varepsilon)(y - x_1 - x + x_1)}{y - x} \\ &\leq \underbrace{\frac{f(y) - f(x)}{y - x}}_{\substack{\text{reversed triangle ineq.}}} - (L + \varepsilon) \end{aligned}$$

Because  $F_\varepsilon(y) > \gamma = F_\varepsilon(x)$ ,  $\varphi(y) > 0$  for  $y > x$ . So,

$$\frac{|f(y) - f(x)|}{y - x} \geq L + \varepsilon$$

$$|f'(x)| = \lim_{y \rightarrow x+} \left| \frac{f(y) - f(x)}{y - x} \right| \geq L + \varepsilon$$

This contradicts with the boundedness of the derivative by  $L$  and  $f$  is in  $x \notin A$  differentiable.



So, inequality (4) does not hold. Therefore  $\forall x_1, x_2$  with  $x_1 < x_2$  in  $[a, b]$  and  $\forall \varepsilon > 0$ ,

$$\begin{aligned} |f(x_2) - f(x_1)| &\leq (L + \varepsilon) |x_2 - x_1| \\ \implies |f(x_2) - f(x_1)| &\leq L |x_2 - x_1| \end{aligned}$$

□

**Corollary** (Corollary to Lemma 4.4). *Let  $f, g : [a, b] \rightarrow \mathbb{R}$  differentiable for all points  $x \in [a, b] \setminus A$  and  $A$  is countable. Furthermore let  $f'(x) = g'(x) \forall x \notin A$ . Then there exists  $K \in \mathbb{R}$  such that  $f(x) = g(x) + K \forall x \in [a, b]$ .*

*Proof.* Let  $h = f - g$ . Then

$$h'(x) = f'(x) - g'(x) = 0 \forall x \in [a, b] \setminus A$$

By Lemma 4.4 with  $L = 0$ , it follows that

$$\begin{aligned} |h(x_1) - h(x_2)| &\leq 0 \cdot |x_1 - x_2| = 0 \\ \implies h(x_1) &= h(x_2) \forall x_1, x_2 \in [a, b] \end{aligned}$$

Hence,  $h(x) = K \in \mathbb{R}$ .

$$\implies f(x) = g(x) + h(x) = g(x) + K$$

□

↓ This lecture took place on 2018/04/19.

By reference (\*),  $\gamma \in [0, \eta]$  (uncountable) and  $\gamma \notin f(A)$  (countable).

$$\implies \forall u \in [x_1, b] \text{ with } F_\varepsilon(u) = \gamma$$

it holds that  $u \notin A$ , hence  $f$  is differentiable in  $u$ .

*Proof of Theorem 4.4.* Let  $f \in \mathcal{R}[a, b]$  and let  $\phi$  be an antiderivative of  $f$ , hence  $\phi'_+ = f_+$ ,  $\phi'_- = f_-$ . Let  $\alpha \in [a, b]$  be arbitrary. By the Theorem variant 1,  $F(x) = \int_\alpha^x f d\xi$  is also an antiderivative of  $f$ . By Lemma 4.4,  $\exists K \in \mathbb{R} : F(x) = \int_\alpha^x f d\xi = \phi(x) + K$ . Determine  $K$ : Let  $x = \alpha \implies F(\alpha) = \int_\alpha^\alpha f dx = 0 = \phi(\alpha) - K$  hence  $K = \phi(\alpha)$ . Hence,

$$\int_\alpha^x f d\xi = \phi(x) - \phi(\alpha)$$

Let  $x = \beta$ .

□

**Remark 4.9** (Remark for the previous corollary).  $F, \phi$  are differentiable on all points  $x$  for which  $f$  is continuous (all of them except for countable many). For those  $x$ ,  $F'(x) = \phi'(x) = f(x)$ .

**Remark 4.10** (Notation). Let  $f \in \mathcal{R}[a, b]$ . Then

$$\int f \, dx$$

- is some particular antiderivative of  $f$  (usually some arbitrary chosen)
- the set of all antiderivatives of  $f$

$$\int f \, dx = \{F : F \text{ is antiderivative of } f\}$$

If  $F_0$  is some fixed antiderivative, then

$$\int f \, dx = \{F_0 + K : K \in \mathbb{R}\}$$

Then  $\int f \, dx$  is the so-called indefinite integral of  $f$ . Notation:

$$\int x^k \, dx = \frac{x^{k+1}}{k+1} + c \quad (k \neq -1)$$

$f$	$F$	remark
$x^\alpha$	$\frac{x^{\alpha+1}}{\alpha+1} + c$	$\alpha \in \mathbb{R} \setminus \{-1\}$ ; restrict $x$ such that $x^\alpha$ and $x^{\alpha+1}$ are defined
$x^{-1}$	$\ln x + c \, (x > 0)$	
$\left(\frac{1}{-x}\right) \cdot (-1) = x^{-1}$	$\ln -x + c \, (x < 0)$	
$e^x$	$e^x$	
$\sin x$	$-\cos x$	
$\cos x$	$\sin x$	
$\sinh x$	$\cosh x$	
$\cosh x$	$\sinh x$	
$\frac{1}{1+x^2}$	$\arctan x$	$ x  < 1$
$\frac{1}{\sqrt{1-x^2}}$	$\arcsin x$	
$-\frac{1}{\sqrt{1-x^2}}$	$\arccos x$	

Table 1: Table of antiderivatives

## 4.5 Integration methods

In this chapter, we discuss how to determine the antiderivative of a function. Usually they are composites of basic functions. Some of these are given in Table 1.

**Remark 4.11.** Let  $F, G : [a, b] \rightarrow \mathbb{R}$  in  $x \in [a, b)$  right-sided differentiable. Then also  $F \cdot G$  in  $x$  is right-sided differentiable and

$$(F \cdot G)'_+(x) = F'_+(x) \cdot G(x) + F(x) \cdot G'_+(x)$$

hence the product law holds.

Analogously, the same holds for the left-sided derivative.

Look up the proof in the course Analysis 1.

### 4.5.1 Partial integration

**Definition 4.4** (Partial integration). Let  $f, g$  be given. Let  $F, G$  be its antiderivatives respectively. Then  $F \cdot G$  is an antiderivative of  $F \cdot g + f \cdot G$ .

This is immediate, because

$$(F \cdot G)'_+ = F'_+ \cdot G + F \cdot G'_+ = f_+ \cdot G + F \cdot g_+ = f_+ G_+ + F_+ \cdot g_+$$

Hence,

$$\int_a^b (Fg + fG) dx = \underbrace{F(b) \cdot G(b) - F(a)G(a)}_{=: F \cdot G|_a^b}$$

Usually, this is rewritten as

$$\int_a^b F \cdot g dx = F \cdot G|_a^b - \int_a^b fG dx$$

If  $F = u$  is continuously differentiable and  $G = v$  as well, then  $f = u'$  and  $g = v'$  and the law has the structure

$$\int_a^b uv' dx = u \cdot v|_a^b - \int_a^b u'v dx$$

**Example 4.1.** Let  $a \neq -1$  and  $x > 0$ .

$$\int \underbrace{x^a}_{v'} \cdot \underbrace{\ln x}_u dx = \underbrace{\left[ \begin{array}{ll} u = \ln x & u' = \frac{1}{x} \\ v' = x^a & v = \frac{x^{a+1}}{a+1} \end{array} \right]}_{\text{scribble notes}} \frac{x^{a+1}}{a+1} \cdot \ln x - \int \frac{1}{x} \cdot \frac{x^{a+1}}{a+1} dx$$

$$= \frac{x^{\alpha+1}}{\alpha+1} \cdot \ln x - \frac{1}{\alpha+1} \int x^\alpha dx = \frac{x^{\alpha+1}}{\alpha+1} \cdot \ln x - \frac{1}{(\alpha+1)^2} x^{\alpha+1}$$

**Example 4.2.** Let  $k \in \{2, 3, 4, \dots\}$ .

$$\int \cos^k(x) dx = \left| \begin{array}{ll} u = \cos^{k-1}(x) & u' = (k-1) \cdot \cos^{k-2}(x) \cdot (-\sin x) \\ v' = \cos x & v = \sin x \end{array} \right|$$

$$\cos^{k-1}(x) \sin x + (k-1) \int \cos^{k-2}(x) \cdot \underbrace{\sin^2(x)}_{(1-\cos^2 x)} dx$$

$$= \cos^{k-1}(x) \cdot \sin(x) + (k-1) \int \cos^{k-2}(x) dx - (k-1) \int \cos^k(x) dx$$

Then we can use the following identity:

$$k \int \cos^k(x) dx = \cos^{k-1}(x) \cdot \sin(x) + (k-1) \int \cos^{k-2}(x) dx$$

This gives a recursive formula:

$$\int \cos^k(x) dx = \frac{1}{k} \cos^{k-1}(x) \cdot \frac{k-1}{k} \sin(x) + (k-1) \int \cos^{k-2}(x) dx$$

Analogously,

$$\int \sin^k(x) dx = -\frac{1}{k} \sin^{k-1}(x) \cdot \cos(x) + \frac{k-1}{k} \int \sin^{k-2}(x) dx$$

Let  $c_m = \int_0^{\frac{\pi}{2}} \cos^m(x) dx$ . Then the following formula holds:

$$\begin{aligned} c_{2n} &= \frac{2n-1}{2n} \cdot \frac{2n-3}{2n-2} \cdot \frac{2n-5}{2n-4} \cdots \frac{1}{2} \cdot \frac{\pi}{2} \\ &= \prod_{k=1}^n \frac{2k-1}{2k} \cdot \frac{\pi}{2} \\ c_{2n+1} &= \prod_{k=1}^n \frac{2k}{2k+1} \end{aligned}$$

*Proof by induction.* Let  $n = 1$ .

$$\begin{aligned}
c_2 &= \int_0^{\frac{\pi}{2}} \cos^2 x \, dx = \frac{1}{2} \cos x \sin x \Big|_0^{\frac{\pi}{2}} + \frac{1}{2} \int_0^{\frac{\pi}{2}} 1 \, dx = 0 - 0 + \frac{\pi}{4} \\
&= \underbrace{\prod_{k=1}^1 \frac{2k-1}{2k}}_{\frac{1}{2}} \cdot \frac{\pi}{2} \\
c_1 &= \int_0^{\frac{\pi}{2}} \cos x \, dx = \sin x \Big|_0^{\frac{\pi}{2}} = 1 - 0 = 1 \\
\underbrace{\prod_{k=1}^0 \frac{2k}{2k+1}}_{\text{empty product}} &= 1
\end{aligned}$$

We make the induction step  $n \rightarrow n+1$ :

$$\begin{aligned}
c_{2(n+1)} &= \frac{1}{2n+2} \cdot \underbrace{\cos^{2n+1}(x)}_{=0 \text{ for } x=\frac{\pi}{2}} \cdot \underbrace{\sin(x)}_{=0 \text{ for } x=0} \Big|_0^{\frac{\pi}{2}} + \frac{2n+1}{2n+2} \int_0^{\frac{\pi}{2}} \cos^{2n}(x) \, dx \\
&= \frac{2n+1}{2n+2} \prod_{k=1}^n \frac{2k-1}{2k} \cdot \frac{\pi}{2} = \prod_{k=1}^{n+1} \frac{2k-1}{2k} \cdot \frac{\pi}{2}
\end{aligned}$$

$c_{2(n+1)+1}$  analogously. □

**Theorem 4.5** (Wallis product). *John Wallis (1616–1703), result from 1655*

Let  $w_n = \prod_{k=1}^n \frac{(2k)^2}{(2k-1)(2k+1)} = \frac{2 \cdot 2}{1 \cdot 3} \cdot \frac{4 \cdot 4}{3 \cdot 5} \dots$ . Then  $\lim_{n \rightarrow \infty} w_n = \frac{\pi}{2}$ .

*Proof.*

$$\frac{\pi}{2} \cdot \frac{c_{2n+1}}{c_{2n}} = \frac{\pi}{2} \cdot \prod_{k=1}^n \frac{\frac{2k}{2k+1}}{\prod_{k=1}^n \frac{2k-1}{2k} \cdot \frac{\pi}{2}} = \prod_{k=1}^n \frac{(2k)^2}{(2k-1)(2k+1)} = w_n$$

It remains to show that  $\lim_{n \rightarrow \infty} \frac{c_{2n+1}}{c_{2n}} = 1$  in  $[0, \frac{\pi}{2}]$  it holds that  $0 \leq \cos x \leq 1$ .

$$\implies \cos^{2n+2}(x) \leq \cos^{2n+1}(x) \leq \cos^{2n}(x)$$

So,  $c_{2n+2} \leq c_{2n+1} \leq c_{2n}$  for  $n \geq 1$ .

$$1 \geq \frac{c_{2n+1}}{c_{2n}}$$

$$\begin{aligned} \implies 1 &\geq \frac{c_{2n+1}}{c_{2n}} \geq \frac{c_{2n+2}}{c_{2n}} = \frac{\prod_{k=1}^{n+1} \frac{2k-1}{2k} \frac{\pi}{2}}{\prod_{k=1}^n \frac{2k-1}{2k} \frac{\pi}{2}} \\ &= \frac{2n+2-1}{2n+2} \rightarrow 1 \text{ for } n \rightarrow \infty \end{aligned}$$

Because of the sandwich lemma for convergent sequences, the intermediate expression must also converge to 1, hence

$$\lim_{n \rightarrow \infty} \frac{c_{2n+1}}{c_{2n}} = 1 \quad \wedge \quad \frac{\pi}{2} \cdot \lim_{n \rightarrow \infty} \frac{c_{2n+1}}{c_{2n}} = \underbrace{\lim_{n \rightarrow \infty} w_n}_{=1}$$

□

#### 4.5.2 Integration by substitution

**Definition 4.5** (Integration by substitution). Let  $f : [a, b] \rightarrow \mathbb{R}$  be continuous. Let  $t : [\alpha, \beta] \rightarrow [a, b]$  be continuously differentiable. Let  $F$  be an antiderivative of  $f$  ( $F$  is therefore continuously differentiable). Then  $F \circ t : [\alpha, \beta] \rightarrow \mathbb{R}$  is also continuously differentiable and the chain rule holds:

$$(F \circ t)' = (F' \circ t) \cdot t' = (f \circ t) \cdot t'$$

Hence  $F \circ t$  is an antiderivative of  $(f \circ t) \cdot t'$ . We apply it to integration:

$$\int_{\alpha}^{\beta} (f \circ t)(u) \cdot t'(u) du = (F \circ t)(\beta) - (F \circ t)(\alpha) = F(t(\beta)) - F(t(\alpha)) = \int_{t(\alpha)}^{t(\beta)} f(x) dx$$

Then we get the substitution integration method:

$$\int_{t(\alpha)}^{t(\beta)} f(x) dx = \int_{\alpha}^{\beta} f(t(u)) \cdot t'(u) du$$

**Remark 4.12** (Mnemonic). Consider the left-hand side and right-hand side simultaneously. Let  $x = t(u)$  (expressions inside parentheses). Then  $dx = t'(u) \cdot du$  (expressions on the right). Let  $u = \alpha \implies x = t(\alpha)$  and  $u = \beta \implies x = t(\beta)$  (interval boundaries).

**Example 4.3.**

$$\int_0^1 2x \sqrt{1-x^2} dx$$

Usually we have some expression, we want to substitute with  $u$ .

$$1 - x^2 = u \quad x = \sqrt{1-u} = t(u)$$

$$\begin{aligned}
x = 0 &= t(1) & x = 1 &= t(0) \\
dx &= \frac{1}{2} \cdot \frac{1}{\sqrt{1-u}} \cdot (-1) du \\
\int_0^1 2x \sqrt{1-x^2} dx &= \int_1^0 2 \cdot \sqrt{1-u} \cdot u \cdot \frac{1}{2} (-1) \frac{1}{\sqrt{1-u}} du = \int_0^1 \sqrt{u} du = \left. \frac{u^{\frac{3}{2}}}{\frac{3}{2}} \right|_0^1 = \frac{2}{3}
\end{aligned}$$

$$\int_0^1 2x \sqrt{\underbrace{1-x^2}_u} dx = \left| \begin{array}{ll} u = 1-x^2 & \\ x = 0 & \Leftrightarrow u = 1 \\ x = 1 & \Leftrightarrow xu = 0 \\ 1 \cdot du & = -2x dx \end{array} \right| = - \int_1^0 \sqrt{u} du = \int_0^1 \sqrt{u} du$$

In general: we set  $h(u) = g(x)$ , then  $h'(u) du = g'(x) dx$ .

**Theorem 4.6.** Let  $f, \tilde{f} \in \mathcal{R}[a, b]$  and  $A \subseteq [a, b]$  countable. Furthermore  $f(x) = \tilde{f}(x) \forall x \in [a, b] \setminus A$ . Then

$$\int_a^b |f - \tilde{f}| dx = 0$$

Then it follows especially that

$$\int_a^b f dx = \int_a^b \tilde{f} dx$$

↓ This lecture took place on 2018/04/24.

*Proof.* Show:  $r \in \mathcal{R}[a, b], r \geq 0$ .  $\int_a^b r dx = 0$  and  $r(x) = 0$  for  $x \in [a, b] \setminus A$ . Then  $\int_a^b r dx = 0$ . Let  $r$  be as above. First, we show:  $r_+(x) = \lim_{\xi \rightarrow x^+} r(\xi) = 0 \forall x \in [a, b]$  and also  $r_-(x) = 0 \forall x \in (a, b]$ .

Proof of that: Let  $x \in [a, b]$  and  $y = r_+(x)$  (exists because  $r \in \mathcal{R}[a, b]$ ). Choose  $\delta_n = \frac{1}{n}$ .  $(x, x + \frac{1}{n}) \cap [a, b]$  is an open interval with uncountable many points, so there is certainly one point in  $A$ . So there exists  $\xi_n \in ((x, x + \frac{1}{n}) \cap [a, b]) \setminus A$  and  $|\xi_n - x| < \delta_n = \frac{1}{n}$ . Hence,  $\lim_{n \rightarrow \infty} \xi_n = x$  and  $r(\xi_n) = 0$ . Therefore,  $\lim_{n \rightarrow \infty} r(\xi_n) = 0$  where  $r(\xi_n) = y = r_+(x)$ .

Analogously,  $r_-(x) = 0$  on  $(a, b]$ .

Let  $\varepsilon > 0$  be arbitrary. We let  $A_\varepsilon = \{w \in [a, b] \mid r(w) > \varepsilon\}$ . We show:  $A_\varepsilon$  is finite.

Assume  $A_\varepsilon$  would have infinitely many points. Choose a sequence  $(w_n)_{n \in \mathbb{N}}$  with  $w_n \in A_\varepsilon$  and  $w_n \neq w_m$  for  $n \neq m$  (works because  $A_\varepsilon$  is infinite).  $(w_n)_{n \in \mathbb{N}}$  is bounded, hence there exists a convergent subsequence  $(w_{n_k})_{k \in \mathbb{N}}$  with  $x = \lim_{k \rightarrow \infty} w_{n_k} \in [a, b]$  and  $w_{n_k} \in [a, b]$ .

Either  $(w_{n_k})$  contains infinitely many sequence element  $w_{n_k} < x$  (variant (a)) or infinitely many  $w_{n_k} > x$  (variant (b)). Let variant b hold without loss of generality.

Combine all  $w_{n_k} > x$  to one subsequence  $(w_{n_{k_l}})_{l \in \mathbb{N}}$ . This gives  $\lim_{l \rightarrow \infty} w_{n_{k_l}} = x$  and  $w_{n_{k_l}} > x$ , thus  $\lim_{l \rightarrow \infty} \underbrace{r(w_{n_{k_l}})}_{\geq \varepsilon \text{ because } w_{n_{k_l}} \in A_\varepsilon} = r_+(x) = 0$ . This gives a contradiction.

$A_\varepsilon$  must be finite.

Consider

$$A_{\frac{1}{n}} = \{w_1^n, \dots, w_{m_n}^n\}$$

finite. Let  $\varphi_n = \sum_{k=1}^{m_n} r(w_k^n) \cdot \chi_{\{w_k^n\}} \in \tau[a, b]$ .

For  $x = w_k^n \in A_{\frac{1}{n}}$

$$\varphi_n(w_k^n) = \sum_{k=1}^{m_n} r(w_k^n) \cdot \underbrace{\chi_{\{w_k^n\}}(w_j^n)}_{\delta_{jk}} = r(w_j^n)$$

so  $|\varphi_n(x) - r(x)| = 0 \forall x \in A_{\frac{1}{n}}$ . Let  $x \in [a, b] \setminus A_{\frac{1}{n}}$ . Then it holds  $0 \leq r(x) < \frac{1}{n}$  and for  $x \notin A_{\frac{1}{n}}$ ,  $\varphi(x) = 0$ . Therefore,

$$|r(x) - \varphi(x)| = r(x) < \frac{1}{n}$$

hence  $\|r - \varphi_n\|_\infty < \frac{1}{n}$ . This means that  $\varphi_n \rightarrow r$  uniformly on  $[a, b]$ . Therefore

$$\lim_{n \rightarrow \infty} \underbrace{\int_a^b \varphi_n dx}_{=0} = \int_a^b r dx = 0$$

Now we want to finish the proof of our theorem: Let  $r(x) = |f(x) - \tilde{f}(x)| \geq 0$  and  $r(x) = 0$  for  $x \notin A$ . So,  $\int_a^b |f - \tilde{f}| dx = 0$  (first part proven).

$$\begin{aligned} \left| \int_a^b f dx - \int_a^b \tilde{f} dx \right| &= \left| \int_a^b (f - \tilde{f}) dx \right| \leq \int_a^b |f - \tilde{f}| dx = 0 \\ \implies \int_a^b f dx &= \int_a^b \tilde{f} dx \end{aligned}$$

Second part proven. □

**Lemma 4.5.** Let  $f \in \mathcal{R}[a, b]$ . Then  $f_+ \in \mathcal{R}[a, b]$  and also  $f_- \in \mathcal{R}[a, b]$ .






Figure 23:  $\xi$  must be sufficiently close enough to  $z$  such that  $|f(\xi) - f_+(z)| \leq \frac{\varepsilon}{2}$ .




Figure 24:  $\xi$  and  $z$

*Proof.* Only for  $f_+$ : First, we show: Let  $x \in [a, b)$ .

$$f_+(x) = \lim_{\xi \rightarrow x_+} f(\xi) = \lim_{\xi \rightarrow x_+} f_+(x)$$

(the plus is important on the right-hand side!).

Proof of this: Let  $\varepsilon > 0$  be arbitrary. Then there exists  $\delta > 0$  such that  $\forall \xi \in (x, x + \delta)$ :  $|f(\xi) - f_+(x)| < \frac{\varepsilon}{2}$ . Now let  $z \in (x, x + \delta)$  be arbitrary chosen. For  $z$  there exists  $\xi \in (z, x + \delta)$  because  $f_+(z)$  exists. Compare with Figure 23.

$$|f_+(z) - f_+(x)| \leq |f_+(z) - f(\xi)| + |f(\xi) - f_+(x)| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2}$$

so  $f_+(x) = \lim_{z \rightarrow x_+} f_+(z)$ .

It remains to show:  $f_+$  has left-sided limits. Let  $x \in (a, b]$  be arbitrary and  $f_-(x) = \lim_{\xi \rightarrow x_-} f(\xi)$ . We show:  $f_-(x) = \lim_{\xi \rightarrow x_-} f_+(x)$  (again: the plus is important).

Let  $\varepsilon > 0$  be arbitrary. Choose  $\delta > 0$  such that  $\forall z \in (x - \delta, x)$ ,  $|f(z) - f_-(x)| < \frac{\varepsilon}{2}$ .

Now let  $\xi \in (x - \delta, x)$  (compare with Figure 24) and choose  $x > z > \xi$  with the property that  $|f(z) - f_+(\xi)| < \frac{\varepsilon}{2}$  (feasible because  $f$  in  $\xi$  has a right-sided limit):

$$|f_+(\xi) - f_-(x)| \leq \underbrace{|f_+(\xi) - f(z)|}_{< \frac{\varepsilon}{2}} + \underbrace{|f(z) - f_-(x)|}_{< \frac{\varepsilon}{2}}$$

because of the choice of  $\delta$  and  $z \in (\xi, x) \subseteq (x - \delta, x)$ .

Hence,  $\lim_{\xi \rightarrow x_-} f_+(\xi) = f_-(x)$ . Analogously for  $f_-$

□

**Remark 4.13.**

$$\lim_{\xi \rightarrow x_+} f_+(\xi) = f_+(x)$$

$$\lim_{\xi \rightarrow x_-} f_-(\xi) = f_-(x)$$

from the proof. So  $f_+$  is right-sided continuous and  $f_-$  is left-sided continuous.

**Lemma 4.6.** Let  $f \in \mathcal{R}[a, b]$ . Then

$$\int_a^b f \, dx = \int_a^b f_+ \, dx = \int_a^b f_- \, dx$$

*Proof.* For  $f_+$ :

$$f, f_+ \in \mathcal{R}[a, b]$$

$\forall x \in [a, b]$  with  $f$  is continuous in  $x$ ,

$$f(x) = \lim_{\xi \rightarrow x} f(\xi) = \lim_{\xi \rightarrow x_+} f(\xi) = f_+(x)$$

$f$  has at most countable many discontinuity points. By Satz 4.6,

$$\int_a^b |f - f_+| \, dx = 0 \quad \text{and accordingly} \quad \int_a^b f \, dx = \int_a^b f_+ \, dx$$

□

## 4.6 Improper integrals

Let  $I$  be an interval in  $\mathbb{R}$  with marginal points  $a$  and  $b$  with  $-\infty \leq a < b \leq +\infty$ . Let  $f$  be a regulated function on  $I$ . We define

1. If  $I = [a, b)$ ,  $\int_a^b f \, dx = \lim_{\beta \rightarrow b_-} \int_a^\beta f \, dx$
2. If  $I = (a, b]$ ,  $\int_a^b f \, dx = \lim_{\alpha \rightarrow a_+} \int_\alpha^b f \, dx$
3. If  $I = (a, b)$ ,  $\int_a^b f \, dx = \lim_{\alpha \rightarrow a_+} \int_\alpha^c f \, dx + \lim_{\beta \rightarrow b_-} \int_c^\beta f \, dx$

for an arbitrarily chosen  $c \in (a, b)$  under the constraint that the corresponding limits in  $\mathbb{R}$  exist.

Standard examples will follow:




Figure 25: The function  $\frac{1}{x^s}$  for  $s > 1$

**Example 4.4.** Let  $s > 1$ .

$$\begin{aligned} \int_1^\infty x^{-s} dx &= \lim_{\beta \rightarrow \infty} \int_1^\beta x^{-s} dx = \lim_{\beta \rightarrow \infty} \left( \frac{1}{-s+1} x^{-s+1} \right) \Big|_1^\beta \\ &= \frac{1}{1-s} \cdot \underbrace{\lim_{\beta \rightarrow \infty} \frac{1}{s-1}}_{=0} - \frac{1}{1-s} \cdot 1 = \frac{1}{s-1} \end{aligned}$$

Compare with Figure 25.

**Example 4.5.** Let  $s < 1$ .

$$\begin{aligned} \int_0^1 x^{-s} dx &= \lim_{\alpha \rightarrow 0^+} \int_\alpha^1 x^{-s} dx = \lim_{\alpha \rightarrow 0^+} \frac{1}{-s+1} x^{-s+1} \Big|_\alpha^1 \\ &= \frac{1}{1-s} - \frac{1}{1-s} \cdot \underbrace{\lim_{\alpha \rightarrow 0^+} \alpha^{1-s}}_{=0} = \frac{1}{1-s} \end{aligned}$$

Compare with Figure 26.

For  $s = 1$ , neither  $\int_0^1 \frac{1}{x} dx$  nor  $\int_1^\infty \frac{1}{x} dx$  exists.

**Example 4.6.** For  $c > 0$ ,

$$\int_0^\infty e^{-cx} dx = \lim_{\beta \rightarrow \infty} \int_0^\beta e^{-cx} dx = \lim_{\beta \rightarrow \infty} \left( -\frac{1}{c} \right) \cdot e^{-cx} \Big|_0^\beta - \frac{1}{c} \cdot \underbrace{\lim_{\beta \rightarrow \infty} e^{-c\beta}}_{=0} + \frac{1}{c} = \frac{1}{c}$$




Figure 26: The function  $x^s$  for  $s < 1$

**Theorem 4.7** (Direct comparison test for improper integrals). *In German, “Majorantenkriterium für uneigentliche Integrale”.*

Let  $f, g$  be regulated functions on  $I$  and

$$|f(x)| \leq g(x) \forall x \in I$$

Assume  $\int_a^b g \, dx$  exists as improper integral. Then also the following improper integrals exist:

$$\int_a^b |f| \, dx \text{ and } \int_a^b f \, dx$$

In German,  $g$  is called Majorante of  $f$  (there is no equivalent terminology in English).

*Proof.* Without loss of generality, let  $I = [a, b)$ . Let  $G(\beta) = \int_a^\beta g \, dx$ . We know that  $\lim_{\beta \rightarrow b^-} G(\beta)$  exists. By Lemma 3.6 (Cauchy criterion for existence of limits): Let  $\varepsilon > 0$  be arbitrary, then there exists a left-sided neighborhood  $U$  of  $b$  ( $U = (b - \delta, b)$  if  $b < \infty$  and  $U = (M, \infty)$  if  $b = \infty$ ) with  $u, v \in U$ , then  $|G(v) - G(u)| < \varepsilon$ .

$$|G(v) - G(u)| = \left| \int_a^v g \, dx - \int_a^u g \, dx \right| = \left| \int_u^v g \, dx \right|$$

Let  $F(\beta) = \int_a^\beta |f| \, dx$ . Analogously as for  $G$ ,  $F(v) - F(u) = \int_u^v |f| \, dx$ . Let  $u, v \in U$ . Then

$$|F(v) - F(u)| = \left| \int_u^v |f| \, dx \right| \leq \left| \int_u^v g \, dx \right| = |G(v) - G(u)| < \varepsilon$$

hence by the Cauchy criterion for  $F$ :  $\lim_{\beta \rightarrow b^-} F(\beta)$  exists, so there exists  $\int_a^b |f| \, dx$  as improper integral. The same applies for the existence of  $\int_a^b f \, dx$ .  $\square$

**Example 4.7.** The cardinal sine function is defined as

$$\text{sinc}(x) = \frac{\sin x}{x}$$

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1 \quad \text{sinc}(0) = 1$$

So  $\text{sinc}(x)$  is continuous on  $\mathbb{R}$ .

$$\int_0^\infty \frac{\sin x}{x} dx = \underbrace{\int_0^1 \frac{\sin x}{x} dx}_{\text{continuous exists}} + \int_1^\infty \frac{\sin x}{x} dx$$

How about  $\int_1^\infty \frac{\sin(x)}{x} dx$ ?

$$\lim_{\beta \rightarrow \infty} \int_1^\beta \frac{\sin x}{x} dx = \left| \begin{array}{ll} u = \frac{1}{x} & u' = -\frac{1}{x^2} \\ v' = \sin x & v = -\cos x \end{array} \right| = \lim_{\beta \rightarrow \infty} \left[ -\frac{1}{x} \cos x \right]_1^\beta - \int_1^\beta \frac{\cos x}{x^2} dx$$

$$= \cos(1) - \lim_{\beta \rightarrow \infty} \int_1^\beta \frac{\cos(x)}{x^2} dx$$

$$\left| \frac{\cos(x)}{x^2} \right| \leq \frac{1}{x^2} \text{ on } [1, \beta]$$

and  $\int_1^\infty \frac{1}{x^2} dx$  exists. So  $g(x) = \frac{1}{x^2}$  is a majorant of  $\frac{\cos(x)}{x^2}$  and by Theorem 4.7,  $\lim_{\beta \rightarrow \infty} \int_1^\beta \frac{\cos(x)}{x^2} dx$  exists.

Attention!  $\int_0^\infty \left| \frac{\sin(x)}{x} \right| dx$  does not exist. Is not Lebesgue integrable.

**Definition 4.6.** Let  $x > 0$ . We call  $\Gamma$  Euler's Gamma function.

$$\Gamma(x) := \int_0^\infty t^{x-1} e^{-t} dt$$

**Remark 4.14.** The improper integral in the definition of the  $\Gamma$ -function exists for all  $x > 0$ .

↓ This lecture took place on 2018/04/26.

Euler's  $\Gamma$ -function exists for all  $x > 0$ :

$$\int_0^1 t^{x-1} e^{-t} dt$$

for  $0 < x \leq 1$ :

$$t^{x-1} \cdot \underbrace{e^{-t}}_{\leq 1} \leq t^{x-1} \text{ and } \underbrace{\int_0^1 \overbrace{t^{x-1}}^{\in(-1,0]} dt}_{\text{exists}}$$

also exists  $\int_0^1 t^{x-1} e^{-t} dt$  because of the direct comparison criterion. For  $x \geq 1$ ,  $t^{x-1} e^{-t}$  is continuous on  $[0, 1]$ , hence  $\int_0^1 t^{x-1} e^{-t} dt$  exists.

**Claim.** Let  $x > 0$  be fixed.  $\exists c > 0$  such that

$$t^{x-1} e^{-t} \leq c \cdot e^{-\frac{t}{2}} \forall t \in [0, \infty)$$

*Proof.*

$$\lim_{t \rightarrow \infty} \underbrace{t^{x-1}}_{\text{polynomially in } t} \cdot \underbrace{e^{-t}}_{\text{exponentially } \rightarrow 0} = 0$$

Also there exists  $L > 1$ , such that  $\forall x > L : t^{x-1} e^{-t/2} < 1$  on  $[1, L]$  (which is a compact interval) continuous. So there exists  $M > 0$  such that  $t^{x-1} e^{-\frac{t}{2}} \leq M \forall t \in [1, L]$ . Let  $c = \max\{M, 1\}$ . Therefore it holds on  $[1, L]$  and also on  $(L, \infty)$ .

$$t^{x-1} e^{-\frac{t}{2}} \leq c$$

Multiply with  $e^{-\frac{t}{2}} > 0$ , then  $t^{x-1} \cdot e^{-t} \leq c e^{-\frac{t}{2}} \forall t \in [1, \infty)$ .

$$c \int_1^\infty e^{-\frac{t}{2}} dt$$

exists. By the direct comparison test, we get  $\int_1^\infty t^{x-1} e^{-t} dt$  exists.  $\square$

**Lemma 4.7.** For all  $x > 0$ ,

$$\Gamma(x+1) = x \cdot \Gamma(x) \quad (\text{functional equation of the } \Gamma\text{-function})$$

*Epecially with  $\Gamma(1) = 1$  it holds that  $\Gamma(n+1) = n!$  for all  $n \in \mathbb{N}_0$ .*

*Proof.*

$$\begin{aligned} \Gamma(x+1) &= \int_0^\infty t^{x+1-1} e^{-t} dt = \int_0^\infty t^x e^{-t} dt \\ &= \left| \begin{array}{ll} u = t^x & u' = x \cdot t^{x-1} \\ v' = e^{-t} & v = -e^{-t} \end{array} \right| \\ &= \underbrace{-t^x \cdot e^{-t}}_{\substack{=0 \text{ on the upper bound} \\ =0 \text{ on the lower bound}}} \Big|_0^\infty + \int_0^\infty x \cdot t^{x-1} \cdot e^{-t} dt = x \int_0^\infty t^{x-1} e^{-t} dt = x \Gamma(x) \end{aligned}$$




Figure 27:  $\Gamma$  on  $\mathbb{C}$

$$\Gamma(1) = \int_0^\infty \underbrace{t^{1-1}}_{=1} \cdot e^{-t} dt = -e^{-t} \Big|_0^\infty = 1$$

$$\Gamma(n+1) = n \cdot \Gamma(n) = n \cdot (n-1) \Gamma(n-1) = n \cdot (n-1) \cdot \dots \cdot 1 \cdot \underbrace{\Gamma(1)}_{=1} = n!$$

□

**Remark 4.15.** There exists a power series  $\Gamma(x) = \sum_{n=0}^\infty a_n(x-x_0)^n$ .  $\Gamma(z)$  is also defined for  $z \in \mathbb{C}$  with  $\Re z > 0$ . Compare with Figure 27.

## 4.7 Young's inequality

Some important inequalities in integration theory follow.

**Theorem 4.8** (Young's inequality). Let  $f : [0, \infty) \rightarrow [0, \infty)$  be continuously differentiable, strictly monotonically increasing with  $f(0) = 0$  and  $f$  is unbounded. Then  $f : [0, \infty) \rightarrow [0, \infty)$  bijective and  $f^{-1} : [0, \infty) \rightarrow [0, \infty)$  is strictly monotonically increasing and continuous. Let  $a, b \geq 0$  be given. Then

$$ab \leq \int_0^a f(x) dx + \int_0^b f^{-1}(y) dy$$




Figure 28: Young's inequality visualized. The blue area denotes  $\int_0^a f(x) dx$  and  $\int_0^b f^{-1}(y) dy$  is the green area.

Equality is given if and only if,  $b = f(a)$  or  $a = f^{-1}(b)$ . Compare with Figure 28.

**Remark 4.16.** The inverse function  $f^{-1}$  can be retrieved by considering reflection along  $f(x) = x$ .

*Proof.* Let  $f : [0, \infty) \rightarrow [0, \infty)$  be as above. Let  $x_1 \neq x_2$ . Without loss of generality  $x_1 < x_2$ . Then  $f(x_1) < f(x_2) \implies f$  is injective. Surjectivity:  $f(0) = 0$ , hence  $0 \in f([0, \infty))$ . Let  $\eta > 0$  be arbitrary. Because  $f$  is unbounded, there exists  $z \in (0, \infty)$  with  $f(z) > \eta$ .  $f(0) = 0 < \eta < f(z)$ .

By the Intermediate Value Theorem ( $f$  is continuous), there exists  $\xi \in (0, z)$  with  $f(\xi) = \eta$ . So  $f$  is surjective.

$$f^{-1} : [0, \infty) \rightarrow [0, \infty)$$

*Monotonicity:* Let  $y_1 < y_2$ . Then  $x_1 = f^{-1}(y_1) < x_2 = f^{-1}(y_2)$ . If this would not be true (hence,  $x_2 \leq x_1$ ) then  $y_2 = f(x_2) \leq y_1 = f(x_1)$  gives a contradiction.

*Continuity of  $f^{-1}$ :* Let  $\varepsilon > 0$  be arbitrary. Let  $y \in (0, \infty)$  be chosen arbitrarily. We show  $f^{-1}$  is continuous in  $y$ . Let  $x = f^{-1}(y) > 0$  and choose  $\hat{\varepsilon} = \min\left\{\frac{x}{2}, \frac{\varepsilon}{2}\right\}$ .

$$x_1 = x - \hat{\varepsilon} > 0 \quad x_2 = x + \hat{\varepsilon} > 0$$

Let  $y_1 = f(x_1)$ ,  $y_2 = f(x_2)$ ,  $x_1 = f^{-1}(y_1)$  and  $x_2 = f^{-1}(y_2)$ . By monotonicity of  $f$ :  $x_1 < x < x_2 \implies y_1 < y < y_2$ .

Choose  $\delta = \min\{y - y_1, y_2 - y\} > 0$  (compare with Figure 29). Hence  $(y - \delta, y + \delta) \subseteq (y_1, y_2) \forall \eta \in (y - \delta, y + \delta)$ ,

$$f^{-1}(\eta) < f^{-1}(y + \delta) < f^{-1}(y_2) = x_2 = x + \hat{\varepsilon}$$






Figure 29:  $\delta$ ,  $y$ ,  $y_1$  and  $y_2$

$$f^{-1}(\eta) < f^{-1}(y - \delta) < f^{-1}(y_1) = x_1 = x - \varepsilon$$

So  $f^{-1}(\eta) \in (x - \varepsilon, x + \varepsilon)$ , and accordingly

$$|\eta - y| < \delta \implies \left| f^{-1}(\eta) - \underbrace{f^{-1}(y)}_{=x} \right| < c \leq \frac{\varepsilon}{2} < \varepsilon$$

So  $f^{-1}$  is continuous in  $y$  and  $f^{-1}$  is continuous in  $y_0$  analogously.  $\square$

Consider

$$\begin{aligned} \int_0^b f^{-1}(y) dy &= \left| \begin{array}{ll} y & = f(x) \\ dy & = f'(x) dx \\ y=0 & \implies x = f^{-1}(0) = 0 \\ y=b & \implies x = f^{-1}(b) \end{array} \right| \\ &= \int_0^{f^{-1}(b)} \underbrace{f^{-1}(f(x))}_{=x} \cdot f'(x) dx = \int_0^{f^{-1}(b)} x \cdot f'(x) dx \\ &\underbrace{=}_{\text{integration by parts}} x \cdot f(x) \Big|_0^{f^{-1}(b)} - 0 \int_0^{f^{-1}(b)} 1 \cdot f(x) dx \\ &= f^{-1}(b) \cdot b - \int_0^{f^{-1}(b)} f(x) dx \end{aligned}$$

So

$$\begin{aligned} I &:= \int_0^a f(x) dx + \int_0^b f^{-1}(y) dy = \int_{f^{-1}(b)}^0 f(x) dx + b \cdot f^{-1}(b) \\ &= \int_{f^{-1}(b)}^a f(x) dx + b \cdot f^{-1}(b) \end{aligned}$$

**Case 1**  $a = f^{-1}(b)$

$$\implies I = \underbrace{\int_a^a f(x) dx}_{=0} + b \cdot a$$

**Case 2**  $b < f(a)$ , and accordingly  $f^{-1}(b) < a$

$$\Rightarrow \int_{f^{-1}(b)}^a \underbrace{f(x)}_{f(f^{-1}(b)) \text{ for } x > f^{-1}(b)} dx > \overbrace{b}^{\text{minimal value}} \cdot \underbrace{(a - f^{-1}(b))}_{\text{length of integration interval}}$$

Therefore  $I > b(a - f^{-1}(b)) + b \cdot f^{-1}(b) = ab$ .

**Case 3**  $b > f(a)$ , and accordingly  $f^{-1}(b) > a$

$$\begin{aligned} \int_{f^{-1}(b)}^a f(x) dx &= \int_a^{f^{-1}(b)} \underbrace{(-f(x))}_{\substack{\text{monotonically decreasing} \\ > -f(f^{-1}(b)) \forall x \in [a, f^{-1}(b))}} dx > -f(f^{-1}(b)) \cdot (f^{-1}(b) - a) \\ &= -b(f^{-1}(b) - a) \\ I &> -b(f^{-1}(b) - a) + b \cdot f^{-1}(b) = ab \end{aligned}$$

**Remark 4.17.** Young's inequality also holds without requiring differentiability of  $f$  (but the proof is more complex).

**Lemma 4.8** (Special case of Young's inequality). Let  $A, B \geq 0$  and  $p, q > 1$  such that  $\frac{1}{p} + \frac{1}{q} = 1 \iff p + q = p \cdot q$ . Then  $p$  and  $q$  are called conjugate exponents. Then  $AB \leq \frac{A^p}{p} + \frac{B^q}{q}$ .

*Proof.*

$$\begin{aligned} f(x) &= x^{p-1} \text{ in Young's inequality} \\ y &= x^{p-1} \iff x = y^{\frac{1}{p-1}} \\ \frac{1}{p-1} &= q-1 \text{ is immediate, because} \\ \frac{1}{p-1} &= q-1 \iff 1 = pq - p - q + 1 \iff p + q = pq \end{aligned}$$

So  $f^{-1}(y) = y^{\frac{1}{p-1}} = y^{q-1}$ . By Young's inequality:

$$\begin{aligned} AB &\leq \int_0^A x^{p-1} dx + \int_0^B y^{q-1} dy \\ &= \frac{x^p}{p} \Big|_0^A + \frac{y^q}{q} \Big|_0^B = \frac{A^p}{p} + \frac{B^q}{q} \end{aligned}$$

□

**Remark 4.18.**

$$AB = \frac{A^p}{p} + \frac{B^q}{q}$$

Equality holds if and only if  $B = A^{p-1} \iff B^q = \overbrace{A^{pq} - q}^p = A^p$ .

## 4.8 Hölder's inequality

**Theorem 4.9** (Hölder's inequality). *Let  $I$  be an interval with boundary values  $a$  and  $b$ .  $-\infty \leq a < b \leq +\infty$ . Let  $p$  and  $q$  be conjugate exponents. Let  $f_1$  and  $f_2$  be regulated functions on  $I$  such that*

$$\int_a^b |f_1(x)|^p dx < \infty \quad \int_a^b |f_2(x)|^q dx < \infty$$

*both exist.*

*We let  $\|f_1\|_p := \left( \int_a^b |f_1(x)|^p dx \right)^{\frac{1}{p}}$  and  $\|f_2\|_q := \left( \int_a^b |f_2(x)|^q dx \right)^{\frac{1}{q}}$ . They are called  $L^p$ -norm of  $f_1$  and  $L^q$ -norm of  $f_2$ .*

*Then*

$$\int_a^b |f_1(x) \cdot f_2(x)| dx < \infty$$

*exists and*

$$\int_a^b |f_1(x) \cdot f_2(x)| dx \leq \|f_1\|_p \cdot \|f_2\|_q$$

*Proof.* Assume that  $\|f_1\|_p > 0$  and  $\|f_2\|_q > 0$ . Let  $A = \frac{|f_1(x)|}{\|f_1\|_p}$  and  $B = \frac{|f_2(x)|}{\|f_2\|_q}$ . By Lemma 4.8,

$$\frac{|f_1(x)|}{\|f_1\|_p} \cdot \frac{|f_2(x)|}{\|f_2\|_q} \leq \frac{1}{q} \cdot \frac{|f_1(x)|^p}{\|f_1\|_p^p} + \frac{1}{q} \cdot \frac{|f_2(x)|^q}{\|f_2\|_q^q}$$

We integrate the inequality,

$$\begin{aligned} & \frac{1}{\|f_1\|_p \cdot \|f_2\|_q} \cdot \int_a^b |f_1(x) \cdot f_2(x)| dx \\ & \leq \frac{1}{p} \cdot \frac{1}{\|f_1\|_p^p} \cdot \underbrace{\int_a^b |f_1(x)|^p dx}_{=\|f_1\|_p^p} + \frac{1}{q} \cdot \frac{1}{\|f_2\|_q^q} \underbrace{\int_a^b |f_2(x)|^q dx}_{=\|f_2\|_q^q} = \frac{1}{p} + \frac{1}{q} = 1 \end{aligned}$$

$$\frac{1}{\|f_1\|_p \cdot \|f_2\|_q} \cdot \int_a^b |f_1(x) \cdot f_2(x)| dx \leq 1 \implies \int_a^b |f_1(x) f_2(x)| dx \leq \|f_1\|_p \cdot \|f_2\|_q$$

Special case: Let  $\|f_1\|_p = 0$

$$\implies \left( \int_a^b |f_1(x)|^p dx \right)^{\frac{1}{p}} = 0 \implies \int_a^b \underbrace{|f_1(x)|^p}_{\geq 0} dx = 0$$

By Theorem 4.6,  $f_1(x) = 0 \forall x \in [a, b] \setminus A$  and  $A$  is at most countable.

$$\implies f_1(x) \cdot f_2(x) = 0 \forall x \in [a, b] \setminus A$$

$$\implies \int_a^b |f_1(x) \cdot f_2(x)| dx = 0$$

$$\implies 0 = 0 \text{ in Hölder's inequality}$$

□

**Remark 4.19** (Special case of Hölder's inequality). Let  $p = q = 2$ ,  $\frac{1}{2} + \frac{1}{2} = 1$ .

$$\int_a^b |f_1(x) \cdot f_2(x)| dx \leq \|f_1\|_2 \|f_2\|_2$$

is called *Cauchy-Schwarz inequality for  $L^2$  functions*.

$$\int_a^b f_1(x)f_2(x) dx = \langle f_1, f_2 \rangle_2 = \langle f_1, f_2 \rangle_{L^2}$$

is an inner product on a proper space of functions.

## 5 Elaboration on differential calculus

We consider a metric space  $X$  and functions  $f : X \rightarrow \mathbb{C}$ . We define a concept of uniform convergence of such sequences:

$$f_n : X \rightarrow \mathbb{C} \quad (n \in \mathbb{N}) \text{ and } f : X \rightarrow \mathbb{C}$$

We say,  $(f_n)_{n \in \mathbb{N}}$  converges uniformly towards  $f$  if  $\forall \varepsilon > 0 \forall N \in \mathbb{N}$  such that  $\forall x \in X$  and  $\forall n \geq N$

$$\underbrace{|f_n(x) - f(x)|}_{\text{absolute value in } \mathbb{C}} < \varepsilon$$

$$\iff \sup \{|f_n(x) - f(x)| : x \in X\} < \varepsilon$$

**Remark 5.1.** Do not use  $\|f\|_\infty$  for the definition of uniform convergence, because  $f_n$  and  $f$  must not be necessarily bounded. Hence,

$$\|f\|_\infty = \sup \{|f(x)| : x \in X\}$$

must not be finite.

**Theorem 5.1.** Let  $X$  be a metric space,  $f_n : X \rightarrow \mathbb{C}$  be a sequence of continuous functions and  $f : X \rightarrow \mathbb{C}$  such that  $f_n \rightarrow f$  uniform on  $X$ . Then  $f$  is also continuous on  $X$ .




Figure 30: Uniform convergence of  $f_N$  to  $f$

↓ This lecture took place on 2018/05/03.

*Proof.* Let  $\varepsilon > 0$  be arbitrary. Choose  $x \in X$ . Show:  $f$  is continuous in  $x$ .

Compare with Figure 30.

Because of uniform convergence  $f_n \rightarrow f$ , there exists  $N \in \mathbb{N}$  such that  $|f_N(z) - f(z)| < \frac{\varepsilon}{3} \forall z \in X$ . Let  $N$  be fixed. Because  $f_N$  is continuous in  $x$ , there exists  $\delta > 0$  such that  $d(x, \xi) < \delta \implies |f_N(\xi) - f_N(x)| < \frac{\varepsilon}{3}$ .

We consider now  $\xi \in X$  with  $d_X(x, \xi) < \delta$ . Then

$$\begin{aligned}
 |f(x) - f(\xi)| &= |f(x) - f_N(x) + f_N(x) - f_N(\xi) + f_N(\xi) - f(\xi)| \\
 &\leq \underbrace{|f(x) - f_N(x)|}_{< \frac{\varepsilon}{3}} + \underbrace{|f_N(x) - f_N(\xi)|}_{< \frac{\varepsilon}{3}} + \underbrace{|f_N(\xi) - f(\xi)|}_{< \frac{\varepsilon}{3}} \\
 &= \varepsilon
 \end{aligned}$$

by uniform convergence, by continuity and by uniform convergence respectively.

Thus,  $f$  is continuous in  $x$ . □

**Theorem 5.2.** Let  $P(z) = \sum_{k=0}^{\infty} a_k z^k$  be a power series in  $\mathbb{C}$  with convergence radius  $\rho_P > 0$ . Furthermore, let  $0 < r < \rho_P$ . Let  $P_n(z) = \sum_{k=0}^n a_k z^k$  ( $n$ -th partial sum of  $P$ ). Then  $P_n \rightarrow P$  uniformly on  $\overline{K_r(0)}$ .

**Remark 5.2.** See Figure 31.

*Proof.* Approximation theorem for power series. Lettl Analysis 1, lecture notes, section 5, theorem 10.




Figure 31: We cannot make a general statement about convergence/divergence. But on every small closed sphere  $P$  converges absolutely for every  $z$

Let  $0 < r < \rho_P$ . Choose  $\bar{r}$  with  $r < \bar{r} < \rho_P$ . Then it holds for  $z \in \overline{K_r(0)}$  that

$$|P(z) - P_n(z)| < \frac{\bar{r}}{\bar{r} - r} \cdot \left(\frac{r}{\bar{r}}\right)^n$$

$$\frac{r}{\bar{r}} < 1$$

hence  $\left(\frac{r}{\bar{r}}\right)^n$  is arbitrary small, for every  $n$  sufficiently large.

$$\Rightarrow \sup \left\{ |P(z) - P_n(z)| : z \in \overline{K_r(0)} \right\} \leq \underbrace{\frac{\bar{r}}{\bar{r} - r}}_{\text{fixed}} \cdot \underbrace{\left(\frac{r}{\bar{r}}\right)^n}_{\text{arbitrary small for } n \text{ sufficiently large}}$$

Hence,  $P_n \rightarrow P$  uniform on  $\overline{K_r(0)}$ . □

**Corollary.**  $P$  is continuous on  $K_{\rho_P}(0)$ .

**Theorem 5.3.** Let  $I \subseteq \mathbb{R}$  be an interval. Let  $f_n : I \rightarrow \mathbb{R}$  be continuously differentiable on  $I \forall n \in \mathbb{N}$ .

1.  $\exists g : I \rightarrow \mathbb{R}$  such that  $f'_n \rightarrow g$  uniform on  $I$
2.  $\exists f : I \rightarrow \mathbb{R}$  such that  $\forall x \in I : f(x) = \lim_{n \rightarrow \infty} f_n(x)$  ("pointwise convergence").

Then  $f$  is continuously differentiable on  $I$  and  $g = f'$ .

*Proof.*  $g$  is continuous as uniform limit of continuous  $f'_n$  (Theorem 5.1). For  $f_n$ , the Fundamental Theorem of Differential Calculus can be applied ( $f'_n$  is continuous, hence a regulated function). Let  $x_0 \in I$ . Then

$$f_n(x) = f_n(x_0) + \int_{x_0}^x f'_n(\xi) d\xi$$

Convergence for  $n \rightarrow \infty$ :

$$f_n(x) \rightarrow f(x) \quad f_n(x_0) \rightarrow f(x_0)$$

(Pointwise convergence)

$$\int_{x_0}^x f'_n(\xi) d\xi \rightarrow \int_{x_0}^x g(\xi) d\xi$$

Therefore, for  $n \rightarrow \infty$ ,

$$f(x) = f(x_0) + \int_{x_0}^x g(\xi) d\xi$$

The right-hand side is continuously differentiable by  $x$  according to the Fundamental Theorem, variant 1, with

$$\left( f(x_0) + \int_{x_0}^x g(\xi) d\xi \right)'(x) = g(x)$$

Hence, by  $f(x) = f(x_0) + \int_{x_0}^x g(\xi) d\xi$  it follows that

$$f'(x) = g(x) \quad \forall x \in I$$

□

To finish our proof, we need a result we missed in the section about Integrals.

**Lemma 5.1.** *Let  $(f_n)_{n \in \mathbb{N}}$  be a sequence of regulated functions on  $[a, b]$  and  $f_n \rightarrow f$  uniform on  $[a, b]$ . Then*

$$\int_a^b |f_n - f| dx \rightarrow 0 \quad \text{for } n \rightarrow \infty \quad \text{especially } \int_a^b f_n dx \rightarrow \int_a^b f dx$$

*Proof.*  $f$  as a uniform limit of regulated functions is a regulated function. The proof has been done in the practicals.

Let  $N \in \mathbb{N}$  large enough such that

$$\forall n \geq N \forall x \in [a, b] : |f_n(x) - f(x)| < \frac{\varepsilon}{b-a}$$

Then

$$\int_a^b |f_n(x) - f(x)| dx < \int_a^b \frac{\varepsilon}{b-a} dx = \frac{\varepsilon}{b-a}(b-a) = \varepsilon$$

Hence,

$$\lim_{n \rightarrow \infty} \int_a^b |f_n(x) - f(x)| dx = 0$$

$$\underbrace{\left| \int_a^b f_n dx - \int_a^b f dx \right|}_{\Rightarrow \rightarrow 0} \leq \underbrace{\int_a^b |f_n - f| dx}_{\rightarrow 0}$$

So,

$$\int_a^b f dx = \lim_{n \rightarrow \infty} \int_a^b f_n dx$$

□

## 5.1 Multiple times differentiable

**Definition 5.1.** Let  $f : I \rightarrow \mathbb{R}$ .  $I \subseteq \mathbb{R}$  is an interval. We define inductively:

$$f^{(0)}(x) = f(x)$$

Assume  $f^{(n-1)}$  is defined continuously on  $I$  and differentiable in  $x \in I$ . Then we let

$$f^{(n)}(x) = \left( f^{(n-1)} \right)'(x)$$

$f^{(n)}(x)$  is called  $n$ -th derivative of  $f$  in  $x$ .

Notational remark:

$$f^{(0)} = f \quad f^{(1)} = f' \quad f^{(2)} = f'' \quad f^{(3)} = f''' \quad f^{(4)} = f''''$$

Furthermore, we let

$$C^n(I) := \left\{ f : I \rightarrow \mathbb{R} : f^{(k)}(x) \text{ exists } \forall x \in I \text{ and } x \mapsto f^{(k)}(x) \text{ is continuous } \forall 0 \leq k \leq n \right\}$$

We call  $C$  the space of  $n$ -times continuously differentiable functions on  $I$ .

**Remark 5.3.**  $C^n(I)$  is a vector space. If  $I = [a, b]$  is compact, then

$$\|f\|_{C^n} = \max \left\{ \sup |f^{(k)}(x)| : x \in I : 0 \leq k \leq n \right\}$$

defines a norm on  $C^n(I)$  with  $\sup |f^{(k)}(x)| : x \in I = \|f^{(k)}\|_{\infty}$ .

## 5.2 Best polynomial approximation of a function

**Remark 5.4** (New topic). Let  $f \in C^n(I)$  and  $x_0 \in I$ . Find an appropriate polynomial  $T$  which approximated  $f$  in a neighborhood of  $x_0$  in the “best” way.

**Definition 5.2.** Let  $P(x) = \sum_{k=0}^n a_k x^k$  be a polynomial with  $a_n \neq 0$  (hence degree of  $P$  is  $n$ ).

$P \in \mathbb{R}[x] \dots$  set of all polynomials with coefficients in  $\mathbb{R}$

This set of polynomials is a ring.

$x_0 \in \mathbb{R}$  is called  $k$ -times root of  $P$  ( $k \in \mathbb{N}$ ) if  $Q \in \mathbb{R}[x]$  exists such that  $P(x) = (x - x_0)^k Q(x)$  with  $Q(x_0) \neq 0$ .



**Remark 5.5.**  $P(x) = (x - x_0)^k \cdot Q(x)$  means that division of  $P$  by  $(x - x_0)^k$  gives no remainder. Recall that division with remainder means that  $\exists \hat{Q}, \hat{R}$  that are polynomials of degree  $\hat{R} < k$ ,

$$P(x) = (x - x_0)^k \cdot \hat{Q}(x) + \hat{R}(x)$$

$\hat{Q}, \hat{R}$  is unique. If  $P(x) = (x - x_0)^k \cdot Q(x) \implies \hat{R} = 0, \hat{Q} = Q$ .

**Lemma 5.2.** Let  $P(x) = \sum_{l=0}^n a_l x^l$  with  $a_n \neq 0$ . Let  $1 \leq k \leq n$ . Then  $x_0 \in \mathbb{R}$  is a  $k$ -times root of polynomial  $P \iff P^{(j)}(x_0) = 0$  for  $j = 0, \dots, k-1$  and  $P^{(k)}(x_0) \neq 0$ .

*Proof.* Proof by complete induction.

**Induction begin** Consider  $k = 1$ . Direction  $\implies$ .

Let  $x_0$  be a simple root of  $P$ , then  $P(x) = (x - x_0) \cdot Q(x)$  and  $Q(x_0) \neq 0$ . Hence,  $P(x_0) = (x_0 - x_0) \cdot Q(x_0) = 0$  and  $P'(x) = Q(x) + (x - x_0) \cdot Q'(x)$ . Thus,  $P'(x_0) = Q(x_0) + (x_0 - x_0) \cdot Q'(x_0) = Q(x_0) \neq 0$ .

Direction  $\impliedby$ .

Let  $P(x_0) = 0$  and  $P'(x_0) \neq 0$ . Division with remainder:  $P(x) = (x - x_0) \cdot Q(x) + R(x)$  with  $\text{degree}(R) \leq \text{degree}(x - x_0) = 1$ . Thus,  $R$  is constant. We insert  $x_0$ . This gives  $P(x_0) = (x_0 - x_0) \cdot Q(x_0) + R$  with  $P(x_0) = 0$  and  $(x_0 - x_0) = 0$ . Hence,  $R = 0$  is the zero polynomial and  $P(x) = (x - x_0) \cdot Q(x)$ . It remains to show that  $Q(x_0) \neq 0$ .  $P'(x) = 1 \cdot Q(x_0) + (x - x_0) \cdot Q'(x_0)$ . We insert  $x = x_0 \implies 0 \neq P'(x_0) = Q(x_0) + (x_0 - x_0) \cdot Q'(x_0)$ . Thus it holds that  $Q(x_0) = P'(x_0) \neq 0$ .

**Induction step**  $k \rightarrow k+1$

**Claim** (Auxiliary claim). Let  $P(x) = (x - x_0) \cdot \tilde{P}(x)$ . Let  $P, \tilde{P}$  be polynomials. Then it holds  $\forall j \in \mathbb{N}$  that

$$P^{(j)}(x) = (x - x_0) \cdot \tilde{P}^{(j)}(x) + j \cdot \tilde{P}^{(j-1)}(x)$$

*Proof.* Proof by complete induction.

Let  $j = 1$ .

$$P'(x) = 1 \cdot \underbrace{\tilde{P}(x)}_{\tilde{P}^{(0)}(x)} + (x - x_0) \cdot \underbrace{\tilde{P}'(x)}_{\tilde{P}^{(1)}(x)}$$

Consider  $j \rightarrow j+1$ .

$$\begin{aligned} P^{(j+1)}(x) &= \left( P^{(j)}(x) \right)'(x) \\ &= \underbrace{((x - x_0) \cdot \tilde{P}^{(j)}(x))'}_{\text{induction assumption}} \\ &= j \tilde{P}^{(j-1)}(x)'(x - x_0) \tilde{P}^{(j+1)}(x) + \tilde{P}^{(j)}(x) + j \cdot \tilde{P}^{(j)}(x) \\ &= (x - x_0) \tilde{P}^{(j+1)}(x) + (j+1) \cdot \tilde{P}^{(j)}(x) \end{aligned}$$

□

We continue with the induction step after verifying our auxiliary claim.  
Direction  $\implies$ .

Let  $x_0$  be an  $k+1$  times zero of  $P$ . Hence  $P(x) = (x - x_0)^{k+1} \cdot Q(x)$ .  $Q(x_0) \neq 0$ .  
Let  $\tilde{P}(x) = (x - x_0)^k \cdot Q(x)$ . We can apply the induction assumption on  $\tilde{P}$ .  
Hence

$$\tilde{P}^{(j)} = 0 \quad \text{for } j = 0, \dots, k-1 \quad \text{and} \quad \tilde{P}^{(k)}(x_0) \neq 0$$

$$P(x) = (x - x_0) \cdot \tilde{P}(x)$$

By the auxiliary claim,  $P^{(j)}(x) = (x - x_0) \cdot \tilde{P}^{(j)}(x) + j \cdot \tilde{P}^{(j-1)}(x)$ . Therefore

$$P^{(j)}(x_0) = j \cdot \tilde{P}^{(j-1)}(x_0) = \begin{cases} 0 & \text{for } j = 0, \dots, k \\ (k+1)\tilde{P}^{(k)}(x_0) \neq 0 & \text{for } j = k+1 \end{cases}$$

Hence, our claim about the derivatives is true (all derivatives are zero).

Direction  $\impliedby$ .

Let  $P^{(j)}(x_0) = 0$  for  $j = 0, \dots, k$  and  $P^{(k+1)}(x_0) \neq 0$  and induction assumption holds for  $k$ . Division with remainder and  $P^{(0)}(x_0) = 0 \implies P(x) = (x - x_0) \cdot \tilde{P}(x)$ . By our auxiliary claim, we get

$$P^{(j)}(x) = (x - x_0) \cdot \tilde{P}^{(j)}(x) + j\tilde{P}^{(j-1)}(x)$$

we insert  $x = x_0$  and use  $P^{(j)}(x_0) = 0$  for  $j = 0, \dots, k$

$$\implies \tilde{P}^{(j)}(x_0) = 0 \quad \text{for } j = 0, \dots, k-1$$

By the induction assumption,  $\tilde{P}(x) = (x - x_0)^k Q(x)$  with  $Q(x_0) \neq 0$

$$\implies P(x) = (x - x_0) \cdot \tilde{P}(x) = (x - x_0)^{k+1} Q(x)$$

□

↓ This lecture took place on 2018/05/08.

### 5.3 Taylor polynomial

**Remark 5.6.** Brook Taylor (1685–1731)

**Definition 5.3.** Let  $I \subseteq \mathbb{R}$  be an interval,  $f \in C^n(I)$ . We let

$$T_f^n(x; x_0) = \sum_{k=0}^n \frac{1}{k!} f^{(k)}(x_0) (x - x_0)^k$$

$T_f^n(x; x_0)$  is a polynomial in  $x$  with  $\text{degree}(T_f^n) \leq n$ .  $T_f^n(x; x_0)$  is called Taylor polynomial of  $f$  of order  $n$  in  $x_0$ .

**Lemma 5.3.** *The premise is the same like in Definition 5.3. The Taylor polynomial of  $T_f^n(x; x_0)$  is the only polynomial of degree  $\leq n$  which satisfies*

$$(T_f^n)^{(k)}(x_0) = f^{(k)}(x_0) \quad \text{for } k = 0, \dots, n$$

*Proof.* Claim:

$$(T_f^n)^{(k)}(x; x_0) = \sum_{l=k}^n \frac{f^{(l)}(x_0)}{(l-k)!} (x - x_0)^{l-k} \quad \text{for } 0 \leq k \leq n$$

Proof of the claim by complete induction:

**Induction base**  $n = 0$

$$(T_f^n)^{(0)}(x; x_0) = \sum_{l=0}^n \frac{f^{(l)}(x_0)}{l!} (x - x_0)^l$$

**Induction step**  $k \rightarrow k + 1$  Let  $(T_f^n)^{(k)}(x; x_0)$

$$= \sum_{l=k}^n \frac{f^{(l)}(x_0)}{(l-k)!} (x - x_0)^{l-k}$$

by induction hypothesis. Then,

$$\begin{aligned} &= \sum_{l=k+1}^n \frac{f^{(l)}(x_0)}{(l-k)!} (l-k) \cdot (x - x_0)^{l-k-1} \\ &= \sum_{l=k+1}^n \frac{f^{(l)}(x_0)}{(l-(k+1))!} (x - x_0)^{l-(k+1)} \end{aligned}$$

We apply insertion:  $x = x_0$  into  $(T_f^n)^{(k)}(x; x_0)$

$$(T_f^n)^{(k)}(x; x_0) = \sum_{l=k}^n \frac{f^{(l)}(x_0)}{(l-k)!} (x - x_0)^{l-k} = \frac{f^{(k)}(x_0)}{0!} = f^{(k)}(x_0)$$

We need to prove uniqueness: Let  $T, \tilde{T}$  be polynomials with  $T^{(k)}(x_0) = \tilde{T}^{(k)}(x_0) = f^{(k)}(x_0)$  for  $k = 0, \dots, n$ . Assume  $T \neq \tilde{T}$ , hence  $T - \tilde{T} \neq 0$  (where 0 is the zero polynomial). For  $P = T - \tilde{T}$ ,

$$P^{(k)}(x_0) = T^{(k)}(x_0) - \tilde{T}^{(k)}(x_0) = 0 \quad (\text{for } 0 \leq k \leq n)$$

By Lemma 5.2,  $x_0$  is an  $n + 1$ -times root of  $P$ . Thus, there exists a polynomial  $Q \neq 0$  with  $Q(x_0) \neq 0$  such that

$$\underbrace{P(x)}_{\text{degree} \leq n} = \underbrace{(x - x_0)^{(n+1)} \cdot Q(x)}_{\text{degree} \geq n+1}$$

This is a contradiction. Hence,  $T - \tilde{T} = 0$ . □




Figure 32: Visualization of the remainder term of a Taylor polynomial

## 5.4 Integral form of the Taylor remainder term

**Definition 5.4.** Let  $f \in C^n(I)$ ,  $x_0 \in I$ . Furthermore let  $T_f^n(x; x_0)$  be the Taylor polynomial of  $n$ -th degree of  $f$  in  $x_0$ . We let  $R_f^n(x; x_0) = f(x) - T_f^n(x; x_0)$ . We call  $R_f^{n+1}(x; x_0)$  the approximation error of the Taylor polynomial. Also called remainder term of  $n + 1$ -th order. Compare with Figure 32.

**Theorem 5.4.** Let  $f \in C^{(n+1)}$ ,  $x \in I$ ,  $x_0 \in I$ . Then

$$R_f^{n+1}(x; x_0) = \frac{1}{n!} \int_{x_0}^x (x-t)^n f^{(n+1)}(t) dt$$

We call it the integral form of the remainder term.

*Proof.* Complete induction over  $n$ .

**Induction base  $n = 0$**

$$T_f^0(x; x_0) = f(x_0)$$

$$\begin{aligned} R_f^1(x; x_0) &= \underbrace{f(x) - f(x_0)}_{f \in C^1} \\ &= \int_{x_0}^x f'(t) dt \\ &= \frac{1}{0!} \int_{x_0}^x (x-t)^0 f^{(1)}(t) dt \end{aligned}$$

**Induction step**  $n - 1 \rightarrow n$

$$\begin{aligned}
R_f''(x; x_0) &= f(x) - T_f^{n-1}(x; x_0) \\
&\stackrel{\text{ind. hypothesis}}{=} \frac{1}{(n-1)!} \int_{x_0}^x (x-t)^{n-1} f^{(n)}(t) dt \\
&= \left| \begin{array}{l} u' = (x-t)^{n-1} \quad v = f^{(n)}(t) \\ u = -\frac{1}{n}(x-t)^n \quad v' = f^{(n+1)}(t) \end{array} \right| \\
&= \frac{1}{(n-1)!} \left[ \underbrace{-\frac{1}{n}(x-t)^n \cdot f^{(n)}(t)}_{= \frac{1}{n!}(x-x_0)^n \cdot f^{(n)}(x_0)} \right] \Bigg|_{t=x_0}^x \\
&\quad + \underbrace{\frac{1}{(n-1)!} \int_{x_0}^x \frac{1}{n}(x-t)^n \cdot f^{(n+1)}(t) dt}_{= \frac{1}{n!} \int_{x_0}^x (x-t)^n \cdot f^{(n+1)}(t) dt}
\end{aligned}$$

So,

$$\begin{aligned}
&\underbrace{f(x) - T_f^{n-1}(x; x_0) - \frac{1}{n!} f^{(n)}(x_0)(x-x_0)^n}_{-T_f^n(x; x_0)} \\
&= \frac{1}{n!} \int_{x_0}^x (x-t)^n \cdot f^{(n+1)}(t) dt
\end{aligned}$$

Therefore,

$$R_f^{(n+1)}(x; x_0) = \frac{1}{n!} \int_{x_0}^x (x-t)^n f^{(n+1)}(t) dt$$

□

## 5.5 Lagrange form of the Taylor remainder term

**Theorem 5.5** (Lagrange form of the remainder term). *Let  $f \in C^{n+1}(I)$ ,  $n \in \mathbb{N}_0$ ,  $x, x_0 \in I$ ,  $x \neq x_0$ . Then there exists some  $\xi$  between  $x_0$  and  $x$  (hence,  $\xi \in (x_0, x)$  if  $x > x_0$  or  $\xi \in (x, x_0)$  if  $x < x_0$ ) such that*

$$R_f^{(n+1)}(x; x_0) = \frac{1}{(n+1)!} f^{(n+1)}(\xi)(x-x_0)^{n+1}$$

*Proof.* Idea: we apply the Mean Value Theorem for definite integrals on the Taylor remainder.

**Case 1** Let  $x_0 < x$ .

$$\begin{aligned}
R_f^{n+1}(x; x_0) &= \frac{1}{n!} \int_{x_0}^x \underbrace{(x-t)^n}_{\substack{\geq 0 \\ \text{regulated function}}} \underbrace{f^{(n+1)}(t)}_{\text{continuous in } t} dt \\
&\stackrel{\text{MVT}}{=} \frac{1}{n!} f^{(n+1)}(\xi) \cdot \int_{x_0}^x (x-t)^n dt \\
&= \frac{1}{n!} f^{(n+1)}(\xi) \left[ -\frac{1}{n+1} (x-t)^{n+1} \right]_{t=x_0}^x \\
&= \frac{1}{(n+1)!} f^{(n+1)}(\xi) (x-x_0)^{n+1}
\end{aligned}$$

where MVT is the Mean Value Theorem for definite integrals (Theorem 4.2).

**Case 2** Let  $x < x_0$  and  $n$  odd.

$$\begin{aligned}
R_f^{n+1}(x; x_0) &= -\frac{1}{n!} \int_x^{x_0} \underbrace{(x-t)^n}_{= (-1)^n (t-x)^n} \cdot f^{(n+1)}(t) dt \\
&= \frac{1}{n!} \int_x^{x_0} \underbrace{(t-x)^n}_{\geq 0} \cdot \underbrace{f^{(n+1)}(t)}_{\text{continuous}} dt \\
&= \frac{f^{(n+1)}(\xi)}{n!} \int_x^{x_0} (t-x)^n dt \\
&= \frac{f^{(n+1)}(\xi)}{n!} \left[ \frac{1}{n+1} (t-x)^{n+1} \right]_x^{x_0} \\
&= \frac{1}{(n+1)!} f^{(n+1)}(\xi) (x_0-x)^{n+1}
\end{aligned}$$

$n+1$  is even

$$= \frac{1}{(n+1)!} f^{(n+1)}(\xi) (x-x_0)^{n+1}$$

**Case 3** Let  $x < x_0$  and  $n$  even.

$$\begin{aligned}
 R_f^{n+1}(x; x) &= -\frac{1}{n!} \int_x^{x_0} \underbrace{(x-t)^n}_{\geq 0} \cdot \underbrace{f^{(n+1)}(t)}_{\text{continuous}} dt \\
 &= -\frac{1}{n!} f^{(n+1)}(\xi) \cdot \int_x^{x_0} (x-t)^n dt \\
 &= -\frac{1}{n!} f^{(n+1)}(\xi) \cdot \left[ -\frac{1}{n+1} (x-t)^{n+1} \right]_x^{x_0} \\
 &= \frac{1}{(n+1)!} f^{(n+1)}(\xi) (x-x_0)^{n+1}
 \end{aligned}$$

□

Any extreme value satisfies that its derivative is zero. But not every point with derivative zero is an extreme value. We now consider conditions to select extreme values from all value satisfying derivative zero.

## 5.6 Sufficient criteria for extreme values

**Corollary** (Sufficient conditions for existence of extreme values). *Let  $I$  be an open interval. Let  $x_0 \in I$  and  $f \in C^{n+1}(I)$ . Assume*

$$f^{(1)}(x_0) = f^{(2)}(x_0) = \cdots = f^{(n)}(x_0) = 0$$

*and  $f^{(n+1)}(x_0) \neq 0$ . Then  $f$  in  $x_0$  has*

1. *a strict local maximum if  $n$  is odd and  $f^{(n+1)}(x_0) < 0$*
2. *a strict local minimum if  $n$  is odd and  $f^{(n+1)}(x_0) > 0$*
3. *no extreme value in  $x_0$  if  $n$  is even.*

*Proof. Case a* Let  $f^{(n+1)}(x_0) < 0$  and  $f^{(n+1)}$  be continuous, then  $\exists \varepsilon > 0$  such that  $(x_0 - \varepsilon, x_0 + \varepsilon) \subseteq I$  ( $I$  is open) and  $f^{(n+1)}(\xi) < 0 \forall \xi \in (x_0 - \varepsilon, x_0 + \varepsilon)$ . Now let  $x \in (x_0 - \varepsilon, x_0 + \varepsilon)$ . Then by Theorem 5.5,

$$\frac{f^{(n+1)}(\xi)}{(n+1)!} (x-x_0)^{n+1} = R_f^{n+1}(x; x_0) = f(x) - \sum_{k=0}^n \frac{1}{k!} f^{(k)}(x_0) (x-x_0)^k = f(x) - f(x_0)$$

for  $k = 1, \dots, n$ . So,

$$f(x) - f(x_0) = \underbrace{\frac{f^{(n+1)}(\xi)}{(n+1)!}}_{<0} \underbrace{(x-x_0)^{n+1}}_{\substack{\text{even} \\ >0 \text{ for } x \neq x_0}}$$

hence  $f(x) - f(x_0) < 0$ , and accordingly

$$f(x) < f(x_0) \quad \forall x \in (x_0 - \varepsilon, x_0 + \varepsilon), x \neq x_0$$

So  $f$  is a strict local maximum.

**Case b** Analogously.

**Case c** We apply the same idea as in Case a up to the point, where we consider  $f(x) - f(x_0)$ .

$$f(x) - f(x_0) = \frac{f^{(n+1)}(\xi)}{(n+1)!} (x - x_0)^{n+1}$$

$f^{(n+1)}(\xi)$  has the same sign as  $\underbrace{f^{(n+1)}(x_0)}_{\neq 0} \quad \forall \xi \in (x_0 - \varepsilon, x_0 + \varepsilon)$ . This is feasible due to continuity of  $f^{(n+1)}$  for sufficiently small  $\varepsilon$ .

$$f(x) - f(x_0) = \underbrace{\frac{f^{(n+1)}(\xi)}{(n+1)!}}_{\text{has constant sign indep. of } x} \cdot \underbrace{(x - x_0)}_{\text{changes its sign}} \overbrace{n+1}^{\text{odd}}$$

Therefore  $f(x) - f(x_0)$  changes its sign for  $x = x_0$ . Hence  $f$  has no extreme value in  $x = x_0$ .

□

## 5.7 Qualitative Taylor equation

**Theorem 5.6** (Qualitative Taylor equation). *Let  $f \in C^n(I)$ ,  $x, x_0 \in I$ . Then there exists some function  $r \in C(I)$  with  $r(x_0) = 0$  such that*

$$f(x) = T_f^n(x; x_0) + (x - x_0)^n \cdot r(x)$$

and accordingly,

$$R_f^{n+1}(x; x_0) = (x - x_0)^n \cdot r(x)$$

**Remark 5.7.** *For some function  $r$  with  $\lim_{x \rightarrow x_0} r(x) = 0$ , we also denote  $o(x - x_0)$  instead of  $r(x)$ . This general notation is called Landau's Big-Oh notation.*

$$f(x) = T_f^n(x; x_0) + (x - x_0)^n \cdot o(x - x_0)$$

*Proof.* Let  $r(x) = \frac{f(x) - T_f^n(x; x_0)}{(x - x_0)^n}$  for  $x \neq x_0$  and  $r(x_0) := 0$ . Then  $f$  is continuous and  $T_f^n$  is continuous in every point  $x \neq x_0$ . It remains to show that  $r$  is continuous



in  $x = x_0$ .

$$\begin{aligned}
 r(x) &= \frac{1}{(x - x_0)^n} \underbrace{\left( f(x) - T_f^{n-1}(x; x_0) - \frac{1}{n!} f^{(n)}(x_0)(x - x_0)^n \right)}_{R_f^n(x; x_0)} \\
 &\stackrel{\text{Lagrange}}{=} \frac{1}{(x - x_0)} \left[ \frac{1}{n!} (x - x_0)^n \cdot f^{(n-1)}(\xi) - \frac{1}{n!} f^{(n)}(x_0)(x - x_0)^n \right]
 \end{aligned}$$

$\xi \in (x_0, x)$

$$= \frac{1}{n!} [f^{(n)}(\xi) - f^{(n)}(x_0)] \rightarrow 0 \text{ for } x \rightarrow x_0 \text{ because } f^{(n)} \text{ is continuous}$$

as  $x_0 < x < x$ , hence  $\xi \rightarrow x_0$  for  $x \rightarrow x_0$

So  $\lim_{x \rightarrow x_0} r(x) = 0 = r(x_0)$ , so  $r$  in  $x_0$  is continuous.  $\square$

↓ This lecture took place on 2018/05/15.

## 5.8 Taylor series

Assume  $f : I \rightarrow \mathbb{R}$  is infinitely often differentiable on  $I$ ,  $x_0 \in I$ . Then there exists  $T_f^n(x; x_0)$  for arbitrary  $n \in \mathbb{N}$ .

$$T_f(x; x_0) = \sum_{k=0}^{\infty} \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k$$

$T_f(x; x_0)$  defines the *Taylor series* on  $f$  in  $x_0$ . Power series in  $\xi = x - x_0$ .  $T_f$  has a convergence radius,

$$\rho(T_f) = \left[ \limsup_{k \rightarrow \infty} \sqrt[k]{\frac{|f^{(k)}(x_0)|}{k!}} \right]^{-1}$$

If  $\rho(T_f) > 0$ , then

$$f(x) = T_f(x; x_0) = \sum_{k=0}^{\infty} \frac{f^{(k)}(x)}{k!} (x - x_0)^k$$

in  $(x_0 - \rho(T_f), x_0 + \rho(T_f))$ ? Compare with Figure 33.




Figure 33: Taylor series

**Example 5.1** (Counterexample). Let  $f : \mathbb{R} \rightarrow \mathbb{R}$ .

$$f(x) = \begin{cases} e^{-\frac{1}{x}} & \text{if } x > 0 \\ 0 & \text{if } x \leq 0 \end{cases}$$

It holds for  $x > 0$ ,

$$f^{(n)}(x) = \frac{P(x)}{Q(x)} \cdot e^{-\frac{1}{x}}$$

where  $P, Q$  are polynomials. So the function value (of an infinitely often differentiable function) must not equate with its Taylor series value.

*Proof.* Proof by complete induction over  $n$ .

**Case**  $n = 0$  immediate with  $P = Q = 1$ .

**Case**  $n \mapsto n + 1$

$$\begin{aligned} f^{(n+1)}(x) &= \left[ \underbrace{\frac{P(x)}{Q(x)} \cdot e^{-\frac{1}{x}}}_{f^{(n)}(x) \text{ by induction hypothesis}} \right]' \\ &= \frac{P' \cdot Q - Q' \cdot P}{Q^2} \cdot e^{-\frac{1}{x}} + \frac{P}{Q} \cdot \frac{1}{x^2} \cdot e^{-\frac{1}{x}} \\ &= \frac{(P'Q - Q'P)x^2 + PQ}{Q^2x^2} \cdot e^{-\frac{1}{x}} \end{aligned}$$

It holds that  $\lim_{x \rightarrow 0^+} \frac{P(x)}{Q(x)} \cdot e^{-\frac{1}{x}} = 0$ . Immediately,  $\lim_{x \rightarrow 0^-} f^{(n)}(x) = 0$ , hence  $f^{(n)}(0) = 0 \forall n \in \mathbb{N}$ .  $f$  is arbitrarily often continuously differentiable on  $\mathbb{R}$ .

Thus,

$$T_f(x; 0) = \sum_{k=0}^{\infty} \overbrace{\frac{f^{(k)}(0)}{k!}}^{=0} x^k = 0$$

but  $f(x) \neq 0$  on  $\mathbb{R}$ . Thus,  $f \neq T_f(x; 0)$ . But  $R_f = f - T_f(x; 0) = f$ .

$$|R_f(x)| \leq c_n |x|^n \quad \forall n \in \mathbb{N}$$

□

**Theorem 5.7.** Let  $f(x) = \sum_{k=0}^{\infty} a_k(x-x_0)^k$  be an analytical<sup>2</sup> function with convergence radius  $\rho(f) > 0$ . Then  $f$  is infinitely often continuously differentiable on  $I := (x_0 - \rho(f), x_0 + \rho(f))$  and  $a_k = \frac{f^{(k)}(x_0)}{k!}$ , hence the given power series is the Taylor series of the function.

*Proof.* See Analysis 1 lecture notes, chapter 8, theorem 1 by G. Lettl.

$f$  is differentiable on  $I = (x_0 - \rho(f), x_0 + \rho(f))$  and  $f'(x) = \sum_{k=0}^{\infty} k a_k(x-x_0)^{k-1}$ . Thus,  $f'$  is also analytical and the power series of  $f'$  converges on  $K(x_0) \implies \rho(f') \geq \rho(f)$  (if you consider the Cauchy-Hadamard Theorem, then  $\rho(f') = \rho(f)$ ).

Induction:  $f^{(n)}(x)$  is analytical on  $I$  and

$$f^{(n)}(x) = \sum_{k=n}^{\infty} k \cdot (k-1) \dots (k-n+1) \cdot a_k \cdot (x-x_0)^{k-n}$$

We insert:  $x = x_0$

$$f^{(n)}(x_0) = n \cdot (n-1) \dots 1 \cdot a_n \implies a_n = \frac{f^{(n)}(x_0)}{n!}$$

□

Revision: Expansion on a different point ( $\xi_0$  instead of  $x_0$ ):

$$f(z) = \sum_{k=0}^{\infty} a_k(z-x_0)^k = \sum_{k=0}^{\infty} \frac{f^{(k)}(x_0)}{k!} (z-x_0)^k$$

with  $a_k = \frac{f^{(k)}(x_0)}{k!}$ .  $f(z) = \sum_{k=0}^{\infty} \tilde{a}_k(z-\xi_0)^k$  with  $\tilde{a}_k = \frac{f^{(k)}(\xi_0)}{k!}$ . Compare with Figure 34.

---

<sup>2</sup>Reminder: A function is analytical if it is locally given by a convergent power series.




Figure 34: Expansion on a different point

## 6 Multidimensional differential calculus

Let  $V, W$  be vector space over  $\mathbb{K}$  ( $\mathbb{R}, \mathbb{C}$ ).

$$\underbrace{\mathcal{L}(V, W)}_{\text{Hom}(V, W)} = \{\varphi : V \rightarrow W : \varphi \text{ is linear}\}$$

$\text{Hom}(V, W)$  has vector space properties.  $\varphi, \psi \in \mathcal{L}(V, W), \lambda, \mu \in \mathbb{K}$ . Then  $\lambda\varphi + \mu\psi \in \mathcal{L}(V, W)$ . In general, it is feasible that to define a norm on  $\mathcal{L}(V, W)$ . Hence,  $\|\cdot\| : \mathcal{L}(V, W) \rightarrow [0, \infty)$  with

1.  $\|\varphi\| = 0 \iff \varphi = 0$  (zero mapping)
2.  $\forall \lambda \in \mathbb{K}, \varphi \in \mathcal{L}(V, W) : \|\lambda\varphi\| = |\lambda| \cdot \|\varphi\|.$
3.  $\forall \varphi, \psi \in \mathcal{L}(V, W) : \|\varphi + \psi\| \leq \|\varphi\| + \|\psi\|.$

### 6.1 Frobenius and matrix norm

**Example 6.1.** Let  $\mathcal{L}(\mathbb{R}^m, \mathbb{R}^n) \cong \mathbb{R}^{n \times m}$ . (identify linear maps with its matrix representation with respect to the canonical basis)

$$A \in \mathbb{R}^{n \times m} \quad A = (a_{ij})_{\substack{i=1, \dots, n \\ j=1, \dots, m}}$$

$$\|A\|_F = \left( \sum_{i=1}^n \sum_{j=1}^m |a_{ij}|^2 \right)^{\frac{1}{2}} \quad \text{"Frobenius norm"}$$

It basically works by appending the next column to the previous one. Hence, this gives a column vector. We square every entries, sum it up and take its square root (a common norm procedure). A norm on  $\mathbb{R}^{n \times m}$  (a matrix) is called matrix norm ( $\mathbb{C}^{n \times m}$ ).

## 6.2 Operator norm and bounded linear operators

**Definition 6.1.** Let  $V, W$  be normed vector spaces over  $\mathbb{K}$ . A linear map  $\varphi : V \rightarrow W$  is called bounded if  $\exists m \geq 0 : \|\varphi(x)\|_W \leq m \cdot \|x\|_V$  for all  $x \in V$  (we call this the boundedness criterion).

The set  $\mathcal{L}_b(V, W) = \{\varphi : V \rightarrow W : \varphi \text{ is linear and bounded}\}$  is a subvectorspace of  $\mathcal{L}(V, W)$ . We let

$$\|\varphi\| = \inf \{m \geq 0 : \|\varphi(x)\|_W \leq m \cdot \|x\|_V \forall x \in V\}$$

and call  $\|\varphi\|$  the operator norm on  $\varphi$  with respect to  $\|\cdot\|_V$  and  $\|\cdot\|_W$ .

Regarding the subvector space property:

Let  $\varphi, \psi \in \mathcal{L}_b(V, W)$ ,  $\lambda, \mu \in \mathbb{K}$ . Show that  $\lambda\varphi + \mu\psi \in \mathcal{L}_b(V, W)$ .

$$\begin{aligned} \|(\lambda\varphi + \mu\psi)(x)\|_W &= \|\lambda \cdot \varphi(x) + \mu \cdot \psi(x)\|_W \\ &\leq |\lambda| \|\varphi(x)\|_W + |\mu| \|\psi(x)\|_W \\ &\leq \underbrace{|\lambda| m \|x\|_V + |\mu| m' \|x\|_V}_{\text{because } \varphi, \psi \text{ are bounded}} \\ &= \underbrace{(|\lambda| m + |\mu| m')}_{=:\tilde{m} \geq 0} \|x\|_V \end{aligned}$$

hence  $\lambda\varphi + \mu\psi \in \mathcal{L}_b(V, W)$ .  $\mathcal{L}_b(V, W) \neq \emptyset$ .

**Lemma 6.1.** Let  $V, W$  be normed vector spaces. Then it holds for any  $\varphi \in \mathcal{L}_b(V, W)$

1.  $\|\varphi(x)\|_W \leq \|\varphi\| \cdot \|x\|_V \forall x \in V$ . Hence,  $m = \|\varphi\|$  satisfies the boundedness criterion, hence informally  $\inf$  equals  $\min$  in Definition 6.1.

2.

$$\|\varphi\| = \sup \left\{ \frac{\|\varphi(x)\|_W}{\|x\|_V} : x \in V \setminus \{0\} \right\} = \sup \left\{ \|\varphi(x)\|_W : x \in V \text{ with } \|x\|_V = 1 \right\}$$

3.  $\|\cdot\|$  is a norm on  $\mathcal{L}_b(V, W)$ .

*Proof.* 1. Let  $m_n \geq 0$  with  $m_n$  satisfies the boundedness criterion, hence

$$\|\varphi(x)\|_W \leq m_n \cdot \|x\|_V \forall x \in V$$

and  $m_n \rightarrow \|\varphi\|$ . The inequality retains in the limit. Thus,  $\|\varphi(x)\|_W \leq \|\varphi\| \cdot \|x\|_V$ .

2. Let  $\tilde{m} = \sup \left\{ \frac{\|\varphi(x)\|_W}{\|x\|_V} : x \neq 0 \right\}$ . Hence,  $\frac{\|\varphi(x)\|_W}{\|x\|_V} \leq \tilde{m} \forall x \in V$ , because  $\tilde{m}$  is an upper bound. So,  $\|\varphi(x)\|_W \leq \tilde{m} \|x\|_V$ . Thus  $\tilde{m}$  satisfies the boundedness criterion and  $\|\varphi\| \leq \tilde{m}$ .

On the opposite: Let  $m$  such that the boundedness criterion is satisfied  $\Rightarrow \|\varphi(x)\|_W \leq m \cdot \|x\|_V \forall x \in V, x \neq 0$  and accordingly,  $\frac{\|\varphi(x)\|_W}{\|x\|_V} \leq m$ . Hence,  $m$  is upper bound of  $\left\{ \frac{\|\varphi(x)\|_W}{\|x\|_V} : x \neq 0 \right\}$ , hence  $m \geq \tilde{m} = \sup \{ \cdot \}$ . Hence,  $m \geq \tilde{m} = \sup \left\{ \frac{\|\varphi(x)\|_W}{\|x\|_V} : x \neq 0 \right\}$ . The statement above also holds for the infimum of  $m - s$ , hence  $\|\varphi\| \geq \tilde{m}$ , hence  $\|\varphi\| = \tilde{m} = \sup \left\{ \frac{\|\varphi(x)\|_W}{\|x\|_V} : x \neq 0 \right\}$ . Because  $\{x \in V : \|x\|_V = 1\} \subseteq \{x \in V : x \neq 0\}$ ,  $\sup \left\{ \frac{\|\varphi(x)\|_W}{\|x\|_V} : \|x\|_V = 1 \right\} = \sup \left\{ \frac{\|\varphi(x)\|_W}{\|x\|_V} : x \neq 0 \right\} = \|\varphi\|$ .

On the opposite: Let  $x \neq 0$ . Then  $\tilde{x} = \frac{x}{\|x\|_V}$  defines a *unit vector*.

$$\|\tilde{x}\|_V = \left\| \frac{x}{\|x\|_V} \right\|_V = \frac{1}{\|x\|_V} \cdot \|x\|_V = 1$$

and

$$\frac{\|\varphi(x)\|_W}{\|x\|_V} = \frac{1}{\|x\|_V} \|\varphi(x)\|_W = \left\| \frac{1}{\|x\|_V} \varphi(x) \right\|_W \underbrace{=}_{\varphi \text{ is linear}} \left\| \varphi\left(\frac{x}{\|x\|_V}\right) \right\|_W = \|\varphi(\tilde{x})\|_W$$

$$\begin{aligned} \Rightarrow \forall x \neq 0 : \frac{\|\varphi(x)\|_W}{\|x\|_V} &= \|\varphi(\tilde{x})\|_W \leq \sup \left\{ \|\varphi(z)\|_W : \|z\|_V = 1 \right\} \\ \Rightarrow \sup \left\{ \frac{\|\varphi(x)\|_W}{\|x\|_V} : x \neq 0 \right\} &\leq \sup \left\{ \|\varphi(z)\|_W : \|z\|_V = 1 \right\} \end{aligned}$$

3. Show that  $\|\varphi\|$  is a norm.

$$\|\varphi\| = 0 \iff \forall x \in V : \|\varphi(x)\|_W \leq 0 \cdot \|x\|_W$$

hence  $\varphi(x) = 0 \forall x \in V$  and accordingly,  $\varphi = 0$  in  $\mathcal{L}(V, W)$ .

$$\begin{aligned} \|\lambda\varphi\| &= \sup \left\{ \|\lambda\varphi(x)\|_W : \|x\|_V = 1 \right\} = \sup \left\{ |\lambda| \|\varphi(x)\|_W : \|x\|_V = 1 \right\} \\ &= |\lambda| \sup \left\{ \|\varphi(x)\|_W : \|x\|_V = 1 \right\} = |\lambda| \|\varphi\| \end{aligned}$$

Triangle inequality: Let  $\varphi, \psi \in \mathcal{L}_b(V, W)$ .

$$\begin{aligned} \|\varphi(x) + \psi(x)\|_W &\underbrace{\leq}_{\text{triangle inequality in } W} \|\varphi(x)\|_W + \|\psi(x)\|_W \\ &\leq \|\varphi\| \cdot \|x\|_V + \|\psi\| \cdot \|x\|_V = (\|\varphi\| + \|\psi\|) \cdot \|x\|_V \end{aligned}$$

By (1.),  $\|\varphi\| + \|\psi\|$  satisfies the boundedness criterion for the linear map  $\varphi + \psi$ . Hence,  $\|\varphi + \psi\| \leq \|\varphi\| + \|\psi\|$ .

□

**Remark 6.1.** •  $\|A\|_F$  is no operator norm on  $\mathcal{L}(\mathbb{R}^m, \mathbb{R}^n)$ .

- Boundedness of linear mappings is required to define  $\|\varphi\|$ .

We consider special case  $V = \mathbb{R}^m, W = \mathbb{R}^n$ .

$$\|\cdot\|_V = \|\cdot\|_\infty \quad \|\cdot\|_W = \|\cdot\|_\infty$$

Let  $A \in \mathbb{R}^{n \times m} (\mathcal{L}(\mathbb{R}^m, \mathbb{R}^n))$ . Then

$$\begin{aligned} \|Ax\|_\infty &= \max \{ |(Ax)_i| : i = 1, \dots, n \} \\ &= \max \left\{ \left| \sum_{j=1}^m a_{ij} x_j \right| : i = 1, \dots, n \right\} \\ &\leq \max \left\{ \sum_{j=1}^m |a_{ij}| \cdot \underbrace{|x_j|}_{\leq \|x\|_\infty} : i = 1, \dots, n \right\} \\ &\leq \max \left\{ \|x\|_\infty \cdot \sum_{j=1}^m |a_{ij}| : i = 1, \dots, n \right\} \\ &= \max \left\{ \underbrace{\sum_{j=1}^m |a_{ij}|}_{=m} : i = 1, \dots, n \right\} \cdot \|x\|_\infty \\ &= m \cdot \|x\|_\infty \end{aligned}$$

Hence the boundedness criterion is satisfied.  $A$  is bounded with respect to  $\|\cdot\|_\infty$  in the preimage and image space. By the norm equivalence theorem, it follows that  $A$  is bounded with respect to arbitrary norms on  $\mathbb{R}^m$ , and accordingly  $\mathbb{R}^n$ .

↓ This lecture took place on 2018/05/17.

Further remarks:

**Remark 6.2.** A linear map  $A : \mathbb{R}^m \rightarrow \mathbb{R}^n$  is always bounded. Thus,

$$\|Ax_1 - Ax_2\|_{\mathbb{R}^n} = \|A(x_1 - x_2)\|_{\mathbb{R}^n} \leq \|A\| \|x_1 - x_2\|_{\mathbb{R}^m}$$

So every linear map  $A : \mathbb{R}^m \rightarrow \mathbb{R}^n$  is Lipschitz continuous with Lipschitz constant  $\|A\|$ .

The considerations above hold for arbitrary finite-dimensional normed vector spaces  $V$  and  $W$  (over  $\mathbb{K} = \mathbb{R}, \mathbb{C}$ ).

**Lemma 6.2.** *Let  $X, Y$  and  $Z$  be normed vector spaces. Let  $A : X \rightarrow Y$  be linear and bounded. Let  $B : Y \rightarrow Z$  be linear and bounded. Then*

$$B \cdot A : X \rightarrow Z$$

*is also bounded and*

$$\|B \cdot A\| \leq \|B\| \cdot \|A\|$$

*Proof.* Let  $x \in X$  be arbitrary and  $BAx = B(Ax) \in Z$  and

$$\|BAx\|_Z = \|B(Ax)\|_Z \overset{\substack{\leq \\ B \text{ is bounded}}}{\leq} \|B\| \|Ax\|_Y \overset{\substack{\leq \\ A \text{ is bounded}}}{\leq} \|B\| \cdot \|A\| \|x\|_X$$

Hence  $m = \|B\| \|A\|$  satisfies the boundedness criterion for the linear map  $B \cdot A : X \rightarrow Z$ . Because  $\|B \cdot A\|$  is the smallest constant for which the boundedness criterion holds, it follows that  $\|BA\| \leq \|B\| \|A\|$ .  $\square$

### 6.3 Landau notation

**Definition 6.2** (Landau  $O$  symbols). *Let  $h, g : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$  and  $D$  is open,  $a \in D$ .*

1. *We denote  $h = O(g)$  in  $a$  (German pronunciation:  $h$  ist groß  $O$  von  $g$ ) iff  $\exists U \subseteq D$  neighborhood of  $a$  in  $D$  and  $\exists r : U \rightarrow \mathbb{R}$  with  $r$  bounded such that  $h(x) = r(x) \cdot g(x) \forall x \in U$ . Thus,*

$$\left| \frac{h(x)}{g(x)} \right| = |r(x)| \leq M \quad \forall x \in U$$

$$(g(x) = 0 \text{ iff } h(x) = 0)$$

2. *We denote  $h = o(g)$  in  $a$  (German pronunciation:  $h$  ist klein  $o$  von  $g$ ) iff  $\exists U \subseteq D$ , with  $U$  being the neighborhood of  $a$ , and  $r : U \rightarrow \mathbb{R}$  such that  $\lim_{x \rightarrow a} r(x) = 0$  and  $h(x) = r(x) \cdot g(x) \forall x \in U$ . In that sense,*

$$\lim_{x \rightarrow a} \frac{h(x)}{g(x)} = 0$$

*Most often,  $O(\underbrace{\|x - x_0\|^n}_{g(x)})$  will be used (in this context of differentiability here)*

*with  $a = x_0$ .*



## 6.4 Multidimensional derivative of a function

**Definition 6.3** (Definition of the derivative of a function).

$$f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n \quad D \text{ is open, } x_0 \in D$$

$$f(x) = \begin{bmatrix} f_1(x) \\ f_2(x) \\ \vdots \\ f_n(x) \end{bmatrix} \in \mathbb{R}^n$$

$$f_i(x) = f_i \left( \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{bmatrix} \right) = f_i(x_1, x_2, \dots, x_m) \in \mathbb{R}$$

**Remark 6.3.** First trial to define the derivative:

$$f'(x_0) := \lim_{x \rightarrow x_0} \frac{\overbrace{f(x) - f(x_0)}^{\in \mathbb{R}^n}}{\underbrace{x - x_0}_{\in \mathbb{R}^m}}$$

Does not work because of incompatibility of dimensions (and we cannot divide vectors).

**Remark 6.4.** We use Taylor's Theorem to characterize  $f'(x_0)$ . A Taylor polynomial of 1st degree is given by

$$f(x) = f(x_0) + f'(x_0)(x - x_0) + R_f^2(x; x_0)$$

and  $R_f^2(x; x_0) = r(x)(x - x_0)$  with  $\lim_{x \rightarrow x_0} r(x) = 0$ . Hence,

$$f(x) - f(x_0) - f'(x_0)(x - x_0) = O(x - x_0)$$

or we insert the absolute operators:

$$|f(x) - f(x_0) - f'(x_0)(x - x_0)| = |r(x)| \cdot |x - x_0| = o(|x - x_0|)$$

where  $(f(x) - f(x_0)) \in \mathbb{R}^n$  for the multidimensional case and  $(x - x_0) \in \mathbb{R}^m$  for the multidimensional case. Thus,

$$= O(|x - x_0|)$$

**Definition 6.4.** Let  $f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$ ,  $D$  is open and  $x_0 \in D$ . We say, “ $f$  is differentiable in  $x_0$ ” (specifically, “Fréchet differentiable”) if there exists  $A \in \mathbb{R}^{n \times m}$  such that

$$\|f(x) - f(x_0) - A(x - x_0)\|_{\mathbb{R}^n} = o(\|x - x_0\|_{\mathbb{R}^m})$$

We call this condition differentiability condition.  $A$  is a linear approximation of  $f$  in  $x_0$ . Because of norm equivalence on  $\mathbb{R}^n$ , and accordingly  $\mathbb{R}^m$ , it is irrelevant which norm  $\|\cdot\|_{\mathbb{R}^n}$  and  $\|\cdot\|_{\mathbb{R}^m}$  is chosen.

**Lemma 6.3.** *Let  $f$  be, as in Definition 6.4, differentiable in  $x_0$ . Then the linear approximation  $A$ , by the differentiability condition, is uniquely determined.*

*Proof.* Assume  $A, B \in \mathbb{R}^{n \times m}$  satisfy the differentiability condition. Let  $r > 0$  such that  $K_r(x_0) \subseteq D$  (feasible because  $D$  is open and  $x_0 \in D$ ). Furthermore let  $v \in \mathbb{R}^m$  and  $\|v\| < r$ , hence  $x = x_0 + v \in K_r(x_0) \subset D$  and  $v = x - x_0$ .

$$\begin{aligned} \|(A - B)v\| &= \|Av - Bv\| = \|A(x - x_0) - B(x - x_0)\| \\ &= \|f(x) - f(x_0) - B(x - x_0) - (f(x) - f(x_0) - A(x - x_0))\| \\ &\leq \|f(x) - f(x_0) - B(x - x_0)\| + \|f(x) - f(x_0) - A(x - x_0)\| \end{aligned}$$

by the differentiability criterion

$$r(x) \cdot \|x - x_0\| + \tilde{r}(x) \|x - x_0\|$$

with  $\lim_{x \rightarrow x_0} r(x) = \lim_{x \rightarrow x_0} \tilde{r}(x) = 0$ . Hence, for  $\hat{r}(x) = r(x) + \tilde{r}(x)$ ,

$$\|(A - B)v\| \leq \hat{r}(x) \|x - x_0\| = \hat{r}(x) \|v\| = O(\|v\|) \text{ in } x_0$$

(Thus  $\lim_{x \rightarrow x_0} \hat{r}(x) = 0$ )

Show:  $(A - B)w = 0 \forall w \in \mathbb{R}^m$ . Assume  $\exists w \in \mathbb{R}^m, w \neq 0$  with  $(A - B)w \neq 0$ . For  $|\alpha| < \frac{r}{\|w\|}$  (with  $r$  as radius of the sphere),

$$\|\alpha w\| = |\alpha| \|w\| < \frac{r}{\|w\|} \cdot \|w\| = r$$

Let  $v = \alpha w$ . Then

$$\begin{aligned} \|(A - B)v\| &= |\alpha| \|(A - B)w\| \leq \hat{r}(x) \|\alpha w\| = |\alpha| \hat{r}(x) \cdot \|w\| \\ \implies \|(A - B)w\| &\leq \underbrace{\hat{r}(x)}_{\xrightarrow{\text{for } x \rightarrow x_0} 0} \cdot \underbrace{\|w\|}_{\text{constant}} \\ \implies (A - B)w &= 0 \end{aligned}$$

This contradicts with our assumption.

Therefore,  $Aw = Bw \forall w \in \mathbb{R}^m$ , hence  $A = B$ . □

## 6.5 Fréchet derivative

**Definition 6.5** (Part 2 of Definition 6.4). *If  $f$  is differentiable in  $x_0$ , then we call the uniquely determined linear map  $A$  the “Fréchet derivative” of  $f$  in  $x_0$  and denote  $A = Df(x_0)$ . An alternative notations are  $f'(x_0)$  and  $D_{x_0}f$ .*

**Lemma 6.4.** *Let  $f : D \rightarrow \mathbb{R}^n$ ,  $D \subseteq \mathbb{R}^m$  open,  $x_0 \in D$ . If  $f$  is differentiable in  $x_0$ , then  $f$  is also continuous in  $x_0$ .*

*Proof.* Let  $x \in D$ . Then,

$$\begin{aligned}
\|f(x) - f(x_0)\|_{\mathbb{R}^n} &= \|f(x) - f(x_0) - Df(x_0) \cdot (x - x_0) + Df(x_0)(x - x_0)\| \\
&\leq \|f(x) - f(x_0) - Df(x_0)(x - x_0)\| + \|Df(x_0)(x - x_0)\| \\
&\leq r(x) \cdot \underbrace{\|x - x_0\|}_{\rightarrow 0 \text{ for } x \rightarrow x_0} + \underbrace{\left\| \overbrace{Df(x_0)}^{\text{constant}} \right\|}_{\in \mathbb{R}^{n \times m}} \cdot \underbrace{\|x - x_0\|}_{\rightarrow 0 \text{ for } x \rightarrow x_0} \\
&\rightarrow 0 \text{ for } x \rightarrow x_0
\end{aligned}$$

Hence  $f$  is continuous in  $x_0$ .  $\square$

**Lemma 6.5.** Let  $f, g : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$ . Let  $f$  and  $g$  be differentiable in  $x_0 \in D$ . Let  $\lambda \in \mathbb{R}$ . Then,

1.  $f + g$  is differentiable in  $x_0$  with

$$D(f + g)(x_0) = Df(x_0) + Dg(x_0)$$

2.  $\lambda f$  is differentiable in  $x_0$  and

$$D(\lambda f)(x_0) = \lambda Df(x_0)$$

Thus differentiability is a linear operation on the vector space's appropriate differentiable functions.

*Proof.* Let  $F := \|(f + g)(x) - (f + g)(x_0) - [Df(x_0) + Dg(x_0)](x - x_0)\|$  with  $[Df(x_0) + Dg(x_0)] = D(f + g)(x_0)$ . Show that  $F = o(\|x - x_0\|)$ .

$$\begin{aligned}
F &\leq \|f(x) - f(x_0) - Df(x_0)(x - x_0)\| + \|g(x) - g(x_0) - Dg(x_0)(x - x_0)\| \\
&= o(\|x - x_0\|) + o(\|x - x_0\|) \\
&= o(\|x - x_0\|)
\end{aligned}$$

For  $\lambda f$  it holds analogously.  $\square$

**Lemma 6.6.** Let  $C : D \rightarrow \mathbb{R}^n$ .  $C(x) = k \in \mathbb{R}^n$  is constant. Then  $C$  is differentiable in every point  $x_0 \in D$  and  $DC(x_0) = 0 \in \mathbb{R}^{n \times m}$ . Let  $A : \mathbb{R}^m \rightarrow \mathbb{R}^n$  be linear. Then  $A$  is differentiable in every point  $x_0 \in \mathbb{R}^m$  and  $DA(x_0) = A$ .

Let  $f(x) = k + Ax : \mathbb{R}^m \rightarrow \mathbb{R}^n$  be linear affine, then  $f$  is differentiable in every point  $x_0 \in \mathbb{R}^m$  with  $Df(x_0) = A$ .

*Proof.*

$$\|c(x) - c(x_0) - 0 \cdot (x - x_0)\|$$




Figure 35: Chain rule in multiple dimensions

where 0 denotes the zero-matrix.

$$= \|k - k\| = \|0\| = 0 = o(\|x - x_0\|)$$

Hence 0 satisfies the differentiability condition for  $c$ .

$$\implies 0 = Dc(x_0)$$

in the linear case

$$\|Ax - Ax_0 - A(x - x_0)\| = \|Ax - Ax_0 - Ax + Ax_0\| = 0 = o(\|x - x_0\|)$$

hence  $DA(x_0) = A$ . Affine: use Lemma 6.5.

$$D(k + A)(x_0) = \underbrace{Dk(x_0)}_{=0} + \underbrace{DA(x_0)}_{=A} = A$$

This is analogous to the one-dimensional case:

$$(k + ax)' = a$$

□

## 6.6 Chain rule

**Theorem 6.1** (Chain rule in multiple dimensions). *Let  $D \subseteq \mathbb{R}^l$  be open. Let  $E \subseteq \mathbb{R}^m$  be open. Let  $f : D \rightarrow \mathbb{R}^m$  such that  $f(D) \subseteq E$  and  $g : E \rightarrow \mathbb{R}^n$ . Compare with Figure 35.*

*Let  $f$  in  $x_0$  be differentiable and  $g$  in  $y_0 = f(x_0)$  is differentiable. Then also  $g \circ f : D \rightarrow \mathbb{R}^n$  is differentiable in  $x_0$  and*

$$\underbrace{D(g \circ f)(x_0)}_{\in \mathbb{R}^{n \times l}} = \underbrace{Dg(f(x_0))}_{\in \mathbb{R}^{n \times m}} \underbrace{Df(x_0)}_{\in \mathbb{R}^{m \times l}}$$

(The dimensions match.)

*Proof.*  $(g \circ f)(x)$  is differentiable in  $x_0$  iff

$$\lim_{x \rightarrow x_0} \frac{\|g(f(x)) - g(f(x_0)) - Dg(f(x_0)) \cdot Df(x_0) \cdot (x - x_0)\|}{\|x - x_0\|} = 0$$

according to Fréchet differentiability. This is equivalent to

$$\frac{\|g(f(x)) - g(f(x_0)) - Dg(f(x_0)) \cdot Df(x_0) \cdot (x - x_0)\|}{\|x - x_0\|} < \varepsilon$$

for any bounded  $\varepsilon$ . Let  $0 < \varepsilon < 2$ . Show that

$$\frac{1}{\|x - x_0\|} \|g(f(x)) - g(f(x_0)) - Dg(y_0) \cdot Df(x_0)(x - x_0)\| < \varepsilon$$

for sufficiently small  $\|x - x_0\|$  with  $x \neq x_0$ .

$$\begin{aligned} & \frac{1}{\|x - x_0\|} \|g(f(x)) - g(f(x_0)) - Dg(y_0) \cdot Df(x_0)(x - x_0)\| \\ &= \frac{1}{\|x - x_0\|} \|g(f(x)) - g(f(x_0)) - Dg(f(x_0)) \cdot (f(x) - f(x_0)) \\ & \quad + Dg(f(x_0))(f(x) - f(x_0)) - Dg(f(x_0))Df(x_0)(x - x_0)\| \end{aligned}$$

recognize that we have a common factor  $Dg(f(x_0))$

$$\begin{aligned} & \leq \frac{1}{\|x - x_0\|} \|g(f(x)) - g(f(x_0)) - Dg(f(x_0))(f(x) - f(x_0))\| \\ & \quad + \frac{1}{\|x - x_0\|} \|Dg(y_0)\| \|f(x) - f(x_0) - Df(x_0)(x - x_0)\| \\ & =: (I) + (II) \end{aligned}$$

**Remark 6.5.**  $f$  is differentiable iff  $\lim_{x \rightarrow x_0} \frac{\|f(x) - f(x_0) - Df(x_0)(x - x_0)\|}{\|x - x_0\|} = 0$

$$\iff \forall \varepsilon > 0 \exists \delta > 0 : \|x - x_0\| < \delta \implies \|f(x) - f(x_0) - Df(x_0)(x - x_0)\| < \varepsilon$$

Thus, differentiability can also be characterized using  $\varepsilon$ - $\delta$ .

Choose  $\delta_1 > 0$  such that  $\|x - x_0\| < \delta_1 \implies$

$$\frac{1}{\|x - x_0\|} \cdot \|f(x) - f(x_0) - Df(x_0)(x - x_0)\| < \frac{\varepsilon}{2} \frac{1}{\|Dg(y_0)\| + 1}$$

is feasible, because  $f$  is differentiable in  $x_0$ .

$$\frac{\varepsilon}{2} \cdot \frac{1}{\|Dg(y_0)\| + 1} < \frac{2}{2} = 1$$

so it also holds that

$$\|f(x) - f(x_0) - Df(x_0)(x - x_0)\| < 1 \cdot \|x - x_0\|$$

By the reverse triangle inequality,  $\|x - y\| \geq \|x\| - \|y\|$ ,

$$\begin{aligned} \|f(x) - f(x_0) - Df(x_0)(x - x_0)\| &\geq \|f(x) - f(x_0)\| - \|Df(x_0)(x - x_0)\| \\ &\geq \|f(x) - f(x_0)\| - \|Df(x_0)\| \cdot \|x - x_0\| \\ &\implies \frac{\|f(x) - f(x_0)\|}{\|x - x_0\|} \leq \|Df(x_0)\| + 1 \end{aligned}$$

↓ This lecture took place on 2018/05/24.

$$\|f(x) - f(x_0)\| - \|Df(x_0)\| \|x - x_0\| < 1 \|x - x_0\|$$

hence, for  $x \neq x_0$

$$\frac{\|f(x) - f(x_0)\|}{\|x - x_0\|} < \|Df(x_0)\| + 1$$

$g$  is differentiable in  $y_0 = f(x_0)$ . Hence, we can choose  $\delta_g > 0$  such that  $\forall y \in E$  with  $\|y - y_0\| < \delta_g$  it holds that

$$\|g(y) - g(y_0) - Dg(y_0) \cdot (y - y_0)\| < \frac{\varepsilon}{2(\|Df(x_0)\| + 1)} \|y - y_0\|$$

Because  $f$  is continuous in  $x_0$ , there exists  $\delta_2 > 0$  such that  $x \in D$  and  $\|x - x_0\| < \delta_2 \implies \|f(x) - f(x_0)\| < \delta_g$ . Now let  $\delta = \min(\delta_1, \delta_2) > 0$ . Then

$$I := \frac{1}{\|x - x_0\|} \|g(f(x)) - g(f(x_0)) - Dg(f(x_0)) \cdot (f(x) - f(x_0))\|$$

Let  $y = f(x)$ ,  $y_0 = f(x_0)$ . Because  $\|f(x) - f(x_0)\| < \delta_g$  gives  $\|x - x_0\| < \delta_2$

$$\implies I < \frac{\varepsilon}{2(\|Df(x_0)\| + 1)} \underbrace{\frac{\|f(x) - f(x_0)\|}{\|x - x_0\|}}_{< \|Df(x_0)\| + 1} < \frac{\varepsilon}{2}$$

$$\begin{aligned} II &:= \|Dg(y_0)\| \frac{1}{\|x - x_0\|} \|f(x) - f(x_0) - Df(x_0)(x - x_0)\| \\ &< \|Dg(y_0)\| \cdot \frac{\varepsilon}{2} \cdot \frac{1}{\|Dg(y_0)\| + 1} < \frac{\varepsilon}{2} \end{aligned}$$

hence,  $I + II < \varepsilon$  for  $\|x - x_0\| < \delta$ . □

## 6.7 Differentiability on $D$

**Definition 6.6.** Let  $f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$  be differentiable in every point  $x \in D$ . Then we are used to say “ $f$  is differentiable on  $D$ ”.

In this case, we call the map

$$\begin{aligned} x &\mapsto Df(x) \\ D &\subseteq \mathbb{R}^m \rightarrow \mathbb{R}^{n \times m} \end{aligned}$$

the mapping function of  $f$  (dt. Abbildungsfunktion). If this function is continuous (in terms of  $\|\cdot\|_{\mathbb{R}}$  or  $\|\cdot\|_{\mathbb{R}^{n \times m}}$  . . . operator norm), then  $f$  is called continuously differentiable on  $D$ .

**Remark 6.6.** To define the differentiability of  $f$ , we require  $x_0$  to be an accumulation point of  $D$ . So  $x_0$  might also be a point on the boundary of  $D$ .

### 6.7.1 Determination of $Df(x_0) \in \mathbb{R}^{n \times m}$

**Definition 6.7.** Let  $f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$  be given.  $D$  is open,  $x_0 \in D$ ,  $v \in \mathbb{R}^m$  is arbitrary, but  $v \neq 0$ . We consider  $t \mapsto f(x_0 + tv)$  defined on  $(-\frac{r}{\|v\|}, \frac{r}{\|v\|})$  for  $r > 0$  such that  $K_r(x_0) \subseteq D$ .

$$\left(-\frac{r}{\|v\|}, \frac{r}{\|v\|}\right) \subseteq \mathbb{R} \rightarrow \mathbb{R}^n$$

Compare with Figure 36.

We define  $df(x_0; v) = \lim_{t \rightarrow 0} \frac{1}{t}(f(x_0 + tv) - f(x_0))$  if this limit exists.  $df(x_0, v)$  is called directional derivative of  $f$  in  $x_0$  in direction  $v$ . It is also called Gâteaux derivative of  $f$  in  $x_0$  in direction  $v$ .

**Remark 6.7.** How does it go together?

Derivative  $Df(x_0)$  and  $df(x_0; \cdot)$ . Assumption: Let  $f$  be differentiable in  $x_0$ . We define  $l_{x_0, v}(t) = x_0 + tv$  where  $tv$  is the linear part.

$$l_{x_0, v} : \left(-\frac{r}{\|v\|}, \frac{r}{\|v\|}\right) \rightarrow D$$

$l_{x_0, v}$  is linear affine from  $\mathbb{R}$  to  $\mathbb{R}^m$ .

$$Dl_{x_0, v}(0) = V \in \mathbb{R}^{m \times 1}$$

with  $V$  as linear part of  $l_{x_0, v}$ .

$$l_{x_0, v}(0) = x_0$$

$$f(x_0 + tv) = f \circ l_{x_0, v}(t)$$

Therefore ( $\rightarrow$  chain rule)

$$D(f \circ l_{x_0, v})(0) = Df(l_{x_0, v}(0)) \cdot Dl_{x_0, v}(0) = Df(x_0) \cdot v$$




Figure 36: Setting in Definition 6.7. The line is given by  $g = \{x_0 + tv : t \in \mathbb{R}\}$

On the other side,

$$\begin{aligned}
 0 &= \lim_{t \rightarrow 0} \underbrace{\frac{1}{|t|} |f_{x_0,v}(t) - f_{x_0,v}(0) - Df_{x_0,v}(0) \cdot t|}_{=0} \\
 &= \lim_{t \rightarrow 0} \frac{1}{|t|} |f(x_0 + tv) - f(x_0) - Df_{x_0,v}(0) \cdot t| \\
 &= \lim_{t \rightarrow 0} \left| \frac{f(x_0 + tv) - f(x_0)}{t} - Df_{x_0,v}(0) \right| = 0
 \end{aligned}$$

therefore

$$Df_{x_0,v}(0) = \lim_{t \rightarrow 0} \frac{f(x_0 + tv) - f(x_0)}{t} = df(x_0; v)$$

**Lemma 6.7.** Let  $f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$  in  $x_0 \in D$  (Fréchet) differentiable with derivative  $Df(x_0)$ . Then also the directional derivative  $df(x_0; v)$  for every direction  $v \in \mathbb{R}^m \setminus \{0\}$  and

$$df(x_0; v) = Df(x_0) \cdot v$$

**Remark 6.8.**  $v \mapsto df(x_0; v)$  is linear. We can derive the structure of the derivative matrix. Let  $f$  as above. Let  $\mathcal{B} = \{e_1, \dots, e_m\}$  be the canonical basis in  $\mathbb{R}^m$ . Then:  $Df(x_0) \cdot e_j$  is the  $j$ -th column of  $Df(x_0)$  for  $j = 1, \dots, m$ . On the other hand,

$$Df(x_0) \cdot e_j = df(x_0; e_j) = \lim_{t \rightarrow 0} \frac{1}{t} [f(x_0 + te_j) - f(x_0)]$$



$$= \begin{bmatrix} \lim_{t \rightarrow 0} \frac{1}{t} (f_1(x_0 + te_j) - f_1(x_0)) \\ \vdots \\ \lim_{t \rightarrow 0} \frac{1}{t} (f_n(x_0 + te_j) - f_n(x_0)) \end{bmatrix} \text{ for } f(x) = \begin{bmatrix} f_1(x) \\ \vdots \\ f_n(x) \end{bmatrix} \in \mathbb{R}^n$$

**Remark 6.9** (Notation). Consider  $x$  instead of  $x_0$

$$x = \begin{bmatrix} x_1 \\ \vdots \\ x_m \end{bmatrix} = [x_1, \dots, x_m]^t$$

Instead of  $f(x) = f\left(\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_m \end{bmatrix}\right)$  we also write  $f(x_1, x_2, \dots, x_m)$ .

## 6.8 Partial derivative and Jacobi matrix

**Definition 6.8.** Let  $f : D \rightarrow \mathbb{R}^n$ ,  $x \in D$  as above.  $f$  is differentiable in  $x$ . Then we let

$$\begin{aligned} \frac{\partial f}{\partial x_j}(x) &= df(x; e_j) = \lim_{t \rightarrow 0} \frac{1}{t} [f(x + te_j) - f(x)] \\ &= \lim_{t \rightarrow 0} \frac{1}{t} [f(x_1, \dots, x_j + t, \dots, x_m) - f(x_1, \dots, x_j, \dots, x_m)] \end{aligned}$$

and we call  $\frac{\partial f}{\partial x_j}(x)$  the partial derivative of  $f$  of variable  $x_j$  in point  $x$ .

Notations for  $\frac{\partial f}{\partial x_j}$ :

$$f_{x_j} \quad f_j \quad \partial_j f$$

The second notation is ambiguous. We will prefer the last one.

**Remark 6.10.**

$$\frac{\partial f}{\partial x_j}(x_0) = \begin{bmatrix} df_1(x; e_j) \\ df_2(x; e_j) \\ \vdots \\ df_n(x; e_j) \end{bmatrix} = \begin{bmatrix} \frac{\partial f_1}{\partial x_j}(x) \\ \vdots \\ \frac{\partial f_n}{\partial x_j}(x) \end{bmatrix} \in \mathbb{R}^n$$

Because  $\frac{\partial f}{\partial x_j}(x)$  is the  $j$ -th column of  $Df(x_0)$ , we get

$$(Df(x))_{i,j} = \frac{\partial f_i}{\partial x_j}(x) = \partial_j f_i(x)$$

We say,  $Df(x) = (\partial_j f_i(x))_{\substack{i=1, \dots, n \\ j=1, \dots, m}}$  is the Jacobi matrix of  $f$ .

**Remark 6.11.** The Jacobi matrix can exist even though the derivative does not exist. If the derivative exists, the Jacobi matrix exists for sure.

**Remark 6.12.**

$$\frac{\partial f}{\partial x_j} = \lim_{t \rightarrow 0} \frac{1}{t} \left[ f(x_1, \dots, x_j + t, \dots, x_m) - f(x_1, \dots, x_j, \dots, x_n) \right]$$

Thus, consider  $x_j$  as derivation variable and all  $x_k$  for  $k \neq j$  as constant parameters.

**Example 6.2.** Consider  $f : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  with

$$f(x_1, x_2, x_3) = \begin{bmatrix} x_1 x_3^2 + \sin(x_1 x_3) \\ \frac{x_2^2}{x_1^2 + 1} \end{bmatrix}$$

$$\partial_1 f(x_1, x_2, x_3) = \begin{bmatrix} 1 \cdot x_3^2 \\ -x_2^2 (x_1^2 + 1)^{-2} \cdot 2x_1 \end{bmatrix} = \begin{bmatrix} x_3^2 \\ -2 \frac{x_1 x_2^2}{(x_1^2 + 1)^2} \end{bmatrix}$$

$$\partial_2 f(x_1, x_2, x_3) = \begin{bmatrix} x_3 \cos(x_2 x_3) \\ \frac{2x_2}{x_1^2 + 1} \end{bmatrix}$$

$$\partial_3 f(x_1, x_2, x_3) = \begin{bmatrix} 2x_1 x_3 + x_2 \cos(x_2 x_3) \\ 0 \end{bmatrix}$$

Jacobi-Matrix

$$Df(x) = \left[ \frac{\partial f_i}{\partial x_j} \right]_{\substack{i=1, \dots, 2 \\ j=1, \dots, 3}} = \begin{bmatrix} x_3^2 & x_3 \cos(x_2 x_3) & 2x_1 x_3 + x_2 \cos(x_2 x_3) \\ -\frac{2x_1 x_2^2}{(x_1^2 + 1)^2} & \frac{2x_2}{x_1^2 + 1} & 0 \end{bmatrix}$$

**Remark 6.13.** Existence of partial derivatives of  $f$  does not suffice to ensure Fréchet-differentiability.

↓ This lecture took place on 2018/05/29.

Usually, we always have to point out which norm is used to define differentiability. Of course, in  $\mathbb{R}$  itself, all norms are equivalent.

**Remark 6.14.** Let  $f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$  be given. Let  $\|\cdot\|_{1,m}$  and  $\|\cdot\|_{2,m}$  be two equivalent norms on  $\mathbb{R}^m$  (norm equivalence theorem) and  $\|\cdot\|_{1,n}$  and  $\|\cdot\|_{2,n}$  are equivalent norms on  $\mathbb{R}^n$ .

Let  $f$  in  $x_0 \in D$  be differentiable with respect to  $\|\cdot\|_{1,m}$  and  $\|\cdot\|_{1,n}$ . Then also  $f$  is differentiable in  $x_0$  with respect to  $\|\cdot\|_{2,m}$  and  $\|\cdot\|_{2,n}$ .

Rationale: Let  $c \|x\|_{2,m} \leq \|x\|_{1,m} \leq C \|x\|_{2,m}$  and  $k \|y\|_{2,n} \leq \|y\|_{1,n} \leq K \|y\|_{2,n}$ , then

$$\begin{aligned} & \frac{\|f(x) - f(x_0) - A(x - x_0)\|_{2,n}}{\|x - x_0\|_{2,m}} \\ & \leq \frac{\frac{1}{k} \|f(x) - f(x_0) - A(x - x_0)\|_{1,n}}{\frac{1}{c} \|x - x_0\|_{1,m}} \\ & = \frac{c}{k} \underbrace{\frac{\|f(x) - f(x_0) - A(x - x_0)\|_{1,n}}{\|x - x_0\|_{1,m}}}_{\xrightarrow{x \rightarrow x_0} 0} \end{aligned}$$

Let  $f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$ ,  $f(x) = \begin{bmatrix} f_1(x) \\ \vdots \\ f_n(x) \end{bmatrix}$ . Then  $f$  is differentiable in  $x_0 \in D \iff f_k : D \rightarrow \mathbb{R}$  is differentiable for all  $k \in \{1, \dots, n\}$ .

Rationale: Let  $f$  be differentiable in  $x_0$ . Let  $A = \begin{bmatrix} a_1 \\ \vdots \\ a_n \end{bmatrix}$  be the derivative of  $f$ . Let  $a_k$  be the rows of  $A$ .  $f$  is differentiable in  $x_0$ , so

$$\begin{aligned} & \frac{\|f(x) - f(x_0) - A(x - x_0)\|_{\infty}}{\|x - x_0\|} \xrightarrow{x \rightarrow x_0} 0 \\ \iff & \frac{|f_n(x) - f_k(x) - a_k(x - x_0)|}{\|x - x_0\|} \xrightarrow{x \rightarrow x_0} 0 \end{aligned}$$

where  $a_k$  is a row vector and  $x - x_0$  is a column vector and  $k \in \{1, \dots, n\}$ .

$$\iff f_k \text{ is differentiable in } x_0$$

**Example 6.3** (Counterexample). We define  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ .

$$f(x_1, x_2) = \begin{cases} \frac{x_1^2 x_2}{x_1^2 + x_2^2} & \text{for } \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix} \\ 0 & \text{for } \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \end{cases}$$

Partial derivatives exist in every point  $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ .

$$\partial_1 f(x_1, x_2) = \frac{2x_1 x_2 (x_1^2 + x_2^2) - x_1^2 x_2 2x_1}{(x_1^2 + x_2^2)^2} = \frac{2x_1 x_2^3}{(x_1^2 + x_2^2)^2}$$

$$\partial_2 f(x_1, x_2) = \frac{x_1^2(x_1^2 + x_2^2) - x_1^2 x_2 \cdot 2x_2}{(x_1^2 + x_2^2)^2} = \frac{x_1^2(x_1^2 x_2^2)}{(x_1^2 + x_2^2)^2}$$

$$\partial_1 f(0, 0) = \lim_{x_1 \rightarrow 0} \frac{1}{x_1} \underbrace{[f(x_1, 0)]}_{=0} - \underbrace{f(0, 0)}_{=0} = 0$$

$$\partial_2 f(0, 0) = \lim_{x_2 \rightarrow 0} \frac{1}{x_2} \underbrace{[f(0, x_2)]}_{=0} - \underbrace{f(0, 0)}_{=0} = 0$$

If  $f$  would be differentiable in  $\begin{bmatrix} 0 \\ 0 \end{bmatrix}$  then  $Df(0) = [00]$ ,  $df(0, v) = Df(0) \cdot v$ . Choose  $v = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \implies df(0; v) = [00] \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 0$ . But!

$$\begin{aligned} df(0; v) &= \lim_{t \rightarrow 0} \frac{1}{t} [f(tv) - f(0)] = \lim_{t \rightarrow 0} \frac{1}{t} [f(t, t) - f(0, 0)] \\ &= \lim_{t \rightarrow 0} \frac{1}{t} \left[ \frac{t^2 \cdot t}{t^2 + t^2} - 0 \right] = \lim_{t \rightarrow 0} \frac{t^3}{2t^3} = \frac{1}{2} \end{aligned}$$

## 6.9 Total differentiability, nabla operator and gradients

**Remark 6.15.** Notation:  $f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}$ . Often, we denote  $df(x_0)$  ( $\in \mathbb{R}^{1 \times m}$ , row vector) instead of  $Df(x_0)$  and we call  $df(x_0)$  the total derivative of  $f$ . We let

$$\nabla f(x_0) = df(x_0)^t = [Df(x_0)]^t = \begin{bmatrix} \partial_1 f(x_0) \\ \partial_2 f(x_0) \\ \vdots \\ \partial_n f(x_0) \end{bmatrix}$$

$\nabla f(x_0)$  is called gradient of  $f$  in  $x_0$ . We call  $\nabla$  the nabla operator. It is also denoted  $\text{grad}(f)$  instead of  $\nabla f$ .

$$df(x_0; v) = df(x_0) \cdot v = \nabla f(x_0)^t \cdot v = \langle \nabla f(x_0), v \rangle_{\mathbb{R}^m}$$

**Remark 6.16.** Let  $f : D \subseteq \mathbb{R}^2 \rightarrow \mathbb{R}$  continuously differentiable be given on  $D$ . Consider  $\Gamma_s = \{x \in D \mid f(x) = s\}$ . Compare with Figure 37.

Assume  $\Gamma_s$  is a graph of a family of curves (dt. Parametrisierte Kurve)  $\gamma_s : I \rightarrow D$ . Let  $I$  be an interval.  $\Gamma_s = \{\gamma_s(t) \mid t \in I\}$ . We assume, that  $\gamma_s$  is regular, hence  $\gamma$  is differentiable and  $\gamma'_s(t) = \begin{bmatrix} \gamma'_1(t) \\ \gamma'_2(t) \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix} \forall t \in I$ .  $\gamma'_s(t)$  is tangential vector on  $\Gamma_s$  in point  $x = \gamma_s(t)$ .

$$\langle \nabla f(\gamma_s(t)), \gamma'_s(t) \rangle = \langle \nabla f(x), v \rangle = 0$$




Figure 37:  $\Gamma_s$  is the niveau level of  $f$

$$\underbrace{f(\gamma_s(t))}_{\in \Gamma_s} = s \quad \dots \text{constant}$$

$$\implies \frac{d}{dt}[f(\gamma_s(t))] = 0$$

where

$$\begin{aligned} \frac{d}{dt}f(\gamma_s(t)) &= \underbrace{Df(x)}_{\nabla f(x)^t} \cdot \underbrace{D\gamma_s(x)}_{\gamma'_s(t)} = \nabla f(x)^t \cdot \gamma'_s(t) \\ &= \langle \nabla f(x), \gamma'_s(t) \rangle \end{aligned}$$

Compare with Figure 38.

**Theorem 6.2.** This theorem establishes the relation of differentiability and partial derivatives.

Let  $f : D \rightarrow \mathbb{R}$ ,  $D \subseteq \mathbb{R}^m$  be open. Assume all partial derivatives  $\partial_i f(x)$  exist  $\forall x \in D$  for  $j = 1, \dots, m$  and the functions  $x \mapsto \partial_j f(x)$  on  $D \rightarrow \mathbb{R}$  are continuous for  $j = 1, \dots, m$ . Then  $f$  is differentiable in every point  $x_0 \in D$  with  $Df(x_0) = [\partial_1 f(x_0), \dots, \partial_m f(x_0)]$ . Then  $f$  is also continuously differentiable.

*Proof.* Proof idea: We approximate  $x_0$  (starting from  $x_0 \in D$ ) along lines parallel to the coordinate system axes. Compare with Figure 39.




Figure 38:  $Df(x) = \nabla f(x)^T$

$$x_0 := \begin{bmatrix} x_1^0 \\ \vdots \\ x_m^0 \end{bmatrix} \quad x := \begin{bmatrix} x_1 \\ \vdots \\ x_m \end{bmatrix} \quad \xi_0 := x_0 = \begin{bmatrix} x_1^0 \\ x_2^0 \\ \vdots \\ x_m^0 \end{bmatrix} \quad \xi_1 := \begin{bmatrix} x_1 \\ x_2^0 \\ \vdots \\ x_m^0 \end{bmatrix} \quad \xi_2 := \begin{bmatrix} x_1 \\ x_2 \\ x_3^0 \\ \vdots \\ x_m^0 \end{bmatrix}$$

$$\xi_k := \begin{bmatrix} x_1 \\ \vdots \\ x_{k_0} \\ x_{k+1} \\ \vdots \\ x_m^0 \end{bmatrix} \quad \dots \quad \xi_m := \begin{bmatrix} x_0 \\ \vdots \\ x_m \end{bmatrix} = x$$

where  $\xi_i$  are “intermediate” points. It holds that  $\xi_k + (x_{k+1} - x_{k+1}^0) \cdot e_{k+1} = \xi_{k+1}$  for  $k = 0, \dots, m-1$ . Define  $\varphi_k : [0, 1] \rightarrow \mathbb{R}$ .

$$\varphi_k(t) = f(\xi_k + t(x_{k+1} - x_{k+1}^0) \cdot e_{k+1})$$

then  $\varphi_k(0) = f(\xi_k)$ ;  $\varphi_k(1) = f(\xi_{k+1})$ .  $\varphi'_k(t) = ?$

$$\underbrace{f(\xi_k + t(x_{k+1} - x_{k+1}^0) \cdot e_{k+1})}_{=\varphi_k(t)} = f(x_1, \dots, x_k, \underbrace{x_{k+1}^0 + t(x_{k+1} - x_{k+1}^0)}_{(k+1)\text{-th variable}}, x_{k+2}^0, \dots, x_m^0)$$

$$\varphi'_k(t) = \frac{d}{dt} \left[ f(x_1, \dots, x_k, \underbrace{x_{k+1}^0 + t(x_{k+1} - x_{k+1}^0)}_{(k+1)\text{-th variable}}, x_{k+2}^0, \dots, x_m^0) \right]$$




Figure 39:  $x_0$  approximates  $x$  along lines parallel to the coordinate system axes

$$\begin{aligned}
 &= \partial_{k+1} f(x_1, \dots, x_k, x_{k+1}^0 + t(x_{k+1} - x_{k+1}^0), x_{k+2}^0, \dots, x_m^0) \cdot (x_{k+1} - x_{k+1}^0) \\
 &= \partial_{k+1} f(\xi_k + t(x_{k+1} - x_{k+1}^0) \cdot e_{k+1}) \cdot (x_{k+1} - x_{k+1}^0)
 \end{aligned}$$

$\varphi_k$  is continuously differentiable on  $[0, 1]$  because  $\partial_{k+1} f$  is continuous. By the mean value theorem of differential calculus, it follows that some  $\tau_{k+1} \in (0, 1)$  exists such that

$$\begin{aligned}
 \varphi(1) - \varphi(0) &= \varphi'(\tau_{k+1}) \cdot (1 - 0) \\
 \implies f(\xi_{k+1}) - f(\xi_k) &= \partial_{k+1} f(\xi_k + \tau_{k+1}(x_{k+1} - x_{k+1}^0) \cdot e_{k+1}) \cdot (x_{k+1} - x_{k+1}^0)
 \end{aligned}$$

For differentiability, we have to show:

$$\lim_{x \rightarrow x_0} \frac{1}{\|x - x_0\|} \left| f(x) - f(x_0) - [\partial_1 f(x_0), \dots, \partial_m f(x_0)] \begin{bmatrix} x_1 - x_1^0 \\ x_2 - x_2^0 \\ \vdots \\ x_m - x_m^0 \end{bmatrix} \right| = 0 \quad (*)$$

Choose  $\|x\| = \|x\|_\infty$  on  $\mathbb{R}^m$ .

$$\frac{1}{\|x - x_0\|} \left| \underbrace{f(x)}_{\xi_m} - \underbrace{f(x_0)}_{\xi_0} - \sum_{k=1}^m \partial_k f(x_0)(x_k - x_k^0) \right|$$

$$\begin{aligned}
&= \frac{1}{\|x - x_0\|} \left| \underbrace{\sum_{k=1}^m (f(\xi_k) - f(\xi_{k-1}) - \partial_k f(x_0)(x_k - x_k^0))}_{\text{telescoping sum}} \right| \\
&\stackrel{\text{triangle ineq.}}{\leq} \frac{1}{\|x - x_0\|} \sum_{k=1}^m \left| \partial_k f(\xi_{k-1} + \tau_k(x_k - x_k^0) \cdot e_k)(x_k - x_k^0) - \partial_k f(x_0)(x_k - x_k^0) \right| \\
&= \sum_{k=1}^m \underbrace{\frac{|x_k - x_k^0|}{\|x - x_0\|}}_{\leq 1} \cdot \left| \partial_k f(\xi_{k-1} + \tau_k(x_k - x_k^0) \cdot e_k) - \partial_k f(x_0) \right| \quad (\#)
\end{aligned}$$

Choose  $\varepsilon > 0$  arbitrary. By continuity of  $\partial_k f(x)$ , there exists  $\delta > 0$  such that  $\|y - y_0\|_\infty < \delta \implies \|\partial_k f(y) - \partial_k f(x_0)\| < \frac{\varepsilon}{m}$  for every  $k \in \{1, \dots, m\}$ . Choose  $x$  such that  $\|x - x_0\| \leq \delta$ . We consider

$$\begin{aligned}
\xi_{k-1} + \tau_k(x_k - x_k^0) \cdot e_k - x_0 &= \begin{bmatrix} x_1 \\ \vdots \\ x_{k-1} \\ x_k^0 \\ \vdots \\ x_m \end{bmatrix} + \begin{bmatrix} 0 \\ \vdots \\ 0 \\ \tau_k(x_k - x_k^0) \\ 0 \\ \vdots \\ 0 \end{bmatrix} - \begin{bmatrix} x_1^0 \\ \vdots \\ x_{k-1}^0 \\ x_k^0 \\ \vdots \\ x_m^0 \end{bmatrix} = \begin{bmatrix} x_1 - x_1^0 \\ \vdots \\ x_{k-1} - x_{k-1}^0 \\ \tau_k(x_k - x_k^0) \\ 0 \\ \vdots \\ 0 \end{bmatrix} \\
&\implies \|\xi_{k-1} + \tau_k(x_k - x_k^0) \cdot e_k - x_0\|_\infty \\
&= \max \left\{ |x_1 - x_1^0|, |x_2 - x_2^0|, \dots, |x_{k-1} - x_{k-1}^0|, \underbrace{\tau_k}_{\in(0,1)} |x_k - x_k^0| \right\} \\
&\leq \|x - x_0\|_\infty < \delta \\
&\implies \left| \partial_k f(\xi_{k-1} + \tau_k(x - x_0^k)e_k) - \partial_k f(x_0) \right| < \frac{\varepsilon}{m} \\
(\#) &\leq 1 \cdot \sum_{k=1}^m \frac{\varepsilon}{m} = m \cdot \frac{\varepsilon}{m} = \varepsilon
\end{aligned}$$

Thus,  $f$  is differentiable in  $x_0$ .

Because  $Df(x) = [\partial_1 f(x), \dots, \partial_m f(x)]$  depends continuously on  $x$ ,  $f$  is continuously differentiable on  $D$   $\square$



**Corollary.** Let  $f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$ .

$$f(x) = \begin{bmatrix} f_1(x) \\ \vdots \\ f_n(x) \end{bmatrix}$$

such that all partial derivatives  $\partial_k f_i(x)$  exist and are continuous in  $x$  (for  $k = 1, \dots, m$ ;  $i = 1, \dots, n$ ). Then  $f$  is continuously differentiable on  $D$ .

*Rationale.* Use  $f$  is continuously differentiable on  $D \iff f_i$  is continuously differentiable for  $i = 1, \dots, n$  and use Theorem 6.2.  $\square$

**Example 6.4** (Counterexample).

$$f(x_1, x_2) = \frac{x_1^2 x_2}{x_1^2 + x_2^2} \quad \partial_1 f(x) = \frac{2x_1 x_2^3}{(x_1^2 + x_2^2)^2} \quad \partial_2 f(x) = \frac{x_1^2(x_1^2 - x_2^2)}{(x_1^2 + x_2^2)^2}$$

$$\partial_1 f(0) = \partial_2 f(0) = 0$$

$$\partial_1 f\left(\begin{bmatrix} \varepsilon \\ \varepsilon \end{bmatrix}\right) = \frac{2\varepsilon \varepsilon^3}{(\varepsilon^2 + \varepsilon^2)^2} = \frac{2\varepsilon^4}{4\varepsilon^4} = \frac{1}{2} \not\rightarrow 0$$

Therefore,  $\partial_1 f$  is not continuous in 0.

↓ This lecture took place on 2018/06/05.

## 6.10 Optimality criteria for multi-dimensional functions

Necessary (but not sufficient) optimality criteria:

**Definition 6.9.** Let  $D \subseteq \mathbb{R}^n$ ,  $f : D \rightarrow \mathbb{R}$ . We say that  $f$  in  $x_0 \in D$  has a local maximum, if some neighborhood  $U$  of  $x_0$  exists such that

$$\forall x \in D \cap U : f(x) \leq f(x_0)$$

and accordingly for strict maxima:

$$\forall x \in (D \cap U) \setminus \{x_0\} : f(x) < f(x_0)$$

Analogously for a minimum with  $f(x) \geq f(x_0)$  and strict minima  $f(x) > f(x_0)$ .

**Lemma 6.8.** Let  $f : D \rightarrow \mathbb{R}$  be given. Let  $D \subseteq \mathbb{R}^n$  be open. Let  $x_0 \in D$  be a local maximum or a local minimum and let  $f$  in  $x_0$  be differentiable. Then  $Df(x_0) = [0, \dots, 0]$  or  $\nabla f(x_0) = \begin{bmatrix} 0 & 0 & \dots & 0 \end{bmatrix}^T$ .

*Proof.* We consider one-dimensional  $l_{e_k, x_0} : (-\varepsilon, \varepsilon) \rightarrow D$  with  $l_{e_k, x_0}(t) \mapsto x_0 + t \cdot e_k$  and  $l_{e_k, x_0}(0) \mapsto x_0$ .  $f$  has a maximum in  $x_0$  (without loss of generality)  $\implies f \circ l_{e_k, x_0}$  (differentiable in  $t = 0$ ) is a local maximum in  $t = 0$ . This is necessary for the optimality of  $t = 0$  for  $f \circ l_{e_k, x_0}$  that  $(f \circ l_{e_k, x_0})'(0) = 0 \implies Df(x_0) \cdot e_k = \partial_k f(x_0) = 0 \implies Df(x_0) = [0, \dots, 0]$ .  $\square$

**Remark 6.17.** By Lemma 6.8 it follows immediately that  $x_0$  is optimal.

$$\implies \underbrace{Df(x_0) \cdot v}_{=0} = df(x_0; v) = 0 \quad \forall v \in \mathbb{R}^n$$

## 6.11 Diffeomorphism

**Definition 6.10.** Let  $U, V \subseteq \mathbb{R}^n$  be open. We call  $f : U \rightarrow V$  a diffeomorphism if

1.  $f : U \rightarrow V$  is bijective
2.  $f$  is continuously differentiable on  $U$
3.  $f^{-1} : V \rightarrow U$  is continuously differentiable on  $V$

Let  $f : U \rightarrow V$ ,  $x_0 \in U$  and  $y_0 = f(x_0) \in V$ . We say:  $f$  is a local diffeomorphism in  $x_0$  if open neighborhoods  $x_0 \in U' \subset U$  and  $y_0 \in V' \subseteq V$  such that  $f : U' \rightarrow V'$  is a diffeomorphism.

**Lemma 6.9.** Let  $g : D \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^n$  and  $D$  is open.  $g$  is continuously differentiable on  $D$ . Furthermore let  $g : D \rightarrow g(D)$  be bijective,  $g^{-1} : g(D) \rightarrow D$  and  $Dg(x)$  is regular for all  $x \in D$ . Then  $g(D) \rightarrow D$  is continuously differentiable with

$$Dg^{-1}(y) = Dg^{-1}(g(x)) = [Dg(x)]^{-1}$$

*Proof.* Will be provided later on (page 121).  $\square$

## 6.12 Local inversion theorem

**Theorem 6.3** (Local Inversion Theorem, Theorem of Inverse Maps). Let  $D \subseteq \mathbb{R}^n$  be open. Let  $f : D \rightarrow \mathbb{R}^n$  be continuously differentiable. Furthermore let  $Df(x_0)$  be regular for  $x_0 \in D$ . Then  $f$  is a local diffeomorphism in  $x_0$ . Furthermore:

$$Df^{-1}(f(x_0)) = [Df(x_0)]^{-1}$$

*Proof.*

$$\begin{aligned} \tilde{f}(\xi) &= f(\xi + x_0) - f(x_0) \\ \tilde{f} : D - x_0 = \{\xi = x - x_0 : x \in D\} &\rightarrow \mathbb{R}^n \\ \tilde{f}(0) &= f(x_0) - f(x_0) = 0 \\ D\tilde{f}(0) &= Df(x_0) \text{ is regular} \end{aligned}$$

**Claim.**  $f$  is a local diffeomorphism  $\iff \tilde{f}$  is a local diffeomorphism.

*Proof.* **Direction  $\Leftarrow$**  Let  $\tilde{f}$  be a local diffeomorphism.

$$\tilde{f}(\xi) = \eta \iff \underbrace{f(\xi + x_0)}_{=x} = \underbrace{\eta + f(x_0)}_y \iff f(x) = y$$

and  $\xi = \tilde{f}^{-1}(\eta)$ . Also,

$$x = \xi + x_0 = \tilde{f}^{-1}(y - f(x_0)) + x_0$$

Thus,  $f$  is invertible and

$$f^{-1}(y) = \tilde{f}^{-1}(y - f(x_0)) + x_0$$

hence  $f^{-1}$  is continuously differentiable, hence  $f$  is a local diffeomorphism.

**Direction  $\Rightarrow$**  Analogous.

□

$$F(x) = \underbrace{[Df(x_0)]^{-1}}_{\in \mathbb{R}^{n \times n}} \tilde{f}(x)$$

$$v \mapsto [Df(x_0)]^{-1} \cdot v \text{ linear} \implies \text{diffeomorphism}$$

$$[Df(x_0)]^{-1} \cdot \tilde{f} \text{ is a local diffeomorphism} \iff \tilde{f} \text{ is a local diffeomorphism.}$$

$$DF(0) = [Df(x_0)]^{-1} \cdot \underbrace{D\tilde{f}(0)}_{Df(x_0)} = E$$

Thus, it suffices to show that  $F$  is a local diffeomorphism where  $F(0) = 0$  and  $DF(0) = I$  (the unit matrix).

Now the interesting part of the proof begins: Let  $y \in \mathbb{R}^n$  arbitrary. We define

$$\varphi(x) = y + x - F(x)$$

$y = F(x) \iff y + x - F(x) = x$ , hence  $\varphi_Y(x) = x$ , hence  $x$  is a fixed point of  $\varphi_Y$ .

$F$  is continuously differentiable with  $DF(0) = I$ . Thus for every  $\varepsilon > 0 \exists \delta > 0$  such that

$$\|x - 0\| \leq \delta \implies \|DF(x) - DF(0)\| = \|DF(x) - I\| \leq \varepsilon$$

Choose  $r > 0$  such that  $\|x\| \leq 2r \implies \|DF(x) - I\| \leq \frac{1}{2}$ . Additionally let  $r$  be sufficiently small such that

$$\overline{K_{2r}(0)} \subseteq \tilde{D} = D - x_0$$




Figure 40: We consider a sphere with radius  $2r$ . The blue line denotes  $l_{x_1, x_2}(t)$ .  $l_{x_1, x_2}(t) \in \overline{K_{2r}(0)}$  because  $\overline{K_{2r}(0)}$  is convex

Recall that  $\tilde{D}$  is the domain of  $F$ . Let  $x_1, x_2 \in \overline{K_{2r}(0)}$  and let

$$l_{x_1, x_2} : \begin{matrix} t \mapsto (1-t)x_1 + tx_2 \\ [0, 1] \rightarrow \overline{K_{2r}(0)} \end{matrix}$$

Compare with Figure 40.

$$l_{x_1, x_2}(0) = x_1 \quad l_{x_1, x_2}(1) = x_2$$

$$\begin{aligned} \|\varphi_y(x_2) - \varphi_y(x_1)\| &= \|\varphi_y \circ l_{x_1, x_2}(1) - \varphi_y \circ l_{x_1, x_2}(0)\| \\ &= \left\| \int_0^1 \frac{d}{dt} [\varphi_Y \circ l_{x_1, x_2}(t)] dt \right\| \\ &= \left\| \int_0^1 D\varphi_Y(l_{x_1, x_2}(t)) \cdot \underbrace{(x_2 - x_1)}_{\substack{\text{inner derivative} \\ l_{x_1, x_2}(t) \cdot dt}} dt \right\| \\ &\leq \int_0^1 \left\| \underbrace{D\varphi_Y(l_{x_1, x_2}(t))}_{\in \overline{K_{2r}(0)}} \right\| \cdot \underbrace{\|(x_2 - x_1)\|}_{\text{constant}} dt \\ &\leq (*) \end{aligned}$$

$$D\varphi_Y(x) = I - DF(x)$$

hence  $\forall x \in \overline{K_{2r}(0)}, \|D\varphi_Y(x)\| = \|I - DF(x)\| \leq \frac{1}{2}$ .

$$(*) \leq \frac{1}{2} \|x_2 - x_1\| \underbrace{\int_0^1 1 \cdot dt}_{=1} = \frac{1}{2} \|x_2 - x_1\|$$

Hence  $\forall x_1, x_2 \in \overline{K_{2r}(0)}$ ,

$$\|\varphi_Y(x_2) - \varphi_Y(x_1)\| \leq \frac{1}{2} \|x_2 - x_1\|$$

Therefore  $\varphi_Y$  is a contraction with constant  $\nu = \frac{1}{2}$ .

$K_{2r}(0) \subseteq \mathbb{R}^n$  is a complete, metric space. It remains to show:  $\varphi_Y : \overline{K_{2r}(0)} \rightarrow \overline{K_{2r}(0)}$ . This only holds if  $y$  is sufficiently small.

Let  $y \in K_r(0)$  and  $x \in \overline{K_{2r}(0)}$ .

$$\begin{aligned} \|\varphi_Y(x)\| &= \|\varphi_Y(x) - \varphi_Y(0) + \varphi_Y(0)\| \leq \|\varphi_Y(x) - \varphi_Y(0)\| + \|\varphi_Y(0)\| \\ &\leq \underbrace{\frac{1}{2} \|x - 0\|}_{\text{contraction}} + \underbrace{\|y\|}_{\leq 2r} < 2r \end{aligned}$$

So, for  $y \in K_r(0)$  and  $x \in \overline{K_{2r}(0)}$ ,  $\varphi_Y(x) \in K_{2r}(0)$ . By Banach Fixed Point Theorem, there exists a uniquely determined  $x$  such that

$$x = \varphi_Y(x) \iff y = F(x)$$

Let  $U = F^{-1}(K_r(0)) \cap K_{2r}(0)$  where  $F^{-1}(K_r(0))$  is open, because  $F$  is continuous. Hence,  $U$  is open and  $0 \in U$ . Therefore  $U$  is an open neighborhood of 0.

Show:  $F^{-1} : F(U) \rightarrow U$  is continuous <sup>3</sup>: Let  $x_1 = F^{-1}(y_1), x_2 = F^{-1}(y_2); y_1, y_2 \in F(U)$ .

$$\varphi_0(x) = x + 0 - F(x) = x - F(x)$$

$$\begin{aligned} x_2 - x_1 &= \underbrace{x_2 - F(x_2)}_{\varphi_0(x_2)} + \underbrace{F(x_2) - x_1 + F(x_1) - F(x_1)}_{-\varphi_0(x_1)} \\ &= \varphi_0(x_2) - \varphi_0(x_1) + F(x_2) - F(x_1) \\ \|x_2 - x_1\| &\leq \|\varphi_0(x_2) - \varphi_0(x_1)\| + \|F(x_2) - F(x_1)\| \\ &\leq \underbrace{\frac{1}{2} \|x_2 - x_1\|}_{\varphi_0 \text{ is a contraction}} + \|F(x_2) - F(x_1)\| \end{aligned}$$

$$\|F^{-1}(y_2) - F^{-1}(y_1)\| \leq 2 \|y_2 - y_1\|$$

---

<sup>3</sup>This proof was added on 2018/06/07 as we initially forgot this condition

hence  $F^{-1}$  is Lipschitz continuous on  $F(U)$ , so also continuous.

We show:  $DF(x)$  is invertible for all  $x \in U$ .

Let  $v \in \ker(DF(x)) \iff \|v\| = \|DF(x) \cdot v - v\|$  where  $DF(x) \cdot v = 0$ .

$$= \|(DF(x) - I) \cdot v\| \leq \underbrace{\|DF(x) - I\|}_{\leq \frac{1}{2}} \cdot \|v\| \leq \frac{1}{2} \|v\|$$

$$\implies \frac{1}{2} \|v\| \leq 0 \implies \|v\| \leq 0 \implies v = 0$$

Hence  $\ker(DF(x)) = \{0\}$ , so  $DF(x)$  is regular.

$$\forall x \in U = F^{-1}(K_r(0)) \cap K_{2r}(0) : y = F(x) \in K_r(0)$$

On the opposite,  $\forall y \in K_r(0)$  there exists some uniquely determined  $x \in K_{2r}(0) \cap F^{-1}(K_r(0))$  with  $y = F(x)$ . So  $F : U \rightarrow K_r(0)$  is bijective and continuously differentiable. Furthermore,  $DF(x)$  is regular  $\forall x \in U$ . By Lemma 6.9,  $F$  is a local diffeomorphism in  $x = 0$ . Thus,  $f$  is a local diffeomorphism in  $x_0$ .

So the central idea of this proof was to rewrite  $F$  such that we can apply Banach Fixed Point Theorem.  $\square$

## 6.13 Implicit functions

**Theorem 6.4** (Implicit function theorem). *Let  $U \subseteq \mathbb{R}^{n+m} = \mathbb{R}^n \times \mathbb{R}^m$ .  $f : U \rightarrow \mathbb{R}^m$  is continuously differentiable.*

Notation:  $\begin{pmatrix} x \\ y \end{pmatrix} \in U; x \in \mathbb{R}^n, y \in \mathbb{R}^m$ .  $f\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = f(x, y)$ .

For  $(x_0, y_0) \in U$ ,  $f(x_0, y_0) = 0$ . Let

$$Df(x_0, y_0) = \begin{bmatrix} \partial_{x_1} f_1 & \dots & \partial_{x_n} f_1 & \partial_{y_1} f_1 & \dots & \partial_{y_m} f_1 \\ \vdots & & \vdots & \vdots & & \vdots \\ \partial_{x_1} f_m & \dots & \partial_{x_n} f_m & \partial_{y_1} f_m & \dots & \partial_{y_m} f_m \end{bmatrix}$$

where the left half is given by  $D_x f(x_0, y_0) \in \mathbb{R}^{m \times n}$  and the right half is given by  $D_y f(x_0, y_0) \in \mathbb{R}^{m \times m}$ .

*Assumption: Let  $D_y f(x_0, y_0)$  be regular.*

*Then there exists some neighborhood  $D$  of  $x_0$  in  $\mathbb{R}^n$  and a function  $g : D \rightarrow \mathbb{R}^m$ .  $E$  is a neighborhood of  $y_0$  in  $\mathbb{R}^m$  such that  $D \times E \subseteq U$  and  $f(x, y) = 0 \iff y = g(x)$  for  $(x, y) \in D \times E$ . Hence,  $f(x, g(x)) = 0 \forall x \in D$ .*

↓ This lecture took place on 2018/06/07.

**Remark 6.18.** So it holds:  $g(x_0) = y_0$ .

*Proof.*

$$F : U \subset \mathbb{R}^{n+m} \rightarrow \mathbb{R}^{n+m}$$

$$F\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x & (\in \mathbb{R}^n) \\ f(x, y) & (\in \mathbb{R}^m) \end{bmatrix} \in \mathbb{R}^{n+m}$$

$$DF\left(\begin{bmatrix} x_0 \\ y_0 \end{bmatrix}\right) = \begin{bmatrix} I & 0 \\ D_x f(x_0, y_0) & D_y f(x_0, y_0) \end{bmatrix} \in \mathbb{R}^{(n+m) \times (n+m)}$$

is the Jacobi matrix of  $f$ .

$$\det\left(DF\left(\begin{bmatrix} x_0 \\ y_0 \end{bmatrix}\right)\right) = \underbrace{\det(I)}_{=1} \cdot \underbrace{\det(D_y f(x_0, y_0))}_{\neq 0} \neq 0$$

so  $DF\left(\begin{bmatrix} x_0 \\ y_0 \end{bmatrix}\right)$  is regular. By the local inversion theorem (Theorem 6.3),  $F$  is a local Diffeomorphism. Thus,  $\exists V : \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} \in V \subseteq D$  such that  $F : V \rightarrow F(V)$  is a diffeomorphism.  $V$  is open, hence  $\exists r, r' > 0$  such that

$$(x_0, y_0) \in K_r(x_0) \times K_{r'}(y_0) \subseteq V$$

here  $\mathbb{R}^{n+m}$  is identified as  $\mathbb{R}^n \times \mathbb{R}^m$ .

$$F^{-1} : F(V) \rightarrow V \quad F^{-1}\left(\begin{bmatrix} \xi \\ \eta \end{bmatrix}\right) = ?$$

$$F\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} \xi \\ \eta \end{bmatrix} \iff \begin{bmatrix} x \\ y \end{bmatrix} = F^{-1}\left(\begin{bmatrix} \xi \\ \eta \end{bmatrix}\right)$$

$$F\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{pmatrix} x \\ f(x, y) \end{pmatrix}$$

hence  $x = \xi$  and  $F^{-1}\left(\begin{bmatrix} \xi \\ \eta \end{bmatrix}\right) = \begin{bmatrix} \xi \\ G(\xi, \eta) \end{bmatrix}$ . Define  $g(x) := G(x, 0)$ .  $g : K_r(x_0) \rightarrow \mathbb{R}^m$ .

$$\begin{bmatrix} x \\ 0 \end{bmatrix} = F\left(F^{-1}\left(\begin{bmatrix} x \\ 0 \end{bmatrix}\right)\right) = F\left(\begin{bmatrix} x \\ G(x, 0) \end{bmatrix}\right) = F\left(\begin{bmatrix} x \\ g(x) \end{bmatrix}\right) = \begin{bmatrix} x \\ f(x, g(x)) \end{bmatrix}$$

$$\implies f(x, g(x)) = 0 \forall x \in K_r(x_0)$$

Uniqueness: Let  $(x, y) \in K_r(x_0) \times K_{r'}(y_0) \subset V$  and  $f(x, y) = 0$ . Hence

$$\begin{aligned} F\begin{pmatrix} x \\ y \end{pmatrix} &= \begin{bmatrix} x \\ f(x, y) \end{bmatrix} = \begin{bmatrix} x \\ 0 \end{bmatrix} \\ \Rightarrow [xy] &= F^{-1}(F\begin{pmatrix} x \\ y \end{pmatrix}) = F^{-1}\begin{pmatrix} x \\ 0 \end{pmatrix} = \begin{bmatrix} x \\ G(x, 0) \end{bmatrix} = \begin{bmatrix} x \\ g(x) \end{bmatrix} \\ &\Rightarrow y = g(x) \end{aligned}$$

□

**Example 6.5.**

$$f(x, y) = x^2y - x^3y^2 - 2$$

$$f(-1, 1) = 1 + 1 - 2 = 0$$

Hence  $(x_0, y_0) = (-1, 1)$  is root of  $F$ .

$$Df(x, y) = \begin{bmatrix} \underbrace{2xy - 3x^2y^2}_{D_x f} & \underbrace{x^2 - 2x^3y}_{D_y f} \end{bmatrix}$$

$$D_y f\left(\begin{bmatrix} -1 \\ 1 \end{bmatrix}\right) = 3 \neq 0$$

hence  $f(x, y)$  is in neighborhood of  $(-1, 1)$  resolvable by  $y$ .

$$x^2y - x^3y^2 - 2 = 0$$

$$x^3y^2 - x^2y + 2 = 0$$

$$y = \frac{x^2 \pm \sqrt{x^4 - 8x^3}}{2x^3} = \frac{x^2(1 \pm \sqrt{1 - \frac{8}{x}})}{2x^3} = \frac{1}{2x} \left( 1 \pm \sqrt{1 - \frac{8}{x}} \right)$$

It has to hold that  $g(-1) = 1$ , thus

$$\frac{1}{-2} \left( 1 + \sqrt{1 - \frac{8}{-1}} \right) = -\frac{1}{2}(1 + 3) = -2$$

This is apparently wrong, thus we consider the second result for  $y$ :

$$\frac{1}{-2} \left( 1 - \sqrt{1 - \frac{8}{-1}} \right) = -\frac{1}{2}(1 - 3) = -1$$

So  $g(x) = \frac{1}{2x} \cdot \left( 1 - \sqrt{1 - \frac{8}{x}} \right)$  is the desired function.  $D_x f(-1, 1) = -2 - 3 = -5$ , hence  $f(x, y) = -x^3y^2 + x^2y - 2$  is also uniquely resolvable by  $x$  in a neighborhood of  $(-1, 1)$ .



*Proof of Lemma 6.9.* Choose  $x_0 \in D$  and show that  $g^{-1}$  is differentiable in  $y_0 = g(x_0)$ . We use the same construction as in Proof 6.13 (Implicit function theorem proof). Without loss of generality:  $x_0 = 0; g(x_0) = y_0 = 0$  and  $Dg(0) = I$ . Let  $v \in \mathbb{R}^n$  be sufficiently small such that  $w = g^{-1}(v)$  is defined.

By differentiability of  $g$ ,

$$g(w) = \underbrace{Dg(0)}_I \cdot w + \underbrace{R(w)}_{o(\|w\|)} = w + R(w)$$

$$g^{-1}(v) = w = g(w) - R(w) = g(g^{-1}(v)) - R(g^{-1}(v)) = v + R^*(v)$$

with  $R^*(v) = -R(g^{-1}(v))$ . Show that  $R^*(v) = o(\|v\|)$ .

$$g^{-1}(v) = v + R^*(v) \tag{5}$$

By differentiability of  $g$ ,

$$\exists r > 0 : \|R(w)\| \leq \frac{1}{2} \|w\|$$

for all  $\|w\| \leq r$  (because  $\frac{\|R(w)\|}{\|w\|} \rightarrow 0$  for  $w \rightarrow 0$ ). Continuity of  $g^{-1}$  combined with  $g^{-1}(0) = 0$

$$\implies \forall \|v\| < \delta : \|g^{-1}(v)\| = \|g^{-1}(v) - g^{-1}(0)\| \leq r$$

$$\implies \|R^*(v)\| = \|R(g^{-1}(v))\| \leq \frac{1}{2} \|g^{-1}(v)\|$$

for all  $v$  with  $\|v\| \leq \delta$ .

By Equation (5),

$$\|g^{-1}(v)\| = \|v + R^*(v)\| \leq \|v\| + \|R^*(v)\| \leq \|v\| + \frac{1}{2} \|g^{-1}(v)\|$$

If  $\|v\| \leq \delta$

$$\|g^{-1}(v)\| \leq 2 \|v\| \quad \forall \|v\| \leq \delta$$

So,

$$\frac{\|R^*(v)\|}{\|v\|} \leq 2 \frac{\|R(g^{-1}(v))\|}{\|g^{-1}(v)\|} = 2 \frac{\|R(w)\|}{\|w\|}$$

$$v \rightarrow 0 \implies w = g^{-1}(v) \rightarrow 0$$

because  $g^{-1}$  is continuous, hence

$$v \rightarrow 0 : \lim_{v \rightarrow 0} \frac{\|R(g^{-1}(v))\|}{\|g^{-1}(v)\|} = \lim_{w \rightarrow 0} \frac{R(w)}{\|w\|} = 0$$

because  $R = o(\|w\|)$ . Thus,  $R^*(v) = o(\|v\|)$ , so  $g^{-1}$  is differentiable in  $y_0 = 0$ .

We know that  $g^{-1}$  is differentiable in every point  $y \in g(D)$ .

$$\underbrace{y}_{= \text{id}(y)} = g(g^{-1}(y)) \forall y \in E = g(D)$$

We derive both sides of the equation and apply the chain rule:

$$\begin{aligned} D \text{id}(y) &= I = Dg(g^{-1}(y)) \cdot Dg^{-1}(y) \\ \implies Dg^{-1}(y) &= [Dg(\underbrace{g^{-1}(y)}_{\text{continuous}})]^{-1} \\ &\quad \underbrace{\hspace{1.5cm}}_{\text{continuous}} \end{aligned}$$

Is the inverse also continuous (does the inverse depend continuously on the coefficients? Yes, you can see it by considering Cramer's Rule which provides a formula with a sum)? So the inverse is a continuous operation on  $GL(n)$ . Therefore  $Dg^{-1}(y)$  depends continuously on  $y$  and  $Dg^{-1}(y) = [Dg(x)]^{-1}$  with  $x = g^{-1}(y)$  and accordingly,  $y = g(x)$ .  $\square$

## 6.14 Partial derivatives and Hessian Matrix

**Remark 6.19** (Concept idea). Let  $D \subseteq \mathbb{R}^n$  be open. Let  $f : D \rightarrow \mathbb{R}$  be continuously differentiable. Hence  $\partial_{x_i} f(x) \in \mathbb{R}$  and  $\partial_{x_i} f : D \rightarrow \mathbb{R}$  is continuous. If  $\partial_{x_i} f$  is also (continuously) differentiable, then its partial derivatives can be determined. In this case, we define

$$\partial_{x_j, x_i} f := \partial_{x_j} [\partial_{x_i} f]$$

Continuation for further higher derivatives:

$$\partial_{x_{i_k}, x_{i_{k-1}}, \dots, x_{i_1}} f = \partial_{x_{i_k}} (\partial_{x_{i_{k-1}} \dots x_{i_1}} f)$$

The index  $k$  in  $\partial_{x_{i_k}, x_{i_{k-1}} \dots x_{i_1}}$  is the order of the partial derivative.

**Example 6.6.**

$$\begin{aligned}
f(x, y) &= x^2y - x^3y^2 - 2 \\
\partial_x f &= 2xy - 3x^2y^2 \\
\partial_y f &= x^2 - 2x^3y \\
\partial_{yx} f &= 2x - 6x^2y \\
\partial_{xy} f &= 2x - 6x^2y = \partial_{yx} \\
\partial_{xx} f &= 2y - 6xy^2 \\
\partial_{yy} f &= -2x^3 \\
\partial_{xxx} f &= -6y^2 \\
\partial_{yyy} f &= 0 \\
\partial_{xyx} f &= 2 - 12xy \\
\partial_{xxy} f &= 2 - 12xy = \partial_{xyx} \\
\partial_{yxy} f &= -6x^2 \\
\partial_{xyy} f &= -6x^2 = \partial_{yxy}
\end{aligned}$$

*It seems that the derivative is independent of the order of the variables.*

**Definition 6.11** (Hesse matrix). *Specifically for second derivatives:*

$$D^2 f(x) = \begin{bmatrix} \partial_{x_1, x_1} f & \partial_{x_2, x_1} f & \dots & \partial_{x_n, x_1} f \\ \partial_{x_1, x_2} f & \partial_{x_2, x_2} f & \dots & \partial_{x_n, x_2} f \\ \vdots & \vdots & \ddots & \vdots \\ \partial_{x_1, x_n} f & \partial_{x_2, x_n} f & \dots & \partial_{x_n, x_n} f \end{bmatrix}$$

*is called Hessian matrix. Named after Otto Hesse (1811–1874).*

**Definition 6.12.** *Let  $f : D \rightarrow \mathbb{R}$  and let all partial derivatives of  $f$  up to order  $k \in \mathbb{N}$  exist and be continuous functions in  $x \in D$ . Then  $f$  is called  $k$ -times continuously differentiable and*

$$C^k(D) := \{f : D \rightarrow \mathbb{R} : f \text{ is } k\text{-times continuously differentiable}\}$$

$C^k(D)$  *is a vector space.*

## 6.15 Schwarz' Theorem

**Remark 6.20.** *Hermann Amadeus Schwarz (1843–1921)*

**Theorem 6.5** (Symmetry of second derivatives, Schwarz' theorem, Clairaut's theorem, Young's theorem). Let  $f \in C^2(D)$ . Let  $D \subseteq \mathbb{R}^n$  be open. Then,

$$\partial_{x_i x_j} f = \partial_{x_j x_i} f \text{ on } D$$

↓ This lecture took place on 2018/06/12.

*Proof.* Let  $x_0 \in D$  and  $\overline{K_r(x_0)} \subseteq D$ . Let  $h, k \in \mathbb{R}$  with  $|h| \leq \frac{r}{2}$  and  $|k| \leq \frac{r}{2}$ . So  $x_0 + he_i \in \overline{K_{\frac{r}{2}}(x_0)}$  and  $x_0 + ke_j \in \overline{K_{\frac{r}{2}}(x_0)}$  and also  $x_0 + he_i + ke_j \in \overline{K_r(x_0)}$ , because

$$\|x_0 + he_i + ke_j - x_0\| \leq |h| \|e_i\| + |k| \|e_j\| = |h| + |k| \leq \frac{r}{2} + \frac{r}{2} = r$$

$$F : \left[-\frac{r}{2}, \frac{r}{2}\right] \times \left[-\frac{r}{2}, \frac{r}{2}\right] \rightarrow \mathbb{R}$$

$$F(h, k) := f(x_0 + he_i + ke_j) - f(x_0 + he_i) - f(x_0 + ke_j) + f(x_0)$$

$$\varphi : \left[-\frac{r}{2}, \frac{r}{2}\right] \rightarrow \mathbb{R} \quad k \text{ fixed}$$

$$\varphi(\lambda) := f(x_0 + \lambda e_i + ke_j) - f(x_0 + \lambda e_i)$$

$$\implies \varphi(0) = f(x_0 + ke_j) - f(x_0)$$

$$\varphi(h) = f(x_0 + he_i + ke_j) - f(x_0 + he_i)$$

Therefore,  $F(h, k) = \varphi(h) - \varphi(0)$ .  $\varphi$  is continuously differentiable (because  $f \in C^2$ ). By the Mean Value Theorem of Differential Calculus:

$$\exists \lambda \in (-h, h) : F(h, k) = \varphi(h) - \varphi(0) = \int_0^h \varphi'(x) \cdot 1 \, dx = \varphi'(\lambda) \cdot (h - 0) = \varphi'(\lambda) \cdot h$$

and accordingly  $|\lambda| < |h|$ .

$$\varphi'(\lambda) \cdot h = h \cdot [Df(x_0 + \lambda e_i + ke_j) \cdot e_i - Df(x_0 + \lambda e_i) \cdot e_i]$$

$$Df \cdot e_i = \partial_{x_i} f$$

and also

$$\varphi'(\lambda) \cdot h = h \cdot [\partial_{x_i} f(x_0 + \lambda e_i + k \cdot e_j) - \partial_{x_i} f(x_0 + \lambda e_i)]$$

$$\partial_{x_i} f \text{ is continuously differentiable}$$

$$\psi(\mu) = \partial_{x_i} f(x_0 + \lambda e_i + \mu e_j) \text{ is continuously differentiable}$$

So we apply the Mean Value Theorem:

$$F(h, k) = \varphi'(\lambda) \cdot h = h[\psi(k) - \psi(0)]$$

$$h \cdot k \cdot \psi'(\mu) \text{ with appropriate } \mu \in [-|k|, |k|]$$

$$\begin{aligned}\psi'(\mu) &= \partial_{x_j x_i} f(x_0 + \lambda e_i + \mu e_j) \\ \implies \exists(\lambda, \mu) &\in [-|h|, |h|] \times [-|k|, |k|] \text{ such that} \\ \textcolor{blue}{F(h, k)} &= h \cdot k \cdot \partial_{x_j x_i} f(x_0 + \lambda e_i + \mu e_j)\end{aligned}$$

Analogously,

$$\tilde{\varphi}(\eta) = f(x_0 + h e_i + \eta e_j) - f(x_0 + \eta e_j)$$

$$\text{like above: } F(h, k) = \tilde{\varphi}(k) - \tilde{\varphi}(0) = k \tilde{\varphi}'(\eta)$$

with appropriate  $\eta \in [-|k|, |k|]$ .

$$k \tilde{\varphi}'(\eta) = k \left[ \partial_{x_j} f(x_0 + h e_i + \eta e_j) - \partial_{x_j} f(x_0 + \eta e_j) \right]$$

$$\tilde{\psi}(\xi) = \partial_{x_i} f(x_0 + \xi e_i + \eta e_j) \text{ and Mean Value Theorem}$$

$$\implies \textcolor{red}{F(h, k)} = k \cdot h \cdot \partial_{x_i x_j} f(x_0 + \xi e_i + \eta e_j)$$

with  $(\xi, \eta) \in [-|h|, |h|] \times [-|k|, |k|]$ .

Because  $\textcolor{blue}{F(h, k)} = \textcolor{red}{F(h, k)}$ ,

$$\begin{aligned}hk \cdot \partial_{x_j x_i} f(x_0 + \lambda e_i + \mu e_j) &= kh \cdot \partial_{x_i x_j} f(x_0 + \xi e_i + \mu e_j) \\ \implies \partial_{x_i x_j} f(x_0 + \lambda e_i + \mu e_j) &= \partial_{x_i x_j} f(x_0 + \xi e_i + \eta e_j)\end{aligned}$$

with  $|\lambda| < |h|, |\xi| < |h|$  and  $|\mu| < |k|, |\eta| < |k|$ .  $h \rightarrow 0, k \rightarrow 0 \implies \lambda \rightarrow 0, \mu \rightarrow 0, \xi \rightarrow 0, \eta \rightarrow 0$ .

$$f \in C^2 \implies \partial_{x_i x_j} f \text{ and } \partial_{x_j x_i} f \text{ are continuous}$$

$$\implies \partial_{x_j x_i} f(x_0) = \lim_{\lambda, \mu \rightarrow 0} \partial_{x_j x_i} f(x_0 + \lambda e_i + \mu e_j) = \lim_{\xi, \eta \rightarrow 0} \partial_{x_i x_j} f(x_0 + \xi e_i + \eta e_j) = \partial_{x_i x_j} f(x_0)$$

Counterexample:

$$f(x_1, x_2) = \begin{cases} \frac{x_1 x_2 (x_1^2 - x_2^2)}{x_1^2 + x_2^2} & \text{for } (x_1, x_2) \neq (0, 0) \\ 0 & \text{for } (x_1, x_2) = (0, 0) \end{cases}$$

here it holds that  $\partial_{x_1 x_2} f(0, 0) \neq \partial_{x_2 x_1} f(0, 0)$ . □

**Theorem 6.6** (Generalization of Schwarz' Theorem). *Let  $f \in C^k(D)$ ,  $D \subseteq \mathbb{R}^n$  open and  $x_0 \in D$ . Let  $(i_1, i_2, \dots, i_k) \in \underbrace{\{1, 2, \dots, n\}^k}_{M_n} = M_n^k$ . Furthermore let  $(i'_1, i'_2, \dots, i'_k)$  be*

*a rearrangement of  $(i_1, i_2, \dots, i_k)$ . Then*

$$\partial_{x_{i_1} x_{i_2} \dots x_{i_k}} f(x_0) = \partial_{x_{i'_1} x_{i'_2} \dots x_{i'_k}} f(x_0)$$

## 6.16 Multiindex notation

*Proof.* Proof by complete induction.

**Definition 6.13** (Multiindex notation). Let  $I \in \mathbb{N}^n$ ,  $I = (\alpha_1, \alpha_2, \dots, \alpha_n)$ .  $\alpha_i \geq 0$ . We call  $I$  a multiindex  $k = |I| = \sum_{i=1}^n \alpha_i$  is the order of  $I$ .

$$\underbrace{\partial_{x_i} \partial_{x_i} \dots \partial_{x_i}}_{\alpha \text{ times}} =: \partial_{x_i}^\alpha$$

is called  $\alpha$ -times partial derivative to variable  $x_i$ . By convention,

$$\partial_{x_i}^0 = f$$

We let

$$\partial_I f = \partial_{x_1}^{\alpha_1} \partial_{x_2}^{\alpha_2} \dots \partial_{x_n}^{\alpha_n} f$$

Furthermore, we use

$$\partial_{i_1, i_2, \dots, i_k} \text{ for } \partial_{x_{i_1} x_{i_2} \dots x_{i_k}}$$

hence

$$\underbrace{\partial_{1312}}_{\text{no parentheses}} f = \partial_{x_1 x_3 x_1 x_2} f = \underbrace{\partial_{(2,1,1)}}_{\substack{\text{multiindex} \\ \text{with parentheses}}} f$$

Furthermore we let  $I! = \alpha_1! \alpha_2! \dots \alpha_n! = \prod_{i=1}^n \alpha_i!$  and for  $v \in \mathbb{R}^n : v = [v_1, v_2, \dots, v_n]^T$  we write

$$v^I = \alpha_1^{\alpha_1} v_1^{\alpha_1} \dots v_n^{\alpha_n} = \prod_{i=1}^n v_i^{\alpha_i}$$

The proof is left as an exercise to the reader.  $\square$

**Theorem 6.7** (Multidimensional Taylor's theorem). Let  $f : D \rightarrow \mathbb{R}$ .  $D$  is open. Let  $x_0 \in D$  and  $f \in C^{k+1}(D)$ . Let  $r > 0$  such that  $K_r(x_0) \subseteq D$  and  $v \in \mathbb{R}^n$  with  $\|v\| \leq r$  (hence,  $x_0 + v \in \overline{K_r(x_0)}$ ) and therefore also the connecting line  $[x_0, x_0 + v] = \{x_0 + tv : t \in [0, 1]\} \subseteq \overline{K_r(x_0)} \subseteq D$ . Then there exists  $\vartheta \in (0, 1)$  such that

$$\begin{aligned} f(x_0 + v) &= f(x_0) + \underbrace{\sum_{j=1}^k \frac{1}{j!} \sum_{i_1, i_2, \dots, i_j=1}^n \partial_{i_1 i_2 \dots i_j} f(x_0) \cdot v_{i_1} \cdot v_{i_2} \dots v_{i_j}}_{T_f^k(x_0 + v; x_0)} \\ &\quad + \underbrace{\frac{1}{(k+1)!} \sum_{i_1 i_2 \dots i_{k+1}=1}^n \partial_{i_1 i_2 \dots i_{k+1}} f(x_0 + \vartheta v) v_{i_1} v_{i_2} \dots v_{i_{k+1}}}_{R_f^{k+1}(x_0 + v; x_0)} \end{aligned}$$

where  $T_f^k(x_0 + v; x_0)$  is the Taylor polynomial of  $k$ -th order and  $R_f^{k+1}(x_0 + v; x_0)$  is the remainder term. But in this notation some terms occur multiple times. For example,  $\partial_{1121} = \partial_{2111}$ . Alternatively,

$$f(x_0 + v) = f(x_0) + \sum_{j=1}^k \sum_{|I|=j} \frac{1}{I!} \partial_I f(x_0) \cdot v^I + \sum_{|I|=k+1} \frac{1}{I!} \partial_I f(x_0 + \vartheta v) \cdot v^I$$

*Proof.* Consider  $\varphi : [0, 1] \rightarrow \mathbb{R}$ .

$$\varphi(t) = f(x_0 + tv) = f \circ l_{x_0, v}(t)$$

$$f \in C^{(k+1)}(D) \implies \varphi \in C^{(k+1)}([0, 1])$$

**Claim.**

$$\varphi^{(j)}(t) = \sum_{i_1, i_2, \dots, i_j=1}^n \partial_{i_1 i_2 \dots i_j} f(x_0 + tv)$$

*Proof.* Proof by induction over  $j$ .

**Induction base  $j = 1$**

$$\begin{aligned} \varphi(t) &= f \circ l_{x_0, v}(t) \\ \varphi'(t) &\underbrace{=}_{\substack{\text{chain} \\ \text{rule}}} \underbrace{Df(x_0 + tv)}_{\substack{\text{Jacobi matrix} \\ \text{row vector}}} \cdot v \\ &= [\partial_1 f(x_0 + tv), \partial_2 f(x_0 + tv), \dots, \partial_n f(x_0 + tv)] \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix} \\ &= \sum_{i_1=1}^n \partial_{i_1} f(x_0 + tv) \cdot v_{i_1} \end{aligned}$$

**Induction step  $j - 1 \mapsto j$**

$$\begin{aligned} \varphi^{(j)}(t) &= (\varphi^{(j-1)})'(t) \\ &\underbrace{=}_{\substack{\text{induction} \\ \text{hypothesis}}} \frac{d}{dt} \left[ \sum_{i_1, i_2, \dots, i_{j-1}=1}^n \underbrace{\partial_{i_1, \dots, i_{j-1}} f(x_0 + tv)}_{\partial_{i_1, \dots, i_{j-1}} f \circ l_{x_0, v}(t)} v_{i_1} \cdots v_{i_{j-1}} \right] \\ &= \sum_{i_1, \dots, i_j=1}^n \partial_{i_1, \dots, i_j} f(x_0 + tv) v_{i_1} \cdots v_{i_j} \end{aligned}$$

□

Taylor's Theorem for  $\varphi(1)$  in the 1-dimensional case:

$$\varphi(1) = \varphi(0) + \sum_{j=1}^n \frac{1}{j!} \varphi^{(j)}(0) \cdot 1^j + \frac{1}{(k+1)!} \varphi^{(k+1)}(\vartheta) \cdot 1$$

with  $\vartheta \in (0, 1)$ . Insertion:

$$\begin{aligned} f(x_0 + v) &= f(x_0) + \sum_{j=1}^n \frac{1}{j!} \sum_{i_1, \dots, i_j=1}^n \partial_{i_1, \dots, i_j} f(x_0) v_{i_1} \dots v_{i_j} \\ &+ \frac{1}{(k+1)!} \sum_{i_1, \dots, i_{k+1}=1}^n \partial_{i_1, \dots, i_{k+1}} f(x_0 + \vartheta v) \cdot v_{i_1} v_{i_2} \dots v_{i_{k+1}} \end{aligned}$$

Alternative notation with multiindices:

$$\partial_{i_1 i_2 \dots i_j} f(x_0) = \partial_{i'_1 i'_2 \dots i'_j} f(x_0)$$

if  $i_1 i_2 \dots i_j$  and  $i'_1 i'_2 \dots i'_j$  only distinguish by the order of indices. How many ways are there to put indices in order?

Let  $I = (\alpha_1, \alpha_2, \dots, \alpha_n)$  be a multiindex

$$\partial_I = \partial_{(\alpha_1, \alpha_2, \dots, \alpha_n)} = \partial_{\underbrace{11 \dots 1}_{\alpha_1 \text{-times}} \underbrace{2 \dots 2}_{\alpha_2 \text{-times}} \underbrace{33 \dots 3}_{\alpha_3 \text{-times}} \dots \underbrace{nn \dots n}_{\alpha_n \text{-times}}}$$

How many ways are there to distribute

$$\underbrace{\underbrace{11 \dots 1}_{\alpha_1} \underbrace{22 \dots 2}_{\alpha_2} \underbrace{33 \dots 3}_{\alpha_3} \dots \underbrace{nn \dots n}_{\alpha_n}}_j$$

is different order? There are  $\binom{j}{\alpha_1}$  different ways to distribute one. There are  $\binom{j-\alpha_1}{\alpha_2}$  different ways to distribute two. There are  $\binom{j-\alpha_1-\alpha_2}{\alpha_3}$  different ways to distribute three. And so on. There are  $\binom{j-\alpha_1-\alpha_2-\dots-\alpha_{n-1}}{\alpha_n}$  different ways to distribute  $n$ .

$$\binom{j}{\alpha_1} \binom{j-\alpha_1}{\alpha_2} \binom{j-(\alpha_1+\alpha_2)}{\alpha_3} \dots \binom{j-(\alpha_1+\dots+\alpha_{n-1})}{\alpha_n} \quad (6)$$

are different arrangements of the index list.

$$\underbrace{11 \dots 1}_{\alpha_1} \underbrace{2 \dots 2}_{\alpha_2} \dots \underbrace{nn \dots n}_{\alpha_n}$$



$$\begin{aligned}
(6) &= \frac{j!}{\alpha_1!(j-\alpha_1)!} \frac{(j-\alpha_1)!}{\alpha_2!(j-(\alpha_1+\alpha_2))!} \frac{(j-(\alpha_1+\alpha_2))!}{\alpha_3!(j-(\alpha_1+\alpha_2+\alpha_3))!} \\
&\quad \cdots \frac{(j-(\alpha_1+\dots, \alpha_{n-1}))!}{\alpha_n!(j-(\alpha_1+\dots+\alpha_n))!} \\
&\quad \underbrace{\hspace{10em}}_{=j} \\
&\quad \underbrace{\hspace{10em}}_{=0!=1} \\
&= \frac{j!}{\alpha_1!\alpha_2!\dots\alpha_n!}
\end{aligned}$$

We merge equal partial derivative

$$\begin{aligned}
f(x_0 + v) &= f(x_0) + \sum_{j=1}^k \frac{1}{j!} \sum_{|l|=j} \frac{j!}{l!} \cdot \partial_l f(x_0) \cdot v^l \\
&\quad + \sum_{|l|=k+1} \frac{1}{(k+1)!} \cdot \frac{(k+1)!}{l!} \cdot \partial_l f(x_0 + \vartheta v) v^l
\end{aligned}$$

□

**Example 6.7.**

$f : \mathbb{R}^2 \rightarrow \mathbb{R}$  arbitrarily often differentiable

$$\begin{aligned}
f(x_0 + v) &= f(x_0) + \partial_1 f(x_0) \cdot v_1 + \partial_2 f(x_0) v_2 \\
&\quad + \frac{1}{2!} \partial_{11} f(x_0) v_1 v_1 + \frac{1}{2!} \partial_{12} f(x_0) v_1 v_2 \\
&\quad + \frac{1}{2!} \partial_{21} f(x_0) v_2 v_1 + \frac{1}{2!} \partial_{22} f(x_0) v_2 v_2 \\
&\quad + \frac{1}{3!} \partial_{111} f(x_0) v_1 v_1 v_1 + \frac{1}{3!} \partial_{211} f(x_0) v_2 v_1 v_1 + \\
&\quad + \frac{1}{3!} \partial_{121} f(x_0) v_1 v_2 v_1 + \frac{1}{3!} \partial_{112} f(x_0) v_1 v_1 v_2 + \\
&\quad + \frac{1}{3!} \partial_{122} f(x_0) v_1 v_2 v_2 + \frac{1}{3!} \partial_{212} f(x_0) v_2 v_1 v_2 + \\
&\quad + \frac{1}{3!} \partial_{221} f(x_0) v_2 v_2 v_1 + \frac{1}{3!} \partial_{222} f(x_0) v_2 v_2 v_2 + R \\
&= f(x_0) + \partial_{(1,0)} f(x_0) \cdot v_1^1 v_2^0 + \partial_{(0,1)} f(x_0) v_1^0 v_2^1 \\
&\quad + \frac{1}{2!0!} \partial_{(2,0)} f(x_0) v_1^2 v_2^0 + \frac{1}{1!1!} \partial_{(1,1)} f(x_0) v_1^1 v_2^1 \\
&\quad + \frac{1}{0!2!} \partial_{(0,2)} f(x_0) v_1^0 v_2^2 + \frac{1}{3!0!} \partial_{(3,0)} f(x_0) v_1^3 v_2^0 \\
&\quad + \frac{1}{2!1!} \partial_{(2,1)} f(x_0) v_1^2 v_2^1 + \frac{1}{1!2!} \partial_{(1,2)} f(x_0) v_1^1 v_2^2 \\
&\quad + \frac{1}{0!3!} \partial_{(0,3)} f(x_0) v_1^0 v_2^3 + R
\end{aligned}$$

↓ This lecture took place on 2018/06/14.

## 6.17 Qualitative Taylor Theorem

**Theorem 6.8** (Qualitative Taylor Theorem). *Let  $f \in C^k(D)$ ,  $D \subseteq \mathbb{R}^n$  open,  $x_0 \in D$ . Let  $r > 0$  such that  $K_r(x_0) \subseteq D$  and  $\|v\| < r$ , hence  $x_0 + v \in K_r(x_0) \subseteq D$ . Then*

$$f(x) = \sum_{j=0}^k \sum_{|I|=j} \frac{1}{I!} \partial_I f(x_0) v^I + o(\|v\|)^k$$

*Pay attention to  $C^k(D)$ ,  $j = 0$  and  $o(\|v\|)^k$ .*

*Proof.* Use the 1-norm  $\|v\|_1 = \sum_{i=1}^n |v_i|$  for the proof. By the equivalence of norms in  $\mathbb{R}^n$ , it holds for every norm in  $\mathbb{R}^n$ .

$$\|v\|_1^k = \left( \sum_{k=1}^n |v_i| \right)^k \underbrace{=}_{\text{Theorem 6.7}} \sum_{|I|=k} \frac{k!}{I!} |v^I|$$

for every multiindex  $I$  of order  $k$ ,  $\frac{1}{I!} |v^I| \leq \frac{1}{k!} \|v\|_1^k$ . By Theorem 6.7,

$$\begin{aligned}
f(x) &= \sum_{j=0}^{k-1} \sum_{|I|=j} \frac{1}{I!} \partial_I f(x_0) v^I + \sum_{|I|=k} \frac{1}{I!} \partial_I f(x_0) v^I \\
&\quad + \sum_{|I|=k} \frac{1}{I!} \partial_I f(x_0 + \vartheta v) \cdot V^I - \sum_{|I|=k} \frac{1}{I!} \partial_I f(x_0) v^I \\
&= \sum_{j=0}^k \sum_{|I|=j} \frac{1}{I!} \partial_I f(x_0) v^I \\
&\quad \underbrace{\sum_{|I|=k} \frac{1}{I!} (\partial_I f(x_0 + \vartheta v) - \partial_I f(x_0)) \cdot v^I}_{\text{to show: } = o(\|v\|_1^k)} \\
&\quad \left\| \sum_{|I|=k} \left| \frac{1}{I!} v^I \cdot (\partial_I f(x_0 + \vartheta v) - \partial_I f(x_0)) \right| \right\| \\
&\leq \sum_{|I|=k} \underbrace{\left| \frac{1}{I!} v^I \right|}_{\leq \frac{1}{k!} \|v\|_1^k} \cdot \|\partial_I f(x_0 + \vartheta v) - \partial_I f(x_0)\| \\
&\leq \underbrace{\frac{1}{k!} \sum_{|I|=k} |\partial_I f(x_0 + \vartheta v) - \partial_I f(x_0)| \|v\|_1^k}_{\xrightarrow{v \rightarrow 0} 0}
\end{aligned}$$

is immediate, because  $f \in C^k(D)$ , hence  $\partial_I f(x_0 + \vartheta v) \rightarrow \partial_I f(x_0)$  for  $v \rightarrow 0$ .  $\square$

Reminder:

$$D_2 f(x_0) = \begin{bmatrix} \partial_{11} f(x_0) & \partial_{12} f(x_0) & \dots & \partial_{1n} f(x_0) \\ \partial_{21} f(x_0) & \partial_{22} f(x_0) & \dots & \vdots \\ \vdots & \ddots & \ddots & \vdots \\ \partial_{n1} f(x_0) & \partial_{n2} f(x_0) & \dots & \partial_{nn} f(x_0) \end{bmatrix}$$

$f \in C^2(D) \implies D_2 f(x_0)$  is symmetrical. Let  $v \in \mathbb{R}^n$ , then

$$v^t D_2 f(x_0) v = \sum_{i=1}^n v_i \sum_{j=1}^n \partial_{ij} f(x_0) \cdot v_j = \sum_{i,j=1}^n \partial_{ij} f(x_0) v_i v_j$$

hence, the Taylor expansion of  $f$  up to the second degree is

$$f(x) = f(x_0) + \sum_{i=1}^n \partial_i f(x_0) v_i + \frac{1}{2} \sum_{i,j=1}^n \partial_{ij} f(x_0) v_i v_j + o(\|v\|^2)$$

$$= f(x_0) + Df(x_0) \cdot v + \frac{1}{2} v^t D_2 f(x_0) v + o(\|v\|^2)$$

## 6.18 Sufficient optimality criteria

**Theorem 6.9** (Sufficient optimality criteria). *Let  $D \subseteq \mathbb{R}^n$  be open. Let  $f \in C^2(D)$ ,  $x_0 \in D$  such that  $Df(x_0) = 0$  ( $\nabla f(x_0) = 0$ ) (hence  $x_0$  is a critical point of  $f$ ).*

1. *Let  $D_2 f(x_0)$  be positive definite. Then  $f$  has a strict local minimum in  $x_0$ .*
2. *Let  $D_2 f(x_0)$  be negative definite. Then  $f$  has a strict local maximum in  $x_0$ .*
3. *Let  $D_2 f(x_0)$  be indefinite. Then  $f$  in  $x_0$  has no local extremum.*

**Remark 6.21** (Reminder). *Let  $M \in \mathbb{R}^{n \times n}$  and  $M$  be symmetrical.  $M$  is called positive definite if  $\forall \lambda \in \mathbb{R}$  eigenvalue of  $M$ :  $\lambda > 0 \iff \forall v \in \mathbb{R}^n \setminus \{0\} : v^t M v > 0$*

*$M$  is called indefinite if  $\exists \lambda$  eigenvalue of  $A : \lambda > 0 \wedge$   
 $\exists \mu$  eigenvalue of  $A : \mu < 0$*

*For corresponding eigenvectors  $v$  (and accordingly  $w$ ),*

$$\begin{aligned} v^t M v &= v^t \lambda v = \lambda \|v\|^2 > 0 \\ w^t M w &= w^t \mu w = \underbrace{\mu}_{<0} \|w\|^2 < 0 \end{aligned}$$

*Proof.* 1. Taylor expansion of 2nd degree with  $Df(x_0) = 0$ :  $f(x) - f(x_0) = \frac{1}{2}(x - x_0)^t D_2 f(x_0)(x - x_0) + g(x - x_0) \cdot \|x - x_0\|^2$  where  $g(x - x_0) \cdot \|x - x_0\|^2$  represents  $o(\|x - x_0\|^2)$  with  $\lim_{v \rightarrow 0} g(v) = 0$ .

$$S^{k-1} = \{v \in \mathbb{R}^n : \|v\| = 1\}$$

is the  $n - 1$  dimensional unit sphere in  $\mathbb{R}^n$ .  $S^{n-1}$  is bounded, closed and therefore compact.  $\forall v \in S^{n-1} : v \neq 0 \implies v^t D_2 f(x_0) v > 0$  by positive definiteness. Let  $m = \min \left\{ \frac{1}{2} v^t D_2 f(x_0) v : v \in S^{n-1} \right\}$  where  $\frac{1}{2} v^t D_2 f(x_0)$  is continuous, has therefore a minimum and this minimum is necessarily positive. So  $m > 0$ . Choose  $\delta > 0$  such that  $\|v\| < \delta \implies |g(v)| < \frac{m}{2}$  (feasible because  $\lim_{v \rightarrow 0} g(v) = 0$ ). For  $x \neq x_0$ ,

$$\begin{aligned} f(x) - f(x_0) &= \underbrace{\left( \frac{1}{2} \frac{(x - x_0)^t}{\|x - x_0\|} D_2 f(x_0) \frac{(x - x_0)}{\|x - x_0\|} \right)}_{\geq m} \cdot \|x - x_0\|^2 + g(x - x_0) \cdot \|x - x_0\|^2 \\ &\geq m \cdot \|x - x_0\|^2 - \underbrace{|g(x - x_0)|}_{\leq \frac{m}{2}} \|x - x_0\|^2 \\ &\geq \frac{m}{2} \|x - x_0\|^2 > 0 \end{aligned}$$

For all  $x \in D$  with  $\|x - x_0\| < \delta$ ,  $x_0$  is a strict local minimum of  $f$ .

2. Follows analogously.

3. Let  $D_2f(x_0)$  be indefinite. Let  $\lambda > 0$  be the eigenvalue of  $D_2f(x_0)$  with eigenvector  $v \neq 0$  and  $\mu < 0$  is negative eigenvalue of  $D_2f(x_0)$  with eigenvector  $w \neq 0$ .

$$\varphi(t) = f(x_0 + tv) \quad t \in (-r, r)$$

$$\varphi'(t) = Df(x_0 + tv) \cdot v = \sum_{i=1}^n \partial_i f(x_0 + tv) \cdot v_i$$

$$\varphi'(0) = 0$$

$$\begin{aligned} \varphi''(t) &= \frac{d}{dt} \left( \sum_{i=1}^n \partial_i f(x_0 + tv) \cdot v_i \right) \\ &= \sum_{i=1}^n \sum_{j=1}^n \partial_{ij} f(x_0 + tv) \cdot v_i v_j \\ &= v^t D_2f(x_0 + tv) \cdot v \\ &\implies \varphi''(0) = v^t D_2f(x_0) v = \lambda \|v\|^2 > 0 \end{aligned}$$

By some previous result establishing the necessary optimality criteria,  $\varphi''(0) > 0 \implies t = 0$  is a strict local minimum of  $\varphi$ . Analogously  $\psi(t) = f(x_0 + tw)$

$$\left. \begin{array}{l} \psi'(0) = 0 \\ \psi''(0) = \mu \|w\|^2 < 0 \end{array} \right\} \implies t = 0 \text{ is strict local maximum for } \psi$$

Let  $\varepsilon > 0$  be sufficiently small such that  $\varphi(t) > \varphi(0) \forall |t| < \varepsilon, t \neq 0$ . and  $\psi(t) < \psi(0) \forall |t| < \varepsilon$  with  $t \neq 0$ . Then  $f(x_0 + tv) = \varphi(t) > \varphi(0) = f(x_0)$  and  $x_0 + tv = x$

$$\text{and } \|x - x_0\| = \|tv\| = |t| \cdot \|v\| = |t| \cdot 1 < \varepsilon$$

$$\text{and } \underbrace{f(x_0 + tw)}_y = \psi(t) < \psi(0) = f(x_0)$$

$$\|y - x_0\| = \|tw\| = |t| \|w\| = |t| \cdot 1 < \varepsilon$$

Thus,  $x_0$  is not a local extreme value. Compare with Figure 41.

□

**Remark 6.22** (Sufficient optimality conditions for the scalar case).

$$f''(x_0) > 0 \vee f''(x_0) < 0$$




Figure 41:  $v$  and  $w$

$$f''(x_0) = 0 \implies \text{no conclusion}$$

$f''(x_0) = 0$  gives a saddle point.

In the 2-dimensional case, we have

$$D_2f(x_0) \text{ symmetrical, positive definite}$$

$$Df(x_0) = 0$$

Compare with Figure 42 (a).

$$D_2f(x_0) \text{ indefinite}$$

$$Df(x_0) = 0$$

$\rightarrow$  no extreme value. Compare with Figure 42 (b).

## 7 Differential geometry of curves

### 7.1 Curves, regular curves and tangential vectors

We consider:




Figure 42: (a) local minimum and (b) saddle point

1. Maps  $\gamma : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^n$  (compare with Figure 43)
2. geometric properties of  $\Gamma = \gamma(I)$ .

**Definition 7.1.** Let  $I \subseteq \mathbb{R}$  be an interval. Let  $\gamma : I \rightarrow \mathbb{R}^n$  be a continuous (or differentiable or smooth) function. Then  $\gamma$  is a continuous (or differentiable or smooth) function curve parameterization. Colloquially, we are used to call  $\gamma$  a parameterized curve (in German:  $\gamma$  is eine parametrisierte Kurve).

$\gamma(t) \in \mathbb{R}^n$  is a curve point.  $t$  is the curve parameter.

$$\Gamma = \gamma(I) = \{\gamma(t) : t \in I\} \subseteq \mathbb{R}^n$$

is called trace of the curve parameterization.

$$\gamma(t) = [\gamma_1(t) \quad \gamma_2(t) \quad \cdots \quad \gamma_n(t)]^T$$

If  $\gamma$  is differentiable, then

$$D\gamma(t) = [\partial_t \gamma_1(t) \quad \partial_t \gamma_2(t) \quad \cdots \quad \partial_t \gamma_m(t)]^T = [\gamma'_1(t) \quad \gamma'_2(t) \quad \cdots \quad \gamma'_n(t)]^T =: \gamma'(t)$$

$\gamma'(t)$  is the derivative vector. Often we also use the notation  $\dot{\gamma}(t)$ .

Let  $\gamma : I \rightarrow \mathbb{R}^n$  be differentiable.  $\gamma$  is called regular curve parameterization if  $\gamma'(t) \neq 0 \forall t \in I$ .

**Definition 7.2.** Let  $\gamma : I \rightarrow \mathbb{R}^n$  be a regular curve parameterization. Let  $T_\gamma(t) := \frac{\gamma'(t)}{\|\gamma'(t)\|_2}$ .  $T_\gamma(t)$  is called unit tangent vector of  $\gamma$  in point  $x = \gamma(t)$ .

$$\tau(\gamma, x) := \{x + \tau \cdot T_\gamma(t) : \tau \in \mathbb{R}\}$$




Figure 43: Maps in differential geometry

is a line in  $\mathbb{R}^n$  with directional vector  $T_\gamma(t)$  through  $x = \gamma(t)$ .  $\tau(\gamma, x)$  is called tangent on  $\Gamma$  in point  $x$ . Compare with Figure 44.

Motivation:

$$\begin{aligned}\gamma(t + \tau) &= \gamma(t) + \gamma'(t) \cdot \tau + o(\tau) \\ &= x + \underbrace{\frac{\tau}{\|\gamma'(t)\|} \cdot T_\gamma(t)}_{\in \tau(\gamma, x)} + o(\tau)\end{aligned}$$

A tangent is the best linear approximation of the curve.

↓ This lecture took place on 2018/06/19.

## 7.2 Reparameterization of parameterized curves

**Definition 7.3.** Let  $\gamma : I \rightarrow \mathbb{R}^n$  be a continuous or differentiable or smooth curve parameterization. Furthermore let  $\sigma : J \rightarrow I$  with  $J \subseteq \mathbb{R}$  as interval.

Let  $\sigma$  be a

- homeomorphism (continuous curve)
- diffeomorphism (differentiable curve)
- $C^\infty$ -diffeomorphism (smooth curve)






Figure 44:  $\tau(\gamma, x)$  is called *tangent* on  $\Gamma$  in point  $x$

Then we call  $\tilde{\gamma} : J \rightarrow \mathbb{R}^n$  with  $\tilde{\gamma} := \gamma \circ \sigma$  a reparameterization of  $\gamma$ . Compare with Figure 45.

On the opposite, let  $\gamma, \tilde{\gamma}$  be a curve parameterization. Then  $\tilde{\gamma}$  is called reparameterization of  $\gamma$  if  $\sigma$  exists as above such that  $\tilde{\gamma} = \gamma \circ \sigma$ .

**Definition 7.4** (Discussion of terminology). Let  $X, Y$  be topological spaces.  $f : X \rightarrow Y$  is called homeomorphism if  $f$  is continuous and bijective and furthermore  $f^{-1} : Y \rightarrow X$  is also continuous.

Equivalently, for  $C^\infty$ -diffeomorphism,  $f$  is smooth and  $f^{-1}$  is smooth.

$\sigma : J \rightarrow I$  is bijective and continuous, hence  $\sigma$  is either strictly monotonically increasing or strictly monotonically decreasing.

**Definition 7.5.** If  $\sigma$  is strictly monotonically increasing, we call a reparameterization orientation preserving. If  $\sigma$  is strictly monotonically decreasing, we call a reparameterization orientation reversing. Compare with Figure 46.

( $\sigma$  is a diffeomorphism,  $\sigma' > 0$  and accordingly  $\sigma' < 0$  defines the orientation.  $\sigma$  is called parameter exchange)

**Claim.** Let  $G = \{\gamma : \gamma \text{ is a regular curve parameterization}\}$ . We define a relation  $\sim_r$  on  $G$ .  $\gamma \sim_r \tilde{\gamma} \iff \tilde{\gamma}$  is a reparameterization of  $\gamma$ . Then  $\sim_r$  is an equivalence relation.

**Proof. Reflexivity**  $\gamma \sim_r \gamma$  with  $\sigma : I \rightarrow I, \sigma = \text{id}$ .

**Symmetry** Let  $\tilde{\gamma} \sim_r \gamma$ , hence  $\tilde{\gamma} = \gamma \circ \sigma \implies \tilde{\gamma} \circ \sigma^{-1} = \gamma \implies \gamma \sim_r \tilde{\gamma}$ .




Figure 45: Reparameterization

**Transitivity** Let  $\tilde{\gamma} \sim_r \tilde{\gamma}$  and  $\tilde{\gamma} \sim_r \gamma$ , hence  $\tilde{\gamma} = \tilde{\gamma} \circ \rho$ .  $\tilde{\gamma} = \gamma \circ \sigma \implies \tilde{\gamma} = \gamma \circ \sigma \circ \rho \implies \tilde{\gamma} \sim_r \gamma$  with  $\sigma \circ \rho$  as diffeomorphism.

□

**Definition 7.6.** Let  $[\gamma]_{\sim_r}$  be an equivalence class of  $\gamma$  with respect to  $\sim_r$ . A size or property, which only depends on  $[\gamma]_{\sim_r}$  but not on a specific representative  $\gamma$  is called geometrical quantity.

**Example 7.1.** Later, we will show: arc length is a geometrical property.

**Definition 7.7.**  $[\gamma]_{\sim_r}$  is called closed curve if  $\gamma(a) = \gamma(b)$  with  $\gamma : [a, b] \rightarrow \mathbb{R}^n$ .

Because  $\gamma(t) = \tilde{\gamma}(\sigma(t))$  with  $\tau = \sigma^{-1}(t)$ , parameterizations have the same trace  $\Gamma = \gamma(I) = \tilde{\gamma}(J)$ . Sometimes we call  $[\gamma]_{\sim_r}$  a curve in  $\mathbb{R}^n$ .

**Example 7.2.** Let  $I = [0, 2\pi]$ .  $\gamma(t) := \begin{pmatrix} \cos(t) \\ \sin(t) \end{pmatrix}$ .  $\gamma : I \rightarrow \mathbb{R}^2$ . It holds  $\gamma(0) = \gamma(2\pi)$ , hence  $\gamma$  is closed. Compare with Figure 47.

$$\sigma : [-2\pi, 0] \rightarrow [0, 2\pi] \quad \sigma : t \mapsto -t$$

$$\hat{\gamma}(t) = \begin{pmatrix} \cos(t) \\ -\sin(t) \end{pmatrix} \text{ with } t \in [0, 2\pi]$$

Why?

$$\hat{\gamma}(t) = \gamma \circ \sigma(t) = \begin{pmatrix} \cos(-t) \\ \sin(-t) \end{pmatrix} = \begin{pmatrix} \cos(t) \\ -\sin(t) \end{pmatrix} = \tilde{\gamma}(t)$$




Figure 46: Orientation

with  $\tilde{\gamma} : [-2\pi, 0] \rightarrow \mathbb{R}^2$  is periodic with period  $2\pi$ . Hence,  $\tilde{\gamma}(t - 2\pi) = \tilde{\gamma}(t) \forall t \in \mathbb{R}$ .

$$\hat{\gamma}(t) := \tilde{\gamma}(t - 2\pi) \quad \hat{\gamma} : [0, 2\pi]$$

$$\hat{\gamma}(t) = \begin{pmatrix} \cos(t) \\ -\sin(t) \end{pmatrix} \quad \hat{\gamma}(t) = \underbrace{\gamma(-t + 2\pi)}_{\sigma(t)}$$

with  $\sigma'(t) = -1$ . Hence reparameterization  $\hat{\gamma}$  is orientation reversing.

**Example 7.3.** Let  $\gamma : \mathbb{R} \rightarrow \mathbb{R}^2$  (see Figure 49).

$$\gamma(t) = \begin{pmatrix} t^2 - 1 \\ t^3 - t \end{pmatrix} \quad \gamma'(t) = \begin{pmatrix} 2t \\ 3t^2 - 1 \end{pmatrix}$$

$$\begin{aligned} \gamma'(t) = 0 &\implies 2t = 0 \text{ hence } t = 0 \\ &\implies 3t^2 - 1 = -1 \neq 0 \end{aligned}$$

As  $\gamma'(t) \neq 0$ , the parameterization is regular. It holds that  $\gamma(1) = \begin{pmatrix} 0 & 0 \end{pmatrix}^T$  and  $\gamma(-1) = \begin{pmatrix} 0 & 0 \end{pmatrix}^T$ .  $\begin{pmatrix} 0 & 0 \end{pmatrix}^T$  lies on the trace of the curve for 2 different parameter values.  $x = 0$  is a double point of the curve.

$$\gamma'(-1) = \begin{pmatrix} -2 \\ 2 \end{pmatrix} \quad T_{\gamma}(-1) = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

$$\gamma'(1) = \begin{pmatrix} 2 \\ 2 \end{pmatrix} \quad T_{\gamma}(1) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

For parameter values  $t = -1$  and  $t = 1$ , tangents exist. But in  $x = 0$ , we get ambiguity with  $\gamma(1) = \gamma(-1)$ . So, no tangent exists.




Figure 47: Example 7.2 defines a curve with  $\gamma(t) := [\cos(t) \quad \sin(t)]^T$




Figure 48: Example 7.2 shows the orientation reversion with  $\sigma'(t) = -1$




Figure 49: Example 7.3 gives two different tangents at the origin




Figure 50: Plot of Neile's parabola (Example 7.4)




Figure 51:  $\Gamma : \gamma(\mathbb{R})$




Figure 52:  $T_{\tilde{\gamma}}(\tau) = -T_{\gamma}(t)$

**Example 7.4** (Neile's parabola). Let  $\gamma : \mathbb{R} \rightarrow \mathbb{R}^2$ .

$$\gamma(t) = \begin{pmatrix} t^2 \\ t^3 \end{pmatrix} \quad \gamma'(t) = \begin{pmatrix} 2t \\ 3t^2 \end{pmatrix} \quad \gamma(0) = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \gamma'(0) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Compare with Figure 50. Curvature parameterization is not regular. For  $t = 0$ , no tangent exists.

**Example 7.5.**

$$\gamma(t) = \begin{pmatrix} t^3 \\ t^3 \end{pmatrix} \quad \gamma'(t) = \begin{pmatrix} 3t^2 \\ 3t^2 \end{pmatrix} \quad \gamma'(0) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$\Rightarrow$  non-regular. Compare with Figure 51.  $\Gamma$  is smooth and has a “geometrical” tangent everywhere.

Let  $\sigma$  be an orientation preserving parameter exchange. Let  $\gamma$  be regular.

$$\tilde{\gamma} = \gamma \circ \sigma \quad \gamma(t) = \tilde{\gamma}(\tau) \quad \text{with } t = \sigma(\tau) \text{ and } \tau = \sigma^{-1}(t)$$

$$T_{\gamma}(t) = \frac{\gamma'(t)}{\|\gamma'(t)\|}$$

$$\tilde{\gamma}'(t) = (\gamma \circ \sigma)'(\tau) = \gamma'(\sigma(\tau)) \cdot \sigma'(\tau) = \gamma'(t) \cdot \underbrace{\sigma'(\tau)}_{>0}$$

$$\Rightarrow \|\tilde{\gamma}'(\tau)\|_2 = \|\gamma'(t) \cdot \sigma'(\tau)\|_2 = \sigma'(\tau) \|\gamma'(t)\|_2$$

$$T_{\tilde{\gamma}}(\tau) = \frac{\tilde{\gamma}'(\tau)}{\|\tilde{\gamma}'(\tau)\|_2} = \frac{\gamma'(t) \cdot \sigma'(\tau)}{\|\gamma'(t)\| \cdot \sigma'(\tau)} = T_{\gamma}(t)$$

Let  $\sigma$  be orientation reversing.

$$\gamma'(\tau) < 0 \quad |\gamma'(\tau)| = -\sigma'(\tau)$$

like above  $\tilde{\gamma}'(\tau) = \gamma'(t) \cdot \sigma'(\tau)$  and  $\|\tilde{\gamma}'(\tau)\|_2 = |\sigma'(\tau)| \cdot \|\gamma'(t)\|_2 = -\sigma'(\tau) \|\gamma'(t)\|_2$ .

$$T_{\tilde{\gamma}}(\tau) = -T_{\gamma}(t)$$

Compare with Figure 52.




Figure 53: Polygonal line

### 7.3 Arc length of parameterized curves

**Definition 7.8.** Let  $I = [a, b]$  be a compact interval. Let  $\gamma : I \rightarrow \mathbb{R}^n$  be a continuous curve. We define a partition  $Z$  of interval  $[a, b]$ :

$$Z := (a = t_0 < t_1 < t_2 < \cdots < t_N = b)$$

$$S_\gamma(Z) := \sum_{k=1}^N \|\gamma(t_k) - \gamma(t_{k-1})\|_2$$

$S_\gamma(Z)$  is the length of the polygonal line through points  $(x_k)_{k=0}^N$  with  $x_k = \gamma(t_k)$ . We let  $S_\gamma = \sup \{S_\gamma(Z) : Z \text{ is partition of } [a, b]\}$ .

If  $S_\gamma < \infty$ , then  $\gamma$  is called rectifiable and  $S_\gamma$  is the arc length of the curve.

**Remark 7.1.** Let  $\tilde{\gamma}$  be a reparameterization of  $\gamma$ . Let  $t = \sigma(\tau)$  be a parameter exchange and  $\sigma : [\alpha, \beta] \rightarrow [a, b]$ .  $t_k = \sigma(\tau_k)$  with  $k = 0, \dots, N$ . If  $\sigma$  is orientation preserving, then  $\tilde{Z} = \{\alpha = \tau_0 < \tau_1 < \cdots < \tau_N = \beta\}$  is a partition of  $[\alpha, \beta]$ . If  $\sigma$  is orientation reversing, then  $\tilde{Z} = \{\alpha = \tau_N < \tau_{N-1} < \cdots < \tau_0 = \beta\}$  is a partition. In every case,

$$S_\gamma(Z) = S_{\tilde{\gamma}}(\tilde{Z})$$

because  $x_k = \gamma(t_k) = \tilde{\gamma}(\tau_k)$  and accordingly  $x_k = \gamma(t_k) = \tilde{\gamma}(\tau_{N-k})$ . So  $S_\gamma$  is a geometrical quantity.

**Lemma 7.1.** Let  $\gamma : [a, b] \rightarrow \mathbb{R}^n$  be Lipschitz continuous, hence  $\exists L \geq 0$  such that  $\|\gamma(s) - \gamma(t)\|_2 \leq L|s - t| \forall s, t \in [a, b]$ . Then  $\gamma$  is rectifiable.

*Proof.* Let  $Z$  be a partition of  $[a, b]$ .  $Z = (t_k)_{k=0}^N$ . Then

$$\begin{aligned}
S_\gamma(Z) &= \sum_{k=1}^N \|\gamma(t_k) - \gamma(t_{k-1})\|_2 \leq \underbrace{\sum_{k=1}^N L}_{\text{by Lipschitz continuity}} \underbrace{|t_k - t_{k-1}|}_{\substack{= t_k - t_{k-1} \\ \text{because } t_k > t_{k-1}}} \\
&= \sum_{k=1}^N L(t_k - t_{k-1}) = L(t_N - t_0) = L(b - a) < \infty \\
&\implies \sup \{S_\gamma(Z) : Z \text{ partition of } [a, b]\} \leq L(b - a) < \infty
\end{aligned}$$

□

**Lemma 7.2.** Let  $\alpha : [a, b] \rightarrow \mathbb{R}^n$ .  $\alpha(t) = [\alpha_1(t), \alpha_2(t), \dots, \alpha_n(t)]^T$ .  $\alpha_n \in \mathcal{R}[a, b]$  for  $k = 1, \dots, n$ .

$$\begin{aligned}
\int_a^b \alpha(t) dt &= \left[ \int_a^b \alpha_1 dt, \int_a^b \alpha_2 dt, \dots, \int_a^b \alpha_n dt \right]^T \\
\text{Then } \left\| \int_a^b \alpha dt \right\|_2 &\leq \int_a^b \|\alpha(t)\|_2 dt
\end{aligned}$$

*Proof.* Approximation by step functions: Let  $\varphi : [a, b] \rightarrow \mathbb{R}^n$ .  $\gamma(t) = [\varphi_1(t), \dots, \varphi_n(t)]^T$ .  $\gamma_i \in \mathcal{R}[a, b]$ . Without loss of generality all  $\varphi_i$  are step functions with respect to the same partition  $Z = (t_k)_{k=0}^N$ <sup>4</sup>.

$$\varphi_i(t) = C_i^k \text{ for } t \in (t_{k-1}, t_k)$$

$$\int_a^b \varphi_i dt = \sum_{k=1}^N c_i^k (t_k - t_{k-1})$$

$\varphi_i$  should approximate  $\alpha_i$  uniformly, hence for  $\varepsilon > 0$  arbitrarily chosen, choose  $\varphi_i$  such that

$$|\varphi_i(t) - \alpha_i(t)| < \frac{\varepsilon}{2\sqrt{n}(b-a)} \quad \forall t \in [a, b]$$

For step function  $\varphi$ ,

$$\left\| \int_a^b \varphi(t) dt \right\|_2 = \left\| \sum_{k=1}^N (t_k - t_{k-1}) [c_1^k, c_2^k, \dots, c_n^k]^T \right\|$$

By the triangle inequality in  $\mathbb{R}^n$ ,

$$\begin{aligned}
&\leq \sum_{k=1}^n (t_k - t_{k-1}) \cdot \underbrace{\left\| [c_1^k, \dots, c_n^k]^T \right\|_2}_{\|\varphi(t)\|_2 \text{ for } t \in (t_{k-1}, t_k)} = \int_a^b \|\varphi(t)\|_2 dt
\end{aligned}$$

<sup>4</sup>If not, take the union of two non-equal-sized intervals and apply the lemma to a refinement of this new interval again. Therefore without loss of generality.

↓ This lecture took place on 2018/06/21.

Let  $\alpha$  as in the lemma statement. Let  $\varepsilon > 0$  be arbitrary and  $\varphi_k \in \tau[a, b]$  such that  $|\alpha_k(t) - \varphi_k(t)| < \frac{\varepsilon}{2\sqrt{n(b-a)}} \forall t \in [a, b] \forall k = 1, \dots, n$ . Hence,

$$\begin{aligned} \int_a^b \|\alpha - \varphi\|_2 dt &= \int_a^b \sqrt{\sum_{k=1}^n |\alpha_k(t) - \varphi_k(t)|^2} dt & \varphi &:= [\varphi_1, \dots, \varphi_n]^n \\ &\leq \int_a^b \sqrt{\sum_{k=1}^n \frac{\varepsilon^2}{4n(b-a)^2}} dt \\ & \quad \underbrace{\hspace{10em}}_{=\frac{\varepsilon^2}{4(b-a)^2}} \\ &= \int_a^b \frac{\varepsilon}{2(b-a)} dt = (b-a) \cdot \frac{\varepsilon}{2(b-a)} = \frac{\varepsilon}{2} \end{aligned}$$

Also,

$$\begin{aligned} \left\| \int_a^b (\alpha - \varphi) dt \right\|_2 &= \left\| \begin{bmatrix} \int_a^b (\alpha_1 - \varphi_1) dt \\ \vdots \\ \int_a^b (\alpha_n - \varphi_n) dt \end{bmatrix} \right\|_2 = \left\| \begin{bmatrix} \left| \int_a^b (\alpha_1 - \varphi_1) dt \right| \\ \vdots \\ \left| \int_a^b (\alpha_n - \varphi_n) dt \right| \end{bmatrix} \right\|_2 \\ &= \left\| \begin{bmatrix} \int_a^b |\alpha_1 - \varphi_1| dt \\ \vdots \\ \int_a^b |\alpha_n - \varphi_n| dt \end{bmatrix} \right\|_2 \leq \left\| \begin{bmatrix} \int_a^b \frac{\varepsilon}{2\sqrt{n(b-a)}} dt \\ \vdots \\ \int_a^b \frac{\varepsilon}{2\sqrt{n(b-a)}} dt \end{bmatrix} \right\|_2 = \left\| \begin{bmatrix} \frac{\varepsilon}{2\sqrt{n}} \\ \vdots \\ \frac{\varepsilon}{2\sqrt{n}} \end{bmatrix} \right\|_2 \\ &= \sqrt{\sum_{k=1}^n \frac{\varepsilon^2}{4n}} = \frac{\varepsilon}{2} \end{aligned}$$

Hence,  $\left\| \int_a^b \alpha dt \right\| = \left\| \int_a^b (\alpha - \varphi) dt + \int_a^b \varphi dt \right\|_2 \leq \left\| \int_a^b (\alpha - \varphi) dt \right\|_2 + \left\| \int_a^b \varphi dt \right\| \leq \frac{\varepsilon}{2} +$



$\int_a^b \|\varphi\|_2 dt$  because  $\varphi \in \tau[a, b]$ .

$$\begin{aligned}
\left\| \int_a^b \alpha dt \right\| &\leq \frac{\varepsilon}{2} + \int_a^b \|\varphi - \alpha + \alpha\|_2 dt \\
&\leq \frac{\varepsilon}{2} + \int_a^b \|\varphi - \alpha\|_2 dt + \int_a^b \|\alpha\|_2 dt \\
&\leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} + \int_a^b \|\alpha\|_2 dt \\
\Rightarrow \left\| \int_a^b \alpha dt \right\|_2 &\leq \varepsilon + \int_a^b \|\alpha\|_2 dt \quad \forall \varepsilon > 0 \\
\Rightarrow \left\| \int_a^b \alpha dt \right\|_2 &\leq \int_a^b \|\alpha\| dt
\end{aligned}$$

□

**Theorem 7.1.** Let  $\gamma : [a, b] \rightarrow \mathbb{R}^n$  be a continuous curve parameterization.  $\gamma(t) = [\gamma_1(t), \dots, \gamma_n(t)]^T$  and let  $\gamma_k$  be an antiderivative of regulated function  $\alpha_k = \gamma'_k(t)$ . Then  $\gamma$  is rectifiable and:

$$S_\gamma = \int_a^b \|\gamma'(t)\|_2 dt$$

*Proof.* Let  $z = (t_i)_{i=0}^N$  be a partition of  $[a, b]$ .

$$\begin{aligned}
S_\gamma(z) &= \sum_{i=1}^N \|\gamma(t_i) - \gamma(t_{i-1})\|_2 \underbrace{=}_{\text{fundamental theorem}} \sum_{i=1}^N \left\| \int_{t_{i-1}}^{t_i} \gamma'(t) dt \right\|_2 \\
&\leq \underbrace{\sum_{i=1}^N \int_{t_{i-1}}^{t_i} \|\gamma'(t)\|_2 dt}_{\text{Lemma 7.2}} = \int_{t_0}^{t_N} \|\gamma'(t)\|_2 dt = \int_a^b \|\gamma'(t)\|_2 dt
\end{aligned}$$

So,  $S_\gamma(Z) \leq \int_a^b \|\gamma'(t)\|_2 dt \forall$  partitions  $Z$ . Therefore,  $S_\gamma \leq \int_a^b \|\gamma'(t)\|_2 dt$  and  $\gamma$  is rectifiable.

Show:  $\forall \varepsilon > 0$ ,  $S_\gamma \geq \int_a^b \|\gamma'(t)\|_2 dt - \varepsilon$ .  $\gamma'$  is a regulated function. Choose  $\varphi = [\varphi_1, \dots, \varphi_n]^T$  with  $\varphi_k \in \tau[a, b]$  with  $\|\varphi(t) - \gamma'(t)\| \leq \frac{\varepsilon}{2(b-a)} \forall t \in [a, b]$ .

$$\begin{aligned}
\varphi(t) &= \begin{bmatrix} c_i^1 & c_i^2 & \dots & c_i^n \end{bmatrix}^T \text{ on } (t_{i-1}, t_i) \\
\Rightarrow \left\| \int_{t_{i-1}}^{t_i} \varphi dt \right\|_2 &= \left\| \begin{bmatrix} c_i^1 \\ \vdots \\ c_i^n \end{bmatrix} \cdot (t_i - t_{i-1}) \right\|_2 = (t_i - t_{i-1}) \left\| \begin{bmatrix} c_i^1 \\ \vdots \\ c_i^n \end{bmatrix} \right\|_2
\end{aligned}$$

$$\int_{t_{i-1}}^{t_i} \|\varphi\| dt = \int_{t_{i-1}}^{t_i} \left\| \begin{bmatrix} c_i^1 \\ \vdots \\ c_i^n \end{bmatrix} \right\|_2 dt = (t_i - t_{i-1}) \left\| \begin{bmatrix} c_i^1 \\ \vdots \\ c_i^n \end{bmatrix} \right\|_2$$

Hence, for  $(t_{i-1}, t_i)$

$$\begin{aligned} \int_{t_{i-1}}^{t_i} \|\varphi\| dt &= \left\| \int_{t_{i-1}}^{t_i} \varphi dt \right\| \\ \|\gamma'(t_i) - \gamma'(t_{i-1})\|_2 &= \left\| \int_{t_{i-1}}^{t_i} \gamma'(t) dt \right\|_2 \\ &= \left\| \int_{t_{i-1}}^{t_i} [\varphi(t) - (\varphi(t) - \gamma'(t))] dt \right\|_2 \geq \left\| \int_{t_{i-1}}^{t_i} \varphi(t) dt \right\|_2 - \left\| \int_{t_{i-1}}^{t_i} (\varphi - \gamma') dt \right\|_2 \end{aligned}$$

Because  $\left\| \int_{t_{i-1}}^{t_i} \varphi(t) dt \right\|_2 = \int_{t_{i-1}}^{t_i} \|\gamma'\|_2 dt$ ,

$$\begin{aligned} &\geq \int_{t_{i-1}}^{t_i} \|\varphi\|_2 dt - \underbrace{\int_{t_{i-1}}^{t_i} \|\varphi - \gamma'\|_2 dt}_{\leq \frac{\varepsilon}{2(b-a)}} \geq \int_{t_{i-1}}^{t_i} \|\varphi\|_2 dt - \frac{\varepsilon(t_i - t_{i-1})}{2(b-a)} \end{aligned}$$

$$\begin{aligned} S_\gamma(z) &= \sum_{i=1}^N \left\| \int_{t_{i-1}}^{t_i} \gamma'(t) dt \right\| \geq \sum_{i=1}^n \int_{t_{i-1}}^{t_i} \|\varphi\|_2 dt - \frac{\varepsilon}{2(b-a)}(t_i - t_{i-1}) \\ &= \int_a^b \|\varphi\|_2 dt - \frac{\varepsilon}{2} = \int_a^b \|\gamma' - (\gamma' - \varphi)\|_2 dt - \frac{\varepsilon}{2} \\ &\geq \int_a^b \|\gamma'\|_2 dt - \int_a^b \left\| \underbrace{\gamma' - \varphi}_{\leq \frac{\varepsilon}{2(b-a)}} \right\|_2 dt - \frac{\varepsilon}{2} \geq \int_a^b \|\gamma'(t)\|_2 dt - \frac{\varepsilon}{2(b-a)} \cdot (b-a) - \frac{\varepsilon}{2} \\ &= \int_a^b \|\gamma'(t)\|_2 dt - \varepsilon \end{aligned}$$

$$\begin{aligned} \implies S_\gamma &\geq S_\gamma(z) \geq \int_a^b \|\gamma'(t)\|_2 dt - \varepsilon \quad \forall \varepsilon > 0 \\ \implies S_\gamma &\geq \int_a^b \|\gamma'(t)\|_2 dt \end{aligned}$$

□

**Remark 7.2.** By definition of  $S_\gamma$ , the arc length is a geometrical quantity.

**Remark 7.3.** Another rationale for the independence of parameterization:

Let  $\gamma : I \rightarrow \mathbb{R}^n$  be a regular  $C^1$ -curve parameterization. Let  $\sigma : J \rightarrow I$  be a diffeomorphism and parameter exchange.  $\tilde{\gamma}(\tau) = \gamma \circ \sigma(\tau)$ .




Figure 54: Traversed path

$\sigma$  is orientation preserving.  $J = [\alpha, \beta], I = [a, b] \implies \sigma(\alpha) = a, \sigma(\beta) = b$ .

$$\begin{aligned}
 S_{\tilde{\gamma}} &= \int_{\alpha}^{\beta} \|\tilde{\gamma}'(\tau)\|_2 d\tau = \int_{\alpha}^{\beta} \|[\gamma'(\sigma(\tau))]\|_2 d\tau = \int_{\alpha}^{\beta} \left\| \gamma'(\sigma(\tau)) \cdot \underbrace{\sigma'(\tau)}_{>0} \right\|_2 d\tau \\
 &= \int_{\alpha}^{\beta} \|\gamma'(\sigma(\tau))\|_2 \cdot \sigma'(\tau) d\tau \stackrel{\text{transformation theorem}}{=} \int_a^b \|\gamma'(t)\|_2 dt = S_{\gamma}
 \end{aligned}$$

Let  $\sigma$  be orientation reversing, hence  $\sigma'(\tau) < 0$ .  $\sigma(\alpha) = b$  and  $\sigma(\beta) = a$ .

$$\begin{aligned}
 S_{\tilde{\gamma}} &= \int_{\alpha}^{\beta} \|\tilde{\gamma}'(\tau)\|_2 d\tau = \int_{\alpha}^{\beta} \left\| \gamma'(\sigma(\tau)) \cdot \underbrace{\sigma'(\tau)}_{<0} \right\|_2 d\tau = - \int_{\alpha}^{\beta} \|\gamma'(\sigma(\tau))\|_2 \underbrace{\sigma'(\tau)}_{=-|\sigma'(\tau)|} dt \\
 &= \int_{\beta}^{\alpha} \|\gamma'(\sigma(\tau))\|_2 \cdot \sigma'(\tau) d\tau = \int_{\sigma(\beta)}^{\sigma(\alpha)} \|\gamma'(t)\|_2 dt = \int_a^b \|\gamma'(t)\|_2 dt
 \end{aligned}$$

Idea: Define  $s = l(t) = \int_{t_0}^t \|\gamma'(\tau)\|_2 d\tau$ . Compare with Figure 54.

$$\text{For } \tilde{t} < t, l(\tilde{t}) = \int_t^{\tilde{t}} \|\gamma'(\tau)\|_2 d\tau = - \underbrace{\int_{\tilde{t}}^t \|\gamma'(\tau)\|_2 d\tau}_{>0} < 0.$$

**Definition 7.9.** We call  $s = l(t) = \int_{t_0}^t \|\gamma'(\tau)\|_2 d\tau$  the arc length parameter of the curve parameterization  $\gamma$ .

Let  $\gamma$  be regular. Then (by the Fundamental Theorem)  $l$  is continuously differentiable and

$$l'(t) = \frac{d}{dt} \left[ \int_{t_0}^t \|\gamma'(\tau)\|_2 d\tau \right] = \|\gamma'(t)\|_2 > 0$$

Hence,  $l$  is strictly monotonically increasing.

Let  $I = [a, b]$ .  $t_0 \in I$  and  $I$  is compact. Then  $l : I \rightarrow J = l(I)$  is strictly monotonically increasing and continuously differentiable.  $J = [\alpha, \beta] = [l(a), l(b)]$ .  $l^{-1} : J \rightarrow I$  is also continuous.  $l^{-1}$  is also continuously differentiable with

$$(l^{-1})'(s) = \frac{1}{l'(l^{-1}(s))} = \frac{1}{\|\gamma'(t)\|_2} > 0$$

with  $t = l^{-1}(s)$  and accordingly  $s = l(t)$ .  $l^{-1}$  is strictly monotonically increasing and continuously differentiable.

Let  $\tilde{\gamma}(s) = \gamma \circ l^{-1}(s)$ . Let  $\tilde{\gamma}$  be the reparameterization of  $\gamma$  by the arc length. Often we write  $\gamma(s)$  instead of  $\tilde{\gamma}(s)$ .  $s$  is a very common variable denoting the arc length parameter. Often:  $s = s(t)$  instead of  $s = l(t)$ ,  $t = s^{-1}(s)$  is inappropriate.

Let  $\tilde{\gamma}$  be the reparameterization of  $\gamma$  by the arc length. Then

$$\tilde{\gamma}'(s) = \frac{d}{ds} [\gamma(l^{-1}(s))] = \gamma'(l^{-1}(s))(l^{-1})'(s) = \gamma'(t) \cdot \frac{1}{l'(t)} = \gamma'(t) \cdot \frac{1}{\|\gamma'(t)\|_2} = T_{\gamma}(t)$$

because for derivatives of inverse functions,  $(f^{-1})'(x) = \frac{1}{f'(f^{-1}(x))}$ . So,  $\|\tilde{\gamma}'(s)\|_2 = 1$ .  $\tilde{\gamma}'(s) = T_{\tilde{\gamma}}(s)$ .

Determine the arc length for  $\tilde{\gamma}$ :

$$\begin{aligned} \tilde{l}(s) &= \int_0^s \underbrace{\|\tilde{\gamma}'(\xi)\|_2}_{=1} d\xi = s \\ \tilde{\gamma}(0) &= \gamma(t_0) \end{aligned}$$

**Example 7.6** (Cycloid).

$$\gamma(t) = \begin{bmatrix} t - \sin(t) \\ 1 - \cos(t) \end{bmatrix}$$

Figure 55 illustrates the trace of a cycloid.

$$\gamma'(t) = \begin{bmatrix} 1 - \cos(t) \\ \sin(t) \end{bmatrix}$$

$$\begin{aligned} s &= \int_0^t \|\gamma'(\tau)\|_2 d\tau = \int_0^t \sqrt{(1 - \cos \tau)^2 + \sin^2 \tau} d\tau \\ &= \int_0^t \sqrt{1 - 2 \cos \tau + 1} d\tau = \int_0^t \sqrt{2} \sqrt{1 - \cos \left( 2 \frac{\tau}{2} \right)} d\tau \end{aligned}$$




Figure 55: Trace of a cycloid

$$\cos\left(\frac{\tau}{2} + \frac{\tau}{2}\right) = \cos\left(\frac{\tau}{2}\right)\cos\left(\frac{\tau}{2}\right) - \sin\left(\frac{\tau}{2}\right)\sin\left(\frac{\tau}{2}\right)$$

$$= \sqrt{2} \int_0^t \underbrace{\sqrt{1 - \cos^2 \frac{\tau}{2} + \sin^2 \frac{\tau}{2}}}_{=\sin^2 \frac{\tau}{2}} d\tau = \sqrt{2} \int_0^t \sqrt{2} \sqrt{\sin^2 \frac{\tau}{2}} d\tau$$

$$t \in [0, 2\pi] \implies \frac{\tau}{2} \in [0, \pi] \implies \sin \frac{\tau}{2} \geq 0. \quad \sqrt{\sin^2 \left(\frac{\tau}{2}\right)} = \sin\left(\frac{\tau}{2}\right)$$

$$= 2 \int_0^t \sin \frac{\tau}{2} d\tau = \left| \begin{array}{l} \text{Integration by substitution} \\ f(x) = \sin(x) \\ \varphi(x) = 2x \\ \varphi'(x) = 2 \end{array} \right| = 2 \int_0^{\frac{t}{2}} \sin(x) \cdot 2 dx$$

$$= -4 \cos(x) \Big|_0^{\frac{t}{2}} = 4 \left(1 - \cos \frac{t}{2}\right) = s$$

$$l(t) = 4 \left(1 - \cos \frac{t}{2}\right)$$

$$l^{-1}(s) = ?$$

$$1 - \cos \frac{t}{2} = \frac{s}{4}$$

$$\cos \frac{t}{2} = 1 - \frac{s}{4} = \frac{1}{4}(4 - s) \in [-1, +1]$$

$$t = 2 \arccos\left(\frac{1}{4}(4 - s)\right)$$

$$\tilde{\gamma}(s) = \left[ \begin{array}{l} 2 \arccos\left(\frac{1}{4}(4 - s)\right) - \sin\left(2 \arccos\left(\frac{1}{4}(4 - s)\right)\right) \\ 1 - \cos\left(2 \arccos\left(\frac{1}{4}(4 - s)\right)\right) \end{array} \right]$$

**Remark 7.4.** The reparameterization by the arc length often do not yield algebraically simpler expression. But it provides the property, that the curve is traversed with uniform speed.




Figure 56: Orthogonal  $\gamma'(t)$  and  $\gamma''(t)$

**Definition 7.10.** Let  $\gamma : I \rightarrow \mathbb{R}^n$  be a regular curve. We call  $\gamma$  parameterized by its arc length, if  $\|\gamma'(t)\|_2 = 1 \forall t \in I$ .

**Remark 7.5.** Let  $\gamma$  be parameterized by an arc length. Then  $s = l(t) = \int_{t_0}^t \|\gamma'(\tau)\|_2 d\tau = \int_{t_0}^t 1 d\tau = t - t_0$ , and accordingly  $t = s + t_0$ . Hence, curve parameter and arc length parameter differ only by constant  $t_0$ . For  $t_0 = 0$ ,  $s = t$ .

Let  $\gamma$  be regular and a  $C^2$  curve and  $\|\gamma'(t)\|_2 = 1 \forall t \in I$ , hence  $\gamma$  is parameterized by the arc length.

Consider  $1 = \|\gamma'(t)\|_2^2 = (\gamma'(t))^T \cdot \gamma'(t)$ . If we derive by  $t$ ,

$$0 = \gamma''(t)^T \cdot \gamma'(t) + \gamma'(t)^T \cdot \gamma''(t) = 2 \langle \gamma'(t), \gamma''(t) \rangle$$

Hence  $\gamma''(t)$  is orthogonal to  $\gamma'(t)$ . Compare with Figure 56.

↓ This lecture took place on 2018/06/26.

## 7.4 Curvature, Torsion, Frenet's formulas

Let  $\gamma : I \rightarrow \mathbb{R}^3, \gamma \in C^3(I, \mathbb{R}^3), \|\gamma'(s)\| = 1$ . Hence  $\gamma$  is parameterized by the arc length.

$$\gamma'(s) = T_\gamma(s) = T_\gamma(s)$$

and  $\langle \gamma'(s), \gamma''(s) \rangle = 0$ , hence  $\gamma''(s) \perp \Gamma(s)$




Figure 57: Orthogonal  $T$  and  $N$

**Definition 7.11.** Let  $\gamma$  be like above. We call  $\kappa(s) = \frac{\|\gamma''(s)\|_2}{\|T'(s)\|_2} = \|T'(s)\|_2$  the curvature of  $\gamma$  in  $x = \gamma(s)$ . If  $\kappa(s) \neq 0$ , hence  $\gamma''(s) \neq 0$ , then we let

$$N = N(s) = \frac{\gamma''(s)}{\|\gamma''(s)\|_2}$$

We call  $N$  the principal orthogonal vector of  $\gamma$  in  $x = \gamma(s)$ .

**Remark 7.6.**

$$N \perp T$$

Compare with Figure 57.  $\kappa(T)$  is the scalar rate of change in direction of  $T$ .

Because  $s = l(t)$  is a geometrical quantity, the reparameterization by the arc length is independent of the original parameterization. Hence  $\kappa, N$  are geometrical quantities. By the definition, it follows that

$$T'(s) = \kappa(s) \cdot N(s)$$

We call this equation the first Frenet formula.

**Definition 7.12.** Let  $\gamma$  be parameterized by the arc length.  $\kappa(s) \neq 0$ , hence  $N$  is defined. We call  $\varepsilon_x = \{x + \xi T + \eta N : \xi, \eta \in \mathbb{R}\}$  the osculating plane of  $\gamma$  in  $x$  (dt. Schmieg Ebene). Compare with Figure 58.

$m = m(s) = x + \frac{1}{\kappa} \cdot N$  is called curvature center of  $\gamma$  in  $x = \gamma(s)$ .

$$K_X(\sigma) = m + \frac{1}{\kappa} (\sin \kappa \sigma \cdot T - \cos \kappa \sigma \cdot N)$$

is called circle of curvature of  $\gamma$  in  $x = \gamma(s)$ .




Figure 58: The osculating plane  $\varepsilon_X$

**Remark 7.7.**

$$m = x + \frac{1}{\kappa} \cdot N \in \varepsilon_X$$

$m$  is in the osculating plane  $K_x(\sigma) \in \varepsilon_x \forall \sigma \in \mathbb{R}$ . Compare with Figure 59. In the osculating plane, the red vertical line is given by  $\frac{1}{\kappa} \cdot \sin \kappa \sigma$  and the horizontal orange line is given by  $\frac{1}{\kappa} \cos \kappa \sigma$ .

$$\begin{aligned} \tilde{\gamma}(\sigma) &= \gamma(s + \sigma) \\ \tilde{\gamma}(0) &= \gamma(s) = x \\ \|\tilde{\gamma}'(\sigma)\| &= \|\gamma'(s + \sigma)\| = 1 \end{aligned}$$






Figure 59: Circle of curvature with radius  $\frac{1}{\kappa}$ , center  $m$  and  $x \in K_x$  for  $\sigma = 0$

Hence  $\tilde{\gamma}$  is also parameterized by the arc length.

$$\begin{aligned}
 \tilde{\gamma}(0) &= x \\
 K_x(0) &= m + \frac{1}{\kappa} (\underbrace{(\sin \kappa \cdot 0)}_{=0} T - \underbrace{(\cos \kappa \cdot 0)}_{=1} \cdot N) \\
 &= x + \frac{1}{\kappa} N - \frac{1}{\kappa} N = x \\
 \tilde{\gamma}'(0) &= \gamma'(s) = T \\
 K'_x(\sigma) &= \frac{1}{\kappa} [(\kappa \cos \kappa \sigma \cdot T + (\kappa \cdot \sin \kappa \sigma) \cdot N)] \\
 K'_x(0) &= \frac{1}{\kappa} \cdot \kappa \cdot T = T \\
 \tilde{\gamma}''(0) &= \gamma''(s) = T'(s) = \kappa \cdot N \\
 K''_x(\sigma) &= -(\kappa \cdot \sin \kappa \sigma) \cdot T + (\kappa \cdot \cos \kappa \sigma) N \\
 K''_x(0) &= \kappa \cdot N
 \end{aligned}$$

$\tilde{\gamma}$  and  $K_x$  correspond in the derivatives up to the second degree for  $\sigma = 0$ . Taylor expansion of 2nd degree for  $\tilde{\gamma} - K_x$ :

$$\|\tilde{\gamma}(\sigma) - K_x(\sigma)\| = o(\sigma^2)$$




Figure 60: Setting of  $B$ ,  $T$  and  $N$  with circle of curvature  $K_x$

Hence  $K_x$  approximates  $\tilde{\gamma}$  in  $x$  up second degree precisely. Compare with Figure 60.

We let  $B = T \times N$  be the normal vector to the osculating plane  $\epsilon_x$ .  $B = T \times N$  is called *binormal vector* of  $\gamma$  in  $x = \gamma(s)$ . Consider  $N^T(s) \cdot N(s) = 1$ . Derive it.

$$\implies N'^T \cdot N + N^T \cdot N' = 2 \langle N'(s), N(s) \rangle = 0$$

hence  $N'(s) \perp N(s)$  hence  $N' = \alpha T + \beta B$ .

$$\langle T, N \rangle = 0 \xrightarrow{\text{derive}} \langle T', N \rangle + \langle T, N' \rangle = 0$$

$T' = \kappa \cdot N$ .  $\langle \kappa \cdot N, N \rangle + \langle T, N' \rangle = 0$ .  $\langle T, N' \rangle = -\kappa$ . Derive  $B = T \times N$ .

$$B' = (T \times N)' = T' \times N + T \times N' = \kappa \underbrace{(N \times N)}_0 + T \times N' = T \times N'$$

Hence  $B' \perp T$  and  $B' \perp N'$ .  $\implies B' = \tilde{\alpha}N + \tilde{\beta}B$ .  $\langle T, N \rangle = 0 \implies \langle T', N \rangle + \langle T, N' \rangle = 0$ , so  $\langle N', T \rangle = -\langle T', N \rangle$ .

Because  $T$  and  $N$  are orthonormal,  $\alpha = \langle T, N' \rangle = -\kappa$  and  $\beta = \langle B, N' \rangle$ . So,

$$N' = -\kappa T + \langle B, N' \rangle \cdot B.$$

$$B' = \tilde{\alpha} \cdot N + \tilde{\beta} \cdot B$$

$$\tilde{\alpha} = \langle N, B' \rangle$$

$$\tilde{\beta} = \langle B, B' \rangle$$

because  $\langle B, B \rangle = 1 \implies \langle B, B' \rangle = 0$ . Hence

$$B' = \langle N, B' \rangle \cdot N$$

We let  $\tau = \tau(s) = \langle N, B' \rangle$ . So,  $B' = \tau \cdot N$ , which we call *third Frenet formula*.  $\langle B, N \rangle = 0 \implies \langle B, N' \rangle + \langle B', N \rangle = 0$ , hence  $\beta = \langle B, N' \rangle = -\langle B', N \rangle = -\tau$ . We call  $N' = -\kappa T - \tau B$  the *second Frenet formula*.

Summary: Let  $\gamma$  be parameterized by the arc length. Then (with  $\gamma \in \mathbb{C}^3$ )

$$\begin{aligned} T' &= \kappa N \\ N' &= -\kappa T - \tau B \\ B' &= \tau N \end{aligned}$$

are called Frenet's formulas.

$$\begin{bmatrix} T \\ N \\ B \end{bmatrix}' = \begin{bmatrix} 0 & \kappa I & 0 \\ -\kappa I & 0 & -\tau I \\ 0 & \tau I & 0 \end{bmatrix} \begin{bmatrix} T \\ N \\ B \end{bmatrix}$$

$[T(s), N(s), B(s)]$  define for every  $s$  an orthonormal basis in  $\mathbb{R}^3$ . The “accompanying triple” of the curve (dt. das begleitende Dreibein der Kurve).

$B' = \tau \cdot N$ . Compare with Figure 61.  $B'$  is the rate of change of the normal vector of the osculating plane  $\varepsilon_x$ .  $\tau$  is the oriented rate of change of the normal vector of the osculating plane.  $\tau(s)$  is called *torsion* of a curve in  $x = \gamma(s)$ .

**Example 7.7** (Determination of the curvature of an arbitrary parameterized curve  $\gamma(t)$ ). Let  $\tilde{\gamma}(s) = \gamma(t(s))$ .




Figure 61: Curve triple  $B, T, N$  and  $B'$

Prerequisite:  $\alpha : I \rightarrow \mathbb{R}^3$  is differentiable.  $\alpha(t) \neq 0 \forall t \in I$ .

$$\frac{d}{dt} [\|\alpha(t)\|] = \frac{d}{dt} \left[ \left( \sum_{i=1}^3 \alpha_i^2(t) \right)^{\frac{1}{2}} \right] = \frac{1}{2} \left( \sum_{i=1}^3 \alpha_i^2(t) \right)^{-\frac{1}{2}} \left( 2 \cdot \sum_{i=1}^3 \alpha_i \cdot \alpha'_i \right) = \frac{\langle \alpha(t), \alpha'(t) \rangle}{\|\alpha(t)\|}$$

$$s(t) = \int_{t_0}^t \|\gamma'(\tau)\| d\tau$$

$$s'(t) = \|\gamma'(t)\|$$

$$t'(s) = \frac{1}{\|\gamma'(t(s))\|}$$

$$\Gamma(s) = \frac{\gamma'(t(s))}{\|\gamma'(t(s))\|}$$

$$\begin{aligned} \frac{d}{ds} \Gamma(s) &= \frac{1}{\|\gamma'(t(s))\|} \cdot \gamma''(t(s)) \cdot \underbrace{t'(s)}_{= \frac{1}{\|\gamma'(t(s))\|}} \\ &= \frac{1}{\|\gamma'(t(s))\|^2} \cdot \frac{\langle \gamma'(t(s)), \gamma''(t(s)) \rangle}{\|\gamma'(t(s))\|} \cdot \underbrace{t'(s)}_{= \frac{1}{\|\gamma'(t(s))\|}} \cdot \gamma'(t(s)) \\ &= \frac{1}{\|\gamma'(t(s))\|^2} \cdot \gamma''(t(s)) - \frac{\langle \gamma'(t(s)), \gamma''(t(s)) \rangle}{\|\gamma'(t(s))\|^4} \cdot \gamma'(t(s)) \end{aligned}$$

Omit  $t(s)$ .

$$\begin{aligned}
 \kappa^2(s) &= \|T'(s)\|^2 = \left\langle \frac{1}{\|\gamma''\|^2} \gamma'' - \frac{\langle \gamma', \gamma'' \rangle}{\|\gamma'\|^4} \gamma', \frac{1}{\|\gamma''\|^2} \gamma'' - \frac{\langle \gamma', \gamma'' \rangle}{\|\gamma'\|^4} \gamma' \right\rangle \\
 &= \frac{\|\gamma''\|^2}{\|\gamma'\|^4} - 2 \cdot \frac{\langle \gamma', \gamma'' \rangle^2}{\|\gamma'\|^6} + \frac{\langle \gamma', \gamma'' \rangle^2 \cdot \|\gamma'\|^2}{\|\gamma'\|^8} \\
 &= \frac{1}{\|\gamma'\|^6} \left( \|\gamma''\|^2 \cdot \|\gamma'\|^2 - \langle \gamma', \gamma'' \rangle^2 \right)
 \end{aligned}$$

$$\Rightarrow \kappa(t) = \frac{1}{\|\gamma'\|^3} \left( \|\gamma''\|^2 \|\gamma'\|^2 - \langle \gamma', \gamma'' \rangle^2 \right)^{\frac{1}{2}}$$

Left as an exercise to the reader:

$$\begin{aligned}
 \|\gamma''\|^2 \cdot \|\gamma'\| - \langle \gamma', \gamma'' \rangle^2 &= \left( \|\gamma''\|^2 \cdot \|\gamma'\|^2 - \|\gamma''\| \cdot \|\gamma'\| \cdot \cos^2(\alpha) \right) \\
 &= \|\gamma''\|^2 \cdot \|\gamma'\|^2 \cdot \sin^2(\alpha) = \|\gamma' \times \gamma''\|^2
 \end{aligned}$$

$$\Rightarrow \kappa(t) = \frac{\|\gamma' \times \gamma''\|}{\|\gamma'\|^3}$$

↓ This lecture took place on 2018/06/28.


Let  $\gamma(t)$  be a helical curve (compare with Figure 62).

$$\gamma(t) = \begin{bmatrix} \cos t \\ \sin t \\ t \end{bmatrix}$$

The pitch (dt. Ganghöhe) is given by  $h$ :

$$\gamma'(t) = \begin{bmatrix} -\sin(t) \\ \cos(t) \\ h \end{bmatrix}$$

$$\|\gamma'(t)\| = \sqrt{1 + h^2} = A$$



**Example 7.8** (helical curve).

Figure 62: Helical curve with pitch  $h$

$$s(t) = \int_0^t \|\gamma'(\tau)\| d\tau = \sqrt{1+h^2} \cdot t = A \cdot t$$

$$t(s) = \frac{1}{A} \cdot s$$

*Reparameterization by arc length:*

$$\tilde{\gamma}(s) = \begin{bmatrix} \cos \frac{s}{a} \\ \sin \frac{s}{a} \\ h \cdot \frac{s}{a} \end{bmatrix}$$

$$T(s) = \tilde{\gamma}'(s) = \frac{1}{A} \cdot \begin{bmatrix} -\sin \frac{s}{a} \\ \cos \frac{s}{a} \\ h \end{bmatrix}$$

$$T'(s) = \underbrace{\frac{1}{A^2}}_{\|T'(s)\|=\kappa(s)} \underbrace{\begin{bmatrix} -\cos \frac{s}{a} \\ -\sin \frac{s}{a} \\ 0 \end{bmatrix}}_{N(s)}$$

$$\left\| -\cos \frac{s}{a} - \sin \frac{s}{a} 0 \right\| = 1$$

$$\Rightarrow \kappa = \frac{1}{1+h^2} \quad \dots \text{ constant}$$




Figure 63: Curve triple

$$\begin{aligned}
 B &= T \times N = \frac{1}{A} \cdot \begin{bmatrix} -\sin \frac{s}{a} \\ \cos \frac{s}{a} \\ h \end{bmatrix} \times \begin{bmatrix} -\cos \frac{s}{a} \\ -\sin \frac{s}{a} \\ 0 \end{bmatrix} \\
 &= \frac{1}{A} \begin{bmatrix} h \cdot \sin \frac{s}{a} \\ -h \cos \frac{s}{a} \\ 1 \end{bmatrix} \\
 B' &= \frac{h}{A^2} \begin{bmatrix} \cos \frac{s}{A} \\ \sin \frac{s}{A} \\ 0 \end{bmatrix} = -\frac{h}{A^2} \cdot N
 \end{aligned}$$

3rd Frenet's formula:  $B' = \tau \cdot N$ .

$$\Rightarrow \tau = -\frac{h}{A^2} = -\frac{h}{1+h^2}$$

Compare with Figure 63.

## 7.5 Curvatures of plane curves

Let  $\gamma : I \rightarrow \mathbb{R}^2$ .

$$\|\gamma'(s)\| = 1$$

Then

$$\gamma'(s) = \begin{bmatrix} \cos v(s) \\ \sin v(s) \end{bmatrix} \in S^1 = \{x \in \mathbb{R}^2 \mid \|x\|_2 = 1\}$$

$\vartheta(s)$  is the continuously differentiable angle parameter.

$$T(s) = \gamma'(s) = \begin{bmatrix} \cos \vartheta(s) \\ \sin \vartheta(s) \end{bmatrix}$$

$$T'(s) = \vartheta'(s) \begin{bmatrix} -\sin \vartheta(s) \\ \cos \vartheta(s) \end{bmatrix} := \vartheta'(s) \cdot N(s)$$

$$N(s) = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \cdot \begin{bmatrix} \cos \vartheta \\ \sin \vartheta \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \cdot T(s)$$

Compare with Figure 64. The curvature changes the sign on change of the traversing direction.

$$T' = \kappa \cdot N$$

$$N(s) = \begin{bmatrix} -\sin \vartheta(s) \\ \cos \vartheta(s) \end{bmatrix}$$

$$N'(s) = \vartheta'(s) \begin{bmatrix} -\cos \vartheta(s) \\ -\sin \vartheta(s) \end{bmatrix} = -\kappa T$$

2 Frenet formulas:

$$T' = \kappa N \quad N' = -\kappa T$$



Determination of  $\kappa$  in case of arbitrary parameterization:

$$\gamma(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

$$T(t) = \frac{1}{\sqrt{x'^2 + y'^2}} \begin{bmatrix} x' \\ y' \end{bmatrix}$$

$s(t)$  ... arc position with

$t(s)$  ... change of parameter to arc length

$$t'(s) = \frac{dt}{ds} = \frac{1}{(x'^2 + y'^2)^{\frac{1}{2}}}$$

$$\frac{d}{ds} [T(t(s))] = \underbrace{\frac{d}{dt} T(t)}_{T'(t)} \cdot \frac{dt}{ds}$$

$$\begin{aligned} T'(t) &= \frac{1}{(x'^2 + y'^2)^{\frac{1}{2}}} \begin{bmatrix} x'' \\ y'' \end{bmatrix} - \frac{1}{(x'^2 + y'^2)^{\frac{3}{2}}} (x'x'' + y'y'') \begin{bmatrix} x' \\ y' \end{bmatrix} \\ &= \frac{y''x' - x''y'}{(x'^2 + y'^2)^{\frac{3}{2}}} \cdot \begin{bmatrix} -y' \\ x' \end{bmatrix} \end{aligned}$$

$$\frac{d}{ds} [T(s)] = \frac{y'' \cdot x' - x'' \cdot y'}{(x'^2 + y'^2)^{\frac{3}{2}}} \cdot \underbrace{\frac{1}{(x'^2 + y'^2)} \cdot \begin{bmatrix} -y' \\ x' \end{bmatrix}}_N$$

$$T' = \underbrace{\frac{y''x' - x''y'}{(x'^2 + y'^2)^{\frac{3}{2}}}}_{=\kappa} \cdot N$$

$$\kappa = \frac{y''x' - x''y'}{(x'^2 + y'^2)^{\frac{3}{2}}}$$

Let  $\gamma : I \rightarrow \mathbb{R}^2$  and  $\gamma$  is parameterized by the arc length.  $\mu(s)$  is center of the circle of curvature of  $\gamma$  by change of parameter  $s$ .

$s \mapsto \mu(s)$  is called *evolute* of  $\gamma$ , a curve in the plane.

$$\mu(s) = \gamma(s) + \frac{1}{\kappa(s)} \cdot N(s)$$

$$\mu'(s) = \gamma'(s) + \frac{1}{\kappa(s)} \cdot N'(s) - \frac{\kappa'(s)}{\kappa(s)^2} \cdot N(s) = T(s) + \frac{1}{\kappa} (-\kappa T) - \frac{\kappa'}{\kappa^2} N = -\frac{\kappa'}{\kappa^2} \cdot N$$

Hence, if  $\kappa \neq 0$ , then  $N(s)$  is a tangential vector of  $\mu$  in  $\mu(s)$ . The normal lines at the curve corresponds to the tangent at the evolute.




Figure 64: We call  $\kappa(s) = \vartheta'(s)$ , thus  $T' = \kappa \cdot N$ , the *signed curvature*. It can have positive or negative sign.




Figure 65: Evolute




Figure 66: The family of normal lines yield a curve  $\mu(t)$

The normal lines to  $\gamma$  give a family of lines such that every line is a tangent at  $\mu$ .  $\mu$  is the *envelope of the family of curves of  $\gamma$*  (dt. Einhüllende der Schar der Normalen an  $\gamma$ ).

**Example 7.9.**

$$\begin{aligned}
 \gamma(t) &= \begin{bmatrix} t \\ t^2 \end{bmatrix} \quad \dots \text{parabola} \\
 \gamma'(t) &= \begin{bmatrix} 1 \\ 2t \end{bmatrix} \\
 T(t) &= \frac{1}{\sqrt{1+4t^2}} \begin{bmatrix} 1 \\ 2t \end{bmatrix} \quad N(t) = \frac{1}{\sqrt{1+4t^2}} \begin{bmatrix} -2t \\ 1 \end{bmatrix} \\
 \kappa(t) &= \frac{2}{(1+4t^2)^{\frac{3}{2}}} \\
 \mu(t) &= \gamma(t) + \frac{1}{\kappa(t)} \cdot N(t) = \begin{bmatrix} t \\ t^2 \end{bmatrix} + \frac{(1+4t^2)^{\frac{3}{2}}}{2} \cdot \frac{1}{(1+4t^2)^{\frac{1}{2}}} \begin{bmatrix} -2t \\ 1 \end{bmatrix} = \begin{bmatrix} -4t^3 \\ \frac{1}{2} + 3t^2 \end{bmatrix} = \underbrace{\frac{1}{2} \begin{bmatrix} 0 \\ 1 \end{bmatrix} + \begin{bmatrix} -4t^3 \\ 3t^2 \end{bmatrix}}_{\text{Neile's parabola}} \\
 \mu'(t) &= 0 \text{ for } t = 0 \quad (\kappa' = 0 \text{ hence } \kappa \text{ has maximum})
 \end{aligned}$$

## Study hints by students

### Exam questions 2018/07/06

1. Let  $f : X \rightarrow Y$  be continuous. Let  $X$  be compact. Prove that  $f(X)$  is compact. Give any corollary of this proof.

2. Give and prove Hölder's inequality.
3. Prove: If  $f(x) = \sum_{k=0}^{\infty} a_k \cdot (x - x_0)^k$  with  $\rho_f > 0$  is analytical. Show that,
  - (a)  $a_k = f^{(k)}(x_0) \cdot \frac{1}{k!}$
  - (b)  $f$  is arbitrary often continuously differentiable in  $(x_0 - \rho_f, x_0 + \rho_f)$
4.  $f : D \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^n$  is differentiable in  $x_0$  with  $Df(x_0)$ . Show that  $\forall v \in \mathbb{R}^m \setminus \{0\} : df(x_0, v) = Df(x_0) \cdot v$
5. Let  $\gamma : I \rightarrow \mathbb{R}^3$  be regular, smooth and  $\|\gamma'(t)\| = 1 \ \forall t \in I$ . Define its curvature, circle of curvature, osculating plane and principal normal vector. Show  $\langle T, N \rangle = 0$ . Give the approximating behavior of curvature and the circle of curvature.

### Example questions 2018/07/31

1. Give and prove Banach Fixed Point Theorem. Give another important theorem, which can be proved using Banach Fixed Point Theorem.
2. Prove that a regulated function  $f \in \mathcal{R}[a, b]$  is non-continuous at at most countable many points.
3. Let  $\gamma : I \rightarrow \mathbb{R}^n$  be a rectifiable curve with arc length  $S_\gamma$ . Assume  $\gamma$  is continuously differentiable and let  $I = [a, b]$ . Prove that  $S_\gamma = \int_a^b \|\gamma'(t)\|_2 \, dt$ .
4. Let  $P(z) = \sum_{k=0}^{\infty} a_k z^k$  be a power series with convergence radius  $\rho_P > 0$ . Furthermore let  $0 < r < \rho_P$ . Prove that the partial sums  $P_n(z) = \sum_{k=0}^n a_k z^k$  of the series on  $\overline{K_r(0)}$  converge *uniformly* to  $P(z)$ . If you use an already-proven theorem, name it.
5. Let  $X$  and  $Y$  be normed vector spaces and let  $A : X \rightarrow Y$  be bounded and linear. Give the definition of *boundedness* and the definition of the operator norm of  $A$ . Let  $Z$  be also a normed vector space and let  $B : Y \rightarrow Z$  be also linear and bounded. Prove that

$$\|BA\| \leq \|B\| \|A\|$$

holds for operator norms.

### List of 12 major theorems

1. Bolzano-Weierstrass Theorem in  $\mathbb{R}^n$
2. Norm equivalence in  $\mathbb{R}^n$
3. Banach Fixed Point Theorem

4. Mean Value Theorem of Integration Calculus
5. Fundamental Theorem of differential/integration calculus, variant 1+2
6. Young's inequality
7. Hölder's inequality
8. Multidimensional Taylor Theorem
9. Chain rule in multidimensional calculus
10. Local Inversion Theorem
11. Implicit Function Theorem
12. Schwarz' Theorem

### **Sample exam by a student**

1. Give and prove the Implicit Function Theorem
2. Give at least two equivalent statements about regulated functions. Prove at least one of them.
3. Prove the Cauchy criterion for limits of functions
4. Let  $A \subseteq X$  be closed. Show that equivalence with the following statement:  $A^c = O \subseteq X$  open in  $X$ . Which corollary can be made regarding intersection and union of open sets?

### **Another sample exam by a student**

1. Prove that all norms are equivalent in  $\mathbb{R}^n$  and show that norm equivalence is an equivalence relation.
2. Prove the approximation theorem for regulated functions.
3. Prove: Let  $f$  be a regulated function on  $[a, b]$ . Then there exists an at most countable set  $A \subseteq [a, b]$  such that  $f$  is continuous in every point  $x \in [a, b] \setminus A$ .
4. Prove the fundamental theorem of differential/integration calculus, variant 1.
5. Prove the multidimensional Chain Rule.

## Index

- $L^p$ -norm, 75
- $C^\infty$ -Homeomorphism, 137
- $k$ -times continuously differentiable, 123
- Accumulation point, 7, 8
- Antiderivative, 54
- Approximation error of a Taylor polynomial, 84
- Arc length of a curve, 142
- Arc length parameter, 147
- Binormal vector, 154
- Boundedness, 19
- Cauchy sequence, 6
- Cauchy-Schwarz inequality, 76
- Characteristic function, 33
- Circle of curvature, 151
- Clairaut's theorem, 123
- Closed curve, 138
- Compactness, 18
- Complete metric space, 7
- Conjugate exponents, 74
- Contact point, 9
- Continuity on a set, 14
- Contraction, 15
- Convergence, 6
- Curvature, 135
- Curvature of a curve in a point, 150
- Diffeomorphism, 114
- Differentiable on  $D$ , 103
- Directional derivative, 103
- Distance function, 3
- Envelope, 161
- Euclidean norm, 5, 7
- Euler's Gamma function, 69
- Evolute, 161
- First Frenet formula, 151
- Fixed point, 15
- Fréchet derivative, 98
- Frobenius norm, 92
- Gamma function, 69
- Geometrical quantity, 138
- Gradient of a function, 108
- Hölder's inequality, 75
- Half-sided derivative, 52
- Hessian matrix, 123
- Homeomorphism, 137
- Implicit function, 118
- Indefinite integral, 58
- Indicator function, 33
- Inner point, 11
- Integral form of the Taylor remainder, 84
- Jacobi matrix, 105
- Lagrange form of the remainder, 85
- Left-sided derivative, 52
- Limit point, 35
- Lipschitz continuity, 15
- Local Inversion Theorem, 114
- Manhattan metric, 8
- Matrix norm, 92
- Metric, 3
- Metric space, 3
- Multiindex notation, 126
- Nabla operator, 108
- Norm, 5, 7
- Normed vector space, 5
- Open sphere, 6
- Operator norm, 93
- Order of a partial derivative, 122
- Orientation preserving reparameterization, 137
- Orientation reversing reparameterization, 137
- Osculating plane, 151

- Parameter exchange, 137
- Parameterization by a curve's arc length, 150
- Parameterized curve, 135
- Partial derivative, 105
- Partial integration, 59
- Partition of an interval, 28
  
- Qualitative Taylor equation, 88
  
- Rectifiable curve, 142
- Regular curve parameterization, 135
- Regulated function, 38
- Relative topology, 23
- Remainder of a Taylor polynomial, 84
- Reparameterization, Homeomorphism and orientation, 136
- Right-sided derivative, 52
  
- Schwarz' theorem, 123
- Second Frenet formula, 155
- signed curvature, 162
- Step function, 28
- Subspace topology, 23
- Symmetry of second derivatives, 123
  
- Tangent of a curve, 135
- Taylor polynomial of  $f$  of order  $n$  in  $x_0$ , 82
- Taylor series, 89
- Taylor's theorem in the multidimensional case, 126
- Third Frenet formula, 155
- Topological space, 13
- Topology, 13
- Torsion of a curve, 155
- Total boundedness, 20
- Total derivative, 108
- Totally differentiable, 108
- Trace topology, 23
  
- Uniformly continuous, 15
- Unit tangent vector, 135
  
- Young's inequality, 71
- Young's theorem, 123