

Linear Algebra 2

Lecture notes, University (of Technology) Graz
based on the lecture by Franz Lehner

Lukas Prokop

September 19, 2018

Contents

5.1	Lecture	3
6	Linear algebra 1	3
7	Determinants	4
7.1	Definition	4
7.2	Properties	4
7.3	Determinant form	9
7.4	Permutations and transpositions	10
7.5	Leibniz formula for determinants	16
7.6	On determinants, invertibility and linear independence	22
7.7	Laplace expansion	33
8	Inner products	38
8.1	Law of cosines	41
8.2	Outer product	42
8.3	Inner products and positive definiteness	45
8.4	Cauchy-Bunyakovsky-Schwarz inequality	49
8.5	Congruence of matrices	53
8.6	Gram-Schmidt process	85
8.7	Riesz representation theorem	88
8.8	Adjoint maps	91

8.9	The linear adjoint map is the complex transpose	93
8.10	Unitary transformations and self-adjoint matrices	95
8.11	Unitary matrices and orthogonal matrices	96
8.12	Quaternions	103
9	Polynomials and algebras	104
9.1	The greatest common divisor of polynomials	115
10	Eigenvectors and eigenvalues	115
10.1	Eigenspace	116
10.2	Characteristic polynomial	120
10.3	Symmetrical minor	120
10.4	Diagonalizable matrix	123
10.5	Fibonacci sequence and golden ratio	126
10.6	Multiplicities of eigenvalues	129
11	Jordan Normal Form (JNF)	132
11.1	Invariant subspaces	132
11.2	Fitting lemma	134
11.3	Main space	138
11.4	Nilpotent matrix	145
11.5	Jordan's normal form	146
11.6	Jordan block	152
11.7	Matrix exponentiation	157
11.8	Minimal polynomial and annihilator	160
11.9	Cayley-Hamilton Theorem	161
11.10	Spectrum mapping theorem	165
12	Normal matrices	166
12.1	Schur's decomposition, QR decomposition	169
12.2	Application: Conic section	173
12.3	Cholesky decomposition	177
12.4	Singular value decomposition	183
13	Eigenvalue estimates	184

13.1 Geršgorin theorem	188
14 Matrix norms	191
15 Non-negative matrices	203

↓ *This lecture took place on 2018/03/05.*

5.1 Lecture

- Mon, 08:15–09:45, lecture
- Wed, 08:15–09:45, lecture
- Mon, 16:00–18:00, tutorial, AE01
- Mon, 13:15–14:00, conversatorium (BE01)

6 Linear algebra 1

Gottfried Wilhelm von Leibniz (1646–1716). Results from 1693:

- Vector spaces (first definition in 1880)
- Matrices and linear maps

From now, it will be more specific (matrices). In general, we discuss “when is a matrix invertible”?

$$\begin{aligned} ax + by &= e \\ cx + dy &= f \end{aligned}$$

We need to invert the matrix

Assuming $a \neq 0$. We multiply the first row with $\frac{1}{a} \cdot (-c)$.

$$\begin{array}{cc|cc} a & b & 1 & 0 \\ c & d & 0 & 1 \\ \hline 0 & d - \frac{c}{a} \cdot b & -\frac{c}{a} & 1 \end{array}$$

We then divide by $d - \frac{c}{a}b$ if $\neq 0$.

If $a = 0$ and $c = 0$, rank is certainly not 2.

If $a = 0$ and $c \neq 0$, we multiply with $\frac{1}{c}(-a)$.

$$\begin{array}{cc} a & b \\ c & d \\ \hline 0 & b - \frac{ad}{c} \end{array}$$

we divide $b - \frac{ad}{c}$ if $\neq 0$.

When does such a system have a non-trivial solution? There is a non-trivial solution iff $ad - bc \neq 0$.

$ad - bc \neq 0$ iff $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is invertible.

Leibniz was not the first discovering it. The result was found before 1685 by Seki Takahazu.

7 Determinants

7.1 Definition

$$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} =: ad - bc =: \begin{vmatrix} a & b \\ c & d \end{vmatrix}$$

is called *determinant of matrix* $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$.

7.2 Properties

- The determinant is linear in every row and every column. For fixed b and d , it is

$$\begin{pmatrix} x \\ y \end{pmatrix} \mapsto \det \begin{pmatrix} x & b \\ y & d \end{pmatrix} = dx - by \quad \text{is linear} \\ \mathbb{K}^2 \rightarrow \mathbb{K}$$

$$\begin{aligned} \det \begin{pmatrix} \lambda x + \mu x' & b \\ \lambda y + \mu y' & d \end{pmatrix} &= d \cdot (\lambda x + \mu x') - b \cdot (\lambda y + \mu y') \\ &= \lambda(dx - by) + \mu(dx' - by') \\ &= \lambda \det \begin{pmatrix} x & b \\ y & d \end{pmatrix} + \mu \det \begin{pmatrix} x' & b \\ y' & d \end{pmatrix} \end{aligned}$$

The determinant is bilinear in rows and columns.

$$\det(\lambda v + \mu v', w) = \lambda \det(v, w) + \mu \det(v', w)$$

Let $v = \begin{pmatrix} a \\ c \end{pmatrix}$.

$$\det(v, \lambda w + \mu w') = \lambda \det(v, w) + \mu \det(v, w')$$

Let $w = \begin{pmatrix} b \\ d \end{pmatrix}$. Follows analogously.

- If two rows are the same, then $\det(M) = 0$.

$$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ab - ba = 0$$

$$\det \begin{pmatrix} a & a \\ c & c \end{pmatrix} = ac - ca = 0$$

- The determinant of the unit matrix is one.

$$\det \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = 1$$

Theorem 7.1 (properties 1–3 characterize the determinant). *If $\varphi : \mathbb{K}^2 \times \mathbb{K}^2 \rightarrow \mathbb{K}$ and φ satisfies the properties 1–3, φ is the determinant.*

bilinear¹

$$\varphi(\lambda v + \mu v', w) = \lambda \varphi(v, w) + \mu \varphi(v', w)$$

$$\forall v, w, v', w' : \mu(v, \lambda w + \mu w') = \lambda \varphi(v, w) + \mu \varphi(v, w')$$

$$\forall v : \varphi(v, v) = 0$$

$$\implies \varphi = \det$$

$$\varphi(e_1, e_2) = 1$$

Proof.

$$v = \begin{pmatrix} a \\ c \end{pmatrix} = a \cdot e_1 + c \cdot e_2$$

$$w = \begin{pmatrix} d \\ b \end{pmatrix} = b \cdot e_1 + d \cdot e_2$$

$$\begin{aligned} \varphi(v, w) &= \varphi(a \cdot e_1 + c \cdot e_2, b \cdot e_1 + d \cdot e_2) \\ &= a \cdot \varphi(e_1, b \cdot e_1 + d \cdot e_2) + c \cdot \varphi(e_2, b \cdot e_1 + d \cdot e_2) \\ &= ab \cdot \underbrace{\varphi(e_1, e_1)}_{=0} + ad \cdot \varphi(e_1, e_2) + cb \cdot \varphi(e_2, e_1) + cd \cdot \underbrace{\varphi(e_2, e_2)}_{=0} \end{aligned}$$




Figure 1: Geometric interpretation of determinants

Is zero, because of property 3.

$$\begin{aligned}
 &= ad \cdot \underbrace{\varphi(e_1, e_2)}_{=1} + cb \cdot \varphi(e_2, e_1) \\
 0 &= \varphi(e_1 + e_2, e_1 + e_2) = \underbrace{\varphi(e_1, e_1)}_{=0} + \underbrace{\varphi(e_1, e_2)}_{=1} + \varphi(e_2, e_1) + \underbrace{\varphi(e_2, e_2)}_{=0} \\
 &\implies \varphi(e_2, e_2) = -1
 \end{aligned}$$

□

Corollary 7.1.

$$\forall v, w : \varphi(v, w) = -\varphi(w, v)$$

Corollary 7.2 (Geometrical interpretation). *See Figure 1. The determinant $\det(v, w)$ is the area of the spanned parallelogram. We denote F as the function returning the area of a geometric object.*

Proof. $\text{area}(v, w)$ satisfies properties (i) – (iii).

Consider orthogonal e_1 and e_2 . $F = 1 = \det(e_1, e_2)$. $\det(e_2, e_1) = -1$.

The sign indicates the orientation of the area.

□

By property 2, if $v = w$, then $F = 0$. By property 1,

1. If v and w are *linear dependent*², then

$$\lambda v + \mu w = 0 \quad (\lambda, \mu) \neq (0, 0)$$

Without loss of generality, $\mu \neq 0 \implies w = -\frac{\lambda}{\mu} \cdot v$.

2. To show:

$$F(\lambda v, w) = \lambda \cdot F(v, w)$$

$$F(v + v', w) = F(v, w) + F(v', w)$$

Let $\lambda \in \mathbb{N}$. We multiply the area n times.


$$F(n \cdot v, w) = n \cdot F(v, w)$$

- 3.

$$F\left(\frac{1}{n} \cdot v, w\right) = \frac{1}{n} F(v, w)$$

follows from $F(\lambda v, w) = \lambda \cdot F(v, w)$, because $v = n \cdot (\frac{1}{n}v)$:

$$F\left(n\left(\frac{1}{n}v\right), w\right) = n \cdot F\left(\frac{1}{n}v, w\right)$$



- 4.

Figure 2: The sign changes if the orientation changes

If we combine (2) and (3),

$$F\left(\frac{m}{n}v, w\right) = \frac{m}{n} F(v, w)$$

See Figure 2.

²Hence, one vector is a multiple of the other

5. By continuity, $F(\lambda v, w) = \lambda F(v, w) \forall \lambda \in \mathbb{R}_+^3$. If the orientation changes, the sign changes. By this property, this actually holds for \mathbb{R} , not only \mathbb{R}_+ .

Analogously:

$$F(v, \lambda w) = \lambda F(v, w) \forall \lambda \in \mathbb{R} \forall v, w \in \mathbb{R}^2$$

6. To show: $F(v + v', w) = F(v, w) + F(v', w)$

If v and w are linear independent, then $F(v + w, w) = F(v, w)$. In general, for a parallelogram of height h and vector w , it holds that

$$F = |w| \cdot h$$

The height of the parallelogram stays the same.

$$F(v, w) = F(v + w, w)$$

- 7.

$$F(\lambda v + \mu w, w) = \lambda F(v, w)$$

Case $\mu = 0$ Already shown, $F(\lambda v, w) = \lambda F(v, w) \forall \lambda \in \mathbb{R}$.

Case $\mu \neq 0$ $F(\lambda v + \mu w, w) = \frac{1}{\mu} F(\lambda v + \mu w, \mu w) = \frac{1}{\mu} F(\lambda v, \mu w) = F(\lambda v, w) = \lambda F(v, w)$

8. Let v and w be linear independent, then they define a basis of \mathbb{R}^2 .

$$v_1 = \lambda_1 v + \mu_1 w$$

$$v_2 = \lambda_2 v + \mu_2 w$$

$$\begin{aligned} \rightarrow F(v_1 + v_2, w) &= F(\lambda_1 v + \mu_1 w + \lambda_2 v + \mu_2 w, w) \\ &= F((\lambda_1 + \lambda_2)v + (\mu_1 + \mu_2)w, w) \\ &= F((\lambda_1 + \lambda_2)v, w) \\ &= (\lambda_1 + \lambda_2)F(v, w) \\ &= \lambda_1 F(v, w) + \lambda_2 F(v, w) \\ &= F(\lambda_1 v, w) + F(\lambda_2 v, w) \\ &= F(\lambda_1 v + \mu_1 w, w) + F(\lambda_2 v + \mu_2 w, w) \\ &= F(v_1, w) + F(v_2, w) \end{aligned}$$

This shows that additivity is given.

³By the way, how are real numbers defined?

7.3 Determinant form

Definition 7.1. Let V be an n -dimensional vector space over \mathbb{K} . A determinant form is a map

$$\begin{aligned}\Delta : V^n &\rightarrow \mathbb{K} \\ (a_1, \dots, a_n) &\mapsto \Delta(a_1, \dots, a_n)\end{aligned}$$

Let $n = 2$.

$$\Delta : \left(\begin{pmatrix} a \\ c \end{pmatrix}, \begin{pmatrix} b \\ d \end{pmatrix} \right) \mapsto \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

It satisfies the properties of *multilinearity*:

1. $\Delta(a_1, \dots, \lambda a_k, \dots, a_n) = \lambda \Delta(a_1, \dots, a_n)$
2. $\Delta(a_1, \dots, a_k + v, \dots, a_n) = \Delta(a_1, \dots, a_k, \dots, a_n) + \Delta(a_1, \dots, a_{k-1}, v, a_{k+1}, \dots, a_n)$

Multilinearity is given, if linearity is given in every component. Hence, if $a_1, \dots, a_{k-1}, a_{k+1}, \dots, a_n$ are fixed, then

$$\begin{aligned}V &\rightarrow \mathbb{K} \\ v &\mapsto \Delta(a_1, \dots, a_{k-1}, v, a_{k+1}, \dots, a_n) \text{ linear}\end{aligned}$$

Furthermore, it satisfies the following property:

3. $\Delta(a_1, \dots, a_n) = 0$
if $\exists k \neq l : a_k = a_l$. If $\Delta \neq 0$, then Δ is called *non-trivial*.

Corollary 7.3. 4. $\Delta(a_1, \dots, a_k + \lambda a_i, \dots, a_n) = \Delta(a_1, \dots, a_k, \dots, a_n) \forall \lambda \in \mathbb{K}, \forall i \neq k$

5. $\Delta(a_1, \dots, a_i, \dots, a_j, \dots, a_n) = -\Delta(a_1, \dots, a_j, \dots, a_i, \dots, a_n)$

Proof. 1.

$$\begin{aligned}\Delta(a_1, \dots, a_k + \lambda a_i, \dots, a_n) &= \Delta(a_1, \dots, a_k, \dots, a_n) + \Delta(a_1, \dots, a_{k-1}, \lambda a_i, a_{k+1}, \dots, a_n) \\ &= \Delta(a_1, \dots, a_n) + \lambda \Delta(a_1, \dots, a_{k-1}, a_i, a_{k+1}, \dots, a_n) \\ &= 0 \quad \text{because } a_i \text{ occurs twice}\end{aligned}$$

2.

$$\begin{aligned}0 &= \Delta(a_1, \dots, a_i + a_j, \dots, a_i + a_j, \dots, a_n) \\ &= \Delta(a_1, \dots, a_i, \dots, a_i, \dots, a_n) \\ &\quad + \Delta(a_1, \dots, a_i, \dots, a_j, \dots, a_n) \\ &\quad + \Delta(a_1, \dots, a_j, \dots, a_i, \dots, a_n) \\ &\quad + \Delta(a_1, \dots, a_j, \dots, a_j, \dots, a_n)\end{aligned}$$

The first and last term are zero. Multilinearity is given:

$$\lambda(a_1, \dots, \lambda a_k, \dots, a_n) = \lambda \Delta(a_1, \dots, a_n)$$

$$\lambda(a_1, \dots, \lambda a_k + v, \dots, a_n) = \lambda \Delta(a_1, \dots, a_n) + \Delta(a_1, \dots, a_{k-1}, v, a_{k+1}, \dots, a_n)$$

□

↓ This lecture took place on 2018/03/07.

Determinant form: $\dim V = n$

$$\Delta : V^n \rightarrow \mathbb{K}$$

1. $\Delta(a_1, \dots, a_{k-1}, \lambda a_k, a_{k+1}, \dots, a_n) = \lambda \Delta(a_1, \dots, a_n)$
2. $\Delta(a_1, \dots, a_{k-1}, a_k + v, a_{k+1}, \dots, a_n) = \Delta(a_1, \dots, a_k, \dots, a_n) + \Delta(a_1, \dots, v, \dots, a_n)$
3. $\Delta(a_1, \dots, a_n) = 0$ if $\exists i \neq j : a_i = a_j$

Multilinearity is given by the first two properties.

$\Delta \neq 0$

Then the fourth property follows:

4. $\Delta(a_1, \dots, a_k + \lambda a_i, \dots, a_n) = \Delta(a_1, \dots, a_n) \forall i \neq k \forall \lambda \in \mathbb{K}$
5. $\Delta(a_1, \dots, a_i, \dots, a_j, \dots, a_n) = -\Delta(a_1, \dots, a_j, \dots, a_i, \dots, a_n)$

Example 7.1. Let $n = 2$, $V = \mathbb{K}^2$.

$$\Delta\left(\begin{pmatrix} a \\ c \end{pmatrix}, \begin{pmatrix} b \\ d \end{pmatrix}\right) = ad - bc = \det\begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

7.4 Permutations and transpositions

Definition 7.2. A permutation is a bijective map $\sigma : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$. σ_n is the set of all permutations.

$$|\sigma_n| = n!$$

Remark 7.1. σ_n in regards of composition defines a group with neutral element id and is called symmetric group.

Remark 7.2. For $n \geq 3$, it is non-commutative.

Example 7.2. *Permutations:*

$$\begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 1 & 3 & 2 \end{pmatrix} \circ \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 3 & 4 & 2 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 3 & 2 & 1 \end{pmatrix}$$

So, e.g. 2 is mapped to 3 (right side of \circ) and 3 is mapped to 3 (left side of \circ). Hence 2 is mapped to 3 (right-hand side of $=$).

$$\begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 1 & 3 & 2 \end{pmatrix}^{-1} = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 4 & 3 & 1 \end{pmatrix}$$

Definition 7.3. A transposition is a permutation exchanging exactly 2 elements.

$$\tau_{ij} : \begin{cases} i \mapsto j \\ j \mapsto i \\ k \mapsto k \forall k \notin \{i, j\} \end{cases}$$

$$\tau_{ij}^{-1} = \tau_{ij}$$

Remark 7.3. Every permutation $\sigma \in \sigma_n$ with $\sigma \neq \text{id}$ can be denoted as product of transpositions.

Proof.

$$\sigma = \begin{pmatrix} 1 & 2 & \dots & n \\ \sigma(1) & \sigma(2) & \dots & \sigma(n) \end{pmatrix}$$

Example:

$$\sigma = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 3 & 5 & 4 & 7 & 6 & 2 \end{pmatrix}$$

□

Find transpositions τ_1, \dots, τ_k such that $\sigma = \tau_1 \circ \tau_2 \circ \dots \circ \tau_k$.

If $\sigma = \text{id}$, then $k = 0$.

If $\sigma \neq \text{id}$,

$$k_1 := \min \{i \mid \sigma(i) \neq i\} \neq \emptyset$$

$$\tau_1 := \tau_{k_1 \sigma(k_1)}$$

$$\sigma_1 := \tau_1 \circ \sigma$$

if $\sigma_i = \text{id}$, then $\tau_1 \circ \sigma = \text{id}$. Then $\sigma = \tau_1^{-1} = \tau_1$.

$$k_2 := \min \{i \mid \sigma_1(i) \neq i\}$$

$$\tau_2 := \tau_{k_2 \sigma_1(k_2)}$$

$$\sigma_2 := \tau_2 \circ \sigma_1$$

Example 7.3. Let $k_1 = 2$.

$$\tau_1 = \tau_{23}$$

$$\begin{aligned}\sigma_1 &= \tau_{23} \circ \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 3 & 5 & 4 & 7 & 6 & 2 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 2 & 5 & 4 & 7 & 6 & 3 \end{pmatrix}\end{aligned}$$

$k_2 = 3$.

$$\tau_2 = \tau_{35}$$

$$\sigma_2 = \tau_2 \circ \sigma_1 = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 2 & 3 & 4 & 7 & 6 & 5 \end{pmatrix}$$

$k_3 = 5$.

$$T_3 = T_{57}$$

$$\begin{aligned}\sigma_3 &= \tau_3 \circ \sigma_2 = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 2 & 3 & 4 & 5 & 6 & 7 \end{pmatrix} \\ &= \text{id}\end{aligned}$$

$$\tau_3 \circ \tau_2 \circ \tau_1 \circ \sigma = \text{id}$$

$$\implies \tau_2 \circ \tau_1 \circ \sigma = T_3^{-1} \circ \text{id} = \tau_3$$

$$\tau_1 \circ \sigma = \tau_2^{-1} \circ T_3 = \tau_2 \circ \tau_3$$

$$\sigma = \tau_1 \circ \tau_2 \circ \tau_3$$

and so on and so forth.

$$\tau_k$$

$$\sigma_k = \tau_k \circ \tau_{k-1} \circ \cdots \circ \tau_i \circ \sigma = \text{id}$$

$$\implies \sigma = \tau_1 \circ \tau_2 \circ \cdots \circ \tau_k$$

Remark 7.4. This decomposition is not unique.

Definition 7.4. Let $\pi \in \sigma_n$ be a permutation. A malposition (dt. Fehlstand) of π is a pair (i, j) such that $i < j$ and $\pi(i) > \pi(j)$.

$$f_\pi := \left| \left\{ (i, j) \mid (i, j) \text{ is malposition of } \pi \right\} \right|$$

$$\text{sign}(\pi) := (-1)^{f_\pi} =: (-1)^\pi$$

is called signature of π

Example 7.4.

$$\pi = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 3 & 5 & 4 & 7 & 6 & 2 \end{pmatrix}$$

Malpositions:

$$\{(2, 7), (3, 4), (3, 7), (5, 6), (5, 7), (4, 7), (6, 7)\}$$

$$2 < 7$$

$$\pi(2) - 3 > 2 = \pi(7)$$

$$f_\pi = 7$$

Theorem 7.2.

$$\text{sign}(\pi) = \prod_{\substack{i, j \\ i < j}} \frac{\pi(j) - \pi(i)}{j - i}$$

1. $\binom{n}{2}$ factors

2. for transposition, $\text{sign } \tau = -1$.

Proof.

$$\prod_{i < j} \frac{\pi(j) - \pi(i)}{j - i} = \frac{\prod_{i < j} (\pi(j) - \pi(i))}{\prod_{i < j} (j - i)}$$

π is bijective in $\{1, \dots, n\}$. Hence, every difference (the expression $j - i$) occurs exactly one time in the numerator and the denominator with sign ± 1 depending on whether (i, j) is a malposition or not (statement 1). In other words, the term in the numerator and denominator cancel out if they are not malpositions.

$$\text{sign}(\pi(j) - \pi(i)) = \begin{cases} +1 & \pi(j) > \pi(i) \\ -1 & \pi(j) < \pi(i) \text{ hence malposition} \end{cases}$$

Consider any transposition, let k be the first index with $k \neq \pi(k)$ and let l be the last index with $l \neq \pi(l)$. $(k, k+1), (k, k+2), \dots, (k, l-1)$ are $(l-1-k+1+1)$ malpositions. $(k+1, l), (k+2, l), \dots, (l-1, l)$ are $(l-1-k+1+1)$ malpositions. The sum gives an even number. Additional, we have malposition (k, l) , thus an odd number of malpositions is given. Thus $\text{sign } \tau = -1$ (statement 2). \square

Example 7.5.

$$\pi = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 3 & 5 & 4 & 7 & 6 & 2 \end{pmatrix}$$

Malposition:

$$\{(2, 7), (3, 4), (3, 7), (5, 6), (5, 7), (4, 7), (6, 7)\}$$

$$2 < 7$$

$$\pi(2) - 3 > 2 = \pi(7)$$

$$f_\pi = 7$$

$$\frac{\prod_{i < j} (\pi(j) - \pi(i))}{\prod_{i < j} (j - i)} = \frac{\prod_{i < j} (j - i) \cdot (-1)^{f_\pi}}{\prod_{i < j} (j - i)} = \text{sign } \pi$$

$$\pi = \begin{pmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$

$$\begin{aligned} \prod_{i < j} \frac{\pi(j) - \pi(i)}{j - i} &= \frac{\pi(2) - \pi(1)}{2 - 1} \cdot \frac{\pi(3) - \pi(1)}{3 - 1} \cdot \frac{\pi(3) - \pi(2)}{3 - 2} \\ &= \frac{(2 - 3) \cdot (1 - 3) \cdot (1 - 2)}{(2 - 1)(3 - 1)(3 - 2)} \\ &= (-1)^3 = -1 \end{aligned}$$

Malpositions:

1. (1, 2)
2. (1, 3)
3. (2, 3)

Transposition: Let $k < \tau(k)$.

$$\tau = \begin{pmatrix} 1 & 2 & \dots & k-1 & k & k+1 & \dots & \tau(k) & \tau(k+1) & \dots & n \\ 1 & 2 & \dots & k-1 & \tau(k) & k+1 & \dots & k & \tau(k+1) & \dots & n \end{pmatrix}$$

Malpositions (denoted F_τ):

$$F_\tau = \begin{cases} (k, k+1), \dots, (k, \tau(k)) \\ (k+1, \tau(k)), (k+2, \tau(k)), \dots, (\tau(k)-1, \tau(k)) \end{cases}$$

Let us count on a specific example:

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 2 & 6 & 4 & 5 & 3 & 7 \end{pmatrix}$$

$$\begin{cases} (3, 4), (3, 5), (3, 6) \\ (4, 6), (5, 6) \end{cases}$$

$$|F_\tau| = (\tau(k) - k) + ((\tau(k) - 1) - k) = 2\tau(k) - 2k - 1 = 2(\tau(k) - k) - 1 \text{ even}$$

Theorem 7.3. 1. $\text{sign}(\text{id}) = 1$

2. $\text{sign}(\pi \circ \sigma) = \text{sign}(\pi) \circ \text{sign}(\sigma)$
Hence, $\text{sign} \sigma_n \rightarrow \{\pm 1\}$ is a homomorphism.
 $(\{+1, -1\}, \cdot)$ is a group $\cong (\mathbb{Z}_2, +)$

$$+1 \rightarrow [0]_2$$

$$-1 \rightarrow [1]_2$$

3. $\text{sign}(\pi^{-1}) = \text{sign}(\pi)$

Proof. 1. obvious, because there are no malpositions

2.

$$\text{sign}(\pi \circ \sigma) = \prod_{i < j} \frac{(\pi \circ \sigma(j) - \pi \circ \sigma(i))}{j - i} \prod_{i < j} \frac{\sigma(j) - \sigma(i)}{\sigma(j) - \sigma(i)}$$

because of bijectivity

$$= \underbrace{\prod_{i < j} \frac{\pi(\sigma(j)) - \pi(\sigma(i))}{\sigma(j) - \sigma(i)}}_{\text{sign } \pi} \cdot \underbrace{\prod_{i < j} \frac{\sigma(j) - \sigma(i)}{j - i}}_{\text{sign } \pi}$$

3. Homomorphism

$$\text{sign}(\pi^{-1} \circ \pi) = 1 \iff \text{sign}(\pi^{-1}) \circ \text{sign}(\pi) = 1 \iff \text{sign}(\pi^{-1}) = \text{sign}(\pi)^{-1}$$

□

Remark 7.5. Recall that the kernel of a homomorphism defines a subgroup.

Corollary 7.4. 1. If $\pi = \tau_1 \circ \dots \circ \tau_k$ is a product of transpositions, then $\text{sign}(\pi) = (-1)^k$

2. $\mathfrak{a}_n = \{\pi \in \sigma_n \mid \text{sign}(\pi) = +1\} = \ker(\{\text{sign} : \sigma_n \rightarrow \{\pm 1\}\})$ is a subgroup of σ_n , the so-called alternating group

$$|\mathfrak{a}_n| = \frac{n!}{2}$$

Corollary 7.5.

$$\dim V = n$$

$$\Delta : V^n \rightarrow \mathbb{K} \quad \text{determinant form}$$

then it holds that $\forall \sigma \in \sigma_n : \Delta(a_{\sigma(1)}, \dots, a_{\sigma(n)}) = \text{sign}(\sigma) \cdot \Delta(a_1, \dots, a_n)$

Proof. If $\sigma = \tau$ is a transposition, the fourth property:

$$\Delta(a_{\tau(1)}, \dots, a_{\tau(n)}) = -\Delta(a_1, \dots, a_n)$$

and $\text{sign}(\tau) = -1$.

The general case: $\sigma = \tau_1 \circ \dots \circ \tau_k$ and $\sigma = \tau_1 \circ \sigma_1$.

$$\begin{aligned} \Delta(a_{\sigma(1)}, \dots, a_{\sigma(n)}) &= \Delta(a_{\tau_1(\sigma_1(1))}, \dots, a_{\tau_1(\sigma_1(n))}) \\ &= -\Delta(a_{\sigma_1(1)}, \dots, a_{\sigma_1(n)}) \end{aligned}$$

$$\sigma_1 = \tau_2 \circ \sigma_2$$

$$\begin{aligned} &= \text{and so on and so forth} \\ &= (-1)^2 \Delta(a_{\sigma_2(1)}, \dots, a_{\sigma_2(n)}) \\ &= (-1)^k \Delta(a_1, \dots, a_n) \\ &= \text{sign } \sigma \Delta(a_1, \dots, a_n) \end{aligned}$$

□

7.5 Leibniz formula for determinants

Definition 7.5 (Definition with theorem). Define the determinant of matrix A .

$$\Delta(a_1, \dots, a_n) = \Delta(b_1, \dots, b_n) \cdot \det A$$

if $a_j = \sum_{i=1}^n a_{ij} b_i$. Hence,

$$\begin{pmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{nj} \end{pmatrix} = \Phi_B(a_j)$$

Let $\dim V = n$. Let $B = (b_1, \dots, b_n)$ be a basis of V . $a_1, \dots, a_n \in V$ with coordinates

$$\Phi_B(a_j) = \begin{pmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{nj} \end{pmatrix} \quad A := \begin{bmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{n1} & \dots & a_{nn} \end{bmatrix}$$

Then $\Delta(a_1, \dots, a_n) = \det(A) \cdot \Delta(b_1, \dots, b_n)$ where

$$\det(A) := \sum_{\pi \in \mathcal{S}_n} \text{sign}(\pi) a_{1\pi(1)} a_{2\pi(2)} \dots a_{n\pi(n)}$$

is called determinant of A

This formula was discovered by Leibniz.

Example 7.6. Consider $n = 2$.

$$\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = \underbrace{a_{11}a_{22}}_{\pi=\text{id}} - \underbrace{a_{12}a_{21}}_{\pi=\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}}$$

Proof.

$$a_j = \sum_{i=1}^n a_{ij} b_i$$

$$\Delta(a_1, \dots, a_n) = \Delta \left(\sum_{i_1=1}^n a_{i_1,1} b_{i_1}, \sum_{i_2=1}^n a_{i_2,2} b_{i_2}, \dots, \sum_{i_n=1}^n a_{i_n,n} b_{i_n} \right)$$

because it is multilinear

$$= \sum_{i_1=1}^n \sum_{i_2=1}^n \cdots \sum_{i_n=1}^n a_{i_1,1} a_{i_2,2} \cdots a_{i_n,n} \cdot \Delta(b_{i_1}, b_{i_2}, \dots, b_{i_n})$$

where Δ is zero if $b_i = b_j$.

$$\Rightarrow i_1, \dots, i_n \text{ are all different elements in } \{1, \dots, n\}$$

$$\Rightarrow \text{every element occurs exactly once}$$

$$i_1, \dots, i_n \text{ is permutation of } 1, \dots, n$$

$$\exists \sigma \in \sigma_n : i_1 = \sigma(1), \dots, i_n = \sigma(n)$$

$$= \sum_{\sigma \in \sigma_n} a_{\sigma(1)1} a_{\sigma(2)2} \cdots a_{\sigma(n)n} \underbrace{\Delta(b_{\sigma(1)} \cdots b_{\sigma(n)})}_{\text{sign } \sigma \Delta(b_1, \dots, b_n) \text{ because of Corollary 7.5}}$$

$$= \sum_{\pi \in \sigma_n} a_{1\pi(1)} \cdots a_{n\pi(n)} \cdot \text{sign}(\pi) \Delta(b_1, \dots, b_n)$$

□

Corollary 7.6. A determinant form is uniquely defined by the value $\Delta(b_1, \dots, b_n)$ on a basis. Especially, $\Delta(b_1, \dots, b_n) \neq 0 \iff \Delta(b_1, \dots, b_n) \neq 0$ [for any basis]
 $\iff \Delta(b_1, \dots, b_n) \neq 0$ [for every basis].

Assume $\Delta(b_1, \dots, b_n) = 0$ for any basis. Every other basis can be expressed by b_1, \dots, b_n and the formula gives $\Delta(a_1, \dots, a_n) = 0 \quad \forall a_1, \dots, a_n$.

□

Remark 7.7 (Summary). 1. The determinant form defines a 1-dimensional vector space.

2. There exists a non-trivial determinant form. Given a basis b_1, \dots, b_n

$$\Delta(b_1, \dots, b_n) = 1$$

By Theorem 7.5, $\Delta(a_1, \dots, a_n) = \det(\Phi_B(a_1), \dots, \Phi_B(a_n))$.

Proof of Theorem 7.5. 1.

$$\begin{aligned} \Delta(a_1, \dots, \lambda a_k, \dots, a_n) &= \sum_{\pi \in \sigma_n} (-1)^\pi a_{\pi(1)1} \dots \lambda a_{\pi(k)k} \dots a_{\pi(n)n} \\ &= \lambda \cdot \sum_{\pi \in \sigma_n} (-1)^\pi a_{\pi(1)1} \dots a_{\pi(n)n} \\ &= \lambda \cdot \Delta(a_1, \dots, a_n) \end{aligned}$$

2.

$$\begin{aligned} \Delta(a_1, \dots, a_k + v, \dots, a_n) &= \sum_{\pi \in \sigma_n} (-1)^\pi a_{\pi(1)1} \dots (a_{\pi(k)k} + v_{\pi(k)}) \dots a_{\pi(n)n} \\ &= \sum_{\pi \in \sigma_n} (-1)^\pi a_{\pi(1)1} \dots a_{\pi(k)k} \dots a_{\pi(n)n} \\ &\quad + \sum_{\pi \in \sigma_n} (-1)^\pi a_{\pi(1)1} \dots v_{\pi(k)} \dots a_{\pi(n)n} \\ &= \Delta(a_1, \dots, a_k, \dots, a_n) + \Delta(a_1, \dots, v, \dots, a_n) \end{aligned}$$

This proves multilinearity.

3. Let $a_k = a_l$, $a_{ik} = a_{il} \forall i = 1, \dots, n$. Without loss of generality, $k < l$.

$$\Delta(a_1, \dots, a_k) = \sum_{\pi \in \sigma_n} (-1)^\pi a_{\pi(1)1} \dots a_{\pi(k)k} \dots a_{\pi(l)l} \dots a_{\pi(n)n} = \text{(ref *)}$$

Let $\tau = \tau_{kl}$, exchange of k and l .

Claim.

$$\begin{aligned} \sigma_n &= \underbrace{\mathcal{A}_n}_{\substack{\text{alternating group} \\ = \{ \pi \mid \text{sign}(\pi) = +1 \}}} \cup \underbrace{\mathcal{A}_n \cdot \tau}_{= \{ \pi \circ \tau \mid \pi \in \mathcal{A}_n \}} \end{aligned}$$

The set of all permutations is the set of even permutations unified with the set of even permutations with one transposition applied. Thus, one transposition suffices to turn even permutations into odd permutations.

Proof. Direction \Leftarrow . Let $\text{sign}(\pi) = -1$.

$$\Rightarrow \pi = (\pi \circ \tau) \circ \underbrace{\tau}_{=\text{id}}$$

$$\sigma = \pi \circ \tau \text{ has } \text{sign}(\sigma) = \text{sign}(\pi \circ \tau) = \text{sign}(\pi) \cdot \text{sign}(\tau) = (-1) \cdot (-1) = 1.$$

$$\sigma \in \mathcal{A}_n \text{ and } \pi = \sigma \circ \tau$$

□

$$\begin{aligned} (\text{ref } *) &= \sum_{\pi \in \mathcal{A}_n} \underbrace{(-1)^\pi}_{=+1} a_{\pi(1)1} \dots a_{\pi(n)n} \\ &+ \sum_{\substack{\pi \in \mathcal{A}_n \\ \pi = \sigma \circ \tau}} \underbrace{(-1)^{\text{sign}(\pi)}}_{=-1} a_{\pi(1)1} \dots a_{\pi(n)n} \\ &= \sum_{\pi \in \mathcal{A}_n} a_{\pi(1)1} \dots a_{\pi(n)n} - \sum_{\sigma \in \mathcal{A}_n} \underbrace{a_{\sigma \circ \tau(1)1} \dots a_{\sigma \circ \tau(k)2} \dots a_{\sigma \circ \tau(l)l} \dots a_{\sigma \circ \tau(n)n}}_{\substack{a_{\sigma(1)1} \dots \underbrace{a_{\sigma(l)k}}_{=a_{\sigma(l)l}} \dots \underbrace{a_{\sigma(k)l}}_{=a_{\sigma(k)k}} \dots a_{\sigma(n)n}}} = 0 \end{aligned}$$

□

This previous part, beginning with the reference from 2018/03/12, was actually added on 2018/03/14, because we skipped it by accident.

$$\Delta(a_1, \dots, a_n)$$

Determinant form \Longleftrightarrow

$$\textbf{multilinear} \quad \Delta(a_1, \dots, \lambda a_k + \mu a'_k, \dots, a_n) = \lambda \Delta(a_1, \dots, a_k, \dots, a_n) + \mu \Delta(a_1, \dots, a'_k, \dots, a_n)$$

$$\textbf{anti-symmetrical} \quad \Delta(a_1, \dots, a_k, \dots, a_l, \dots, a_n) = -\Delta(a_1, \dots, a_l, \dots, a_k, \dots, a_n)$$

$$\Delta(a_{\pi(1)}, \dots, a_{\pi(n)}) = (-1)^\pi \Delta(a_1, \dots, a_n)$$

$$\text{where } (-1)^\pi := \text{sign}(\pi) = (-1)^{F(\pi)}$$

$$F(\pi) = \left\{ (i, j) \mid i < j \wedge \pi(i) > \pi(j) \right\}$$

$$\text{sign}(\pi \circ \sigma) = \text{sign}(\pi) \cdot \text{sign}(\sigma)$$

Basis b_1, \dots, b_n .

$$\Delta\left(\sum_{i=1}^n a_{i1}b_i, \dots, \sum_{i=1}^n a_{in}b_i\right) = \det A \cdot \Delta(b_1, \dots, b_n)$$

$$\det(A) = \sum_{\pi \in \sigma_n} (-1)^\pi a_{1\pi(1)} \dots a_{n\pi(n)} = \sum_{\pi \in \sigma_n} (-1)^\pi a_{\pi(1)1} \dots a_{\pi(n)n}$$

Lemma 7.1. Let V, W be vector spaces over \mathbb{K} with $\dim V = \dim W = n$. Let $\Delta : W^n \rightarrow \mathbb{K}$ be a determinant form and $f : V \rightarrow W$ linear.

$$V \xrightarrow{f} W$$

$$V^n \xrightarrow{f^{(n)}} W^n \xrightarrow{\Delta} \mathbb{K}$$

$$(v_1, \dots, v_n) \mapsto (f(v_1), \dots, f(v_n))$$

$$\implies \Delta^f : V^n \rightarrow \mathbb{K}$$

$$\Delta^f(v_1, \dots, v_n) = \Delta(f(v_1), \dots, f(v_n))$$

is a determinant form on V .

Proof. 1. Multilinear

$$\begin{aligned} & \Delta^f(v_1, \dots, \lambda v_k + \mu v'_k, \dots, v_n) \\ &= \Delta(f(v_1), \dots, f(\lambda v_k + \mu v'_k), \dots, f(v_n)) \\ &= \Delta(f(v_1), \dots, \lambda f(v_k) + \mu f(v'_k), \dots, f(v_n)) \\ &= \lambda \Delta(f(v_1), \dots, f(v_k), \dots, f(v_n)) + \mu \Delta(f(v_1), \dots, f(v'_k), \dots, f(v_n)) \\ &= \lambda \Delta^f(v_1, \dots, v_k, \dots, v_n) + \mu \Delta^f(v_1, \dots, v'_k, \dots, v_n) \end{aligned}$$

□

Corollary 7.8. Let $V = W$, $\Delta : V^n \rightarrow \mathbb{K}$ determinant form.

$$f : V \rightarrow V \text{ linear}$$

$$\implies \Delta^f \text{ is determinant form}$$

Because there is (except for one factor) only one determinant form:

$$\exists C_f \in \mathbb{K} : \Delta^f(v_1, \dots, v_n) = C_f \cdot \Delta(v_1, \dots, v_n) \forall v_1, \dots, v_n \in V$$

$$\det(f) := C_f \text{ is called determinant on } f$$

Proof. Let Δ_1, Δ_2 be two determinant forms. Let b_1, \dots, b_n be a basis.

$$\Delta_1(v_1, \dots, v_n) = \det A \cdot \Delta_1(b_1, \dots, b_n)$$

$$\Delta_2(v_1, \dots, v_n) = \det A \cdot \Delta_2(b_1, \dots, b_n)$$

$$v_j = \sum_{i=1}^n a_{ij} b_i$$

$$\implies \Delta_2(v_1, \dots, v_n) = \frac{\Delta_2(b_1, \dots, b_n)}{\Delta_1(b_1, \dots, b_n)} \cdot \Delta_1(v_1, \dots, v_n)$$

$$\implies C_f = \frac{\Delta^f(b_1, \dots, b_n)}{\Delta(b_1, \dots, b_n)} = \det(f)$$

□

7.6 On determinants, invertibility and linear independence

Corollary 7.9. $B = (b_1, \dots, b_n)$ is basis of V . $\phi_B^B(f)$ is matrix representation of linear f and $\det(f) = \det \phi_B^B(f)$ (LHS by Corollary 7.8, RHS by Definition 7.5 $\sum_{\pi} (-1)^{\pi} \dots$).

Proof.

$$\det(f) = \frac{\Delta(f(b_1), \dots, \Delta(f(b_n)))}{\Delta(b_1, \dots, b_n)}$$

$$\begin{aligned} f(b_j) &= \sum_{i=1}^n \phi_B(f(b_j))_i \cdot b_i \\ &= \sum_{i=1}^n (\phi_B^B(f))_{ij} b_i \end{aligned}$$

with $\phi_B^B(f)_{ij} = \phi_B(f(b_j))_i$.

$$\det f = \frac{\det \phi_B^B(f) \cdot \Delta(b_1, \dots, b_n)}{\Delta(b_1, \dots, b_n)}$$

□

Theorem 7.6. $f : V \rightarrow V$ is invertible $\iff \det(f) \neq 0$.

Proof. \implies Let Δ be a non-trivial determinant form.

$$B = (b_1, \dots, b_n) \text{ is a basis } \implies \Delta(b_1, \dots, b_n) \neq 0$$

$$\det(f) = \frac{\Delta(f(b_1), \dots, f(b_n))}{\Delta(b_1, \dots, b_n)}$$

$\Leftarrow \det(f) \neq 0 \Rightarrow (f(b_1), \dots, f(b_n))$ is basis $\iff f$ is invertible. If f is invertible, then $(f(b_1), \dots, f(b_n))$ is basis.

$$\Rightarrow \Delta(f(b_1), \dots, f(b_n)) \neq 0 \Rightarrow \det(f) \neq 0$$

If f is not invertible, then

$$\Rightarrow f(b_1) \dots f(b_n) \text{ is linear dependent}$$

$$\exists k : f(b_k) = \sum_{i \neq k} \lambda_i f(b_i)$$

Without loss of generality: $k = n$

$$\begin{aligned} \Delta(f(b_1), \dots, f(b_n)) &= \Delta(f(b_1), \dots, f(b_{n-1}), \sum_{i=1}^{n-1} \lambda_i f(b_i)) \\ &= \sum_{i=1}^n \lambda_i \underbrace{\Delta(f(b_1), \dots, f(b_{n-1}), f(b_i))}_{=0 \forall i \in \{1, \dots, n-1\}} \\ &= 0 \end{aligned}$$

□

Corollary 7.10. For a matrix $A \in \mathbb{K}^{n \times n}$ it holds that $\det A \neq 0 \iff A$ has full rank.

Proof. \Rightarrow If A is invertible $\ker(A) = \emptyset$, so A has full rank.

\Leftarrow If A does not have full rank, consider $x \neq 0$ with $x \in \ker(A)$ then $Ax = 0$ and $A(2x) = 0$. Thus it is not injective and therefore not invertible.

If A has full rank it is surjective (column space spans all n dimensions) and injective ($x \neq y \Rightarrow Ax \neq Ay$). Thus invertible.

□

Theorem 7.7. $f, g : V \rightarrow V$ linear.

$$\Rightarrow \det(f \circ g) = \det(f) \cdot \det(g)$$

for a matrix: $\det(A \cdot B) = \det(A) \cdot \det(B)$

Proof. If f and g are invertible

$$\det(f) = \frac{\Delta(f(b_1), \dots, f(b_n))}{\Delta(b_1, \dots, b_n)}$$

for arbitrary bases (b_1, \dots, b_n) of V .

$$\begin{aligned}\det(f \circ g) &= \frac{\Delta(f(g(b_1)), \dots, f(g(b_n)))}{\Delta(b_1, \dots, b_n)} \cdot \frac{\Delta(g(b_1), \dots, g(b_n))}{\Delta(g(b_1), \dots, g(b_n))} \\ &= \underbrace{\frac{\Delta(f(g(b_1)), \dots, f(g(b_n)))}{\Delta(g(b_1), \dots, g(b_n))}}_{\det(f) \neq 0} \cdot \underbrace{\frac{\Delta(g(b_1), \dots, g(b_n))}{\Delta(b_1, \dots, b_n)}}_{\det(g) \neq 0}\end{aligned}$$

If f or g is not invertible

$$f \text{ is not invertible} \implies \det(f) = 0$$

Same for g .

Claim. $f \circ g$ invertible $\iff f$ invertible and g invertible.

$f \circ g$ invertible $\implies f \circ g$ surjective $\implies f$ surjective $\implies (\dim V < \infty)$ f is bijective.

$f \circ g$ invertible $\implies f \circ g$ injective $\implies g$ injective $\implies g$ bijective.

$$\implies f \circ g \text{ is not invertible}$$

$$\det(f \circ g) = 0 = \det(f) \cdot \det(g)$$

□

Corollary 7.11. For $A, B \in \mathbb{K}^{n \times n}$ it holds that

1. $\det(A \cdot B) = \det(A) \cdot \det(B)$
2. $\det(A^{-1}) = \frac{1}{\det(A)}$ if invertible
3. $\det(A) = 0 \iff \text{rank}(A) < n$
4. $\det(A^t) = \det(A)$

Proof of Corollary 7.11. 1. $\det(A \cdot B) = \det(f_A \circ f_B) = \det(f_A) \cdot \det(f_B) = \det(A) \cdot \det(B)$ (compare with Corollary 7.9)

2. $A \cdot A^{-1} = I$ and $1 = \det(A \cdot A^{-1}) = \det(A) \cdot \det(A^{-1})$

Remark 7.8 (From the practicals).

$$\det(A) = \det(f_A)$$

Shown so far:

$$\det f = \det(\phi_B^B(f))$$

$$A = \phi_B^B(f_A)$$

for $B = (e_1, \dots, e_n)$

□

Direct proof of Corollary 7.11 (1).

$$A = \begin{bmatrix} s_1 & \cdots & s_n \\ \vdots & & \vdots \end{bmatrix}$$

s_i are column vectors of A . Let Δ be the uniquely defined determinant form by $\Delta(e_1, \dots, e_n) = 1$.

$$\begin{aligned} A \cdot B &= \begin{bmatrix} s_1 & \cdots & s_n \\ \vdots & & \vdots \end{bmatrix} \cdot \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ \vdots & & & \vdots \\ b_{n1} & & & b_{nn} \end{bmatrix} \\ &= \begin{bmatrix} s_1 b_{11} + s_2 b_{21} + \cdots + s_n b_{n1} & s_1 b_{12} + s_2 b_{22} + \cdots + s_n b_{n2} & \cdots & s_1 b_{1n} + s_2 b_{2n} + \cdots + s_n b_{nn} \\ \vdots & \vdots & & \vdots \end{bmatrix} \\ \det(A \cdot B) &= \frac{\Delta(s_1(A \cdot B), \dots, s_n(A \cdot B))}{\Delta(e_1, \dots, e_n)} = \Delta \left(\sum_{i_1=1}^n s_{i_1} b_{i_1 1}, \sum_{i_2=1}^n s_{i_2} b_{i_2 2}, \dots, \sum_{i_n=1}^n s_{i_n} b_{i_n n} \right) \\ &= \sum_{i_1=1}^n \cdots \sum_{i_n=1}^n b_{i_1 1} b_{i_2 2} \cdots b_{i_n n} \underbrace{\Delta(s_{i_1}, \dots, s_{i_n})}_{=0} \end{aligned}$$

if one index occurs twice. It suffices to consider \sum_{i_1, \dots, i_n} such that all i_j are difference. If all are difference, then all occur exactly once. Hence, i_1, \dots, i_n is permutation of $1, \dots, n$.

$$\begin{aligned} &= \sum_{\pi \in \sigma_n} b_{\pi(1)1} \cdots b_{\pi(n)n} \Delta(s_{\pi(1)} \cdots s_{\pi(n)}) \\ &= \sum_{\pi \in \sigma_n} \underbrace{(-1)^\pi b_{\pi(1)1} \cdots b_{\pi(n)n}}_{\det B} \underbrace{\Delta(s_1, \dots, s_n)}_{=\det(A)} = \det(B) \cdot \det(A) \end{aligned}$$

□

Proof of Corollary 7.11 (3). A is invertible $\iff f_A$ is invertible. $\implies \det(A) = 0 \iff \det(f_A) = 0 \iff f_A$ is not bijective $\iff \text{rank}(A) < n$. □

Proof of Corollary 7.11 (4).

$$\begin{aligned} \det(A^t) &= \sum_{\pi \in \sigma_n} (-1)^\pi (A^t)_{\pi(1)1} \cdots (A^t)_{\pi(n)n} \\ &= \sum_{\pi \in \sigma_n} (-1)^\pi a_{1\pi(1)} \cdots a_{n\pi(n)} \end{aligned}$$

Remark 7.9.

$$\begin{aligned}\sigma_n &\rightarrow \sigma_n \\ \pi &\mapsto \pi^{-1} \text{ is bijective} \\ \text{injective: } \pi^{-1} = \sigma^{-1} &\implies \pi = \sigma \\ \text{surjective: } \pi &= (\pi^{-1})^{-1}\end{aligned}$$

$$= \sum_{\pi \in \sigma_n} (-1)^{\pi^{-1}} a_{1\pi^{-1}(1)} \dots a_{n\pi^{-1}(n)}$$

Every index i occurs once on the left side and once on the right side. i occurs right

$$\pi^{-1}(j) = i \iff j = \pi(i)$$

$$= \sum_{\pi \in \sigma_n} (-1)^{\pi} a_{\pi(1)1} \dots a_{\pi(n)n}$$

$$\begin{aligned}\text{sign}(\pi \circ \pi^{-1}) &= 1 \\ &= \text{sign}(\pi) \cdot \text{sign}(\pi^{-1})\end{aligned}$$

Remark 7.10 (A small exercise).

$$\begin{aligned}\det(A) &= \det(f_A) \\ \prod_{j=1}^n a_{j,\pi^{-1}(j)} &= \prod_{i=1}^n a_{\pi(i),\pi^{-1}(\pi(i))} = \prod_{i=1}^n a_{\pi(i),i} \\ j &= \pi(i)\end{aligned}$$

□

Definition 7.6.

$$\text{perm}(A) := \sum_{\pi \in \sigma_n} a_{\pi(1)1} \dots a_{\pi(n)n}$$

is called permanent of A .

Open problem: for which matrix does $\text{perm}(A) = 0$ hold?

Example 7.8 (Computation of the determinant).

$$\begin{aligned}\dim &\leq 3 \\ n = 2 : \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} &= a_{11}a_{22} - a_{12}a_{21}\end{aligned}$$

$$n = 3 : \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \sum_{\sigma \in \sigma_n} (-1)^\pi a_{\pi(1)1} a_{\pi(2)2} a_{\pi(3)3}$$

TODO drawing cayley graph

By the Cayley-Graph of group σ_3 we can see that $\sigma_3 = \left\langle (\underline{12}), (\underline{\underline{23}}) \right\rangle = -1$.

$$= a_{11}a_{22}a_{33} + a_{21}a_{32}a_{13} + a_{31}a_{12}a_{23}$$

TODO drawing tic tac toe

$$-a_{21}a_{12}a_{33} - a_{11}a_{32}a_{23} - a_{31}a_{22}a_{13}$$

TODO drawing tic tac toe

$$\begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$

Rule of Sarrus holds only for $n = 2$ or $n = 3$.

↓ This lecture took place on 2018/03/14.

Example 7.9 (Rule by Sarrus). Let $n = 2$:

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

Let $n = 3$:

$$\begin{vmatrix} 1 & 2 & 5 & 1 & 2 \\ 2 & 5 & 14 & 2 & 5 \\ 5 & 14 & 42 & 5 & 14 \end{vmatrix} = 1$$

$$\begin{aligned} & 1 \cdot 5 \cdot 42 + 2 \cdot 14 \cdot 5 + 5 \cdot 2 \cdot 14 - 5 \cdot 5 \cdot 5 - 1 \cdot 14 \cdot 14 - 2 \cdot 2 \cdot 42 \\ &= 14 \cdot (1 \cdot 5 \cdot 3 + 2 \cdot 5 + 5 \cdot 2) - 125 - 14 \cdot (14 + 2 \cdot 2 \cdot 3) \\ &= 14 \cdot 35 - 125 - 14 \cdot 26 \\ &= 14 \cdot 9 - 125 = 1 \end{aligned}$$

An error in the computation will be enhanced.

Let $n = 4$. $|\sigma_n| = 24$ makes consideration of all permutations impractical.

Lemma 7.2. Let A be an upper triangular matrix, hence $a_{ij} = 0$ if $i > j$.

$$\implies \det(A) = a_{11}a_{22} \dots a_{nn}$$

Proof.

$$\det(A) = \sum_{\pi \in \sigma_n} (-1)^\pi a_{\pi(1)1} \dots a_{\pi(n)n}$$

such that $\pi(j) \leq j \forall j$.

$$\implies \text{id}$$

$$\pi(j) \leq j \forall j \implies \pi(1) \leq 1 \implies \pi(1) = 1$$

$$\pi(2) \leq 2 \implies \pi(2) = 2$$

$$\pi(3) \leq 3 \implies \pi(3) = 3$$

...

$$\pi(n) \leq n \implies \pi(n) = n$$

□

Theorem 7.8. Let $A = (a_{ij})$ be a $n \times n$ matrix.

1. Let z_1, \dots, z_n be row vectors of A . Then

$$\det \begin{bmatrix} z_1 & \dots \\ \vdots & \\ z_n & \dots \end{bmatrix} = \det \begin{bmatrix} z_1 & \dots \\ z_i + \lambda z_j & \dots \\ \vdots & \\ z_n & \dots \end{bmatrix} \forall i \neq j, \lambda \in \mathbb{K}$$

2. Let S_1, \dots, S_n be columns of A . Then,

$$\det \begin{pmatrix} S_1 & \dots & S_n \\ \vdots & & \vdots \end{pmatrix} = \det \begin{pmatrix} S_1 & \dots & S_i + \lambda S_j & \dots & S_j & \dots & S_n \\ \vdots & & \vdots & & \vdots & & \vdots \end{pmatrix}$$

Proof for column i.

$$\begin{aligned} \Delta(s_1, \dots, s_n) &= \Delta(s_1, \dots, s_i + \lambda s_j, \dots, s_n) \\ &= \Delta(s_1, \dots, s_i, \dots, s_n) + \lambda \underbrace{\Delta(s_1, \dots, s_j, \dots, s_j, \dots, s_n)}_{=0} \end{aligned}$$

□

Second proof. Row form is multiplication from left with matrix of structure

$$\det((I + \lambda E_{ij})A) = \underbrace{\det(I + \lambda E_{ij})}_{\text{triangular matrix}=1} \cdot \det(A)$$

□

Example 7.10.

$$\begin{vmatrix} 1 & 2 & 5 \\ 2 & 5 & 14 \\ 5 & 14 & 42 \end{vmatrix} = \begin{vmatrix} 1 & 2 & 5 \\ 0 & 1 & 4 \\ 0 & 4 & 17 \end{vmatrix} = \begin{vmatrix} 1 & 2 & 5 \\ 0 & 1 & 4 \\ 0 & 0 & 1 \end{vmatrix} = 1$$

Example 7.11.

$$\begin{aligned} & \begin{vmatrix} 1 & 0 & 3 & -2 \\ 2 & 6 & 4 & 1 \\ 3 & 3 & -1 & -1 \\ -1 & 2 & 4 & 1 \end{vmatrix} = \begin{vmatrix} 1 & 0 & 3 & -2 \\ 0 & 6 & -2 & 5 \\ 0 & 3 & -10 & 5 \\ 0 & 2 & 7 & -1 \end{vmatrix} \\ &= \frac{1}{3} \frac{1}{2} \begin{vmatrix} 1 & 0 & 3 & -2 \\ 0 & 6 & -2 & 5 \\ 0 & 6 & -20 & 10 \\ 0 & 6 & 21 & -3 \end{vmatrix} = \frac{1}{6} \begin{vmatrix} 1 & 0 & 3 & -2 \\ 0 & 6 & -2 & 5 \\ 0 & 0 & -18 & 5 \\ 0 & 0 & 23 & -8 \end{vmatrix} = \frac{1}{6} \cdot 6 \begin{vmatrix} 1 & 0 & 3 & -2 \\ 0 & 1 & -2 & 5 \\ 0 & 0 & -18 & 5 \\ 0 & 0 & 23 & -8 \end{vmatrix} \\ &= \begin{vmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 8 & 5 \\ 0 & 0 & -8 & 5 \\ 0 & 0 & 7 & -8 \end{vmatrix} = \begin{vmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 8 & 5 \\ 0 & 0 & -8 & 5 \\ 0 & 0 & -1 & -3 \end{vmatrix} = \begin{vmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 8 & 5 \\ 0 & 0 & 0 & 29 \\ 0 & 0 & -1 & -3 \end{vmatrix} \\ &= - \begin{vmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 8 & 5 \\ 0 & 0 & -1 & -3 \\ 0 & 0 & 0 & 29 \end{vmatrix} = 29 \end{aligned}$$

Remark 7.11 (Laws, discussed so far).

$$\begin{aligned} & \begin{vmatrix} z_1 & \dots \\ \lambda \cdot z_1 & \dots \\ z_n & \dots \end{vmatrix} = \lambda \begin{vmatrix} z_1 & \dots \\ z_k & \dots \\ z_n & \dots \end{vmatrix} \\ & \begin{vmatrix} z_1 & \dots \\ z_1 + \lambda z_j & \dots \\ z_n & \dots \end{vmatrix} = \begin{vmatrix} z_1 & \dots \\ z_i & \dots \\ z_n & \dots \end{vmatrix} \quad (i \neq j) \\ & \begin{vmatrix} z_1 & \dots \\ \vdots & \vdots \\ z_i & \dots \\ \vdots & \vdots \\ z_j & \dots \\ \vdots & \vdots \\ z_n & \dots \end{vmatrix} = - \begin{vmatrix} z_1 & \dots \\ \vdots & \vdots \\ z_j & \dots \\ \vdots & \vdots \\ z_i & \dots \\ \vdots & \vdots \\ z_n & \dots \end{vmatrix} \end{aligned}$$

$$\begin{vmatrix} a_{11} & \dots & & \\ & a_{22} & \dots & \\ & & a_{33} & \dots \\ & & & \ddots \\ 0 & & & & a_{nn} \end{vmatrix} = a_{11} \cdot a_{nn}$$

Lemma 7.3. 1.

$$\begin{vmatrix} a_{11} & * & * & * \\ 0 & & & \\ 0 & & B & \\ 0 & & & \end{vmatrix} = a_{11} \cdot \det(B)$$

2.

$$\begin{vmatrix} & & & 0 \\ & B & & 0 \\ & & & 0 \\ * & * & * & a_{nn} \end{vmatrix} = \det(B) \cdot a_{nn}$$

3. If there are individual square matrices (A_1, A_2, \dots, A_k) along the diagonal of a matrix, the determinant of the matrix is the product of the determinant of the submatrices.

$$\det(A) = \det(A_1) \cdot \det(A_2) \cdot \dots \cdot \det(A_k)$$

Proof. We only prove the second property.

All permutations which do not map index 1 to 1, introduce a factor zero making the product zero. If index 1 is mapped to 1, the product in Leibniz' formula is multiplied with a_{11} in all permutations. We can extract factor a_{11} and get the determinant of B multiplied with a_{11} .

$$\begin{aligned} \begin{vmatrix} & & & 0 \\ & B & & \vdots \\ & & & 0 \\ a_{n,1} & \dots & a_{n,n-1} & a_{n,n} \end{vmatrix} &= \sum_{\pi \in \sigma_n} (-1)^\pi a_{\pi(1)1} \dots a_{\pi(n)n} \\ &= \sum_{\pi' \in \sigma_{n-1}} (-1)^{\pi'} a_{\pi'(1)1} \dots a_{\pi'(n-1)n-1} \cdot a_{nn} \\ &= \det(B) \cdot a_{nn} \end{aligned}$$

$$\{\pi \in \sigma_n \mid \pi(n) = n\}$$

$$\pi(n) = n$$

$$B = \begin{pmatrix} a_{11} & \dots & a_{1,n-1} \\ \vdots & & \\ a_{n-1,1} & \dots & a_{n,n-1} \end{pmatrix}$$

Same idea: If

$$A = \begin{bmatrix} 0 \\ \vdots \\ 0 \\ \vdots \\ a_{ij} \\ 0 \\ \vdots \\ 0 \end{bmatrix}$$

Exchange the i -th row with the last row.

$$= \pm 1 \begin{bmatrix} 0 \\ \vdots \\ 0 \\ \vdots \\ 0 \\ \vdots \\ 0 \\ \vdots \\ a_{ij} \end{bmatrix}$$

□

Definition 7.7.

$$A \in \mathbb{K}^{n \times n}$$

$A_{k,l}$ is an $(n-1) \times (n-1)$ matrix, that is created by omitting the k -th row and l -th column.

$$\begin{bmatrix} a_{1,1} & \dots & a_{1,l-1} & a_{1,l+1} & \dots & a_{1,n} \\ \vdots & & & & & \vdots \\ a_{k-1,1} & \dots & a_{k-1,l-1} & a_{k-1,l+1} & \dots & a_{k-1,n} \\ a_{k+1,1} & \dots & a_{k+1,l-1} & a_{k+1,l+1} & \dots & a_{k+1,n} \\ \vdots & & & & & \vdots \\ a_{n,1} & \dots & a_{n,l-1} & a_{n,l+1} & \dots & a_{n,n} \end{bmatrix}$$

Pierre-Simon Laplace (1749–1827)

Definition 7.8 (Laplace expansion). *In German, this theorem is called Entwicklungssatz von Laplace*

Let l be fixed.

$$\det(A) = \sum_{k=1}^n a_{kl} (-1)^{k+l} \det(A_{kl})$$

“Expansion along column l ”.

Let k be fixed.

$$\det(A) = \sum_{l=1}^n a_{kl}(-1)^{k+l} \det(A_{kl})$$

“Expansion along row k ”.

Example 7.12.

$$\begin{aligned} \begin{vmatrix} 1 & 2 & 5 \\ 2 & 5 & 14 \\ 5 & 14 & 42 \end{vmatrix} &= \sum_{l=1}^3 (-1)^{1+l} \det(A_{1l}) \quad \text{for } k = 1 \text{ fixed} \\ &= 1 \begin{vmatrix} 5 & 14 \\ 14 & 42 \end{vmatrix} - 2 \begin{vmatrix} 2 & 14 \\ 5 & 42 \end{vmatrix} + 5 \begin{vmatrix} 2 & 5 \\ 5 & 14 \end{vmatrix} \\ &= 1 \cdot (5 \cdot 42 - 14 \cdot 14) - 2(2 \cdot 42 - 5 \cdot 14) + 5 \cdot (2 \cdot 14 - 5 \cdot 9) \\ &= 1 \cdot (5 \cdot 3 \cdot 14 - 14 \cdot 14) - 2 \cdot (2 \cdot 3 \cdot 13 - 5 \cdot 14) \\ &= 14 - 2 \cdot 14 + 5 \cdot 15 = 1 \end{aligned}$$

Consider $k = 2$.

$$\begin{aligned} &-2 \cdot \begin{vmatrix} 2 & 5 \\ 14 & 42 \end{vmatrix} + 5 \cdot \begin{vmatrix} 1 & 5 \\ 5 & 42 \end{vmatrix} - 14 \cdot \begin{vmatrix} 1 & 2 \\ 5 & 14 \end{vmatrix} \\ &= -2(3 \cdot 14 \cdot 2 - 14 \cdot 5) + 5 \cdot (42 - 25) - 14 \cdot (14 - 10) \\ &= -2 \cdot 14 + 5 \cdot 17 - 4 \cdot 14 = -28 + 85 - 56 = 85 - 84 = 1 \end{aligned}$$

↓ This lecture took place on 2018/03/19.

Review:

- Determinants are multilinear (in rows and columns)
- Determinants switches its sign if two rows or row columns are exchanged
- $\Delta(s_1, \dots, s_n) = (-1)^\pi \Delta(s_{\pi(1)}, \dots, s_{\pi(n)})$ where s_i are column vectors
-

$$\begin{vmatrix} a_{11} & 0 & \dots & 0 \\ * & & & \\ \vdots & & B & \\ * & & & \end{vmatrix} = a_{11} \cdot \det B$$

$B = A_{11}$

where A_{kl} is the $(n-1) \times (n-1)$ matrix created by removal of the k -th row and l -th column. This is a special case of Laplace expansion.

7.7 Laplace expansion

$$\begin{aligned}\det A &= \sum_{k=1}^n (-1)^{k+l} a_{kl} \cdot \det A_{kl} && \text{for fixed } l \in \{1, \dots, n\} \\ &= \sum_{l=1}^n (-1)^{k+l} a_{kl} \cdot \det A_{kl} && \text{for fixed } k \in \{1, \dots, n\}\end{aligned}$$

So in the case of (a very classic example)

$$\begin{vmatrix} a_{11} & 0 & \dots & 0 \\ * & & & \\ \vdots & & B & \\ * & & & \end{vmatrix} = a_{11} \cdot (-1)^{1+1} \cdot \det A_{11}$$

for fixed $k = 1$:

$$\sum_{l=1}^n (-1)^{1+l} \underbrace{a_{1l}}_{=0 \text{ for } l>1} \det A_{1l}$$

Proof. Let $l \in \{1, \dots, n\}$ be fixed. Let e_k be a unit vector. For the l -th column,

$$s_l = \sum_{k=1}^n a_{kl} e_k = \begin{pmatrix} a_{1l} \\ a_{2l} \\ \vdots \\ a_{nl} \end{pmatrix}$$

$$\det(A) = \Delta(s_1, s_2, \dots, s_{l-1}, \sum_{k=1}^n a_{kl} \cdot e_k, s_{l+1}, \dots, s_n)$$

$$= \sum_{k=1}^n a_{kl} \Delta(s_1, \dots, s_{l-1}, e_k, s_{l+1}, \dots, s_n)$$

$$= \sum_{k=1}^n a_{kl} \begin{vmatrix} a_{11} & a_{12} & \vdots & a_{1,l-1} & 0 & a_{1,l+1} & \dots & a_{1n} \\ a_{21} & a_{22} & \vdots & a_{2,l-1} & \vdots & \vdots & & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\ \vdots & \vdots & \vdots & \vdots & 0 & \vdots & & \vdots \\ \vdots & \vdots & \vdots & \vdots & 1 & \vdots & & \vdots \\ \vdots & \vdots & \vdots & \vdots & 0 & \vdots & & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \vdots & a_{n,l-1} & 0 & a_{n,l+1} & \dots & a_{nn} \end{vmatrix}$$

Recognize the one in row k . We consecutively exchange row k with the row above until it becomes row 1. This gives $k - 1$ exchanges. Hence a cycle $(1 \dots k)$. This gives $\text{sign} = (-1)^{k-1}$.

$$= \sum_{k=1}^n a_{kl} (-1)^{k-1} \begin{vmatrix} a_{k1} & a_{k2} & \dots & a_{k,l-1} & 1 & a_{k,l+1} & \dots & a_{kn} \\ a_{11} & a_{12} & \dots & & 0 & & & a_{1n} \\ \vdots & \vdots & \dots & & 0 & & & \vdots \\ a_{k-1,1} & a_{k-1,2} & \dots & & 0 & & & a_{k-1,n} \\ a_{k+1,1} & a_{k+1,2} & \dots & & 0 & & & a_{k+1,n} \\ \vdots & \vdots & \dots & & 0 & & & \vdots \\ a_{n1} & a_{n2} & \dots & & 0 & & & a_{nn} \end{vmatrix}$$

Now we can do $l - 1$ column exchange to move the one into the first column. This gives a cycle $(1, 2, \dots, l)$ and $\text{sign} = (-1)^{l-1}$

$$= \sum_{k=1}^n a_{kl} (-1)^{k-1} (-1)^{l-1} \begin{vmatrix} 1 & a_{k1} & a_{k2} & \dots & a_{k,l-1} & a_{k,l+1} & \dots & a_{kn} \\ 0 & a_{11} & a_{12} & \dots & a_{1,l-1} & a_{1,l+1} & \dots & a_{1n} \\ 0 & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & a_{2,n} \\ 0 & a_{k-1,1} & a_{k-1,2} & \dots & a_{k-1,l-1} & a_{k-1,l+1} & \dots & a_{k-1,n} \\ 0 & a_{k+1,1} & a_{k+1,2} & \dots & a_{k+1,l-1} & a_{k+1,l+1} & \dots & a_{k+1,n} \\ 0 & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & a_{2,n} \\ 0 & a_{n1} & a_{n2} & \dots & a_{nl-1} & a_{nl+1} & \dots & a_{nn} \end{vmatrix}$$

where the k -th row and l -th column is removed

$$= \sum_{k=1}^n (-1)^{k+l} a_{kl} \det A_{kl}$$

□

Example 7.13. Let $A = (a_{kl})_{\substack{1 \leq k \leq n \\ 1 \leq l \leq n}} = (-1)^{k+l}$.

$$A = \begin{pmatrix} +1 & -1 & +1 \\ -1 & +1 & -1 \\ +1 & -1 & +1 \end{pmatrix}$$

Theorem 7.9. $\hat{a}_{kl} := (-1)^{k+l} \det A_{lk}$ is called cofactor.

$$\hat{A} = [\hat{a}_{kl}]_{k,l=1}^n$$

is called complementary matrix or adjugate matrix of A .

$$\begin{aligned} \hat{a}_{kl} &= (-1)^{k+l} \det (\text{the matrix without row } l \text{ and column } k) \\ &= (-1)^{k+l} \det A_{lk} = \frac{\partial}{\partial a_{lk}} \det A \end{aligned}$$

Then it holds that

$$A^{-1} = \frac{1}{\det A} \hat{A}$$

Proof. Show that $\hat{A} \cdot A = I \cdot \det(A)$. Let $B = \hat{A} \cdot A$.

$$b_{kl} = \sum_{i=1}^n \hat{a}_{ki} \cdot a_{il} = \sum_{i=1}^n (-1)^{k+i} \det A_{ik} \cdot a_{il}$$

Case $k = l$

$$b_{ll} = \sum_{i=1}^n (-1)^{l+i} \det A_{il} \cdot a_{il} \quad \underbrace{\quad}_{\substack{\text{Laplace expansion} \\ \text{with } l\text{-th column}}} = \det A$$

Case $k \neq l$ Without loss of generality: $k < l$.

$$\begin{aligned} b_{kl} &= \sum_{i=1}^n \det(A_{ik}) (-1)^{k+i} a_{il} \\ &= \det \begin{bmatrix} a_{11} & \dots & a_{1l} & \dots & a_{1l} & \dots & a_{1n} \\ \vdots & & \vdots & & \vdots & & \vdots \\ a_{n1} & \dots & a_{nl} & \dots & a_{nl} & \dots & a_{nn} \end{bmatrix} \\ &\quad \underbrace{\quad}_{\substack{\text{two equal} \\ \text{columns}}} = 0 \end{aligned}$$

(i.e. matrix A with k -th column replaced by l -th column) expanded by k -th row.

$$\det A = \sum_{i=1}^n (-1)^{k+i} \det(A_{ik}) \cdot a_{ik}$$

\tilde{A} = (matrix A replacing k -th column with l -th column)

$$\det \tilde{A} = \sum_{i=1}^n (-1)^{k+i} \det(A_{ik}) \cdot a_{il}$$

□

Example 7.14 (Small inverse matrices). Let $n = 2$.

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} = \frac{1}{ad - bc} \cdot \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$\hat{a}_{11} = (-1)^{1+1} \cdot \det A_{11} \quad \hat{a}_{21} = (-1)^{2+1} \cdot \det A_{12}$$

$$\hat{a}_{12} = (-1)^{1+2} \cdot \det A_{21} \quad \hat{a}_{22} = (-1)^{2+2} \cdot \det A_{22}$$

Remark 7.12 (Cayley 1855).

$$A^{-1} = \frac{1}{\nabla} \begin{bmatrix} \partial_a \nabla & \partial_c \nabla \\ \partial_b \nabla & \partial_d \nabla \end{bmatrix}$$

$$\nabla f = \begin{bmatrix} \frac{\partial f}{\partial x_1} \\ \vdots \\ \frac{\partial f}{\partial x_n} \end{bmatrix}$$

Example 7.15. Let $n = 3$.

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}^{-1} = \frac{1}{\det(A)} \begin{bmatrix} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} & -\begin{vmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{vmatrix} & \begin{vmatrix} a_{12} & a_{13} \\ a_{22} & a_{23} \end{vmatrix} \\ -\begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} & \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} & -\begin{vmatrix} a_{11} & a_{13} \\ a_{21} & a_{23} \end{vmatrix} \\ \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} & -\begin{vmatrix} a_{11} & a_{12} \\ a_{31} & a_{32} \end{vmatrix} & \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} \end{bmatrix}$$

Corollary 7.12. Let $A \in \mathbb{Z}^{n \times n}$. If $\det A = 1 \implies A^{-1} \in \mathbb{Z}^{n \times n}$.

Let $A \in \mathbb{Z}^{n \times n}$ and $\det A = 1$. Let $B \in \mathbb{Z}^{n \times n}$ and $\det B = 1$.

$$\implies \det(A \cdot B) = 1 \quad \implies \det(A^{-1}) = 1$$

Definition 7.9. Integer matrices with $\det = 1$ define a group called special linear group.

$$\text{SL}(n, \mathbb{Z}) = \{A \in \mathbb{Z}^{n \times n} \mid \det A = 1\}$$

Or in general for a ring R :

$$\text{SL}(n, R) = \{A \in R^{n \times n} \mid \det A = 1\}$$

Theorem 7.10 (Cramer's Rule). Gabriel Cramer (1704–1752)

Show by Cramer in 1750, by McLaurin 1748 for $n \leq 3$.

Let A be a regular matrix with column vectors a_1, \dots, a_n . Then the solution $Ax = b$ ($\implies x = A^{-1}b$ has a unique solution) is given by

$$x_i = \frac{\Delta(a_1, \dots, a_{i-1}, b, a_{i+1}, \dots, a_n)}{\Delta(a_1, \dots, a_n)}$$

$$= \frac{\det \left(\begin{bmatrix} a_1 & \dots & a_{i-1} & b & a_{i+1} & \dots & a_n \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \end{bmatrix} \right)}{\det A}$$

$n + 1$ determinants of form $n \times n$. In practice infeasible except for small matrices.

Geometrical proof for $n = 2$.

$$A = \begin{pmatrix} a_1 & a_2 \\ \vdots & \vdots \end{pmatrix}$$

$$Ax = b \quad a_1 \cdot x + a_2 \cdot x_2 = b$$

$$\Delta(a_1, a_2) = A(a_1, a_2)$$

where A is the area function.

TODO drawing parallelogram

$$\Delta(b, a_2) = A(b, a_2) = \Delta(x_1 \cdot a_1, a_2) = x_1 \cdot \Delta(a_1, a_2)$$

$$\implies x_1 = \frac{\Delta(b, a_2)}{\Delta(a_1, a_2)}$$

□

Generic proof. Let $x = A^{-1} \cdot b = \frac{1}{\det A} \cdot \hat{A} \cdot b$.

$$\begin{aligned} x_i &= \frac{1}{\det A} \cdot \sum_{k=1}^n \hat{a}_{ik} b_k \\ &= \frac{1}{\det A} \sum_{k=1}^n (-1)^{i+k} \det A_{ki} \cdot b_k \\ &\quad \underbrace{=}_{\substack{\text{see proof of} \\ \text{Laplace expansion}}} \frac{1}{\det A} \sum_{k=1}^n \Delta(a_1, \dots, a_{i-1}, e_k, a_{i+1}, \dots, a_n) b_k \\ &= \frac{\Delta(a_1, \dots, a_{i-1}, b, a_{i+1}, \dots, a_n)}{\det A} \end{aligned}$$

□

Example 7.16.

$$2x_1 + x_2 = 7$$

$$x_1 - 3x_2 = 0$$

$$A = \begin{bmatrix} 2 & 1 \\ 1 & -3 \end{bmatrix}$$

$$\det(A) = 2 \cdot (-3) - 1 = -7$$

$$x_1 = -\frac{1}{7} \begin{vmatrix} 7 & 1 \\ 0 & -3 \end{vmatrix} = 3$$

$$x_2 = -\frac{1}{7} \begin{vmatrix} 2 & 7 \\ 1 & 0 \end{vmatrix} = 1$$

Remark 7.13. For large n (hence $n \geq 4$), Cramer's Rule is impractical (tiresome and unstable). But it helps with theoretical considerations.

1. The map $A \mapsto \det A$ is continuous and differentiable.
2. if $\det A \neq 0 \implies$ the set of invertible matrices is open⁴
3. The solution of system $Ax = b$ depends continuously on a_{ij} and b_i ⁵

8 Inner products

Definition 8.1.

$$\mathbb{R}^3 : \left\| \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \right\| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

By Pythagorem Theorem

Pythagorem Theorem. Claim: $a^2 + b^2 = c^2$

TODO




Figure 3: Proof construction of the Pythagorem Theorem

□

⁴Hence for all invertible A , there exists some neighborhood such that all matrices in this neighborhood are invertible.

$$\text{e.g. } d(A, B) = \max_{i,j} |a_{ij} - b_{ij}|$$

⁵ This justifies why Computational Mathematics (dt. Numerik) is practical and interesting

$$\forall \varepsilon \exists \delta : d(b, b') < \delta \implies d(x, x') < \varepsilon$$

↓ This lecture took place on 2018/03/21.

The norm is given by

$$\left\| \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \right\| = \sqrt{a_1^2 + a_2^2 + a_3^2}$$

Definition 8.2 (Scalar product in $\mathbb{R}^2/\mathbb{R}^3$).

$$\langle a, b \rangle := \|a\| \cdot \|b\| \cdot \cos \theta$$

where θ is the angle between vector a and b .

Theorem 8.1.

$$\langle a, a \rangle = \|a\|^2$$

Remark 8.1. Recall that

$$\cos 0 = 1 \quad \cos \frac{\pi}{2} = 0 \quad \cos \pi = -1 \quad \cos \frac{3}{2}\pi = 0$$

$$\sin 0 = 0 \quad \sin \frac{\pi}{2} = 1 \quad \sin \pi = 0 \quad \sin \frac{3}{2}\pi = -1$$

$$\sin \theta = \cos(\theta - \frac{\pi}{2})$$

$$\cos(\pi - \theta) = -\cos(\theta)$$

$$\sin(-\theta) = -\sin(\theta)$$

$$\sin(\pi - \theta) = \sin(\theta)$$

$$\sin(-\theta) = -\sin(\theta)$$

Theorem 8.2. 1. $\langle a, a \rangle = \|a\|^2$

$$2. \langle a, a \rangle = 0 \iff a = 0$$

$$3. \langle a, b \rangle = 0 \iff a = 0 \vee b = 0 \vee \theta = \frac{\pi}{2} \vee \theta = \frac{3}{2}\pi, \text{ hence orthogonal}$$

$$4. \langle a, b \rangle > 0 \iff \text{acute angle}$$

$$5. \langle a, b \rangle < 0 \iff \text{obtuse angle (dt. stumpf)}$$

Theorem 8.3. 1. $\langle a, b \rangle = \langle b, a \rangle$

$$2. \langle \lambda a, b \rangle = \lambda \cdot \langle a, b \rangle = \langle a, \lambda \cdot b \rangle$$

$$3. \langle a + b, c \rangle = \langle a, c \rangle + \langle b, c \rangle$$

Thus, linear in a and b . Thus, bilinear.

Proof. 2. Assume $\lambda > 0$. Angle stays the same.

$$\langle \lambda a, b \rangle = \|\lambda a\| \cdot \|b\| \cdot \cos \theta = \lambda \cdot \|a\| \cdot \|b\| \cdot \cos \theta$$

Assume $\lambda < 0$. θ becomes $\pi - \theta$.

$$\langle \lambda a, b \rangle = \|\lambda a\| \cdot \|b\| \cdot \cos(\pi - \theta) = |\lambda| \cdot \|a\| \cdot \|b\| \cdot (-\cos(\theta)) = \lambda \cdot \|a\| \cdot \|b\|$$

3. Let $\|c\| = 1$. $\langle a, c \rangle = \|a\| \cdot \cos \theta$.

$$\langle a + b, c \rangle = \langle a, c \rangle + \langle b, c \rangle$$

Projections will add up.

In the generic case:

$$\begin{aligned} \langle a + b, c \rangle &= \left\langle a + b, \|c\| \cdot \frac{c}{\|c\|} \right\rangle \\ &= \underbrace{\|c\|}_{\text{by (2.)}} \left\langle a + b, \frac{c}{\|c\|} \right\rangle \end{aligned}$$

□

Theorem 8.4.

$$\left\langle \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix}, \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} \right\rangle = a_1 b_1 + a_2 b_2 + a_3 b_3$$

Proof.

$$\begin{aligned} \langle a, b \rangle &= \langle a_1 e_1 + a_2 e_2 + a_3 e_3, b \rangle \\ &= a_1 \langle e_1, b \rangle + a_2 \langle e_2, b \rangle + a_3 \langle e_3, b \rangle \\ &= a_1 b_1 + a_2 b_2 + a_3 b_3 \\ \langle e_i, b \rangle &= \langle e_i, b_1 e_1 + b_2 e_2 + b_3 e_3 \rangle \\ &= b_1 \langle e_i, e_1 \rangle + b_2 \langle e_i, e_2 \rangle + b_3 \langle e_i, e_3 \rangle \\ &= b_1 \delta_{i1} + b_2 \delta_{i2} + b_3 \delta_{i3} \quad \text{with } \delta \text{ as Kronecker delta} \\ &= b_i \end{aligned}$$

□

In this chapter, we will talk about vector spaces in which we will discuss scalar products with properties 1–3 from Theorem 8.3.

$$\text{in } \mathbb{R}^n : \quad \langle x, y \rangle = \sum_{i=1}^n x_i y_i$$

$$\text{in } V \subseteq \mathbb{R}^\infty : \quad \langle x, y \rangle = \sum_{i=1}^{\infty} x_i y_i$$

if convergent! For this space, $(e_i)_{i \in \mathbb{N}}$ is a basis.

$$\text{in } C[a, b] \quad \langle f, g \rangle = \int f(x)g(x) dx$$

is the Delta function.

Or better: $(\sin nx)_{n \in \mathbb{N}} \cup (\cos nx)_{n \in \mathbb{N}}$.

$$\int_0^{2\pi} \sin(nx) \cos(mx) dx = 0 \forall m, n$$

$$\int_0^{2\pi} \sin(nx) \sin(mx) dx = 0 \text{ if } m \neq n$$

1768/03/21 J. Fourier

Theorem 8.5 (1822 Fourier). *Every function f in $[0, 2\pi]$ can be denoted as*

$$f(x) = \sum_{n=0}^{\infty} a_n \cos(nx) + \sum_{n=1}^{\infty} b_n \sin(nx)$$

$$a_n = \langle f, \cos(nx) \rangle = \int_0^{2\pi} f(x) \cos(nx) dx$$

$$b_n = \langle f, \sin(nx) \rangle = \int_0^{2\pi} f(x) \sin(nx) dx$$

This theorem cannot be proven, because it depends on the definition of “function”. The answer to the question, which functions satisfy this theorem, is an open research topic.

8.1 Law of cosines

Theorem 8.6 (Law of cosines). *In German, “Kosinussatz”.*

$$c^2 = a^2 + b^2 - 2ab \cos \gamma$$

$$\begin{aligned} \|\vec{c}\|^2 &= \|\vec{b} - \vec{a}\|^2 \\ &= \langle \vec{b} - \vec{a}, \vec{b} - \vec{a} \rangle \\ &= \langle \vec{b}, \vec{b} \rangle - \langle \vec{a}, \vec{b} \rangle - \langle \vec{b}, \vec{a} \rangle + \langle \vec{a}, \vec{a} \rangle \\ &= \|\vec{b}\|^2 - 2\|\vec{a}\| \|\vec{b}\| \cos \gamma + \|\vec{a}\|^2 \end{aligned}$$

$$\|a\| \cdot \|b\| \cdot \sin \theta = \text{area of the spanned parallelogram}$$

How to find an orthogonal vector?

Remark 8.2 (Orthogonal vector in \mathbb{R}^2). Find \vec{b} such that $\langle \vec{a}, \vec{b} \rangle = 0$, $a_1 b_1 + a_2 b_2 = 0$. For example, $b_1 = a_2$ and $b_2 = -a_1$.

$$\vec{a} = \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \quad \vec{b} = \begin{pmatrix} a_2 \\ -a_1 \end{pmatrix}$$

8.2 Outer product

Definition 8.3. Called outer product (only in \mathbb{R}^3) or cross product.

Let $a, b \in \mathbb{R}^3$ and $a \times b$ is the vector which

1. $\|a \times b\| = \|a\| \cdot \|b\| \cdot \sin \theta$ is the area of the spanned parallelogram.
2. $a \times b \perp a$ and $a \times b \perp b \iff \langle a \times b, a \rangle = 0$ and $\langle a \times b, b \rangle = 0$
3. $(a, b, a \times b)$ is clockwise.

When does $a \times b = 0$ hold? $a = 0, b = 0, \sin \theta = 0$, hence $\theta = 0 \vee \theta = \pi$

$$\iff a, b \text{ are linear independent}$$

Theorem 8.7. • $b \times a = -a \times b$


- $(\lambda a) \times b = \lambda(a \times b) = a \times (\lambda b)$
- $(a + b) \times c = a \times c + b \times c$

Proof. • Orientation swaps. Consider the right-hand rule. If you assign b to your index finger, a to your middle finger, you retrieve direction $b \times a$ with the thumb. Now assign a (index finger) and b (middle finger) and you retrieve the opposite direction, namely $-(b \times a)$.

- If $\lambda > 0$, it follows immediate. If $\lambda < 0$, lengths stay the same, but orientation swaps.
- If $c = 0$, it is trivial. If $c \neq 0$, E is the plane orthogonal to c . a' and b' are projections of a and b to E .

1. $(a + b)' = a' + b'$
2. $a \times c = a' \times c$.

$$\begin{aligned} \|a \times c\| &= \|a\| \|c\| \cdot \sin \theta \\ &= \|a'\| \cdot \|c\| \\ &= \|a' \times c\| \end{aligned}$$



- Orientation of $a \times c$ and $a' \times c$ is the same
- The plane, spanned by c and a , is also spanned by c and a'

$$\|a'\| = \|a\| \cdot \underbrace{\cos\left(\frac{\pi}{2} - \theta\right)}_{=\sin \theta}$$

Hence,

$$(a + b) \times c = (a + b)' \times c = (a' + b') \times c \stackrel{!}{=} a' \times c + b' \times c = a \times c + b \times c$$

$$(a' + b') \times c = a' \times c + b' \times c$$

rotated by 90° multiplied by $\|c\|$

$$a' \times c = a'$$

rotated by 90° multiplied by $\|c\|$

$$a' \times c + b' \times c = (a' + b') \times c$$

The relation $u + v = w$ will be preserved under rotation by 90° and multiplication with λ .

□

Corollary 8.1. *The cross product is a map of $\mathbb{R}^3 \times \mathbb{R}^3 \rightarrow \mathbb{R}^3$ such that*

- *bilinear*
- *antisymmetrical, $a \times b = -b \times a$*
- *$e_1 \times e_2 = e_3, e_2 \times e_3 = e_1, e_3 \times e_1 = e_2$*

$$e_i \times e_j = e_k \cdot \text{sign } \pi \quad \pi = \begin{pmatrix} 1 & 2 & 3 \\ i & j & k \end{pmatrix}$$

Corollary 8.2.

$$\begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} \times \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix} = \begin{bmatrix} a_2 b_3 - a_3 b_2 \\ a_3 b_1 - a_1 b_3 \\ a_1 b_2 - a_2 b_1 \end{bmatrix} = \begin{bmatrix} \begin{vmatrix} a_2 & b_2 \\ a_3 & b_3 \end{vmatrix} \\ - \begin{vmatrix} a_1 & b_1 \\ a_3 & b_3 \end{vmatrix} \\ \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} \end{bmatrix} \underbrace{=}_{\substack{\text{by Laplace expansion along} \\ \text{the third column}^6}} \begin{vmatrix} a_1 & b_1 & e_1 \\ a_2 & b_2 & e_2 \\ a_3 & b_3 & e_3 \end{vmatrix}$$

Proof.

$$\begin{aligned} & (a_1 e_1 + a_2 e_2 + a_3 e_3) \times (b_1 e_1 + b_2 e_2 + b_3 e_3) \\ &= a_1 b_1 e_1 \times e_1 + a_1 b_2 e_1 \times e_2 + a_1 b_3 e_1 \times e_3 \\ &+ a_2 b_1 e_2 \times e_1 + a_2 b_2 e_2 \times e_2 + a_2 b_3 e_2 \times e_3 \\ &= a_3 b_1 e_3 \times e_1 + a_3 b_2 e_3 \times e_2 + a_3 b_3 e_3 \times e_3 \\ &= a_1 b_2 e_3 - a_1 b_3 e_2 - a_2 b_1 e_3 + a_2 b_3 e_1 + a_3 b_1 e_2 - a_3 b_2 e_1 \\ &= (a_2 b_3 - a_3 b_2) e_1 + (a_3 b_1 - a_1 b_3) e_2 + (a_1 b_2 - a_2 b_1) e_3 \end{aligned}$$

□

Theorem 8.8.

$$\langle a \times b, c \rangle = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

This corresponds to the volume of the spanned parallelepiped (dt. “Spat”). $\|a \times b\|$ is the area of the parallelogram and $\|c\|$ its height.

Equivalently, $\begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix}$ is the area of the parallelogram.


Proof. Laplace expansion in third column

□

Example 8.1. *Let planes in \mathbb{R}^3 be given.*

$$E = \{x_0 + \lambda a + \mu b \mid \lambda, \mu \in \mathbb{R}\}$$

$$c = a \times b = \left\{ x \in \mathbb{R}^3 \mid x - x_0 \perp c \right\} = \left\{ x \in \mathbb{R}^3 \mid \langle x - x_0, c \rangle = 0 \right\}$$



8.3 Inner products and positive definiteness

From now on \mathbb{K} will be \mathbb{R} or \mathbb{C} .

Definition 8.4. An inner product on a vector space V is a map

$$V \times V \rightarrow \mathbb{K}$$

$$(x, y) \mapsto \langle x, y \rangle$$

1. $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle \forall x, y, z \in V$
2. $\langle \lambda x, y \rangle = \lambda \langle x, y \rangle \forall \lambda \in \mathbb{K} \forall x, y \in V$
3. $\langle y, x \rangle = \overline{\langle x, y \rangle} \forall x, y \in V$

where $\overline{\langle x, y \rangle}$ denotes the complex conjugate.

$$\langle x, \lambda y \rangle \underbrace{= \overline{\langle \lambda y, x \rangle}}_{\text{by (3)}} = \underbrace{\overline{\lambda \langle y, x \rangle}}_{\text{by (2)}} = \overline{\lambda} \overline{\langle y, x \rangle} = \overline{\lambda} \langle x, y \rangle$$

Linear in x , semi-linear in y . Sesquilinear⁷.

⁷In Latin, sesqui means 1.5

In physics, the notation is different:

$$\begin{aligned}\langle x|y\rangle & \quad \langle \lambda x|y\rangle = \bar{\lambda} \langle x|y\rangle & \quad \langle x|\lambda y\rangle = \lambda \langle x|y\rangle \\ |y\rangle \dots \text{ket} & \quad \langle x| \dots \text{bra} \\ \langle x|y\rangle & \quad \text{bracket}\end{aligned}$$

The inner product is called positive-semidefinite, if

$$\langle x, x \rangle \geq 0 \forall x \in X$$

if additionally $\langle x, x \rangle = 0 \iff x = 0$, then \langle, \rangle is called positive definite.

↓ This lecture took place on 2018/04/09. Easter holidays finished.

Lemma 8.1. 1. $\langle x, y + z \rangle = \langle x, y \rangle + \langle x, z \rangle$

2. $\langle x, \lambda y \rangle = \bar{\lambda} \cdot \langle x, y \rangle$

3. $\langle x, 0 \rangle = 0$

Definition 8.5. An inner product is positive semidefinite, if $\langle x, x \rangle \geq 0$. Is positive definite, if $\langle x, x \rangle > 0$ for all $x \neq 0$. Is negative definite, if $\langle x, x \rangle < 0$ for all $x \neq 0$. Is indefinite, if neither positive nor negative semidefinite.

A positive definite inner product is called scalar product. A positive definite inner product is in Hermitian form, if $\mathbb{K} = \mathbb{C}$. A positive definite inner product is also called unitary product, if $\mathbb{K} = \mathbb{C}$.

So quadratic form over \mathbb{R} and Hermitian form over \mathbb{C} .

Remark 8.3. For example, the expression $\langle x, x \rangle > 0$ requires that $x \in \mathbb{R}$, but we defined the inner product over \mathbb{C} as well. In Euclidean spaces, $\langle x, x \rangle \in \mathbb{R}$ anyhow, but more generally, the condition should require the absolute value: $|\langle x, x \rangle| > 0$.

Example 8.2. • Let $V = \mathbb{R}^n$.

$$\left\langle \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix} \right\rangle = \sum_{i=1}^n x_i y_i$$

Let $V = \mathbb{C}^n$.

$$\left\langle \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, \begin{pmatrix} y_1 \\ \vdots \\ y_n \end{pmatrix} \right\rangle = \sum_{i=1}^n x_i \bar{y}_i \implies \langle x, x \rangle = \sum_{i=1}^n x_i \bar{x}_i = \sum_{i=1}^n |x_i|^2 \geq 0$$

→ positive definite.

- Another example: let $A \in \mathbb{R}^{n \times n}$. Let $x, y \in \mathbb{R}^n$.

$$\begin{aligned}\langle x, y \rangle_A &= x^t \cdot A \cdot y \quad \text{is bilinear} \\ &= \sum_{i=1}^n x_i \sum_{j=1}^n a_{ij} y_j = \sum_{i,j=1}^n a_{ij} x_i y_j\end{aligned}$$

hence $\langle x, y \rangle_A = \langle y, x \rangle_A$. It must hold that

$$\sum_{i,j=1}^n a_{ij} x_i y_j = \sum_{i,j=1}^n a_{ij} y_i x_j \quad \forall x, y$$

We let $x = e_k$ and $y = e_l$.

$$\implies a_{kl} = a_{lk} \quad \forall k, l$$

Hence $A = A^T$. A is symmetrical.

Let $A \in \mathbb{C}^{n \times n}$. Let $x, y \in \mathbb{C}^n$.

$$\langle x, y \rangle_A = \sum_{i=1}^n \sum_{j=1}^n x_i a_{ij} - i j \overline{y_j}$$

$$\langle x, y \rangle_A = \langle y, x \rangle_A \quad \forall x, y$$

$$\iff A^T = \overline{A} \quad \text{is in Hermitian form}$$

$$a_{ji} = \overline{a_{ij}} \quad \forall i, j$$

•

$$V = C[a, b] = \{f : [a, b] \rightarrow \mathbb{K} \text{ continuous}\}$$

$$\langle f, g \rangle = \int_a^b f(t) \overline{g(t)} dt \quad \text{is a scalar product}$$

$$\langle f, f \rangle = \int_a^b |f(t)|^2 dt \geq 0$$

- Consider $V = l_2$ (\mathbb{R}^∞ would be too large) where $l_2 = \{(x_n)_{n \in \mathbb{N}} \mid x_n \in \mathbb{R}, \sum_{n=1}^\infty x_n^2 < \infty\}$.

$$\langle x, y \rangle = \sum_{n=1}^\infty x_n y_n \quad \text{is a scalar product}$$

Does it converge? This is not obvious.

Fourier claimed that this example (4) and example (3) are the same. He claimed every function can be written as $f(x) = \sum_{n=0}^\infty a_n e^{inx}$.

$$x \cdot x = \langle x, x \rangle = \sum_{i=1}^n x_i^2 = \|x\|^2$$

Definition 8.6. Let V be a vector space. A norm on V is a map $\|\cdot\| : V \rightarrow [0, \infty[$ such that

1. $\|x\| \geq 0$ and $\|x\| = 0 \iff x = 0$
2. $\|\lambda \cdot x\| = |\lambda| \cdot \|x\| \quad \forall \lambda \in K, \forall x \in V$
3. $\|x + y\| \leq \|x\| + \|y\|$ is the triangle inequality

Remark 8.4. Every norm is a metric with $d(x, y) = \|x - y\|$.

d is translation invariant. $d(x + x_0, y + x_0) = d(x, y)$. This is compatible to a vector space.

In a black hole (\rightarrow physics), you have a different metric in every point (Riemannian geometry): $\langle x, y \rangle_{A(x,y)}$.

Example 8.3. Let $V = \mathbb{R}^n$.

- $\|x\|_2 = \left(\sum_{i=1}^n x_i^2\right)^{\frac{1}{2}}$ is called euclidean norm.
- $\|x\|_1 = \sum_{i=1}^n |x_i|$ is called l^1 norm or Manhattan norm.
- $\|x\|_\infty = \max \{|x_i| \mid i = 1, \dots, n\}$

Let $V = C[a, b]$.

•

$$\|f\|_1 = \int_a^b |f(t)| dt$$

L^1 -norm, gives rise to the Lebesgue integral.

•

$$\|f\|_\infty = \max_{t \in [a,b]} |f(t)| \quad \text{is a } L^\infty\text{-norm}$$

•

$$\|f\|_2 = \left(\int_a^b |f(t)|^2 dt \right)^{\frac{1}{2}}$$

Theorem 8.9. Let \langle, \rangle be a scalar product in V (hence, positive-definite inner product). Then $\|x\| = \sqrt{\langle x, x \rangle}$ is a norm on V .

Proof. • $\|x\| \geq 0, \|x\| = 0 \iff \langle x, x \rangle = 0 \iff x = 0$

- $\|\lambda x\| = \sqrt{\langle \lambda x, \lambda x \rangle} = \sqrt{\lambda \cdot \bar{\lambda} \cdot \langle x, x \rangle} = \sqrt{\lambda^2 \cdot \langle x, x \rangle} = |\lambda| \cdot \sqrt{\langle x, x \rangle}$
- Triangle inequality

□

8.4 Cauchy-Bunyakovsky-Schwarz inequality

Lemma 8.2 (Cauchy-Bunyakovsky-Schwarz inequality). *Cauchy (1789–1857) for \mathbb{R}^n , Bunyakovsky (1804–1889) for infinite dimensions, Schwarz (1843–1921) generically.*

$$|\langle x, y \rangle| \leq \|x\| \cdot \|y\|$$

Hence, l^2 if $\sum_{n=1}^{\infty} x_n^2 < \infty$ and $\sum_{n=1}^{\infty} y_n^2 < \infty$. $\langle x, x \rangle < \infty$ and $\langle y, y \rangle < \infty$.

$$\Rightarrow \sum x_n y_n \leq \sqrt{\sum x_n^2} \sqrt{\sum y_n^2}$$

If $|\langle x, y \rangle| = \|x\| \cdot \|y\| \iff x, y$ are linear dependent.

Proof. Now we can continue with part 3 of the proof of Theorem 8.9. Triangle inequality:

$$\begin{aligned} \|x + y\|^2 &= \langle x + y, x + y \rangle \\ &= \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle \\ &\leq \|x\|^2 + 2|\langle x, y \rangle| + \|y\|^2 \\ &\leq \|x\|^2 + 2\|x\|\|y\| + \|y\|^2 \\ &= (\|x\| + \|y\|)^2 \end{aligned}$$

□

Proof of CBS inequality, Lemma 8.2.

Case 1: $y = 0$ trivial

Case 2: $y \neq 0$ Let $\lambda \in \mathbb{K}$ be arbitrary.

$$\begin{aligned} 0 &\leq \langle x - \lambda y, x - \lambda y \rangle \\ &= \langle x, x \rangle - \langle x, \lambda y \rangle - \langle \lambda y, x \rangle + \langle \lambda y, \lambda y \rangle \\ &= \langle x, x \rangle - \bar{\lambda} \langle x, y \rangle - \lambda \langle y, x \rangle + |\lambda|^2 \langle y, y \rangle \end{aligned}$$

This holds for all λ , hence also for $\lambda = \frac{\langle x, y \rangle}{\langle y, y \rangle}$. Because $y \neq 0 \implies \langle y, y \rangle > 0$, we can divide.

$$\begin{aligned}
&= \langle x, x \rangle - \frac{\overline{\langle x, y \rangle}}{\langle y, y \rangle} \cdot \langle x, y \rangle - \frac{\langle x, y \rangle}{\langle y, y \rangle} \cdot \langle y, x \rangle + \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle^2} \cdot \langle y, y \rangle \\
&= \langle x, x \rangle - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} - \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} + \frac{|\langle x, y \rangle|^2}{\langle y, y \rangle} \\
&= \|x\|^2 - \frac{|\langle x, y \rangle|^2}{\|y\|^2} \\
&\implies \|x\|^2 \cdot \|y\|^2 - |\langle x, y \rangle|^2 \geq 0
\end{aligned}$$

□

Alternative proof of CBS inequality in \mathbb{R}^n .

$$\begin{aligned}
0 &\leq \sum_{i=1}^n \sum_{j=1}^n (x_i y_j - x_j y_i)^2 \\
&= \sum_{i,j=1}^n (x_i^2 y_j^2 - 2x_i y_j x_j y_i + x_j^2 y_i^2) \\
&= \sum_{i,j} x_i^2 y_j^2 - 2 \sum_{i,j} x_i x_j y_i y_j + \sum_{i,j} x_j^2 y_i^2 \\
&= 2 \sum_i x_i^2 \sum_j y_j^2 - 2 \sum_i x_i y_i \sum_j x_j y_j \\
&= 2 \|x\|^2 \|y\|^2 - 2 \langle x, y \rangle^2 \\
&\leadsto \|x\|^2 \|y\|^2 = \langle x, y \rangle^2 + \frac{1}{2} \sum_i \sum_j (x_i y_j - x_j y_i)^2
\end{aligned}$$

So for $n = 3$, $\|x\|^2 \|y\|^2 = \langle x, y \rangle^2 + \|x \times y\|^2$. Hence, equality is given iff x and y are linear dependent.

In the general case: If $|\langle x, y \rangle| = \|x\| \cdot \|y\|$. From the proof, it follows that $\exists \lambda : \langle x - \lambda y, x - \lambda y \rangle = 0$

$$\implies x - \lambda y = 0 \implies x, y \text{ are linear independent}$$

□

Theorem 8.10. Let V be a vector space over $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . Let $B = \{b_1, \dots, b_n\}$ is a basis. \langle, \rangle is an inner product. What does \langle, \rangle look like in regards of the coordinate?

There exists a unique matrix A in Hermitian form (hence, $a_{ij} = \overline{a_{ji}}$, $A = \overline{A^T}$) such that $\forall x, y \in V : \langle x, y \rangle = \Phi_B(x)^T \cdot A \cdot \overline{\Phi_B(y)}$. If \langle, \rangle is positive definite, A is regular.

Remark 8.5.

$$\langle x, y \rangle = \sum x_i \overline{y_i}$$

corresponds to $A = I$.

$$x^T \cdot I \cdot \overline{y} = x^T \cdot \overline{y}$$

How about $A = -I$.

$$\langle x, y \rangle_A = - \sum x_i \overline{y_i}$$

This is not a scalar product (because of negative definiteness).

Proof. Let $x = \sum_{i=1}^n \xi_i b_i$, $y = \sum_{j=1}^n \eta_j b_j$.

$$\begin{aligned} \langle x, y \rangle &= \left\langle \sum_{i=1}^n \xi_i b_i, \sum_{j=1}^n \eta_j b_j \right\rangle \\ &= \sum_{i=1}^n \xi_i \sum_{j=1}^n \overline{\eta_j} \underbrace{\langle b_i, b_j \rangle}_{=: a_{ij} \text{ is unique } a_{ij} = \langle b_i, b_j \rangle} \\ &= \sum_{i=1}^n \sum_{j=1}^n \xi_i a_{ij} \overline{\eta_j} \\ &= \xi^T \cdot A \cdot \overline{\eta} \\ &= \Phi_B(x)^T \cdot A \cdot \Phi_B(y) \\ a_{ji} &= \langle b_j, b_i \rangle = \overline{\langle b_i, b_j \rangle} = \overline{a_{ij}} \end{aligned}$$

Show: If \langle, \rangle is positive definite, then A is regular. It suffices to show that $\ker A = \{0\}$.

Assume: $A \cdot \xi = 0 \implies \xi^T \cdot A \cdot \xi = 0$. Let $x = \sum_{i=1}^n \xi_i b_i \implies \langle x, x \rangle = 0 \implies x = 0 \implies \xi = \Phi_B(x) = 0$ \square

Definition 8.7. Let $A \in \mathbb{C}^{n \times n}$. The matrix $A^* := \overline{A^T}$ ($(A^*)_{ij} = \overline{a_{ji}}$) is called conjugate transpose.

A is called self-adjoint if $A = A^*$ (dt. selbst-adjungiert). A is called symmetrical if $A = \overline{A}$ and $\mathbb{K} = \mathbb{R}$ or A is called Hermitian if $A = A^*$ and $\mathbb{K} = \mathbb{C}$.

$A = A^*$ is called (positive/negative) (semidefinite/definite) if the corresponding sesquilinear form

$$\langle \xi, \eta \rangle_A = \xi^T \cdot A \cdot \overline{\eta}$$

Hence, $\xi^T A \bar{\xi} \geq 0 \forall \xi \neq 0$ is positive definite, has the corresponding property or $\xi^T A \bar{\xi} > 0 \forall \xi \neq 0$ is positive semidefinite, has the corresponding property.

$\xi^T A \bar{\xi} \leq 0 \forall \xi \neq 0$ is negative definite or $\xi^T A \bar{\xi} < 0 \forall \xi \neq 0$ is negative semidefinite.

If $\exists \xi : \xi^T A \bar{\xi} > 0$ and $\exists \eta : \eta^T A \bar{\eta} < 0$, then A is called indefinite.

↓ This lecture took place on 2018/04/11.

Inner product: $\langle x, y \rangle$

- $\forall x : \langle x, x \rangle \geq 0$ positive semi-definite
- $\forall x \neq 0 : \langle x, x \rangle > 0$ positive definite

in regards of basis b_1, \dots, b_n .

$$\langle x, y \rangle = \sum a_{ij} \xi_i \bar{\eta}_j$$

$$a_{ij} = \langle b_i, b_j \rangle$$

Remark 8.6. $A = A^*$ is called positive semidefinite if $A \geq 0$ if $\forall \xi : \xi^T A \bar{\xi} \geq 0$.

$A = A^*$ is called positive definite $\iff A > 0 \iff \forall \xi \in \mathbb{K}^n \setminus \{0\} : \xi^T A \bar{\xi} > 0$ with $\xi^T A \bar{\xi} = \sum_{i=1}^n \sum_{j=1}^n a_{ij} \xi_i \bar{\xi}_j$.

Example 8.4.

$$A = I > 0$$

$$\xi^T I \bar{\xi} = \sum_{i=1}^n \xi_i \bar{\xi}_i = \sum |\xi_i|^2 > 0 \quad \text{if } \xi \neq 0$$

$A = -I < 0$ is negative definite

$$A = \begin{bmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & & -1 & \\ & & & & \ddots \\ & & & & & -1 \end{bmatrix}$$

is indefinite:

$$e_1^T A e_1 > 0 \quad e_n^T A e_n < 0$$

Remark 8.7. For a diagonal matrix

$$A = \begin{bmatrix} a_1 & & 0 \\ & \ddots & \\ 0 & & a_n \end{bmatrix}$$

$A = A^* \iff a_i = \bar{a}_i$, hence for all $a_i \in \mathbb{R}$.

For a diagonal matrix it holds that

$$A > 0 \text{ if all } a_i > 0 : \xi^T A \bar{\xi} = \sum_{i=1}^n a_i |\xi_i|^2 \geq 0$$

$$A \leq 0 \text{ if all } a_i \geq 0 \text{ if } \xi^T A \bar{\xi} = 0 \implies \text{all } a_i \cdot |\xi_i|^2 = 0$$

$$A < 0 \text{ if all } a_i < 0$$

$$A \leq 0 \text{ if all } a_i \leq 0$$

$$\text{indefinite if } \exists i : a_i > 0 \exists j : a_j < 0$$

Remark 8.8. Remember, that the rank of matrix satisfies:

$$\exists P, Q \in \text{GL}(n) : PAQ = \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \ddots & \\ & & & 0 \end{pmatrix}$$

$A \sim PAQ$ is equivalent

8.5 Congruence of matrices

Definition 8.8 (Congruence). Consider two self-adjoint matrices $A, B \in \mathbb{K}^{n \times n}$ are called congruent (denoted $A \cong B$) if $\exists C \in \text{GL}(n, \mathbb{K})$ such that $C^* A C = B$.

Remark 8.9. C is invertible, hence C^T is invertible.

$$(C^T)^{-1} = (C^{-1})^T \quad (C^{-1})^T \cdot C^T = (C \cdot C^{-1})^T = I^T = I$$

$$(\overline{A^{-1}}) = \overline{A}^{-1}$$

$$(AB)^* = \overline{(AB)}^T = \overline{B^T A^T} = \overline{B}^T \cdot \overline{A}^T = B^* \cdot A^*$$

$C^* A C$ is self-adjoint.

$$(C^* A C)^* = C^* \cdot A^* \cdot (C^*)^* = C^* \cdot A \cdot C$$

Theorem 8.11. Every Hermitian matrix is congruent to a diagonal matrix D of structure:

$$\text{diag}(D) = (1, 1, \dots, 1, -1, \dots, -1, 0, \dots, 0)$$

Proof. The proof is given by an algorithm.

We construct matrix C inductively such that

$$C^*AC = \text{diag}(\pm 1, \dots, 0)$$

Consider $n = 1$.

$$A = [a_{11}]$$

If $a_{11} = 0$ where $a_{11} \in \mathbb{R}$, we don't have to do anything. If $a_{11} \neq 0$ and $a_{11} \in \mathbb{R}$ (because A is self-adjoint),

$$C = \left[\frac{1}{\sqrt{|a_{11}|}} \right]$$

$$C^*AC = \left[\frac{1}{\sqrt{|a_{11}|}} \cdot a_{11} \cdot \frac{1}{\sqrt{|a_{11}|}} \right] = [\text{sign}(a_{11})]$$

Example 8.5.

$$A = \begin{bmatrix} 0 & 1 & i \\ 1 & 0 & 1 \\ -i & 1 & 0 \end{bmatrix}$$

Remark 8.10. It seems we need to take the absolute value in the complex numbers:
Let $a = 3 + 4i$. $|a| = 5$.

$$C^*AC = \left[\frac{1}{\sqrt{|a_{11}|}} \cdot |a_{11}| \cdot \frac{1}{\sqrt{|a_{11}|}} \right] = \left[\frac{1}{\sqrt{5}} \cdot 5 \cdot \frac{1}{\sqrt{5}} \right] = [1]$$

Then $n - 1 \rightarrow n$:

Case 1: $A = 0$ nothing to do.

Case 2: $a_{11} = 0$ **Case 2a:**

$$\exists j : a_{jj} \neq 0 : \begin{bmatrix} 0 & & & & \\ & a_{jj} & & & \\ & & & & \\ & & & & \\ & & & & \end{bmatrix}$$

$$T_{(1,j)} = \begin{bmatrix} 0 & & & & & & 1 \\ & 1 & & & & & \\ & & \ddots & & & & \\ & & & 1 & & & \\ & & & & 0 & & \\ & & & & & 1 & \\ & & & & & & \ddots \\ 1 & & & & & & & 1 \end{bmatrix} = T_{(ij)}^*$$

Permutation matrix that swaps 1 with j .

$$T_{(1j)}^* A T_{(1j)} = \begin{bmatrix} a_{ji} & \dots & \dots \\ \vdots & \ddots & \\ \vdots & & 0 \end{bmatrix}$$

where $T_{(1j)}^*$ exchanges j -th and first row and $T_{(1j)}$ exchanges j -th and first column.

Case 2b : all $a_{jj} = 0$. Choose i, j such that $a_{ij} \neq 0$. Let E_{ij} be a zero matrix with 1 at row i and column j .

$$C = I + E_{ij}e^{i\theta}$$

where θ such that $a_{ij} = e^{i\theta}|a_{ij}|$.

Example 8.6. $a_{12} \neq 0$

$$C_1 = \begin{bmatrix} 1 & 1 & \\ & 1 & \\ & & 1 \end{bmatrix}$$

$$C_1^* A C_1 = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 0 & 1 & i \\ 1 & 0 & 1 \\ -i & 1 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 1 & i \\ 1 & 2 & 1+i \\ -i & 1-i & 0 \end{bmatrix}$$

In the general case:

$$C^* A C = (I + E_{ji}e^{-i\theta})A(I + E_{ij}e^{i\theta})$$

$$\begin{aligned} (C^* A C)_{jj} &= (A + E_{ji}e^{-i\theta}A + AE_{ij}e^{i\theta} + E_{ji}AE_{ij})_{jj} \\ &= \underbrace{a_{jj}}_{=0} + \underbrace{(E_{ji}e^{-i\theta}A)_{jj}}_{e^{-i\theta}a_{ij}=|a_{ij}|} + \underbrace{(AE_{ij}e^{i\theta})_{jj}}_{a_{ji}e^{i\theta}=\overline{a_{ij}}e^{i\theta}=|a_{ij}|} + \underbrace{a_{ii}}_{=0} \\ &= 2|a_{ij}| \end{aligned}$$

Case 2a is shown.

Example 8.7.

$$C_2 = \begin{bmatrix} 0 & 1 & \\ 1 & 0 & \\ & & 1 \end{bmatrix} = T_{(12)}$$

$$A_2 = C_2^* A_1 C_2 = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 0 & 1 & i \\ 1 & 2 & i+1 \\ -i & 1-i & 0 \end{bmatrix} \cdot \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 1 & 1+i \\ 1 & 0 & i \\ 1+i & -i & 0 \end{bmatrix}$$

Case 3 $a_{11} \neq 0$

$$C = \begin{bmatrix} 1 & -\frac{a_{12}}{a_{11}} & -\frac{a_{13}}{a_{11}} & \cdots & -\frac{a_{1n}}{a_{11}} \\ & 1 & \cdots & 0 & 0 \\ & \vdots & 1 & & 0 \\ & 0 & & \ddots & \\ & 0 & 0 & \cdots & 1 \end{bmatrix}$$

Example 8.8.

$$C_3 = \begin{bmatrix} 1 & -\frac{1}{2} & -\frac{1+i}{2} \\ & 1 & \\ & & 1 \end{bmatrix}$$

$$A_3 = C_3^* A_2 C_3 = \begin{bmatrix} 1 & 0 & 0 \\ -\frac{1}{2} & 1 & 0 \\ -\frac{1-i}{2} & 0 & 1 \end{bmatrix} \cdot \begin{bmatrix} 2 & 1 & 1+i \\ 1 & 0 & i \\ 1-i & -i & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & -\frac{1}{2} & -\frac{1+i}{2} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 1 & 1+i \\ 0 & -\frac{1}{2} & \frac{1}{2}(-i+i) \\ 0 & \frac{1}{2}(-1-i) & -1 \end{bmatrix} \cdot \begin{bmatrix} 2 & 0 & 0 \\ 0 & -\frac{1}{2} & -\frac{1+i}{2} \\ 0 & -\frac{1-i}{2} & -1 \end{bmatrix}$$

$$C^* A C = \begin{bmatrix} a_{11} & 0 & \cdots & 0 \\ 0 & & & \\ \vdots & & \tilde{A} & \\ 0 & & & 1 \end{bmatrix}$$

$$\tilde{A} \in \mathbb{K}^{(n-1) \times (n-1)}$$

$$\tilde{A} = \tilde{A}^*$$

$$C' = \begin{bmatrix} \frac{1}{\sqrt{|a_{11}|}} & & 0 \\ & 1 & \\ & & \ddots \\ 0 & & & 1 \end{bmatrix}$$

$$(C')^* (C^* A C) C' = \begin{bmatrix} \frac{a_{11}}{|a_{11}|} & 0 & 0 \\ 0 & & \\ \vdots & & \tilde{A} \\ 0 & & & 1 \end{bmatrix} \text{ where } \frac{a_{11}}{|a_{11}|} = \pm 1$$

Apply this algorithm to \tilde{A} .

Example 8.9 (Part 4).

$$C_4 = \begin{bmatrix} \frac{1}{\sqrt{2}} & & \\ & 1 & \\ & & 1 \end{bmatrix}$$

$$\begin{aligned}
A_4 &= C_4^* A_3 C_4 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & \frac{-1+i}{2} \\ 0 & \frac{-1-i}{2} & -1 \end{bmatrix} \\
\tilde{A} &= \begin{bmatrix} -\frac{1}{2} & \frac{-1+i}{2} \\ \frac{-1-i}{2} & -1 \end{bmatrix} \\
C_5 &= \begin{bmatrix} 1 & & \\ & 1 & -1+i \\ & 0 & 1 \end{bmatrix} \\
A_5 &= C_5^* A_4 C_5 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -1-i & 1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & \frac{-1+i}{2} \\ 0 & \frac{-1-i}{2} & -1 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1+i \\ 0 & 0 & 1 \end{bmatrix} \\
A_5 &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & \frac{-1+i}{2} \\ 0 & 0 & 0 \end{bmatrix} \cdot \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -1+i \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{2} & 0 \\ 0 & 0 & 0 \end{bmatrix} \\
C_6 &= \begin{bmatrix} 1 & & \\ & \sqrt{2} & \\ & & 1 \end{bmatrix} \\
\sqrt{2} &= \frac{1}{\sqrt{\frac{1}{2}}} \\
C_6^* A_5 C_6 &= \begin{bmatrix} 1 & & \\ & -1 & \\ & & 0 \end{bmatrix} \\
C_6^* \dots C_2^* C_1^* A C_1 C_2 \dots C_6 &= \begin{bmatrix} 1 & & \\ & -1 & \\ & & 0 \end{bmatrix} \Rightarrow \text{indefinite} \\
C &= C_1 C_2 \dots C_6 \\
C^* &= C_6^* C_5^* \dots C_1^*
\end{aligned}$$

□

Remark 8.11. 1. If $A \geq 0$, C arbitrary $\Rightarrow C^* A C \geq 0$.

$$\begin{aligned}
\xi^T (C^* A C) \bar{\xi} &= \underbrace{(\xi^T C^*)}_{\xi^T \overline{C^T} = \overline{\xi^T C^T} = \overline{(C \cdot \bar{\xi})^T} = \bar{\eta}^T} A \underbrace{(C \bar{\xi})}_{\eta} = \bar{\eta}^T A \bar{\eta} \geq 0
\end{aligned}$$

2. If $A > 0$, C invertible

$$\Rightarrow C^* A C > 0$$

$$\text{if } \xi^T C^* A C \bar{\xi} = 0 \Rightarrow \eta = C \bar{\xi} = 0 \text{ because } A > 0$$

$$\Rightarrow \bar{\xi} = 0 \text{ because } C \text{ is invertible}$$

Corollary 8.3. *If we apply the example 8.5 to $A > 0$,*

$$C^*AC = \begin{bmatrix} \pm 1 & & & & \\ & \ddots & & & \\ & & \pm 1 & & \\ & & & \ddots & \\ & & & & 0 \\ & & & & & \ddots \end{bmatrix} \text{ is still positive definite } \implies C^*AC = I$$

Theorem 8.12 (Sylvester's law of inertia). *J. J. Sylvester (1814–1897)*

Let $A \in \mathbb{C}^{n \times n}$ be Hermitian. $C \in \text{GL}(n, \mathbb{C})$ by the algorithm such that

$$C^*AC = \begin{bmatrix} \pm 1 & & & & \\ & \ddots & & & \\ & & \pm 1 & & \\ & & & -1 & \\ & & & & \ddots \\ & & & & & -1 \\ & & & & & & 0 \\ & & & & & & & \ddots \\ & & & & & & & & 0 \end{bmatrix}$$

Then the number of $+1$, -1 and zeros is uniquely determined (it does not depend on the order to the operands).

Proof. C is invertible, hence

$$\text{rank}(A) = \text{rank} \begin{bmatrix} +1 & & & & \\ & \ddots & & & \\ & & +1 & & \\ & & & -1 & \\ & & & & \ddots \\ & & & & & -1 \\ & & & & & & 0 \\ & & & & & & & \ddots \\ & & & & & & & & 0 \end{bmatrix}$$

Let r be the number of $+1$ and s be the number of -1 . The number of $+1$ and -1 is uniquely determined.

Hence, it suffices to show that the number r of $+1$ is uniquely defined.

Let \tilde{C} be another matrix such that

$$\tilde{C}^* A \tilde{C} = \begin{bmatrix} \pm 1 & & & & & & & \\ & \ddots & & & & & & \\ & & \pm 1 & & & & & \\ & & & -1 & & & & \\ & & & & \ddots & & & \\ & & & & & -1 & & \\ & & & & & & 0 & \\ & & & & & & & \ddots \\ & & & & & & & & 0 \end{bmatrix}$$

with \tilde{r} ones and \tilde{s} minus ones.

It suffices to show that $r \leq \tilde{r}$. We know $r + s = \tilde{r} + \tilde{s}$.

C is an invertible matrix, hence a change of basis. In this new basis $B' = \{b_1, \dots, b_n\}$, it holds that

$$x^* A x = \overline{x^T} A x = \overline{\Phi_B(x)^T} \cdot D \cdot \Phi_B(x)$$

$$\begin{aligned} A &= (C^*)^{-1} D C^{-1} \\ \overline{x^T} A x &= \overline{x^T} (C^*)^{-1} D \underbrace{C^{-1} x}_{\overline{C^{-1} x}} \end{aligned}$$

Equivalently, \tilde{C} is a change of basis to basis \tilde{B} such that $x^* A x = \Phi_{\tilde{B}}(x)^* \tilde{D} \Phi_{\tilde{B}}(x)$. For $x \in \mathcal{L}(\{b_1, \dots, b_r\}) \setminus \{0\}$,

$$\Phi_B(x) = \begin{pmatrix} \xi_1 \\ \vdots \\ \xi_r \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

$$\Rightarrow x^* A x = \Phi_B(x)^* D \Phi_B(x)$$

$$= (\overline{\xi_1}, \dots, \overline{\xi_r}, 0, \dots, 0) \begin{bmatrix} +1 & & & & & & & \\ & \ddots & & & & & & \\ & & +1 & & & & & \\ & & & -1 & & & & \\ & & & & \ddots & & & \\ & & & & & -1 & & \\ & & & & & & 0 & \\ & & & & & & & \ddots \\ & & & & & & & & 0 \end{bmatrix} \begin{pmatrix} \xi_1 \\ \vdots \\ \xi_r \\ 0 \\ \vdots \\ 0 \end{pmatrix} = \sum_{i=1}^r |\xi_i|^2 > 0$$

On the other hand, $\forall x \in \mathcal{L}(\tilde{b}_{\tilde{r}+1}, \dots, \tilde{b}_n)$.

$$\Phi_{\tilde{B}}(x) = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ \tilde{\xi}_{\tilde{r}+1} \\ \vdots \\ \tilde{\xi}_n \end{pmatrix}$$

$$x^*Ax = \Phi_{\tilde{B}}(x)^* \tilde{D} \Phi_{\tilde{B}}(x)$$

$$= (0, \dots, 0, \tilde{\xi}_{\tilde{r}+1}, \dots, \tilde{\xi}_n) \begin{bmatrix} +1 & & & & & & \\ & \ddots & & & & & \\ & & +1 & & & & \\ & & & -1 & & & \\ & & & & \ddots & & \\ & & & & & -1 & \\ & & & & & & 0 \\ & & & & & & \ddots \\ & & & & & & & 0 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ \vdots \\ 0 \\ \tilde{\xi}_{\tilde{r}+1} \\ \vdots \\ \tilde{\xi}_n \end{bmatrix} \leq 0$$

$$\Rightarrow \mathcal{L}(b_1, \dots, b_r) \cap \mathcal{L}(\tilde{b}_{\tilde{r}+1}, \dots, \tilde{b}_n) = \{0\}$$

$$\text{dimension } r + (n - \tilde{r}) \leq n \Rightarrow r \leq \tilde{r}$$

□

↓ This lecture took place on 2018/04/16.

$$A = A^*$$

Conjugate complex. The important question: When does it hold that

$$A > 0$$

Hence

$$\forall x \in \mathbb{C}^n : x^*Ax \geq 0$$

$$A > 0 \text{ if } x^*Ax > 0 \forall x \neq 0$$

$$(x^*)_i = \bar{x}_i$$

$$\exists C \in \text{GL}(n, \mathbb{C}) \text{ such that}$$

$$C^*AC \underbrace{=}_{\text{congruence}} \begin{bmatrix} +1 & & & & & & \\ & \ddots & & & & & \\ & & +1 & & & & \\ & & & -1 & & & \\ & & & & \ddots & & \\ & & & & & -1 & \\ & & & & & & 0 \\ & & & & & & & \ddots \\ & & & & & & & & 0 \end{bmatrix}$$

where the number of +1 is r (see Sylvester's Law of inertia).

Definition 8.9. If $A = A^*$ is congruent to

$$\begin{bmatrix} +1 & & & & & & \\ & \ddots & & & & & \\ & & +1 & & & & \\ & & & -1 & & & \\ & & & & \ddots & & \\ & & & & & -1 & \\ & & & & & & 0 \\ & & & & & & & \ddots \\ & & & & & & & & 0 \end{bmatrix}$$

with r occuring +1s and s occuring -1s.

Then $\text{ind}(A) := r$ is called index of A . $\text{sign}(A) := r - s$ is called signature of A .

Corollary 8.4. 1. $A > 0 \iff A \triangleq I \iff \text{ind}(A) = n$

2. $A \geq 0 \iff \text{ind}(A) = \text{sign}(A) = \text{rank}(A)$

3. $A \triangleq B \iff \text{ind}(A) = \text{ind}(B) \wedge \text{sign}(A) = \text{sign}(B)$

It is left as an exercise to the reader that congruence is an equivalence relation.

1. $I \cdot A \cdot I = A$

2. $A \triangleq B \implies C^*AC = B \implies A = (C^*)^{-1}BC^{-1} = (C^{-1})^*BC^{-1} \implies B \triangleq A$

3. $C_1^*A_1C_1 = A_2 \wedge C_2^*A_2C_2 = A_3 \implies \underbrace{C_2^*C_1^*A_1C_1C_2}_{=(C_1C_2)^*A_1(C_1C_2)} = A_3 \implies A_1 \triangleq A_3$

Furthermore it will be shown in the practicals that $A > 0 \iff \exists CA = C^*C$

Remark 8.12 (Idea).

$$\det(C^*AC) = \det \begin{bmatrix} +1 & & & & & & \\ & \ddots & & & & & \\ & & +1 & & & & \\ & & & -1 & & & \\ & & & & \ddots & & \\ & & & & & -1 & \\ & & & & & & 0 & \\ & & & & & & & \ddots & \\ & & & & & & & & 0 \end{bmatrix}$$

$$\det(C^*) \det(A) \det(C) = \begin{cases} 0 & \text{if } \text{rank}(A) < n \\ (-1)^{\text{number of } -1} & \end{cases}$$

$$\overline{\det(C)} \det(A) \det(C)$$

If $A > 0$,

$$|\det(C)|^2 \cdot \det(A) = 1 \implies \det(A) > 0$$

Lemma 8.3. 1.

$$\det(C^*) = \overline{\det(C)}$$

2.

$$A = A^* \implies \det(A) \in \mathbb{R}$$

3.

$$A = A^*, B = B^*, A \hat{=} B \implies \text{sign } \det(A) = \text{sign } \det(B)$$

4.

$$A > 0 \implies \det(A) > 0$$

but not the other way around:

$$\det \begin{bmatrix} -1 & \\ & -1 \end{bmatrix} = 1$$

Proof. 1.

$$\det(C^*) = \sum_{\sigma \in \Sigma_n} (-1)^\sigma \underbrace{(C^*)_{1\sigma(1)} \cdots (C^*)_{n\sigma(n)}}_{\overline{C_{\sigma(1)1}} \cdots \overline{C_{\sigma(n)n}}} = \overline{\sum_{\sigma} (-1)^\sigma C_{\sigma(1)1} \cdots C_{\sigma(n)n}} = \overline{\det(C)}$$

2. immediate

$$3. A\hat{B} \implies C^*AC = B$$

$$\det(C^*AC) = \det(B)$$

$$\underbrace{|\det(C)|^2}_{>0} \cdot \det(A) = \det(B)$$

$$4. A \hat{=} I \implies \text{sign } \det(A) = \text{sign } \det(I) = 1$$

□

Definition 8.10. Let $A \in \mathbb{K}^{m \times n}$, $r \leq \min\{m, n\}$.

$$I = \underbrace{\{i_1 < \dots < i_r\}}_{\subseteq \{1, \dots, m\}} \quad J = \underbrace{\{j_1 < \dots < j_r\}}_{\subseteq \{1, \dots, n\}}$$

Then

$$[A]_{I,J} = \begin{bmatrix} a_{i_1 j_1} & a_{i_1 j_2} & \dots & a_{i_1 j_r} \\ a_{i_2 j_1} & a_{i_2 j_2} & \dots & a_{i_2 j_r} \\ \vdots & \vdots & \ddots & \vdots \\ a_{i_r j_1} & a_{i_r j_2} & \dots & a_{i_r j_r} \end{bmatrix}$$

is called minor of A.

Example 8.10. Let $r = 1$, $I = \{i_1\}$, $J = \{j_1\}$, $[A]_{\{i_1\}, \{j_1\}} = a_{i_1 j_1}$.

Definition 8.11. If $m = n$ with $I = \{1, \dots, r\}$ and $J = \{1, \dots, r\}$, then

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1r} \\ \vdots & \vdots & \ddots & \vdots \\ a_{r1} & a_{r2} & \dots & a_{rr} \end{bmatrix}$$

the first minor of A (Hauptminoren).

$$A < 0 \iff (-A) > 0$$

$$\det(\lambda A) = \lambda^* \det(A)$$

Theorem 8.13. Let $A = A^*$, then it holds that

$$1. A > 0 \iff \text{all first minors satisfy } \det(A_r) > 0$$

$$2. A < 0 \iff (-1)^r \det(A_r) > 0 \forall r \in \{1, \dots, n\}$$

Proof. Direction \implies

For $r = n$: $\det(A_r) = \det(A) > 0$. It suffices to show: the submatrices

$$A_r = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1r} \\ \vdots & & & \\ a_{r1} & & & a_{rr} \end{bmatrix}$$

are positive definite. Hence, $\forall x \in \mathbb{C}^r$ with $x \neq 0 : x^* A_r x > 0$.

$$x \in \mathbb{C}^r \setminus \{0\} : x^* A_r x = \left[x^* \underbrace{0}_{n-r} \right] \cdot A \cdot \begin{bmatrix} x \\ 0 \end{bmatrix} > 0$$

$$= [x^* 0] \begin{bmatrix} A_r & & & * \\ & & & \vdots \\ & & & * \\ * & \dots & * & * \end{bmatrix} \begin{bmatrix} x \\ 0 \end{bmatrix}$$

Remark: every submatrix $\begin{bmatrix} a_{i_1 i_1} & \dots & a_{i_1 i_r} \\ \vdots & \ddots & \vdots \\ a_{i_r i_1} & \dots & a_{i_r i_r} \end{bmatrix}$ of a positive definite matrix is positive definite.

Direction \Leftarrow

Assume all first minors $\det(A_r) > 0$.

We use complete induction:

Let $n = 1$ and $r = 1$ $A = [a_{11}]$ and $\det(A_1) = a_{11}$. $A > 0 \iff a_{11} > 0$.

Consider $n \rightarrow n + 1$ Assume all first minors are greater 0. Then all first minors of matrix A_{n-1} are greater 0.

□

$$A' = \begin{bmatrix} C & \vdots 0 \vdots \\ \dots 0 \dots & 1 \end{bmatrix} A \begin{bmatrix} C & \\ & 1 \end{bmatrix}$$

$$= \begin{bmatrix} C^* & \vdots 0 \vdots \\ \dots 0 \dots & 1 \end{bmatrix} \begin{bmatrix} A_{n-1} & a_{1,n} \\ & a_{2,n} \\ & \vdots \\ & a_{n-1,n} \\ \overline{a_{n,1}} & \overline{a_{n,2}} & \dots & a_{nn} \end{bmatrix} \begin{bmatrix} C & \vdots 0 \vdots \\ \dots 0 \dots & 1 \end{bmatrix}$$

$$= \begin{bmatrix} I & a_{1,n} \\ & a_{2,n} \\ & \vdots \\ \overline{a_{1,n}} & \overline{a_{2,n}} & \dots & \overline{a_{n-1,n}} & a_{n,n} \end{bmatrix}$$

$$C' = \begin{bmatrix} 1 & & 0 & -a_{1,n} \\ & \ddots & & -a_{2,n} \\ & & \vdots & \\ 0 & & -a_{n-1,n} & 1 \end{bmatrix} = \left[\begin{array}{c|c} I & -b \\ \hline 0 & 1 \end{array} \right]$$

with

$$b = \begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{n-1,n} \end{bmatrix}$$

$$(C')^* A' C' = \left[\begin{array}{c|c} I & 0 \\ \hline -b^* & 1 \end{array} \right] \left[\begin{array}{c|c} I & b \\ \hline b^* & a_{n,n} \end{array} \right] \text{TODO}$$

$$\Rightarrow A \hat{=} A' \hat{=} \begin{bmatrix} I & 0 \\ 0 & -b^* b + a_n \end{bmatrix}$$

$$\exists C'' = C \cdot C'$$

such that

$$(C'')^* A C'' = \left[\begin{array}{c|c} I & 0 \\ \hline 0 & a_{n,n} - b^* b \end{array} \right]$$

$$\det(A) \cdot |\det(C'')|^2 = \det \begin{bmatrix} I & 0 \\ 0 & a_{n,n} - b^* b \end{bmatrix} = a_{n,n} - b^* b > 0 \Rightarrow \begin{bmatrix} I & 0 \end{bmatrix}$$

Back to the scalar product:

Definition 8.12. 1. (a) A vector space with a positive definite inner product is called Euclidean space ($K = \mathbb{R}, \dim < \infty$) or unitary space ($K = \mathbb{C}$)

(b) Hilbert space if $\dim = \infty$.

David Hilbert (1862–1943)

$$\|v\| = \sqrt{\langle v, v \rangle}$$

$$\|\lambda v\| = |\lambda| \cdot \|v\|$$

$$\text{in } \mathbb{R}^2: \langle a, b \rangle = \|a\| \|b\| \cos \varphi$$

2. An element $v \in V$ is called **normed** if $\|v\| = 1$ (if not, then $\frac{v}{\|v\|}$ is normed)
3. Let $v, w \in V \setminus \{0\}$. Then the angle spanned between v and w is the angled $\varphi \in [0, \phi]$ such that $\cos \varphi = \frac{\Re \langle v, w \rangle}{\|v\| \|w\|}$
4. Two vectors $v, w \in V$ are **orthogonal** ($v \perp w$) if $\langle v, w \rangle = 0$ (hence $\varphi = \frac{\pi}{2}$)

Theorem 8.14. 1. $\|v + w\|^2 = \|v\|^2 + \|w\|^2 + 2 \|v\| \|w\| \cos \varphi$ (Law of cosines)




Figure 4: Geometrical proof of Theorem 8.14

2. if $v \perp w$: $\|v + w\|^2 = \|v\|^2 + \|w\|^2$ (Pythagorean Theorem)
3. $\|v + w\|^2 + \|v - w\|^2 = 2(\|v\|^2 + \|w\|^2)$ (Parallelogram Law)

$$\begin{aligned}
 e^2 + f^2 &= 2(a^2 + b^2) \\
 e^2 &= (a + \xi)^2 + \eta^2 \\
 f^2 &= (a - \xi)^2 + \eta^2 \\
 e^2 + f^2 &= (a + \xi)^2 + (a - \xi)^2 + 2\eta^2 \\
 &= a^2 + \xi^2 + a^2 + \xi^2 + 2\eta^2 = 2a^2 + 2b^2
 \end{aligned}$$

Proof. 1.

$$\begin{aligned}
 \|v + w\|^2 &= \langle v + w, v + w \rangle = \langle v, v \rangle + \langle v, w \rangle + \langle w, v \rangle + \langle w, w \rangle \\
 &= \|v\|^2 + \langle v, w \rangle + \overline{\langle v, w \rangle} + \|w\|^2 \\
 &= \|v\|^2 + 2 \underbrace{\Re \langle v, w \rangle}_{\cos \varphi \cdot \|v\| \cdot \|w\|} + \|w\|^2
 \end{aligned}$$

2. immediate, $\langle v, w \rangle = 0$

3.

$$\begin{aligned}\|v + w\|^2 + \|v - w\|^2 &= \|v\|^2 + \|w\|^2 + 2\Re \langle v, w \rangle + \|v\|^2 + \|-w\|^2 + 2\Re \langle v, -w \rangle \\ &= 2\|v\|^2 + 2\|w\|^2 + 0\end{aligned}$$

Other norms:

$$\begin{aligned}\left\| \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \right\|_1 &= \sum_1^n |x_i| \\ \left\| \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \right\|_\infty &= \max |x_i|\end{aligned}$$

□

Remark 8.13. You can show (von Neumann did): A norm on \mathbb{R}^n satisfies the Parallelogram Law iff \exists a scalar product on \mathbb{R}^n such that $\|v\| = \sqrt{\langle v, v \rangle}$

Definition 8.13. Let (v, \langle, \rangle) be a vector space with scalar product. A family $(v_i)_{i \in I} \subseteq V$ is called

orthogonal if $\forall i \neq j : \langle v_i, v_j \rangle = 0$

orthonormal if additionally $\|v_i\| = 1 \forall i$
hence $\forall i, j : \langle v_i, v_j \rangle = \delta_{ij}$

orthonormal basis if they are orthonormal and give a basis of V .

Example 8.11. 1. Canonical basis in \mathbb{R}^n in regards of the standard scalar product

$$\langle e_i, e_j \rangle = \delta_{ij}$$

2. Fourier $\{ \sqrt{2} \sin 2\pi x, \sqrt{2} \sin 4\pi x, \dots, \sqrt{2} \sin(2k\pi x), \dots \}$ with $k \in \mathbb{N}$ union with $\{ \sqrt{2} \cos 2\pi x, \sqrt{2} \cos 4\pi x, \dots \} \cup \{g\}$ on $C[0, 1]$.

$$\langle f, g \rangle = \int_0^1 f(x)g(x) dx$$

And this is wrong unless we redefine the term basis (not every function is built using the sine/cosine). A basis here is every function:

$$f(x) = \sum_{k=0}^{\infty} a_k \cos 2k\pi x + \sum_{k=1}^{\infty} b_k \sin 2k\pi x$$

And this is wrong as well unless we define equality more precisely (in the usual sense, it is wrong). Lebesgue did this later.

Remark 8.14. For JPEG compression, Fourier transformation is applied. Hence, we consider the music (amplitudes) as f and

$$f(x) = \sum_{k=0}^n a_k \cos 2k\pi x + \sum_{k=1}^n b_k \sin 2k\pi x$$

with n finite.

Theorem 8.15. Let $(v_i)_{i \in I} \subseteq V$, $v_i \neq 0 \forall i$

1. $(v_i)_{i \in I}$ orthogonal $\iff \left(\frac{v_i}{\|v_i\|} \right)_{i \in I}$ is orthonormal
2. $(v_i)_{i \in I}$ is orthogonal, then $(v_i)_{i \in I}$ is linear independent.

↓ This lecture took place on 2018/04/18.

$$\cos \varphi = \frac{\langle v, w \rangle}{\|v\| \|w\|}$$

$$v \perp w \iff \langle v, w \rangle = 0$$

$(v_i)_{i \in I}$ orthogonal if $\langle v_i, v_j \rangle = 0 \forall i \neq j$

orthonormal: $\langle v_i, v_j \rangle = \delta_{ij}$.

Proof of Theorem 8.15. Let $\sum_{k=1}^n \lambda_k v_{i_k} = 0$.

$$\implies 0 = \left\langle \sum_{k=1}^n \lambda_k v_{i_k}, v_i \right\rangle = \sum_{k=1}^n \lambda_k \langle v_{i_k}, v_i \rangle$$

$\forall l \in \{1, \dots, n\}$: Let $i = i_l$.

$$\begin{aligned} i_l &= \sum_{k=1}^n \lambda_k \left\langle \underbrace{v_{i_k}, v_{i_l}}_{\substack{0 & i_k \neq i_l \\ \|v_{i_l}\|^2 & i_k = i_l}} \right\rangle \\ &= \lambda_l \cdot \|v_{i_l}\|^2 \implies \lambda_l = 0 \end{aligned}$$

□

Theorem 8.16. Let $B = (b_1, \dots, b_n)$ is an orthonormal basis of a finite dimensional

vector space over \mathbb{K} . For $v \in V$, let $\Phi_B(v) = \begin{pmatrix} \lambda_1 \\ \vdots \\ \lambda_n \end{pmatrix}$. For $w \in V$, let $\Phi_B(w) = \begin{pmatrix} \mu_1 \\ \vdots \\ \mu_n \end{pmatrix}$.

1. $\lambda_i = \langle v, b_i \rangle$
2. $\langle v, w \rangle = \sum_{i=1}^n \lambda_i \overline{\mu_i}$

Proof. 1.

$$\begin{aligned}
 \langle v, b_i \rangle &= \left\langle \sum_{j=1}^n \lambda_j b_j, b_i \right\rangle \\
 &= \sum_{j=1}^n \lambda_j \cdot \underbrace{\langle b_j, b_i \rangle}_{=\delta_{ji}} \\
 &= \lambda_i
 \end{aligned}$$

2.

$$\begin{aligned}
 \langle v, w \rangle &= \left\langle \sum_{i=1}^n \lambda_i b_i, \sum_{j=1}^n \mu_j b_j \right\rangle \\
 &= \sum_{i=1}^n \lambda_i \sum_{j=1}^n \overline{\mu_j} \underbrace{\langle b_i, b_j \rangle}_{\delta_{ij}} \\
 &= \sum_{i=1}^n \lambda_i \cdot \overline{\mu_i}
 \end{aligned}$$

Compare: B is an arbitrary basis:

$$\begin{aligned}
 \langle v, w \rangle &= \Phi_B(v)^T \cdot A \cdot \overline{\Phi_B(w)} \\
 a_{ij} &= \langle b_i, b_j \rangle = \delta_{ij} \\
 A &= I \\
 \rightarrow \langle v, w \rangle &= \Phi_B(v)^T \cdot \overline{\Phi_B(w)}
 \end{aligned}$$

□

Definition 8.14. Let V be a vector space with a scalar product. Let $v \in V$, then

$$v^\perp = \{w \in V \mid \langle v, w \rangle = 0\}$$

For $M \subseteq V$: $M^\perp = \{w \in V \mid \forall u \in M : \langle u, w \rangle = 0\}$ is called orthogonal complement of v or orthogonal complement of M




Figure 5: Orthogonal complement

Compare with Figure 5

in \mathbb{R}^n :

$$\begin{aligned} & \{w \mid \langle v, w \rangle = 0\} \\ &= \left\{ \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \mid \sum_{i=1}^n a_i x_i = 0 \right\} \end{aligned}$$

if $v = \begin{pmatrix} a_1 \\ \vdots \\ a_n \end{pmatrix}$.

Theorem 8.17. Let V be a vector with scalar product. $M, N \subseteq V$ are partitions.

1. M^\perp is a subspace.
2. $M \subseteq N \implies N^\perp \subseteq M^\perp$
 $(M_1 \cup M_2)^\perp = M_1^\perp \cap M_2^\perp$
3. $\{0\}^\perp = V$
4. $V^\perp = \{0\}$
5. $M \cap M^\perp \subseteq \{0\}$

$$6. M^\perp = \mathcal{L}(M)^\perp$$

$$7. M \subseteq (M^\perp)^\perp$$

Proof. 1.

$$v^\perp = \{w \in V \mid \langle v, w \rangle = 0\}$$

$$T_v : V \rightarrow \mathbb{K} \text{ (linear functional)}$$

$$w \mapsto \langle w, v \rangle$$

$$v^\perp = \{w \mid T_v(w) = 0\} = \ker T_v$$

is a subspace.

$$\begin{aligned} M^\perp &= \bigcap_{v \in M} v^\perp \\ &= \bigcap_{v \in M} \ker(T_v) \end{aligned}$$

is a subspace.

$$2. M \subseteq N \implies N^\perp \subseteq M^\perp$$

$$\begin{aligned} (M_1 \cup M_2)^\perp &= \{w \mid \forall v \in M_1 : \langle w, v \rangle = 0 \wedge \forall v \in M_2 : \langle w, v \rangle = 0\} \\ &= M_1^\perp \cap M_2^\perp \end{aligned}$$

$$3. \text{ trivial: } \forall v \in V : \langle v, 0 \rangle = 0$$

$$4. \text{ Let } w \in V \text{ such that } \langle w, v \rangle = 0 \forall v \in V. \text{ Especially for } v = w.$$

$$\begin{aligned} \implies \underbrace{\langle w, w \rangle}_{\|w\|^2} = 0 &\implies w = 0 \\ \implies V^\perp &= \{0\} \end{aligned}$$

$$5. \text{ Let } w \in M \cap M^\perp, \text{ hence}$$

$$\begin{aligned} \forall v \in M : \langle w, v \rangle &= 0 \\ w \in M &\implies \langle w, w \rangle = 0 \\ &\implies w = 0 \\ \text{or } M \cap M^\perp &= \varphi \end{aligned}$$

$$6.$$

$$M \subseteq \mathcal{L}(M) \underbrace{\implies}_{\text{by point (2.)}} \mathcal{L}(M)^\perp \subseteq M^\perp$$

Show that: $M^\perp \subseteq \mathcal{L}(M)^\perp$. Hence, $\forall v \in M^\perp \implies v \in \mathcal{L}(M)^\perp$. Let $v \in M^\perp$, $w \in \mathcal{L}(M)$.

$$\exists w_1, \dots, w_n \in M : \exists \lambda_1, \dots, \lambda_n \in \mathbb{K} : w = \sum_{i=1}^n \lambda_i w_i$$

$$\begin{aligned} \langle w, v \rangle &= \left\langle \sum_{i=1}^n \lambda_i w_i, v \right\rangle \\ &\stackrel{\text{by linearity in 1st argument}}{=} \sum_{i=1}^n \lambda_i \underbrace{\left\langle \underbrace{w_i}_{\in M}, \underbrace{v}_{\in M^\perp} \right\rangle}_{=0} = 0 \\ &\implies v \perp w \quad \forall w \in \mathcal{L}(M) \end{aligned}$$

7. Show that $\forall v \in M : v \in (M^\perp)^\perp$. Hence, $\forall w \in M^\perp : v \perp w$

$$\begin{aligned} M^\perp &= \{w \mid \forall v \in M : v \perp w\} \\ \implies \forall v \in M \forall w \in M^\perp : v \perp w &\implies \forall w \in M^\perp \forall v \in M, v \in W^\perp \\ &\implies \forall v \in M : v \in \bigcap_{w \in M^\perp} w^\perp = (M^\perp)^\perp \end{aligned}$$

□

Corollary 8.5. Let $U \subseteq V$ be a subspace. By Theorem 8.17 (1), U^\perp is a subspace and $U \cap U^\perp = \{0\}$ because of Theorem 8.17 (5),

$U + U^\perp$ is direct sum

in $\mathbb{R}^n : U + U^\perp = \mathbb{R}^n$.

Remark 8.15. If $\dim(V) = \infty$, it must not hold that $U + U^\perp = V$.

Example 8.12.

$$V = l^2 = \left\{ (x_n)_{n \in \mathbb{N}} \mid \sum |x_n|^2 < \infty \right\}$$

$$\begin{aligned} U &= \mathcal{L}((e_i)_{i \in \mathbb{N}}) \\ &= \{ (x_n)_{n \in \mathbb{N}} \mid x_n = 0 \text{ except for finite many } n \} \end{aligned}$$

$$U^\perp = \{e_i \mid i \in \mathbb{N}\}^\perp = \left\{ (x_n)_{n \in \mathbb{N}} \mid \underbrace{\langle (x_n)_{n \in \mathbb{N}}, e_i \rangle}_{= \{ (x_n)_{n \in \mathbb{N}} \mid \forall i \in \mathbb{N} : x_i = 0 \} = \{0\}} = 0 \forall i \in \mathbb{N} \right\}$$

$$\langle (x_n)_n, (y_n)_n \rangle = \sum_{n=1}^{\infty} x_n \overline{y_n}$$

$$\implies U^\perp = \{0\}$$

but $U + U^\perp \neq l_2$

$U+U^\perp$ is a direct sum.

$$\begin{aligned} v &\in U+U^\perp \\ U &\xrightarrow{\pi_U} U \\ U^\perp &\xrightarrow{\pi_{U^\perp}} U^\perp \end{aligned}$$

Every $v \in U+U^\perp$ has a unique decomposition:

$$v = u + w \quad u \in U, w \in U^\perp$$

Definition 8.15. Let V be a vector space. A subset $K \subseteq V$ is called convex⁸ if

$$\forall \lambda \in [0, 1] : \forall x, y \in K : \lambda x + (1 - \lambda)y \in K$$

Example 8.13. Subspaces are convex.

1.

$$U \subseteq V : \forall x, y \in U \forall \lambda, \mu : \lambda x + \mu y \in U$$

Especially: $\lambda \in [0, 1], \mu = 1 - \lambda$

2. Let $(V, \|\cdot\|)$ be a normed space.

$$B_{\|\cdot\|}(0, 1) = \left\{ x \in V \mid \underbrace{\|x\|}_{\text{unit circle}} < 1 \right\}$$

We discussed three different norms so far. In \mathbb{R}^2 with $\|\cdot\|_2$ (Euclidean norm), the unit circle is a circle of radius 1. In \mathbb{R}^2 with $\left\| \begin{pmatrix} x \\ y \end{pmatrix} \right\|_\infty = \max(|x|, |y|)$ (infinity norm), the unit circle is a square from $(-1, -1)$ to $(1, 1)$. This square contains the circle of radius 1. In \mathbb{R}^2 with $\left\| \begin{pmatrix} x \\ y \end{pmatrix} \right\|_1 = |x| + |y|$ (Manhattan norm), the unit circle is a square rotated by 45 degrees from $(-1, 0)$ to $(1, 0)$. It also contains the circle of radius 1.

Let $x, y \in B(0, 1)$, hence $\|x\| < 1, \|y\| < 1$.

$$\begin{aligned} \left\| \lambda x + (1 - \lambda)y \right\| &\underbrace{\leq}_{\text{by triangle ineq.}} \lambda \|x\| + (1 - \lambda) \|y\| \\ &< \lambda + (1 - \lambda) \\ &= 1 \\ &\implies \lambda x + (1 - \lambda)y \in B(0, 1) \end{aligned}$$

⁸Wide-sighted people with glasses use a glass with convex curvature.

3. *Translation in a convex set gives a convex set. Let K be convex. $K' = x_0 + K = \{x_0 + z \mid z \in K\}$ Let $x', y' \in K' \implies x' = x_0 + x$ and $y' = x_0 + y$.*

$$\begin{aligned} \implies \lambda x' + (1 - \lambda)y' &= \lambda \cdot (x_0 + x) + (1 - \lambda)(x_0 + y) \\ &= x_0 + \underbrace{\lambda x + (1 - \lambda)y}_{\in K} \end{aligned}$$

Especially: linear manifolds are convex. $B(x_0, 1)$ is convex.

4. $K \subseteq V$ convex. $f : V \rightarrow W$ is linear. $\implies f(K)$ is convex.

Optimization: Given a set M and a function $f : M \rightarrow \mathbb{R}$. Find $y \in M$ such that $f(y)$ is minimal.

Find $y \in M$ such that $d(x_0, y)$ is minimal. Compare with Figure 6.




Figure 6: A generic optimization problem

Now if M is convex (consider M convex in $(\mathbb{R}^n, \|\cdot\|_2)$), there exists a unique element $y \in M$ such that $\|x_0 - y\|$ is minimal.

Finite elements (in computational mathematics) is the same idea.

Theorem 8.18. $(V, \langle \cdot, \cdot \rangle)$ is a vector space with scalar product. $K \subseteq V$ is convex. Let $x \in V$ be given. Let $y_0 \in K$. Then the following statements are equivalent:

1. $\forall y \in K : \|x - y_0\| \leq \|x - y\|$
2. $\forall y \in K : \Re \langle x - y_0, y - y_0 \rangle \leq 0$
3. $\forall y \in K \setminus \{y_0\} : \|x - y_0\| < \|x - y\|$

Compare with Figure 7. In the special case if $K = U$ is a subspace, then the following statement is given (equivalent to statement 2)

$$2'. \forall y \in U : \langle x - y_0, y - y_0 \rangle = 0$$




Figure 7: Optimization on a convex set

Proof $1 \rightarrow 2$. Let $y \in K : 1 > \varepsilon > 0$.

$$y_\varepsilon = \underbrace{y_0 + \varepsilon(y - y_0)}_{\varepsilon y + (1-\varepsilon)y_0 \text{ because of convexity}} \in K$$

$$\begin{aligned}
\forall \varepsilon \in]0, 1[: \|x - y_0\|^2 &\leq \|x - y_\varepsilon\|^2 \\
&= \|x - (y_0 + \varepsilon(y - y_0))\|^2 \\
&= \|(x - y_0) - \varepsilon(y - y_0)\|^2 \\
&= \|x - y_0\|^2 - 2\varepsilon \Re \langle x - y_0, y - y_0 \rangle + \varepsilon^2 \|y - y_0\|^2 \\
\implies \forall 0 < \varepsilon < 1 : 0 &\leq -2\varepsilon \Re \langle x - y_0, y - y_0 \rangle + \varepsilon^2 \|y - y_0\|^2 \\
&= \varepsilon \cdot \left(-2\Re \langle x - y_0, y - y_0 \rangle + \varepsilon \|y - y_0\|^2 \right) \\
&\underbrace{\implies}_{\varepsilon \rightarrow 0} 0 \leq -2\Re \langle x - y_0, y - y_0 \rangle
\end{aligned}$$

2 \rightarrow 3.

$$\begin{aligned}
\|x - y\|^2 &= \|(x - y_0) + (y_0 - y)\|^2 \\
&= \|(x - y_0) - (y - y_0)\|^2 \\
&= \|x - y_0\|^2 + \|y - y_0\|^2 \underbrace{- 2\Re \langle x - y_0, y - y_0 \rangle}_{\geq 0} \\
&\geq \|x - y_0\|^2 + \|y - y_0\|^2 \\
&> \|x - y_0\|^2 \\
&y \neq y_0
\end{aligned}$$

3 \rightarrow 1. trivial.

2 \rightarrow 2'. Consider $K = U$ is subspace.

$$\forall y \in Y : \Re \langle x - y_0, y - y_0 \rangle \leq 0$$

U is a subspace.

$$\{y - y_0 \mid y \in U\} = \{z \mid z \in U\} = U - y_0$$

$$\left. \begin{array}{l} \forall z \in U : \Re \langle x - y_0, z \rangle \leq 0 \\ \forall z \in U : \Re \langle x - y_0, -z \rangle \leq 0 \end{array} \right\} \implies \forall z \in U : \Re \langle x - y_0, z \rangle = 0$$

Case $K = \mathbb{C}$:

$$\begin{aligned}
i \cdot U &= U \\
\implies z \in U : \Re \langle x - y_0, iz \rangle &= 0 \\
\Re i \langle x - y_0, z \rangle &= \Im \langle x - y_0, z \rangle
\end{aligned}$$

□

Corollary 8.6. Let (V, \langle, \rangle) be a vector space.

1. $K \subseteq V$ is convex, $x \in V$. Then the optimization problem

$$\begin{cases} \|x - y\| = \min! \\ y \in K \end{cases}$$

has at most one solution.

2. If $K = U$ subspace, then there exists at most one $y_0 \in U$ such that $x - y_0 \in U^\perp$.

↓ This lecture took place on 2018/04/23.

Orthonormalbasis:

$$\begin{aligned} \langle b_i, b_j \rangle &= \delta_{ij} \\ v &= \sum \lambda_i b_i \rightsquigarrow \langle v, b_i \rangle = \lambda_i \end{aligned}$$

Given: an arbitrary basis of a subspace

Find: orthonormal basis of the subspace

TODO sketch drawing (projection and convexity)

$$K \subseteq V \text{ convex}$$

V with scalar product.

Then the optimization problem

$$\|x - y\| = \min \quad y \in K$$

has at most one solution.

y is the solution.

$$\iff \Re \langle x - y_0, y - y_0 \rangle \leq 0 \forall y \in K$$

If K is the subspace U ($x - y_0 \perp U$), then

$$\Re \langle x - y_0, y \rangle = 0 \forall y \in K$$

$$U^\perp = \{ y \mid y \perp U \}$$

is subspace.

$$U \cap U^\perp = \{0\}$$

If $x \in U \cap U^\perp$, then $x \perp x = \langle x, x \rangle = \|x\|^2 = 0$.

Orthogonal complement: $U + U^\perp$ is direct sum.
Every $x \in U + U^\perp$ has a unique decomposition.

$$x = u + v \quad u \in U, v \in U^\perp$$

The maps $x \mapsto u$ and $x \mapsto v$ are linear.

Definition 8.16. Assume $U + U^\perp = V$. Then the projection maps

$$\pi_U : V \rightarrow V \quad \pi_{U^\perp} : V \rightarrow V$$

such that $\pi_U(x) \in U$ and $\pi_{U^\perp}(x) \in U^\perp$ and $x = \pi_U(x) + \pi_{U^\perp}(x)$ are orthogonal projections.

Remark 8.16. 1. $x \in U \iff \pi_U(x) = x \iff \pi_{U^\perp}(x) = 0$

2. $x \in U^\perp \iff \pi_U(x) = 0 \iff \pi_{U^\perp}(x) = x$

3. $\pi_{U^\perp} = \text{id} - \pi_U$

$$\pi_U(x) \in U$$

$$\implies \text{remark (4): } \pi_U(\pi_U(x)) = \pi_U(x)$$

$$(\sim): \pi_U \circ \pi_U = \pi_U \text{ idempotent}$$

$$\pi_U \text{ is linear: } \pi_U \circ \pi_{U^\perp} = 0$$

Theorem 8.19. Let $V = U + U^\perp$.

1. $\forall x, y \in V : \langle x, \pi_{U^\perp}(y) \rangle = \langle \pi_U(x), y \rangle = \langle \pi_U(x), \pi_U(y) \rangle$

2. Compare with Figure 8.

$$\|\pi_U(x)\| \leq \|x\| \wedge \|\pi_{U^\perp}(x)\| = \|x\| \iff x \in U$$

Proof:

(a)

$$x = \pi_U(x) + \pi_{U^\perp}(x) \quad y = \pi_U(y) + \pi_{U^\perp}(y)$$

$$\langle x, \pi_U(y) \rangle = \langle \pi_U(x) + \pi_{U^\perp}(x), \pi_U(y) \rangle = \langle \pi_U(x), \pi_U(y) \rangle + \underbrace{\left\langle \underbrace{\pi_{U^\perp}(x)}_{\in U^\perp}, \underbrace{\pi_U(y)}_{\in U} \right\rangle}_{=0}$$

$$\langle \pi_U(x), y \rangle = \langle \pi_U(x), \pi_U(y) \rangle + \langle \pi_U(x), \pi_{U^\perp}(y) \rangle$$

(b)

$$x = \pi_U(x) + \pi_{U^\perp}(x)$$

$$\implies \|x\|^2 = \|\pi_U(x)\|^2 + \|\pi_{U^\perp}(x)\|^2 \geq \|\pi_U(x)\|^2$$

$$\text{By equality } \iff \|\pi_{U^\perp}(x)\| = 0 \iff x = \pi_U(x) \iff x \in U$$




Figure 8: Projection

Definition 8.17. Jørgen Pederson Gram (1850–1916)

Let $v_1, v_2, \dots \in V$.

$$\text{Gram}(v_1, \dots, v_m) = \begin{bmatrix} \langle v_1, v_1 \rangle & \langle v_1, v_2 \rangle & \dots & \langle v_1, v_m \rangle \\ \langle v_2, v_1 \rangle & \langle v_2, v_2 \rangle & \dots & \langle v_2, v_m \rangle \\ \vdots & \vdots & \vdots & \vdots \\ \langle v_m, v_1 \rangle & \langle v_m, v_2 \rangle & \dots & \langle v_m, v_m \rangle \end{bmatrix}$$

is called Gram matrix of tuple v_1, v_2, \dots, v_m

Remark 8.17. In case $V = \mathbb{C}^n$.

$$\langle v, w \rangle = \overline{w}^T \cdot v = \sum_1^n \lambda_i \overline{\mu_i} = (\overline{\mu_1}, \dots, \overline{\mu_n}) \begin{pmatrix} \lambda_1 \\ \vdots \\ \lambda_n \end{pmatrix}$$

$$v = \begin{pmatrix} \lambda_1 \\ \vdots \\ \lambda_n \end{pmatrix} \quad w = \begin{pmatrix} \mu_1 \\ \vdots \\ \mu_n \end{pmatrix}$$

Hence, if

$$v_i = \begin{pmatrix} \beta_{1i} \\ \vdots \\ \beta_{ni} \end{pmatrix} \quad i = 1, \dots, m$$

$$\begin{aligned} V &= \begin{pmatrix} v_1 & v_2 & \dots & v_m \\ \vdots & \vdots & & \vdots \end{pmatrix} \in \mathbb{C}^{n \times m} \\ &= \begin{pmatrix} \beta_{11} & \beta_{12} & \dots & \beta_{1m} \\ \vdots & \vdots & & \vdots \\ \beta_{n1} & \beta_{n2} & \dots & \beta_{nm} \end{pmatrix} \\ (V^*V)_{ij} &= \sum_{k=1}^n (v^*)_{ik} v_{kj} = \sum_{k=1}^n \overline{\beta_{ki}} \beta_{kj} = \overline{\langle v_i, v_j \rangle} \\ &= \begin{pmatrix} v_1^* & \dots \\ \vdots & \\ v_m^* & \dots \end{pmatrix} \begin{pmatrix} v_1 & \dots & v_m \\ \vdots & & \vdots \end{pmatrix} \\ V^*V &= \overline{\text{Gram}(v_1, \dots, v_m)} \end{aligned}$$

Theorem 8.20. Let $v_1, \dots, v_m \in V$. $G = \text{Gram}(v_1, \dots, v_m)$.

1. $G = G^*$ is Hermitian, positive semidefinite.

$$\xi^T \cdot G \cdot \bar{\xi} = \left\| \sum_{i=1}^m \xi_i v_i \right\|^2 \geq 0$$

2. $\xi \in \ker G \iff \sum_{i=1}^m \bar{\xi}_i v_i = 0$
3. G is positive definite iff (v_1, \dots, v_m) are linear independent.

Proof. 1. $g_{ij} = \langle v_i, v_j \rangle = \overline{\langle v_j, v_i \rangle} = \overline{g_{ji}}$

$$\xi^T \cdot G \cdot \bar{\xi} = \sum_{i=1}^n \sum_{j=1}^n \xi_i g_{ij} \bar{\xi}_j = \sum_{i=1}^n \sum_{j=1}^n \xi_i \bar{\xi}_j \langle v_i, v_j \rangle = \left\langle \sum_{i=1}^n \xi_i v_i, \sum_{j=1}^n \bar{\xi}_j v_j \right\rangle = \left\| \sum_{i=1}^n \xi_i v_i \right\|^2$$

2. Direction \implies . $\xi \in \ker G \implies G\xi = 0 \implies \xi^T \cdot G \cdot \xi = 0$

$$\xi^T \cdot G \cdot \xi = \xi^T \cdot G \cdot \bar{\xi} \underbrace{=}_{(1)} \left\| \sum_{i=1}^m \bar{\xi}_i v_i \right\|^2$$

Direction \Leftarrow . If $\left\| \sum_{i=1}^m \xi_i v_i \right\| = 0$

$$(G \cdot \xi)_i = \sum_{j=1}^n \langle v_i, v_j \rangle \xi_j = \sum_{j=1}^n \langle v_i, \overline{\xi_j} v_j \rangle = \left\langle v_i, \underbrace{\sum_{j=1}^n \overline{\xi_j} v_j}_{=0} \right\rangle = 0$$

$$\Rightarrow G \cdot \xi = 0$$

3. G is positive definite

$$\Longleftrightarrow \forall \xi \neq 0 : \xi^T \cdot G \cdot \xi > 0$$

$$\Longleftrightarrow \forall \xi \neq 0 : \left\| \sum_{i=1}^m \xi_i \cdot v_i \right\|^2 > 0$$

$$\Longleftrightarrow \forall \xi \neq 0 : \sum_{i=1}^m \xi_i v_i \neq 0$$

$$\Longleftrightarrow (v_1, \dots, v_m) \text{ is linear independent}$$

$$\Longleftrightarrow \ker G = \{0\}$$

$$\Longleftrightarrow G \text{ is regular}$$

□

Theorem 8.21. Let $U \subseteq V$ be a subspace. V is a vector space with scalar product.

(u_1, \dots, u_m) is basis of U

$$G = \text{Gram}(u_1, \dots, u_m) = \left[\langle u_i, u_j \rangle \right]_{i,j=1,\dots,m}$$

Then the projection $\pi_U(x) = \sum_{j=1}^m \eta_j u_j$ where

$$\eta = \overline{G}^{-1} \cdot \begin{pmatrix} \langle x, u_1 \rangle \\ \vdots \\ \langle x, u_m \rangle \end{pmatrix}$$

If u_1, \dots, u_m would be an orthonormal basis, then

$$\begin{pmatrix} \langle x, u_1 \rangle \\ \vdots \\ \langle x, u_m \rangle \end{pmatrix}$$

would be the coordinate of x .




Figure 9: Projection

Let $u = \sum_{j=1}^m \eta_j u_j$. Compare with Figure 9. Show that $x - u \in U^\perp = \mathcal{L}(u_1, \dots, u_m)^\perp = \{u_1, \dots, u_m\}^\perp = \bigcap_{i=1}^m u_i^\perp$

Hence, show that $x - u \perp u_i \forall i \in \{1, \dots, m\}$.

$$\begin{aligned}
 \langle u_i, u \rangle &= \left\langle u_i, \sum_{j=1}^m \eta_j u_j \right\rangle \\
 &= \sum_{j=1}^m \langle u_i, u_j \rangle \cdot \overline{\eta_j} \\
 &= \sum_{j=1}^m g_{ij} \overline{\eta_j} \\
 &= (G\overline{\eta})_i &= \langle u_i, x \rangle
 \end{aligned}$$

because

$$\begin{aligned}\overline{G} \cdot \eta &= \begin{pmatrix} \langle x, u_1 \rangle \\ \vdots \\ \langle x, u_m \rangle \end{pmatrix} \\ G \cdot \overline{\eta} &= \begin{pmatrix} \langle x, u_1 \rangle \\ \vdots \\ \overline{\langle x, u_m \rangle} \end{pmatrix} = \begin{pmatrix} \langle u_1, x \rangle \\ \vdots \\ \langle u_m, x \rangle \end{pmatrix}\end{aligned}$$

Hence, $\forall i \in \{1, \dots, m\}$:

$$\langle u_i, u \rangle = \langle u_1, x \rangle \implies \forall i \in \{1, \dots, m\} : \langle u_i, x - u \rangle = 0 \implies x - u \in \{u_1, \dots, u_m\}^\perp$$

Example 8.14. Find polynomial $p(t)$ of degree 2 such that

$$\int_0^1 |t^3 - p(t)|^2 dt \stackrel{!}{=} \min$$

$V = C[0, 1]$, scalar product

$$\langle f, g \rangle = \int_0^1 f(t) \overline{g(t)} dt$$

$U =$ polynomial function of degree ≤ 2

$$x = t \mapsto t^3 \notin U$$

Find $p \in U$ such that $\|x - p\|^2 \stackrel{!}{=} \min$

$$\|x - p\|^2 = \int |x(t) - p(t)|^2 dt$$

Basis of $U = \mathcal{L}(\{1, t, t^2\})$

$$u_i(t) = t^{i-1} \quad i = 1, 2, 3$$

Gram matrix:

$$g_{ij} = \langle u_i, u_j \rangle = \int_0^1 t^{i-1} t^{j-1} dt = \int_0^1 t^{i+j-2} dt = \left. \frac{t^{i+j-1}}{i+j-1} \right|_0^1 = \frac{1}{i+j-1}$$

$$G = \begin{bmatrix} 1 & \frac{1}{2} & \frac{1}{3} \\ \frac{1}{2} & \frac{1}{3} & \frac{1}{4} \\ \frac{1}{3} & \frac{1}{4} & \frac{1}{5} \end{bmatrix}$$

Hilbert matrix:

$$\left[\frac{1}{i+j-1} \right]_{i,j=1,\dots,k}$$

This matrix is very unstable (in the equation system $Gx = b$) and therefore a very important test matrix in computational mathematics (ie. Numerics).

$$u = \sum_{j=1}^3 \eta_j u_j$$

$$\eta = \overline{G}^{-1} \cdot \begin{pmatrix} \langle x, u_1 \rangle \\ \langle x, u_2 \rangle \\ \langle x, u_3 \rangle \end{pmatrix}$$

$$\langle x, u_j \rangle = \int_0^1 x(t) u_j(t) dt = \int_0^1 t^3 \cdot t^{j-1} dt = \int_0^1 t^{2+j} dt = \frac{1}{3+j}$$

$$\eta = \begin{bmatrix} 1 & \frac{1}{2} & \frac{1}{3} \\ \frac{1}{2} & \frac{1}{3} & \frac{1}{4} \\ \frac{1}{3} & \frac{1}{4} & \frac{1}{5} \end{bmatrix}^{-1} \begin{pmatrix} \frac{1}{4} \\ \frac{1}{5} \\ \frac{1}{6} \end{pmatrix} = \begin{bmatrix} 9 & -36 & 30 \\ -36 & 192 & -180 \\ 30 & -180 & 180 \end{bmatrix} \begin{bmatrix} \frac{1}{4} \\ \frac{1}{5} \\ \frac{1}{6} \end{bmatrix} = \begin{bmatrix} \frac{1}{20} \\ -\frac{3}{5} \\ \frac{3}{2} \end{bmatrix}$$

(Assume that we don't know 180 in the bottom-right corner precisely. Consider $180 + \varepsilon$, then this error ε explodes tremendously in the solution).

Corollary 8.7. Special case u_i is orthonormal basis of U ($\rightarrow G = I$) Then it holds that

$$1. \forall v \in V : \pi_U(v) = \sum_{i=1}^m \langle v, u_i \rangle \cdot u_i$$

2.

$$\|v\|^2 \geq \sum_{i=1}^m |\langle v, u_i \rangle|^2 \quad (\text{Bessel's inequality})$$

$$\|v\|^2 = \sum_{i=1}^m |\langle v, u_i \rangle|^2 \iff v \in U \quad (\text{Parseval's identity})$$

$$\eta_j = \overline{G}^{-1} \begin{pmatrix} \langle v, u_1 \rangle \\ \vdots \\ \langle v, u_m \rangle \end{pmatrix}$$

F. Bessel (1784–1846)

M. A. Parseval (1755–1836)

Proof. Gram's matrix = I .

$$\eta_j = \langle v, u_j \rangle$$

□




Figure 10: Projection used in the Gram-Schmidt process

8.6 Gram-Schmidt process

Given: finite, linear independent vectors v_1, \dots, v_m

Find: orthonormal basis of (v_1, \dots, v_m) .

Theorem 8.22 (Gram-Schmidt process for orthogonalization). *Let $(v_1, \dots, v_m) \subseteq V$ be linear independent. Let $U := \mathcal{L}(v_1, \dots, v_m)$. Then $\exists u_1, \dots, u_m$ as orthonormal basis of U , specifically inductive*

$$u_1 = \frac{v_1}{\|v_1\|}$$

and for $k = 2, \dots, m$:

$$\begin{aligned} \tilde{u}_k &= v_k - \sum_{j=1}^{k-1} \langle v_k, u_j \rangle \cdot u_j \\ u_k &= \frac{\tilde{u}_k}{\|\tilde{u}_k\|} \end{aligned}$$

Proof. **Induction base** $k = 1$ is trivial

Induction step $k - 1 \rightarrow k$. Assume

$$\mathcal{L}(u_1, \dots, u_{k-1}) = \mathcal{L}(v_1, \dots, v_{k-1}) =: U_{k-1}$$

$$\tilde{u}_k = v_k - \pi_{U_{k-1}}(v_k) \in U_{k-1}^\perp \text{ because of Theorem 8.7}$$

$$\implies \tilde{u}_k \perp u_1, \dots, u_{k-1} \implies (u_1, \dots, u_{k-1}, \frac{\tilde{u}_k}{\|\tilde{u}_k\|})$$

is an orthonormal basis.

$$\mathcal{L}(u_1, \dots, u_{k-1}, \frac{\tilde{u}_k}{\|\tilde{u}_k\|}) = \mathcal{L}(u_1, \dots, u_{k-1}, v_k)$$

then $\tilde{u}_k - v_k \in \mathcal{L}(u_1, \dots, u_{k-1})$

□

↓ This lecture took place on 2018/04/25.

Gram-Schmidt process:

$$\mathcal{L}(v_1, v_2) = \mathcal{L}(v_2 - p(v_2), v_1) \quad v_2 - p(v_2) \perp v_1$$

Given: v_1, \dots, v_m

$$u_i = \frac{v_i}{\|v_i\|}$$

$$\tilde{u}_k = v_k - \sum_{i=1}^{k-1} \langle v_k, u_i \rangle \cdot u_i$$

$$u_k = \frac{\tilde{u}_k}{\|\tilde{u}_k\|} \quad \frac{\langle v_k, \tilde{u}_i \rangle \tilde{u}_i}{\|\tilde{u}_i\|^2}$$

Example 8.15. Let $V = \mathbb{R}^3$.

$$\langle x, y \rangle = x^t A y$$

$$A = \begin{bmatrix} 1 & -1 & 1 \\ -1 & 3 & -1 \\ 1 & -1 & 2 \end{bmatrix}$$

$v_i =$ standard basis e_i

$$\|v_1\|^2 = \langle v_1, v_1 \rangle = v_1^T A v_1 = a_{11} = 1$$

$$\|v_2\|^2 = \langle v_2, v_2 \rangle = a_{22} = 3$$

$$u_1 = \frac{v_1}{\|v_1\|} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$$

$$\tilde{u}_2 = v_2 - u_1 \langle v_2, u_1 \rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} - \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \cdot (0 \ 1 \ 0) A \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} + \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$$

$$u_2 = \frac{\tilde{u}_2}{\|\tilde{u}_2\|} \quad \|\tilde{u}_2\|^2 = \langle \tilde{u}_2, \tilde{u}_2 \rangle = (1 \ 1 \ 0) \cdot A \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} = 2 \quad u_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}$$

$$\tilde{u}_3 = v_3 - u_1 \langle v_3, u_1 \rangle - u_2 \langle v_3, u_2 \rangle$$

$$= \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} - \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \cdot \underbrace{(0 \ 0 \ 1) \cdot A \cdot \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}}_{a_{31}=1} - \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \cdot \underbrace{(0 \ 0 \ 1) \cdot A \cdot \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}}_{a_{31}+a_{32}=0} \cdot \frac{1}{\sqrt{2}} = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

$$\|\tilde{u}_3\|^2 = (-1 \ 0 \ 1) \cdot A \cdot \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix} = 1 - 1 - 1 + 2 = 1 \quad u_3 = \begin{pmatrix} -1 \\ 0 \\ 1 \end{pmatrix}$$

Remark 8.18. This is an alternative method to build orthogonal projection on subspace $U \subseteq \mathbb{C}^n$ with standard scalar product.

1. Determine an orthonormal basis of U : $u_1, \dots, u_m \in \mathbb{C}^{n \times 1}$
2. $P = \sum_{i=1}^m u_i \cdot u_i^*$

$$P \cdot v = \sum_{i=1}^m u_i \underbrace{u_i^* \cdot v}_{\langle v, v_i \rangle} = \sum_{i=1}^m u_i \langle v, v_i \rangle$$

Gram matrix = I

Example 8.16 (Example 8.14 again).

$$V = C[0, 1] \quad U = \mathcal{L}(1, x, x^2) =: \mathcal{L}(v_1, v_2, v_3)$$

$$\langle f, g \rangle = \int_0^1 f(t) \overline{g(t)} dt$$

Orthonormal basis:

$$\|v_1\|^2 = \int_0^1 1^2 dt = 1$$

$$u_1 = 1$$

$$\tilde{u}_2 = v_2 - u_1 \cdot \langle v_2, u_1 \rangle = x - 1 \cdot \underbrace{\int_0^1 t \cdot 1 dt}_{=\frac{1}{2}} = x - \frac{1}{2}$$

$$\|\tilde{u}_2\|^2 = \int_0^1 \left(t - \frac{1}{2}\right)^2 dt = \left. \frac{(t - \frac{1}{2})^3}{3} \right|_0^1 = \frac{(\frac{1}{2})^3 - (-\frac{1}{2})^2}{3} = \frac{1}{12}$$

$$u_2 = \frac{\tilde{u}_2}{\|\tilde{u}_2\|} = \sqrt{12} \cdot \left(x - \frac{1}{2}\right)$$

$$\begin{aligned}
\tilde{u}_3 &= v_3 - u_1 \langle v_3, u_1 \rangle - u_2 \cdot \langle v_3, u_2 \rangle \\
&= x^2 - 1 \cdot \underbrace{\int_0^1 t^2 \cdot 1 \, dt}_{=\frac{1}{3}} - \sqrt{12} \left(x - \frac{1}{2}\right) \int_0^1 t^2 \sqrt{12} \left(t - \frac{1}{2}\right) dt \\
&= x^2 - \frac{1}{3} - 12 \left(x - \frac{1}{2}\right) \cdot \frac{1}{12} \\
&= x^2 - x + \frac{1}{6}
\end{aligned}$$

Side note:

$$\begin{aligned}
\int_0^1 t^2 \left(t - \frac{1}{2}\right) dt &= \int_0^1 \left(t^3 - \frac{1}{2}t^2\right) dt = \frac{1}{4} - \frac{1}{6} = \frac{1}{12} \\
\|\tilde{u}_3\|^2 &= \int_0^1 \left(t^2 - t + \frac{1}{6}\right)^2 dt = \frac{1}{180} \\
\implies u_3 &= \sqrt{180} \cdot \left(x^2 - x + \frac{1}{6}\right)
\end{aligned}$$

Projection:

$$\int_0^1 (t^3 - p(t))^2 dt = \min!$$

Solution: $\pi_U(x^3) \quad U = \mathcal{L}(1, x, x^2)$

$$\begin{aligned}
\pi_U(x^3) &= u_1 \langle x^3, u_1 \rangle + u_2 \langle x^3, u_2 \rangle + u_3 \langle x^3, u_3 \rangle \\
&= 1 \cdot \int_0^1 t^3 \cdot 1 \, dt + \sqrt{12} \left(x - \frac{1}{2}\right) \int_0^1 t^3 \sqrt{12} \left(t - \frac{1}{2}\right) dt \\
&\quad + \sqrt{180} \left(x^2 - x + \frac{1}{6}\right) \int_0^1 t^3 \sqrt{180} \left(t^2 - t + \frac{1}{6}\right) dt
\end{aligned}$$

Consider $\langle f, g \rangle := \int_{-1}^1 \sqrt{1-t^2} f(t) \overline{g(t)} dt$. Take $1, x, x^2, \dots$ and apply Gram schmidt process to retrieve the Chebyshev polynomials.

$$\begin{aligned}
&\int_0^1 f(t)g(t) \, dt && \text{Laguerre polynomials} \\
&\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{-\frac{t^2}{2}} f(t)g(t) \, dt && \text{Hermite polynomials}
\end{aligned}$$

8.7 Riesz representation theorem

Remark 8.19. *Frigyes Riesz (1880–1956)*

Theorem 8.23. Let $(V, \langle \cdot, \cdot \rangle)$ be a vector space with scalar product $\dim V < \infty$.

V^* is the dual space, namely $\text{Hom}(V, \mathbb{K})$ which is the space of linear functionals. For some fixed $y \in V$, the map $T_y(x) = \langle x, y \rangle$ is linear in x (and therefore $T_y \in V^*$).

Then the map $V \rightarrow V^*$ defined by $Ty : V \rightarrow \mathbb{K}$ is an antilinear isomorphism (antiisomorphism).

This is trivial in \mathbb{R} , but in \mathbb{C} it is much more complex (pun intended).

Hence,

1. For every y it holds that $Ty \in V^*$
2. For every linear functional $f \in V^* : \exists! y \in V : f = T_y$
3. Let $y \mapsto Ty$ is an antilinear map.

$$T_{\lambda y_1 + \mu y_2} = \bar{\lambda} T y_1 + \bar{\mu} T y_2$$

Example 8.17 (For point 2).

$$V = C[0, 1]$$

Scalar product: $\langle f, g \rangle = \int f(t)g(t) dt$. Let $F : C[0, 1] \rightarrow \mathbb{R}$ linear. Then by the Riesz representation theorem, there exists $g \in C[0, 1] : F(f) = \int f(t)g(t) dt$.

For example $f \mapsto f(1)$

$$\exists g(t) : f(1) = \int_0^1 f(t)g(t) dt$$

In physics, e.g. the Dirac delta function.

Proof of point 3. We show linearity.

$$Ty(x) = \langle x, y \rangle \text{ is linear in } x \implies Ty \in V^*$$

$$\begin{aligned} \forall x \in V : T_{\lambda y_1 + \mu y_2}(x) &= \langle x, \lambda y_1 + \mu y_2 \rangle = \bar{\lambda} \langle x, y_1 \rangle + \bar{\mu} \langle x, y_2 \rangle \\ &= \bar{\lambda} T y_1(x) + \bar{\mu} T y_2(x) = (\bar{\lambda} T y_1 + \bar{\mu} T y_2)(x) \\ &\implies T_{\lambda y_1 + \mu y_2} = \bar{\lambda} T y_1 + \bar{\mu} T y_2 \end{aligned}$$

We show injectivity: the map $y \mapsto Ty$ is injective.

Assume: $Ty = 0$ (zero functional). Show $y = 0$. $Ty = 0$ means $\forall x \in V : Ty(x) = 0$, especially for $x = y$, $T_y(y) = \langle y, y \rangle = 0 \implies y = 0$.

We show surjectivity: the map $y \mapsto Ty$ is surjective.

Let u_1, \dots, u_n is an orthonormal basis (exists because of Gram-Schmidt).

Given: $f \in V^*$. Find: y such that $f = Ty$.

$$\text{Hence, } \forall x \in V : f(x) = \langle x, y \rangle \quad \underbrace{\iff}_{\text{by Fortsetzungssatz}} \quad f(u_i) = \langle u_i, y \rangle$$

Let $y = \sum_{j=1}^n \overline{f(u_j)} \cdot u_j$.

$$\implies \langle u_i, y \rangle = \left\langle u_i, \sum_{j=1}^n \overline{f(u_j)} u_j \right\rangle = \sum_{j=1}^n f(u_j) \underbrace{\langle u_i, u_j \rangle}_{\delta_{ij}} = f(u_i)$$

Hence, y satisfies the condition. \square

Remark 8.20. The Riesz representation theorem also holds in infinite dimensions (thus, in Hilbert spaces, a generalization of Euclidean spaces). In those spaces, there exists some Hilbert base:

$$(u_i)_{i \in I} : x = \sum_{i \in I} \langle x, u_i \rangle \cdot u_i \forall x$$

So every x has such a representation and in infinite dimensions, this representation is a series.

Corollary 8.8.

1. $v = 0 \iff \forall w \in V : \langle v, w \rangle = 0$
2. $\|v\| = \sup \{ |\langle v, w \rangle| \mid \|w\| \leq 1 \}$

Equivalently in the dual space:

1. $v = 0 \iff \forall f \in V^* : f(v) = 0$
2. $\|v\| = \sup \{ |f(v)| \mid f \in V^*, \|f\| \leq 1 \}$

holds in general in a normed space.

Remark 8.21. We make a small revision: dual space $V^* = \text{Hom}(V, \mathbb{K})$

$$\begin{aligned} W &\xrightarrow{T} V \xrightarrow{f} \mathbb{K} \\ \implies f \circ T : W &\rightarrow \mathbb{K} \in W^* \end{aligned}$$

is a linear functional on W . Hence, the map $\text{Hom}(V, \mathbb{K}) \rightarrow \text{Hom}(W, \mathbb{K})$ and $f \mapsto f \cdot T$ is linear.

$$(\lambda f + \mu g) \circ T = \lambda \cdot f \circ T + \mu g \circ T \quad \text{“transposed map”}$$

Linear map: $T^* : V^* \rightarrow W^*$.

Let V, W be spaces with a scalar product. Then $V \simeq V^*$ and $W \simeq W^*$ where \simeq means anti-isomorphic. $T : W \rightarrow V \implies T^* : V^* \rightarrow W^*$.

8.8 Adjoint maps

Definition 8.18 (Theorem and definition). Let $(V, \langle \cdot, \cdot \rangle_V)$ and $(W, \langle \cdot, \cdot \rangle_W)$ be spaces with a scalar product. $\dim V, \dim W < \infty$.

$T \in \text{Hom}(W, V)$ hence, $T : W \rightarrow V$ linear

1. For every $v \in V$ the map $w \mapsto \langle T(w), v \rangle_V$ is linear.
2. $\forall v \in V \exists! u \in W \forall w \in W : \langle T(w), v \rangle_V = \langle w, u \rangle_W$ and $T^*(v) = u$.

Hence,

$$\langle T(w), v \rangle_V = \langle w, T^*(v) \rangle_W \quad \forall w \in W \quad \forall v \in V$$

3. The map $T^* : V \rightarrow W$ with $v \mapsto u$ is linear, hence $T^* \in \text{Hom}(V, W)$ and is called adjoint map.
4. The map $\text{Hom}(W, V) \mapsto \text{Hom}(V, W)$ with $T \mapsto T^*$ is antilinear and $T^{**} = T$.

Proof. 1. $\langle T(w), v \rangle = T_V(T(w)) = T_v \circ T(w)$
Composition of linear maps is linear.

2. $T_V \circ T \in W^*$. By Riesz representation theorem, $\exists! u \in W : T_V \circ T(w) = \langle w, u \rangle \forall w \in W = \langle T(w), v \rangle = \langle w, u \rangle$
3. Show that,

$$\forall v_1, v_2 \in V \forall \lambda, \mu : T^*(\lambda v_1 + \mu v_2) = \lambda T^*(v_1) + \mu T^*(v_2)$$

It suffices to show that

$$\langle w, T^*(\lambda v_1 + \mu v_2) \rangle = \langle w, \lambda T^*(v_1) + \mu T^*(v_2) \rangle \quad \forall w \in W$$

Compare with corollary: $w_1 = w_2$ in $W \iff \forall w : \langle w, w_1 \rangle = \langle w, w_2 \rangle$.

$$\begin{aligned} \langle w, T^*(\lambda v_1 + \mu v_2) \rangle &= \langle T(w), \lambda v_1 + \mu v_2 \rangle \\ &= \bar{\lambda} \langle T(w), v_1 \rangle + \bar{\mu} \langle T(w), v_2 \rangle \\ &= \bar{\lambda} \langle w, T^*(v_1) \rangle + \bar{\mu} \langle w, T^*(v_2) \rangle \\ &= \langle w, \lambda T^*(v_1) \rangle + \langle w, \mu T^*(v_2) \rangle \\ &= \langle w, \lambda T^*(v_1) + \mu T^*(v_2) \rangle \end{aligned}$$

4. Show $(\lambda T_1 + \mu T_2)^* = \bar{\lambda} T_1^* + \bar{\mu} T_2^*$.

$$\iff \forall v \in V : (\lambda T_1 + \mu T_2)^* v = (\bar{\lambda} T_1^* + \bar{\mu} T_2^*)(v)$$

$$\forall v \in V \forall w \in W : \langle w, (\lambda T_1 + \mu T_2)^*(v) \rangle = \langle w, (\bar{\lambda} T_1^* + \bar{\mu} T_2^*)(v) \rangle$$

Hence,

$$\begin{aligned}
\langle w, (\lambda T_1 + \mu T_2)^*(v) \rangle &= \langle (\lambda T_1 + \mu T_2)(w), v \rangle \\
&= \lambda \langle T_1(w), v \rangle + \mu \langle T_2(w), v \rangle \\
&= \lambda \langle w, T_1^*(v) \rangle + \mu \langle w, T_2^*(v) \rangle \\
&= \langle w, \bar{\lambda} T_1^*(v) \rangle + \langle w, \bar{\mu} T_2^*(v) \rangle \\
&= \langle w, \bar{\lambda} T_1^*(v) + \bar{\mu} T_2^*(v) \rangle \\
&= \langle w, (\bar{\lambda} T_1^* + \bar{\mu} T_2^*)(v) \rangle
\end{aligned}$$

$$T : W \rightarrow V \quad T^* : V \rightarrow W \quad T^{**} : W \rightarrow V$$

Show that $\forall w \in W : T^{**}(w) = T(w)$. Hence $\forall w \in W \forall v \in V : \langle T^{**}(w), v \rangle_V = \langle T(w), v \rangle_V$

$$\begin{aligned}
\langle T^{**}(w), v \rangle_V &= \overline{\langle v, T^{**}(w) \rangle} = \overline{\langle T^*(v), w \rangle} = \langle w, T^*(v) \rangle \\
&= \langle T(w), v \rangle \\
\langle T w, v \rangle &= \langle w, T^* v \rangle
\end{aligned}$$

If $V = W$, then $T = T^*$.

□

Assume $u = D^*(x)$ exists $\in \mathbb{R}[x]$

$$\begin{aligned}
&\implies M := \max_{t \in [0,1]} |u(t)| < \infty \\
||x^n| D^*(x)| &= \left| \int_0^1 t^n \cdot u(t) dt \right| \leq \int_0^1 t^n \cdot M dt = \frac{M}{n+1} \\
&\implies \frac{n}{n+1} \leq \frac{M}{n+1} \forall n \in \mathbb{N} \\
&\implies u(x) \notin \mathbb{R}[x]
\end{aligned}$$

Example 8.18 (For Definition 8.18, point 3). If $\dim V = \infty$, then not every linear map has an adjoint map!

$$V = \mathbb{R}[x] \quad \langle f, g \rangle = \int_0^1 f(t)g'(t) dt$$

$$D : V \rightarrow V \quad p(x) \mapsto p'(x)$$

Recall: The derivative of a linear combination is the linear combination of derivatives. Assume D has an adjoint D^* .

$$\implies \langle x^n, D^*(x) \rangle = \langle D(x^n), x \rangle = \int_0^1 n t^{n-1} t dt = \frac{n}{n+1}$$

↓ This lecture took place on 2018/05/02.

Riesz representation theorem
 V with scalar product
 $\text{Hom}(V, \mathbb{K}) \simeq V$ where \simeq is antilinear
 $\forall f \in \text{Hom}(V, \mathbb{K}) : \exists! y \in V : f = T_y$

$$T_y(x) = \langle x, y \rangle$$

$$T_{\lambda x + \mu y} = \bar{\lambda} T_x + \bar{\mu} T_y$$

For $f \in \text{Hom}(V, W)$, the map $x \mapsto \langle f(x), y \rangle \in \text{Hom}(V, \mathbb{K})$

$$\implies \exists! z \in V : \forall x \in V : \langle f(x), y \rangle = \langle x, z \rangle$$

$$z =: f^*(y) \dots \text{adjoint map}$$

$$f^* : W \rightarrow V \text{ is linear}$$

$$\text{Hom}(V, W) \rightarrow \text{Hom}(W, V)$$

$$f \mapsto f^*$$

is an antilinear *involution*.

$$f^{**} = f$$

8.9 The linear adjoint map is the complex transpose

Theorem 8.24. Let $B \subseteq V, C \subseteq W$ be orthonormal bases. $f \in \text{Hom}(V, W)$.

$$\Phi_B^C(f^*) = \Phi_C^B(f)^* = \overline{\Phi_C^B(f)}^T$$

Proof.

$$A = \Phi_C^B(f)$$

Column $s_j(A)$ are the coordinates of $b_j \in B$ in regards of basis C .

$$\begin{aligned} a_{ij} &= \text{i-th coordinate of } f(b_j) \\ &= \Phi_C(f(b_j))_i = \langle f(b_j), c_i \rangle \\ &= \langle b_j, f^*(c_i) \rangle = \overline{\langle f^*(c_i), b_j \rangle} \\ &= \text{j-th coordinate of } f^*(c_i) \\ &= \overline{\Phi_B^C(f^*)_{ji}} = \bar{a}_{ji} \end{aligned}$$

$$\text{if } \tilde{A} = \Phi_B^C(f^*)$$

□

Theorem 8.25. *Let U, V, W be finite-dimensional.*

$$U \xrightarrow{f} V \xrightarrow{g} W$$

1. $(g \circ f)^* = f^* \circ g^*$
2. $f^{**} = f$
3. $\ker f = (\text{image } f^*)^\perp$
4. $\text{image } f = (\ker f^*)^\perp$
5. $f \text{ injective} \iff f^* \text{ surjective}$
6. $f \text{ surjective} \iff f^* \text{ injective}$

Proof. 1. Let $u \in U, w \in W$

$$\begin{aligned} \langle (g \circ f)(u), w \rangle_W &= \langle g(f(u)), w \rangle_W \\ &= \langle f(u), g^*(w) \rangle_V \\ &= \langle u, f^*(g^*(w)) \rangle_U \end{aligned}$$

holds $\forall u \in U \forall w \in W$. By definition

$$\langle (g \circ f)(u), w \rangle_W = \langle u, (g \circ f)^*(w) \rangle$$

Hence,

$$\implies (g \circ f)^* = f^* \circ g^*$$

3. Show that

- $\ker f \subseteq (\text{image } f^*)^\perp$
- $(\text{image } f^*)^\perp \subseteq \ker f$

Proof:

- Let $u \in \ker f$. Show that $\forall y \in \text{image } f^* : \langle u, y \rangle = 0$

$$y \in \text{image } f^* \implies \exists v \in V : y = f^*(v)$$

$$\langle u, y \rangle_U = \langle u, f^*(v) \rangle_U = \left\langle \underbrace{f(u)}_{=0}, \underbrace{v}_V \right\rangle = 0$$

- Let $u \in (\text{image } f^*)^\perp$, hence $\forall v \in V : u \perp f^*(v)$. Hence $\forall v \in V :$
 $\langle u, f^*(v) \rangle_U = 0.$

$$\forall v \in V : \langle f(u), v \rangle_V = 0$$

$$\implies f(u) \in V^\perp = \{0\}$$

$$\implies u \in \ker f$$

4. Apply (3) to f^* .

$$\begin{aligned} \ker f^* &= (\text{image } f^{**})^\perp = (\text{image } f)^\perp \\ \implies (\ker f^*)^\perp &= (\text{image } f)^{\perp\perp} \underbrace{=}_{\dim < \infty} \text{image } f \end{aligned}$$

□

Remark 8.22 (Addition to Theorem 8.17). *So, if subspace $U \subseteq V$. Then $U^{\perp\perp} = U$.*

Proof: It holds that $U + U^\perp = V$ and $U^\perp + U^{\perp\perp} = V$. $U \subseteq U^{\perp\perp}$ and $\dim U = \dim U^{\perp\perp} \implies U = U^{\perp\perp}$.

8.10 Unitary transformations and self-adjoint matrices

Definition 8.19. *Let V be a vector space with scalar product.*

1. $f : V \rightarrow V$ is called *self-adjoint*, if $f = f^*$. Hence $\forall x, y \in V : \langle f(x), y \rangle = \langle x, f(y) \rangle \iff \Phi_B^B(f) = \Phi_B^B(f)^*$ if B is orthonormal basis of V .
2. $f \in \text{Hom}(V, W)$ is called *unitary transformation* or *linear isometry* if

$$\forall x, y \in V : \langle f(x), f(y) \rangle = \langle x, y \rangle$$

*esp. $\|f(x)\| = \|x\|$, hence lengths (and also angles) are preserved.
mostly it is additionally required that f is invertible.*

Remark 8.23. 1. *Unitary transformations are injective.*

2. *If $\dim V = \dim W < \infty$ and $f : V \rightarrow W$ is linear and unitary, then f is regular and $f^{-1} = f^*$.*
3. *If $\dim V = \infty$, $f : V \rightarrow V$ is isometry, it does not imply that f is invertible.*

Proof. 1. Recognize that $\langle f(x), f(x) \rangle = \langle x, x \rangle$ implies $\|f(x)\|^2 = \|x\|^2$. Then
 $f(v) = 0 \implies \|f(v)\| = \|v\| = 0 \implies v = 0$

$$\ker f = \{0\}$$

2. f unitary $\stackrel{(1.)}{\implies} f$ injective $\implies f$ surjective.

$$\begin{aligned} \forall x, y \in V : \langle x, y \rangle &= \langle f(x), f(y) \rangle \\ &= \langle x, f^* \circ f(y) \rangle \end{aligned}$$

hence for fixed y , it holds that

$$\begin{aligned} \forall x \in V : \langle x, y \rangle &= \langle x, f^* \circ f(y) \rangle \\ \implies y &= f^* \circ f(y) \text{ for all } y \implies f^* \circ f = \text{id} \end{aligned}$$

$$3. V = l^2 = \left\{ (x_n)_n \mid \sum |x_n|^2 < \infty \right\}$$

$$S : l^2 \rightarrow l^2$$

$$(x_1, x_2, \dots) = (0, x_1, x_2, \dots)$$

$$\|S(x)\| = \|x\|$$

$$\langle S(x), S(y) \rangle = \langle (0, x_1, x_2, \dots), (y_1, y_2, \dots) \rangle$$

$$= 0 + \sum_{i=1}^{\infty} x_i \overline{y_i}$$

$$= \langle x, y \rangle$$

$$\langle x, S^* y \rangle = \langle Sx, y \rangle$$

$$= \langle (0, x_1, x_2, \dots), (y_1, y_2, \dots) \rangle$$

$$= 0 \cdot \overline{y_1} + x_1 \cdot \overline{y_2} + x_2 \cdot \overline{y_3} + \dots$$

$$= \langle (x_1, x_2, \dots), (y_1, y_2, \dots) \rangle$$

$$S^*(y_1, y_2, \dots) = (y_2, y_3, \dots)$$

$$\langle S_x, S_y \rangle = \langle x, S^* S y \rangle \forall x, y$$

$$\implies S^* \circ S = \text{id}$$

$$\text{but } S \circ S^*(x_1, x_2, \dots) = S(x_2, x_3, \dots)$$

$$= (0, x_2, x_3, \dots)$$

$$\implies S \circ S^* \neq \text{id}$$

S is not invertible

This shifting of indices works in a finite number of dimensions, but does not work in infinity (in this case, you miss one dimension).

□

8.11 Unitary matrices and orthogonal matrices

Definition 8.20. 1. A matrix U is called unitary if $U^* U = I$

2. A matrix $U \in \mathbb{R}^{n \times n}$ is called orthogonal if $U^T U = I$

Theorem 8.26. For a matrix $T \in \mathbb{C}^{n \times n}$ it holds equivalently:

1. T is unitary ($T^* \cdot T = I$)

2. $\forall x \in \mathbb{C}^n : \|Tx\| = \|x\|$ (isometry)

3. $\forall x, y \in \mathbb{C}^n : \Re \langle Tx, Ty \rangle = \Re \langle x, y \rangle$

4. $\forall x, y \in \mathbb{C}^n : \langle Tx, Ty \rangle = \langle x, y \rangle$

5. The columns of T define an orthonormal basis of \mathbb{C}^n

Proof. 1. \rightarrow 2.

$$\|Tx\|^2 = \langle Tx, Tx \rangle = \langle x, T^*Tx \rangle = \langle x, Ix \rangle = \|x\|^2$$

2. \rightarrow 3.

$$\begin{aligned} \|T(x+y)\|^2 &= \|x+y\|^2 \\ \|T(x-y)\|^2 &= \|x-y\|^2 \\ \|Tx+Ty\|^2 &= \|Tx\|^2 + 2\Re\langle Tx, Ty \rangle + \|Ty\|^2 \\ \|Tx-Ty\|^2 &= \|Tx\|^2 - 2\Re\langle Tx, Ty \rangle + \|Ty\|^2 \\ \hline \|Tx+Ty\|^2 - \|Tx-Ty\|^2 &= 4\Re\langle Tx, Ty \rangle \\ \text{analogously, } \|x+y\|^2 - \|x-y\|^2 &= 4\Re\langle x, y \rangle \\ \hline \implies \Re\langle Tx, Ty \rangle &= \Re\langle x, y \rangle \end{aligned}$$

3. \rightarrow 4.

$$\Re\langle Tx, Ty \rangle = \Re\langle x, y \rangle \quad \forall x, y \in \mathbb{C}^n$$

also holds for $i \cdot y$ instead of y

$$\Re\langle Tx, iTy \rangle = \Re\langle x, iy \rangle \quad \forall x, y \in \mathbb{C}^n$$

$$\Re(-i\langle Tx, Ty \rangle) = \Re(-i\langle x, y \rangle)$$

$$\Re(-i(a+ib)) = \Re(-ia+b) = b$$

$$\Re(-i \cdot z) = \Im(z)$$

$$\Im\langle Tx, Ty \rangle = \Im\langle x, y \rangle \quad \forall x, y \in \mathbb{C}^n$$

\Re and \Im are equivalent.

$$\implies \langle Tx, Ty \rangle = \langle x, y \rangle \quad \forall x, y$$

(this is a common proof pattern, that you only show it for \Re and \Im follows immediately)

4. \rightarrow 5. e_1, \dots, e_n define some orthonormal basis.

$$\implies (Te_1, \dots, Te_n) \text{ is orthonormal basis}$$

$$u_i = Te_i = \text{i-th column of } T$$

$$\langle u_i, u_j \rangle = \langle Te_i, Te_j \rangle = \langle e_i, e_j \rangle = \delta_{ij}$$

5. \rightarrow 4. $(T^*T)_{ij}$ is the i -th column vector of T^* times the j -th column vector of T .

$$u_j^* \cdot u_j = \langle u_j, u_j \rangle = \delta_{jj}$$

$$\Rightarrow T^*T = \begin{bmatrix} 1 & \dots & 0 \\ & \ddots & \\ 0 & \dots & 1 \end{bmatrix} = I$$

□

What do isometries of \mathbb{R}^n or \mathbb{C}^n look like?

Definition 8.21. An isometry between two metric spaces (M_1, d_1) and (M_2, d_2) . Metric d :

$$d(x, y) \geq 0$$

$$d(x, y) = 0 \iff x = y$$

$$d(x, y) \leq d(x, z) + d(z, y)$$

is a map $f : M_1 \rightarrow M_2$ such that

$$d_2(f(x), f(y)) = d_1(x, y)$$

Every normed space has metric $d(x, y) = \|x - y\|$. An isometry between two spaces is a (not necessarily linear) map $f : V \rightarrow W$ such that $\|f(x) - f(y)\| = \|x - y\|$.

Example 8.19 (Translation).

$$x_0 \in V \quad T_{x_0} : V \rightarrow V \quad x \mapsto x + x_0$$

Translation T_{x_0} is an isometry because $\|T_{x_0}(x) - T_{x_0}(y)\| = \|x + x_0 - (y + x_0)\| = \|x - y\|$. But translation is not unitary because of non-linearity: $T_{x_0}(0) = 0 + x_0 \neq 0$.

Other examples in \mathbb{R}^n :

1. rotation
2. reflection
3. unitary/orthogonal map

Example 8.20 (Rotation in \mathbb{R}^2).

$$U(e_1) = \begin{pmatrix} \cos \alpha \\ \sin \alpha \end{pmatrix}$$

$$U(e_2) = \begin{pmatrix} -\sin \alpha \\ \cos \alpha \end{pmatrix}$$




Figure 11: Rotation in \mathbb{R}^2

Compare with Figure 11.

$$U_\alpha = \begin{bmatrix} \cos \alpha & \dots & -\sin \alpha \\ & \ddots & \\ \sin \alpha & \dots & \cos \alpha \end{bmatrix} = \begin{bmatrix} 1 & \\ & 1 \end{bmatrix} \cdot \cos \alpha + \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \cdot \sin \alpha$$

Tangent a:

$$\begin{aligned} \begin{pmatrix} x(t) \\ y(t) \end{pmatrix} &= \begin{pmatrix} \dot{x}(t) \\ \dot{y}(t) \end{pmatrix} \\ \vec{x}(t) &\perp \dot{\vec{x}}(t) \\ \dot{\vec{x}}(t) &= \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \vec{x}(t) \\ \vec{x}(t) &= e^{\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} t} \cdot \vec{x}_0 \end{aligned}$$

Compare with Figure 12.

$$x'(t) = a \cdot x(t) \implies x(t) = c \cdot e^{at}$$

Example 8.21 (Rotation considered as motion).




Figure 12: Rotation in \mathbb{R}^2 considered as motion. Commonly done by physicists.

$$\begin{aligned}\frac{dx}{dt} &= ax \\ dx &= ax \cdot dt \\ \int \frac{dx}{x} &= \int a \cdot dt \\ \log x &= at + C \\ x &= C_1 \cdot e^{at}\end{aligned}$$

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$e \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}^t = \sum_{n=0}^{\infty} \frac{\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}^n}{n!} t^n$$

$$e^{it} = \cos t + i \cdot \sin t$$

insert $\sum_{n=0}^{\infty} \frac{(it)^n}{n!}$ and split \Re and \Im .

$$\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}^2 = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$$

$$\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}^3 = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}^4 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$i^2 = -1 \quad i^3 = -i \quad i^4 = 1$$

$$\begin{aligned}
e^{\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} t} &= \cos(t) \cdot \begin{bmatrix} 1 & \\ & 1 \end{bmatrix} + \sin(t) \cdot \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \\
U_{\alpha+\beta} &= U_\alpha \cdot U_\beta \\
\begin{bmatrix} \cos(\alpha+\beta) & -\sin(\alpha+\beta) \\ \sin(\alpha+\beta) & \cos(\alpha+\beta) \end{bmatrix} &= \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} \cdot \begin{bmatrix} \cos \beta & -\sin \beta \\ \sin \beta & \cos \beta \end{bmatrix} \\
&= \begin{bmatrix} \cos \alpha \cos \beta - \sin \alpha \sin \beta & -\cos \alpha \sin \beta - \sin \alpha \cos \beta \\ \sin \alpha \cos \beta + \cos \alpha \sin \beta & \sin \alpha \sin \beta + \cos \alpha \cos \beta \end{bmatrix}
\end{aligned}$$

Example 8.22 (Reflection in \mathbb{R}^2).

$$\begin{aligned}
S(e_1) &= \begin{bmatrix} \cos(2\varphi) \\ \sin(2\varphi) \end{bmatrix} \\
S(e_2) &= \begin{bmatrix} \cos(2\varphi - \frac{\pi}{2}) \\ \sin(2\varphi - \frac{\pi}{2}) \end{bmatrix} = \begin{bmatrix} \sin(2\varphi) \\ -\cos(2\varphi) \end{bmatrix} \\
\frac{\pi}{2} - 2\psi &= \frac{\pi}{2} - 2(\frac{\pi}{2} - \varphi) = 2\varphi - \frac{\pi}{2} \\
S &= \begin{bmatrix} \cos(2\varphi) & \sin(2\varphi) \\ \sin(2\varphi) & -\cos(2\varphi) \end{bmatrix}
\end{aligned}$$

↓ This lecture took place on 2018/05/07.

Linear isometries:

Theorem 8.27.

$$O(n) = \{ U \in \mathbb{R}^{n \times n} \mid U^T U = I \} \quad \text{orthogonal group}$$

$$\mathcal{U}(n) = \{ U \in \mathbb{C}^{n \times n} \mid U^* U = I \} \quad \text{unitary group}$$

$$SO(n) = \{ U \in O(n) \mid \det(U) = 1 \} \subseteq O(n) \quad \text{subgroup, special orthogonal group}$$

$$SU(n) = \{ U \in \mathcal{U}(n) \mid \det(U) = 1 \} \subseteq \mathcal{U}(n) \quad \text{subgroup, special unitary group}$$

$$\mathcal{GL}(n, \mathbb{K}) = \{ A \in \mathbb{K}^{n \times n} \mid \text{invertible} \} \quad \text{general linear group}$$

$$SL(n, \mathbb{K}) = \{ A \in GL(n) \mid \det(A) = 1 \} \quad \text{special linear group}$$

Then, e.g. $O(2)$ is the group of rotations and reflections.

Remark 8.24. For $U \in \mathcal{U}(n)$ it holds that $|\det(U)| = 1$. Why?

We know: $U^* U = I \implies \det(U^* U) = \det(I) = 1 = \det(U^*) \det(U) = \det(\overline{U}^T) \det(U) = \overline{\det(U)} \det(U) = |\det(U)|^2 = 1$.

Example 8.23 (Rotation).

$$U = \begin{bmatrix} \cos \varphi & -\sin \varphi \\ \sin \varphi & \cos \varphi \end{bmatrix}$$

$$\det U_\varphi = \cos^2(\varphi) + \sin^2(\varphi) = 1 \quad \implies U_\varphi \in \mathcal{SO}(2)$$

$$S_\varphi = \begin{bmatrix} \cos(2\varphi) & \sin(2\varphi) \\ \sin(2\varphi) & -\cos(2\varphi) \end{bmatrix}$$

$$\det(S_\varphi) = -\cos^2(2\varphi) - \sin^2(2\varphi) = -1$$

General orthogonal matrix in $O(2)$.

$$U = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \text{ with } \bar{U}U = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} a & c \\ b & d \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a^2 + c^2 & ab + cd \\ ab + cd & b^2 + d^2 \end{bmatrix} \stackrel{!}{=} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

Resulting constraints:

$$a^2 + c^2 = 1 \tag{1}$$

$$b^2 + d^2 = 1 \tag{2}$$

$$ab + cd = 0 \tag{3}$$

$$a = \cos \varphi \quad c = \sin \varphi \quad b = \cos \psi \quad d = \sin \psi$$

$$\cos \varphi \cdot \cos \psi + \sin \varphi \cdot \sin \psi = 0$$

$$\begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \beta & -\sin \beta \\ \sin \beta & \cos \beta \end{bmatrix} = \begin{bmatrix} \cos(\alpha + \beta) \\ \sin(\alpha + \beta) \end{bmatrix}$$

$$\cos \alpha \cos \beta - \sin \alpha \sin \beta = \cos(\alpha + \beta)$$

$$\cos \varphi \cdot \cos \psi = \cos(\varphi - \psi)$$

$$\cos \alpha = 0 \text{ for } \alpha = \frac{\pi}{2} + k \cdot \pi = (k + \frac{1}{2})\pi \quad (k \in \mathbb{Z})$$

$$\implies \varphi - \psi = (k + \frac{1}{2})\pi$$

$$\varphi = \psi + (k + \frac{1}{2})\pi$$

$$\cos \varphi = \cos(\psi + (k + \frac{1}{2})\pi) = \cos \psi \cos(k + \frac{1}{2})\pi - \sin \psi \underbrace{\sin(k + \frac{1}{2})\pi}_{\varepsilon \in \{\pm 1\}}$$

$$= -\varepsilon \cdot \sin \psi \implies \sin \psi = -\varepsilon \cos \varphi$$

$$\begin{aligned}
& \sin \alpha \cos \beta + \cos \alpha \sin \beta = \sin(\alpha + \beta) \\
\sin(\varphi) &= \sin(\psi + (k + \frac{1}{2})\pi) = \underbrace{\sin \psi \cos \left(k + \frac{1}{2}\right)\pi}_{=\varepsilon \cdot \cos(\psi)} + \underbrace{\cos \psi \sin \left(k + \frac{1}{2}\right)\pi}_{=0} \\
& \cos \psi = \varepsilon \sin \varphi \\
U &= \begin{bmatrix} \cos \varphi & \varepsilon \cdot \sin(\psi) \\ \sin \varphi & -\varepsilon \cos \varphi \end{bmatrix} = \underbrace{\begin{bmatrix} \cos \varphi & -\sin \varphi \\ \sin \varphi & \cos \varphi \end{bmatrix}}_{\substack{\text{rotation} \\ \det=1}} \cdot \underbrace{\begin{bmatrix} 1 & \\ & -\varepsilon \end{bmatrix}}_{\substack{\varepsilon=1: \\ \text{reflection on } x\text{-axis} \\ \varepsilon=-1: \text{id}}} \\
U_\varphi &= \cos \varphi \begin{bmatrix} 1 & \\ & 1 \end{bmatrix} + \sin \varphi \begin{bmatrix} & -1 \\ 1 & \end{bmatrix}
\end{aligned}$$

Hence, every orthogonal matrix is either a rotation ($\det = 1$) or a reflection ($\det = -1$).

$$\begin{aligned}
SO(2) : \left\{ U_\varphi = \cos \varphi + i \cdot \sin \varphi \quad 1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, i = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \right\} \\
SU(2) : \left\{ a_0 + ia_1 + ja_2 + ka_3 \mid \sum a_i^2 = 1 \right\}
\end{aligned}$$

8.12 Quaternions

William Rowan Hamilton (1805–1865).

Remark 8.25 (Quaternions). *Hamilton defined the complex numbers in the modern sense in 1833.*

$$\begin{aligned}
C &= \{(a, b) \mid a, b \in \mathbb{R}\} \\
(a, b) \cdot (c, d) &= (ac - bd, ad + bc)
\end{aligned}$$

He tried to invent them over 10 years for the third dimension. He failed. On 1843/10/16, he invented the quaternions next to a bridge. It works on four dimensions, but it is non-commutative. It is a screw field (dt. Schiefkörper).

$$ij = k \quad jk = i \quad ki = j \quad ji = -k \quad kj = -i \quad ik = -j$$

anti-commutative.

$$\begin{aligned}
i^2 &= j^2 = k^2 = -1 \\
(a_0 + a_1i + a_2j + a_3k)(b_0 + b_1i + b_2j + b_3k) &\quad \text{linear} \\
(a_0 + \vec{a})(b_0 + \vec{b}) &= a_0b_0 + a_0\vec{b} + b_1\vec{a} + \vec{a} \times \vec{b}
\end{aligned}$$

$$\begin{aligned}
SO(2) &\approx \left\{ \cos \varphi + i \cdot \sin \varphi \mid \varphi \in [0, 2\pi] \right\} = \{z \in \mathbb{C} \mid |z| = 1\} = \mathcal{T} \text{ Torus} \\
SU(2) &= \left\{ a_0 + ia_1 + ja_2 + ka_3 \mid \sum a_i^2 = 1 \right\} \\
SO(2) &\approx \left\{ \cos \varphi + i \sin \varphi \mid \varphi \in [0, 2\pi] \right\}
\end{aligned}$$

9 Polynomials and algebras

Definition 9.1 (Algebra). Let \mathbb{K} be a field, a \mathbb{K} algebra, a vector space \mathcal{A} over \mathbb{K} with a multiplication operator $*$: $\mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$ with $(a, b) \rightarrow a * b$ such that

1. $a * (b + c) = a * b + a * c$ (distributive law, $a, b, c \in \mathcal{A}$)
2. $(a + b) * c = a * c + b * c$
3. $\lambda \cdot (a * b) = (\lambda \cdot a) * b = a * (\lambda \cdot b)$ ($a, b \in \mathcal{A}, \lambda \in \mathbb{K}$, associativity)

Remark 9.1. **Associativity** is not generally required.

$$a * (b * c) = (a * b) * c$$

If satisfied, it is called associative algebra.

Commutativity is not generally required.

$$a * b = b * a$$

If satisfied, it is called commutative algebra.

Example 9.1. 1. $(\mathbb{K}, +, * = \cdot)$ is a one-dimensional \mathbb{K} algebra.

2. $(\mathbb{K}^{n \times n}, +, * = \text{matrix multiplication})$ is an associative non-commutative algebra where $\mathbb{K}^{n \times n} \simeq \text{Hom}(V, V)$ and $f * g = f \circ g$.
3. $\mathbb{K}^X = \{f : X \rightarrow \mathbb{K}\}$. Let X be an arbitrary set.

$$(\lambda f + \mu g)(x) = \lambda \cdot f(x) + \mu \cdot g(x)$$

$$(f * g)(x) = f(x) \cdot g(x)$$

$(\mathbb{K}^X, +, *)$ is an associative, commutative algebra.

4. \mathbb{R}^3 with $a \times b$ is an algebra.

$$a \times b = -b \times a$$

is non-commutative and also non-associative:

$$a \times (b \times c) \neq (a \times b) \times c$$

Jacobian identity:

$$a \times (b \times c) + b \times (c \times a) + c \times (a \times b) = 0$$

5. $\mathcal{A} = \mathbb{K}^{n \times n}$

$$A * B = [A, B] = A \cdot B - B \cdot A \quad \text{“commutator”}$$

is an algebra with Jacobian identity. Lie algebra:

$$[A, [B, C]] + [B, [C, A]] + [C, [A, B]] = 0$$

$$[A, B] = -[B, A]$$

The so-called Lie groups (like $O(n)$, $U(n)$, $SO(n)$, $SU(n)$).

6. $\mathcal{A} = \mathbb{K}^{n \times n}$

$$A * B = A \cdot B + B \cdot A$$

is associative. It is an Jordan algebra. Pascual Jordan (1902–1980)⁹.

Remark 9.2. Oskar Perron (1880/05/07–1975)

Definition 9.2.

$$\mathbb{K}^\infty = \{(a_0, a_1, a_2, \dots) \mid a_i \in \mathbb{K}\}$$

$$P_{\mathbb{K}} = \{(a_0, a_1, \dots, a_n, 0, \dots) \mid n \in \mathbb{N}, a_i \in \mathbb{K}\}$$

Cauchy product:

$$(a_n)_{n \geq 0} * (b_n)_{n \geq 0} = (c_n)_{n \geq 0}$$

$$c_n = \sum_{k=0}^n a_k b_{n-k}$$

Lemma 9.1. 1. $(P_{\mathbb{K}}, *)$ is a commutative, associative algebra with one-element $(1, 0, \dots)$. The basis is given with $1, x, x^2, \dots$. The algebra is called polynomial algebra

$$\mathbb{K}[x] = \left\{ \sum_{k=0}^n a_k x^k \mid a_k \in \mathbb{K}, n \in \mathbb{N} \right\}$$

2. $(\mathbb{K}^\infty, *)$ is a commutative algebra with one-element $(1, 0, \dots)$ and is called algebra of formal power series¹⁰

$$\mathbb{K}[[x]] = \left\{ \sum_{k=0}^{\infty} a_k x^k \mid a_k \in \mathbb{K} \right\}$$

Proof. Show that $\forall a, b \in P_{\mathbb{K}} : a * b \in P_{\mathbb{K}}$, hence only finitely many c_n are $\neq 0$.

Remark: $a_k = 0 \forall k > m$ and $b_k = 0 \forall k > n$.

⁹Different Jordan than in Gauss-Jordan and different than C. Jordan (19th century) about to come

¹⁰We don't need to consider convergence. This is purely formal object.

Claim.

$$c_k = 0 \quad \forall k > m + n$$

$$\begin{aligned} c_k &= \sum_{l=0}^k a_l b_{k-l} \\ &= \sum_{l=0}^{m-1} a_l b_{k-l} \quad \text{equality if } l > m \implies a_l = 0 \\ &= 0 \end{aligned}$$

$$k > m + n, l < m \implies -l > -m \implies k - l \underbrace{>}_{\implies b_{k-l}=0} m + n - m = n$$

About the Cauchy product:

$$c_n = \sum_{k=0}^n a_k b_{n-k} = \sum_{k'=0}^n a_{n-k'} b_{k'} = (b * a)_n \quad (k' = n - k)$$

Law of distributivity:

$$\begin{aligned} [(a + b) * c]_n &= \sum_{k=0}^n (a + b)_k \cdot c_{n-k} \\ &= \sum_{k=0}^n (a_k c_{n-k}) + (b_k c_{n-k}) \\ &= (a * c)_n + (b * c)_n \end{aligned}$$

□

Definition 9.3. Let $x^0 = (1, 0, \dots)$ and $x^k = (0, \dots, 1, 0, \dots)$ create a basis. The elements of $p(x) = \mathbb{K}[x]$ are called polynomials in the formal variable x

$\deg p(x) = \max \{k \mid a_k \neq 0\}$ is called degree of the polynomial

$$\deg(0) := -\infty$$

Lemma 9.2.

1. $\deg(p(x) \cdot q(x)) = \deg(p(x)) + \deg(q(x))$
2. $\mathbb{K}[x]$ is zero-divisor-free, hence $p(x) \cdot q(x) = 0 \implies p(x) = 0 \vee q(x) = 0$

Definition 9.4. Every polynomial $p(x) \in \mathbb{K}[x]$ induces a polynomial function $p : \mathbb{K} \rightarrow \mathbb{K}$ with $\alpha \mapsto p(\alpha)$ with $p \in \mathbb{K}^{\mathbb{K}}$.

$$\implies (\lambda p + \mu q)(\alpha) = \lambda \cdot p(\alpha) + \mu \cdot q(\alpha)$$

$$(p \cdot q)(\alpha) = p(\alpha) \cdot q(\alpha)$$

The map $\mathbb{K}[x] \rightarrow \mathbb{K}^{\mathbb{K}}$ with $p(x) \mapsto$ polynomial function p is linear and multiplicative (called algebra homomorphism).

Remark 9.3. A polynomial and a polynomial function are not the same. If $|\mathbb{K}| < \infty$, a difference occurs. For example, consider \mathbb{Z}_5 :

$$|\mathbb{Z}_5^{\mathbb{Z}_5}| = 5^5$$

$$|\mathbb{K}[x]| = \infty$$

where $\mathbb{Z}_5^{\mathbb{Z}_5}$ is a set of polynomial functions and $\mathbb{K}[x]$ is a set of polynomials. For example, $\prod_{\alpha \in \mathbb{K}} (x - \alpha)$ corresponds to the polynomial function 0. Consider $\mathbb{K} = \mathbb{Z}_3$. If $\alpha \in \mathbb{Z}_3$ and $x \in \mathbb{Z}_3$, we get $(x - 0)(x - 1)(x - 2)$ and choosing any $x \in \mathbb{Z}_3$ makes at least one factor zero. Thus, the polynomial function 0 is given. Hence the map $\mathbb{K}[x] \rightarrow \mathbb{K}^{\mathbb{K}}$ is surjective but not injective.

On finite fields, every function is a polynomial function.

$$\eta_i = f(\xi_i) \quad \{\xi_1, \dots, \xi_n\} = \mathbb{K}$$

From the practicals, it will follow that there exists a polynomial of degree n such that $p(\xi_i) = \eta_i$.

Definition 9.5. An algebra homomorphism is a linear map between ψ and two \mathbb{K} -algebras \mathcal{A} and \mathcal{B} such that $\forall a, b \in \mathcal{A} : \psi(a * b) = \psi(a) * \psi(b)$.

Example 9.2. 1. $\mathbb{K}[x] \rightarrow \mathbb{K}^{\mathbb{K}}$ with $p(x) \mapsto$ polynomial function

2. Let $\alpha \in \mathbb{K}$ be fixed. $\psi_\alpha : \mathbb{K}[x] \rightarrow \mathbb{K}$ with $p(x) \mapsto p(\alpha)$ is an algebra homomorphism of $\mathbb{K}[x] \rightarrow \mathbb{K}$.

$$\psi_\alpha(\lambda p + \mu q) = (\lambda p + \mu q)(\alpha) = \lambda p(\alpha) + \mu q(\alpha) = \lambda \psi_\alpha(p) + \mu \psi_\alpha(q)$$

3. Consider $\iota : \mathbb{K} \rightarrow \mathbb{K}[x]$ with $\iota : \alpha \mapsto \alpha \cdot x^0$.

$$(\alpha \cdot x^0) \cdot (\beta \cdot x^0) = (\alpha \cdot \beta) \cdot x^0$$

Theorem 9.1 (Insertion theorem, dt. Einsetzungssatz). Let \mathcal{A} be an associative algebra with one-element $\mathbf{1}_A$ and $\iota : \mathbb{K} \rightarrow \mathcal{A}$ with $\alpha \mapsto \alpha \cdot \mathbf{1}_A$ is the insertion of \mathbb{K} .

Then for every $a \in \mathcal{A}$ the map

$$\begin{aligned} \psi_a : \mathbb{K}[x] &\rightarrow \mathcal{A} \\ \sum_{k=0}^n c_k x^k &\mapsto \sum_{k=0}^n c_k a^k \end{aligned}$$

of the unique algebra homomorphism of $\mathbb{K}[x] \rightarrow \mathcal{A}$ with the property $\psi_a(x) = a$. We say, $\mathbb{K}[x]$ is a free, associative algebra over \mathbb{K} . Every algebra homomorphism $\mathbb{K}[x] \rightarrow \mathcal{A}$ has this structure.

↓ This lecture took place on 2018/05/09.

We consider algebras as vector spaces with associative multiplication. For example, matrices and polynomials. An algebra homomorphism is linear and multiplicative.

$$\Phi(a + b) = \Phi(a) * \Phi(b)$$

A is an associative algebra with $\mathbf{1}_A$.

$$l : \mathbb{K} \rightarrow \mathcal{A}$$

$$\alpha \mapsto \alpha \cdot \mathbf{1}_{\mathcal{A}}$$

$a \in \mathcal{A} \implies \mathcal{L}(a^0, a^1, a^2, a^3, \dots) \subseteq \mathcal{A}$ subalgebra.

1. Show linear and multiplicative property.

$$\exists! \Phi_a : \mathbb{K}[x] \rightarrow \mathcal{A} \text{ algebra homomorphism}$$

such that $\Phi_a(x) = a$, namely $\Phi_a\left(\sum_{k=0}^n c_k x^k\right) = \sum_{k=0}^n c_k a^k$.

2. Every homomorphism $\Psi : \mathbb{K}[x] \rightarrow \mathcal{A}$ has this structure.

Proof. Let $a := \Psi(x) \implies \Psi(x^n) = \Psi(x)^n = a^n$ by homomorphism.

$$\Psi \text{ linear} \implies \Psi\left(\sum_{k=0}^n c_k x^k\right) = \sum_{k=0}^n c_k \Psi(x^k) = \sum_{k=0}^n c_k a^k$$

x^0, x^1, \dots give a basis of $\mathbb{K}[x]$. Hence $\Psi = \Phi_a$ with $a = \Psi(x)$. On the opposite (1.): Obviously Φ_a is linear. Multiplicative: Show that

$$\underbrace{\Psi_a(p(x) \cdot q(x))}_{=p(a) \cdot q(a)} \stackrel{!}{=} \underbrace{\Phi_a(p(x)) \cdot \Phi_a(q(x))}_{=p(a) \cdot q(a)}$$

□

Example 9.3. 1. $\mathcal{A} = \mathbb{K}$.

$$\Psi_\alpha : \mathbb{K}[x] \rightarrow \mathbb{K}$$

$$p(x) \mapsto p(\alpha)$$

2. $\mathcal{A} = \mathbb{K}^{n \times n} \approx \text{Hom}(V, V)$

$$A^0 = I \quad A^n = A \cdot A^{n-1}$$

$$l : \mathbb{K} \rightarrow \mathbb{K}^{n \times n}$$

$$\alpha \mapsto \alpha \cdot I$$

$$\Psi_\alpha : \mathbb{K}[x] \rightarrow \mathbb{K}^{n \times n}$$

$$p(x) \mapsto p(A)$$

$$\sum_{k=0}^n c_k x^k \mapsto \sum_{k=0}^n c_k \cdot A^k$$

Remark 9.4. Let $\mathbb{K}[x]$ be a free, associative algebra over \mathbb{K} with a generator. Hence, for all associative algebras \mathcal{A} , given some element $a \in \mathcal{A}$. There exists exactly one homomorphism $\varphi : \mathbb{K}[x] \rightarrow \mathcal{A}$ such that $\varphi(x) = a$.

Compare it with a free group with one generator. Is a group G generated by x such that \forall groups H , if $h \in H$ given, there exists exactly one group homomorphism $\varphi : G \rightarrow H$ such that $\varphi(x) = h$. Namely, $G = (\mathbb{Z}, +)$ is generated by **1**. Given $h \in H \rightarrow \varphi_h : \mathbb{Z} \rightarrow H_k$ and $k \mapsto h$.

Definition 9.6. A root of a polynomial $p(x) \in \mathbb{K}[x]$ is a $\xi \in \mathbb{K}$ such that $p(\xi) = \Psi_\xi(p) = 0$, hence $p(x) \in \ker \Psi_\xi$.

Remark 9.5. 1. $p(x) = c_0$ is a root $\iff c_0 = 0$

2. $p(x) = c_0 + c_1x$ is the only root, $\xi = -\frac{c_0}{c_1}$.

3. $p(x) = c_0 + c_1x + c_2x^2$ has two roots over \mathbb{C}

4. $p(x) = c_0 + c_1x + c_2x^2 + c_3x^3$ has three roots

To find roots, formulas up to fourth degree exist. For degree ≥ 5 , there is no equation.

Paolo Ruffini (1765–1822)

Niels Henrik Abel (1802–1829)

Gerolamo Cardano (1501–1576)

Remark 9.6. Cardano was a polymath.

1. founder of probability theory
2. Liber de ludo aleae: important book on probability
3. Cardan joint (dt. Kardanische Welle)
4. Gimbal (dt. Kardanische Aufhängung)
5. used $\sqrt{-1}$ as a valid expression for the first time
6. published a solution for roots of cubic polynomials (Ars Magna, 1545)

Scipione del Ferro (1465–1526)

1. used a solution for roots of cubic polynomials in competitions, kept it secret
2. came up with the same solution like Tartaglia
3. lost competitions on cubic polynomials to Antonio Fiore, because Ferro's solution was not generic enough

Niccolò Fontana Tartaglia (1500–1557)

1. Cardano cajoled Tartaglia into revealing his solution to the cubic equations by promising not to publish them.

Ludovico Ferrari (1522–1565)

Theorem 9.2 (Method by Cardano/del Ferro).

$$a_0 + a_1x + a_2x^2 + a_3x^3 = 0$$

$$x \rightarrow x + a \quad \text{such that } a_2 = 0$$

$$x^3 + px + q = 0$$

Cubus p.6 rebus aeq 20

$$x^3 + 6x = 20$$

$x = \text{res}$, $x^2 = \text{census}$, $x^3 = \text{cubus}$.

Approach: $x = u + v$.

$$u^3 + 3u^2v + 3uv^2 + v^3 + p(u + v) + q = 0$$

$$u^3 + v^3 + (3uv + p)(u + v) + q = 0$$

Requirement: u and v such that $3uv + p = 0$.

$$\begin{cases} u^3 + v^3 + q = 0 & \implies v^3 = -(q + u^3) \\ 3uv + p = 0 & \implies uv = -\frac{p}{3u} \end{cases}$$

$$u^3 \cdot v^3 = -\frac{p^3}{27}$$

$$-u^3(q + u^3) = -\frac{p^3}{27}$$

$$u^6 + qu^3 - \frac{p^3}{27} = 0$$

$$u^3 = ?$$

Equation for degree 2 by Viète, Francois (1540–1603):

$$(y - \alpha)(x - \beta) = x^2 - (\alpha + \beta)x + \alpha\beta$$

$$x^2 + px + q$$

$$p = -(\alpha + \beta)$$

$$q = \alpha \cdot \beta$$

$$\alpha = \frac{1}{2} \left[(\alpha + \beta) + \sqrt{(\alpha - \beta)^2} \right]$$

$$\beta = \frac{1}{2} \left[(\alpha + \beta) - \sqrt{(\alpha - \beta)^2} \right]$$

$$\frac{\alpha}{\beta} = \frac{1}{2} \left(\alpha + \beta \pm \sqrt{(\alpha - \beta)^2} \right) = \frac{1}{2} \left(\alpha + \beta \pm \sqrt{\underbrace{\alpha^2 + \beta^2 - 2\alpha\beta}_{(\alpha+\beta)^2 - 4\alpha\beta}} \right) = \frac{1}{2} \left(-p \pm \sqrt{p^2 - 4q} \right)$$

Hence,

$$u^3 = \frac{1}{2} \left(-q \mp \sqrt{q^2 + \frac{4p^3}{27}} \right)$$

$$u^3 = \frac{q}{2} \pm \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}$$

$$u = \sqrt[3]{-\frac{q}{2} \pm \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}}$$

$$v^3 = -q - u^3 = -\frac{q}{2} \mp \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}$$

$$x = u + v = \sqrt[3]{-\frac{q}{2} + \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}} + \sqrt[3]{-\frac{q}{2} - \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}}$$

Theorem 9.3 (Division with remainder). $p(x), q(x) \in \mathbb{K}[x]$, $q(x) \neq 0$.

Then there exists exactly one polynomial $s(x), r(x) \in \mathbb{K}[x]$,

$$p(x) = s(x) \cdot q(x) + r(x)$$

with $\deg r(x) < \deg q(x)$.

Proof. Induction over $\deg p(x)$.

Induction base

$$\deg p(x) < \deg q(x) \leadsto p(x) = 0 \cdot q(x) + p(x)$$

If $\deg p(x) \geq \deg q(x)$,

$$p(x) = \sum_{k=0}^n a_k x^k \quad q(x) = \sum_{k=0}^m b_k x^k$$

$$a_n \neq 0 \quad m \leq n \quad b_m \neq 0$$

$$p_1(x) = p(x) - \frac{a_n}{b_m} \cdot q(x) \cdot x^{n-m}$$

cancels the largest term $a_n x^n$ in $p(x)$.

$$\begin{aligned} &= \sum_{k=0}^n a_k x^k - \frac{a_n}{b_m} \sum_{k=0}^m b_k x^{k+n-m} \\ &= a_n x^n + \sum_{k=0}^{n-1} a_k x^k - \frac{a_n}{b_m} b_m \cdot x^{m+n-m} - \frac{a_n}{b_m} \sum_{k=0}^{m-1} b_k x^{k+n-m} \end{aligned}$$

what remains is a polynomial of degree $\deg p_1(x) \leq n-1$.

$$\implies p(x) = \frac{a_n}{b_m} x^{n-m} \cdot q(x) + p_1(x)$$

By induction hypothesis,

$$p_1(x) = s_1(x) \cdot q(x) + r_1(x)$$

Hence,

$$p(x) = \left(\frac{a_n}{b_m} x^{n-m} + s_1(x) \right) q(x) + r_1(x)$$

□

Example 9.4.

$$p(x) = 3x^5 - x^4 + 2x^3 + x^2 + 1$$

$$q(x) = x^2 - 3x + 1$$

$$\left| \begin{array}{ccccccccc} 3x^5 & -x^4 & +2x^3 & +x^2 & +1 & : x^2 & -3x & +1 & = 3x^2 + 8x^2 + 23x + 62 \\ -3x^5 & +9x^4 & -3x^3 & & & & & & \\ 0 & 8x^4 & -x^3 & +x^2 & +1 & & & & \\ & 8x^4 & -24x^3 & +8x^2 & & & & & \\ & 0 & 23x^3 & -7x^2 & +1 & & & & \\ & & 23x^3 & -69x^2 & +23x & & & & \\ & & 0 & 62x^2 & -23x & +1 & & & \\ & & & 62x^2 & -186x & +62 & & & \\ & & & & 163x & -61 & & & \end{array} \right|$$

Hence, $s(x) = 3x^3 + 8x^2 + 23x + 62$ and $r(x) = 163x - 61$.

Definition 9.7. $q(x)$ divides $p(x) \iff$ the remainder is zero \iff there exists $s(x)$ such that $p(x) = s(x) \cdot q(x)$.

Theorem 9.4. 1. If $p(x) = s(x) \cdot (x - \xi) + r$

$$q(x) = x - \xi \quad \implies \quad p(\xi) = r$$

2. ξ is root of $p(x) \implies x - \xi$ divides $p(x)$

Theorem 9.5 (Ruffini-Horner's method). Given $p(x) \in \mathbb{K}[x]$, $\lambda \in \mathbb{K}$. Find $p(\lambda)$.

$$\begin{aligned} p(x) &= a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0 \\ &= a_n \lambda^n + a_{n-1} \lambda^{n-1} + \dots + a_1 \lambda + a_0 \\ &= (a_n \lambda^{n-1} + \dots + a_1) \lambda + a_0 \\ &= ((a_n \lambda^{n-2} + \dots + a_1) \lambda + a_0) \lambda + a_0 \\ &= \vdots \end{aligned}$$

Algorithm:

$$\xi_n = a_n \text{ for } k = n-1, \dots, 0 \quad \xi_k = \lambda \xi_{k+1} + a_k$$

$$p(\lambda) = \xi_0$$

If $p(x) = s(x)(x - \lambda) + r$, $p(\lambda) = r$.

Horner's method provides a more convenient method to evaluate a polynomial for given x than exponentiation by a high degree.

Example 9.5.

$$3x^5 - x^4 + 2x^3 + x^2 + 1$$

$$p(5) = ? \quad \xi_5 = 3$$

$3x^5$	$-x^4$	$+2x^3$	$+x^2$	$+1$	$: (x - 5)$	$=$	$3x^4 + 14x^3 + 72x^2 + 361x + 1805$
$3x^5$	$-15x^4$						
0	$14x^4$	$+2x^3$	$+x^2$	$+1$			
	$14x^4$	$-70x^3$					
	0	$+72x^3$	$+x^2$	$+1$			
		$72x^3$	$-360x^2$				
		0	$+361x^2$	$+1$			
			$361x^2$	$-1805x$			
				$1805x$	$+1$		
				$1805x$	$-5 \cdot 1805$		
					$5 \cdot 1805 + 1$		

$$\xi_5 = 3$$

$$\xi_4 = 5 \cdot \xi_5 + (-1) = 5 \cdot 3 - 1 = 14$$

$$\xi_3 = 5 \cdot 14 + 2 = 72$$

$$\xi_2 = 5 \cdot 72 + 1 = 361$$

$$\xi_1 = 5 \cdot 361 + 0 = 1805$$

$$\xi_0 = 5 \cdot 1805 + 1 = 9026$$

Definition 9.8. A polynomial $p(x) \in \mathbb{K}[x]$ is called *reducible*, if $\exists p_1(x), p_2(x) : \deg p_1(x) < \deg p(x)$ and $p(x) = p_1(x) \cdot p_2(x)$ (is the factorization). $\deg p_2(x) < \deg p(x)$ (proper divisor). Otherwise the polynomial is called *irreducible*.

Remark 9.7. An irreducible polynomial of degree > 1 has no roots.

Example 9.6. • Consider $x^2 = -2$ irreducible over $\mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{R}$. Its roots are $\pm \sqrt{2}$.

It is reducible over \mathbb{R} : $x^2 - 2 = (x - \sqrt{2})(x + \sqrt{2})$.

It is reducible over $\mathbb{Q}(\sqrt{2}) = \{a + b\sqrt{2} \mid a, b \in \mathbb{Q}\}$.

- Consider $x^2 + 1$ irreducible over \mathbb{Q}, \mathbb{R} and reducible over \mathbb{C} . Its roots are $\pm i$.
 $\mathbb{Q}(i) = \{a + bi \mid a, b \in \mathbb{Q}\}$. $x^2 + 1 = (x + i)(x - i)$.
- Consider $\mathbb{K} = \mathbb{Z}_2$ and $p(x) = x^2 + x + 1$. This polynomial has no roots and is irreducible.
- $x^5 + x + 1$ has no roots, is reducible.

$$x^5 + x + 1 = (x^2 + x + 1)(x^3 + x^2 + 1)$$

Is there some field $\mathbb{K} \supseteq \mathbb{Z}_2$ such that $x^3 + x^2 + 1$ has roots?

Yes. Let α be a number such that $\alpha^3 + \alpha^2 + 1 = 0 \implies \alpha^3 = -\alpha^2 - 1 = \alpha^2 + 1$.

$$\mathbb{K} = \mathbb{Z}_2(\alpha) = \{a + b\alpha + c\alpha^2 \mid a, b, c \in \mathbb{Z}_2\}$$

with $\alpha^3 = \alpha^2 + 1$ is a field.

Let i be a number such that $i^2 + 1 = 0$, thus $i^2 = -1$

$$\mathbb{C} = \mathbb{R}(i) = \{a + bi \mid a, b \in \mathbb{R}\}$$

Hence, irreducible is not equivalent to some root exists. The implication works only in one direction. There always exists some field such that roots exist.

Theorem 9.6 (Fundamental theorem of Algebra). \mathbb{C} is algebraically closed, hence every polynomial has a root over \mathbb{C} .

Corollary 9.1. Every polynomial over $\mathbb{C} \dots$

1. has a factorization $p(x) = (x - \xi_1)(x - \xi_2) \dots (x - \xi_n)$.
2. $p(x)$ is irreducible $\iff \deg p(x) \leq 1$.

Remark 9.8. No algebraic proof exists. It is more like a Fundamental Theorem of Calculus over complex numbers. The proof is given by the Lionville theorem (not done here).

Theorem 9.7. For arbitrary fields, it holds that every polynomial has exactly one factorization (except for its order) in irreducible factors.

↓ This lecture took place on 2018/05/14.

9.1 The greatest common divisor of polynomials

The Euclidean algorithm determines the greatest common divisor.

Consider $n = q \cdot m + r$. For the Euclidean algorithm, it holds that $\gcd(n, m) = \gcd(m, r)$. The analogous solution holds for polynomials. Consider $p(x) = s(x) \cdot q(x) + r(x)$. Then the $\gcd(p(x), q(x))$ returns the polynomial of maximum degree that divides the polynomial with leading coefficient 1.

Corollary 9.2. The Euclidean algorithm also works for polynomials.

An application: Find all multiple roots (i.e. roots with multiplicity greater 1).

$$\begin{aligned} (x - \xi)^k | p(x) \\ \implies (x - \xi)^{k-1} | p'(x) \\ p(x) = s(x) \cdot (x - \xi)^k \\ p'(x) = s'(x) \cdot (x - \xi)^k + s(x) \cdot k \cdot (x - \xi)^{k-1} = (s'(x)(x - \xi) + s(x) \cdot k)(x - \xi)^{k-1} \\ \implies (x - \xi)^{k-1} | \gcd(p(x), p'(x)) \end{aligned}$$

10 Eigenvectors and eigenvalues

Given $f : V \rightarrow V$. Find a basis of V such that $\Phi_B^B(f)$ has the simplest possible representation. Hence,

$$A = \Phi_B^B(f) = \begin{bmatrix} a_{11} & & 0 \\ & \ddots & \\ 0 & & a_{nn} \end{bmatrix}$$

$$A \cdot e_i = \lambda_i \cdot e_i$$

Find vector $v \in V$ such that $f(v) = \lambda \cdot v$. 0 can be an eigenvalue, but not an eigenvector. Not every A has λ satisfying $\forall v : A \cdot v = \lambda \cdot v$.




Figure 13: How map f might transform a circle

Definition 10.1. $f \in \text{Hom}(V, V) = \text{End}(V)$. $\lambda \in \mathbb{K}$ is called *eigenvalue* if $\exists v \in V \setminus \{0\} : f(v) = \lambda \cdot v$. Then v is called *eigenvector of eigenvalue λ* . $\text{spec}(f) = \{\text{eigenvalues of } f\}$ is called *spectrum of f* .

In 1925 in quantum mechanisms, it was discovered that the spectrum of light is given as a linear map (spectrum in the mathematical sense).

10.1 Eigenspace

Lemma 10.1. For $\lambda \in \mathbb{K}$, $f \in \text{End}(V)$.

$$\eta_\lambda = \{v \mid f(v) = \lambda \cdot v\}$$

is a subspace and is called *eigenspace of f for eigenvalue λ* .

Proof.

$$f(v) = \lambda \cdot v \iff f(v) - \lambda \cdot v = 0 \iff (f - \lambda \cdot \text{id})(v) = 0 \iff v \in \underbrace{\ker(f - \lambda \cdot \text{id})}_{\text{subspace}}$$

□

Example 10.1. 1. $f = \mu \cdot \text{id}$. $\text{spec}(f) = \{\mu\}$. $f(v) = \mu \cdot v \forall v \in V$. $\eta_\mu = V$.

2. Let b_1, \dots, b_n be a basis of V . Let $\lambda_1, \dots, \lambda_n \in \mathbb{K}$. Then there exists a unique, linear map f such that $f(b_i) = \lambda_i \cdot b_i$. Every b_i is an eigenvector to eigenvalue λ_i .

$$\eta_\lambda = \mathcal{L}(\{b_i \mid \lambda_i = \lambda\})$$

Assume $f(v) = \lambda \cdot v$.

$$\begin{aligned} v &= \alpha_1 \cdot b_1 + \dots + \alpha_n b_n \\ f(v) &= \alpha_1 f(b_1) + \dots + \alpha_n f(b_n) \\ &= \alpha_1 \lambda_1 b_1 + \dots + \alpha_n \lambda_n b_n \\ &= \lambda(\alpha_1 b_1 + \dots + \alpha_n b_n) \end{aligned}$$

$$\implies 0 = \alpha_1(\lambda_1 - \lambda)b_1 + \dots + \alpha_n(\lambda_n - \lambda) \cdot b_n$$

$$\text{linear indep.} \implies \forall i : \alpha_i(\lambda_i - \lambda) = 0$$

hence either $\alpha_i = 0$ or $\lambda_i = \lambda$

$$\implies \text{spec}(f) = \{\lambda_1, \dots, \lambda_n\}$$

$$\Phi_B^B(f) = \begin{bmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix}$$

3. Let $V = C^\infty(\mathbb{R})$.

$$\frac{d}{dx}y(x) = \lambda \cdot y(x) \quad \frac{dy}{dx} = \lambda \cdot y$$

$$\int \frac{dy}{y} = \int \lambda \cdot dx$$

$$\log(y) = \lambda \cdot x + C \quad \text{Eigen function (compare with Fourier analysis)}$$

$$y = C \cdot e^{\lambda x}$$

$$\frac{d}{dx}e^{\lambda x} = \lambda \cdot e^{\lambda x}$$

4. Let $V = C^\infty[0, a]$.

$$\frac{d^2}{dx^2}y(x) = \lambda \cdot y(x)$$

$$\frac{d^2}{dx^2}e^{\lambda x} = \frac{d}{dx}\lambda e^{\lambda x} = \lambda^2 e^{\lambda x}$$

$$\frac{d^2}{dx^2}e^{i\omega x} = -\omega^2 e^{i\omega x}$$

$$\frac{d^2}{dx^2}\sin \omega x = \frac{d}{dx}\omega \cdot \cos(\omega x) = -\omega^2 \cdot \sin(\omega x)$$

$$\frac{d^2}{dx^2}\cos \omega x = \frac{d}{dx}(-\omega) \sin(\omega x) = -\omega^2 \cos(\omega x)$$

$$\begin{aligned}
y(0) = y_0 &\rightarrow y(x) = y_0 \cdot e^{\lambda x} \\
y(0) = y(a) &= 0 \\
y(x) &= \sin(\omega x) \\
\omega a = k \cdot \pi &\implies y(0) = y(a) = \pi \\
\omega &= \frac{k \cdot \pi}{a}
\end{aligned}$$

Eigenvalues of $H = P^2 + Q$ and $PQ - QP = \frac{\hbar}{i}I$. Heisenberg: Quantum mechanics is not commutative (impulses are matrices, not values).

Definition 10.2. Let A be a $n \times n$ matrix. λ is called right-sided eigenvalue if $\exists x \in \mathbb{K}^n \setminus \{0\} : Ax = \lambda \cdot x$. λ is called left-sided eigenvalue if $\exists x \in \mathbb{K}^n \setminus \{0\} : x^T A = \lambda \cdot x^T$. But this definition is satisfied $\iff A^T x = \lambda \cdot x$, hence right-sided eigenvalue of A^T . Thus, these definitions collapse.

Lemma 10.2. Left-sided eigenvalue \iff right-sided eigenvalue. Let λ be a right-sided eigenvalue.

$$\begin{aligned}
Ax = \lambda x &\iff (A - \lambda \cdot I) \cdot x = 0 \\
&\iff \exists x \neq 0 : x \in \ker(A - \lambda I) \\
&\iff \ker(A - \lambda I) \neq \{0\} \\
&\iff \text{rank}(A - \lambda I) < n \\
&\iff \text{rank}(A^T - \lambda I) < n \\
&\iff \ker(A^T - \lambda I) \neq \{0\} \\
&\iff \exists x \neq 0 : A^T x = \lambda \cdot x \\
&\iff \lambda \text{ is a left-sided eigenvalue}
\end{aligned}$$

Example 10.2. For $\dim = \infty$, this must not hold.

$$S : \begin{matrix} \mathbb{K}^\infty \rightarrow \mathbb{K}^\infty \\ (x_1, x_2, \dots) \mapsto (x_2, x_3, \dots) \end{matrix}$$

$$S(1, 0, \dots) = (0, 0, \dots)$$

$$\implies (1, 0, \dots) \text{ is eigenvector for eigenvalue } 0$$

hence, element of $\ker(S)$.

$$S = \begin{bmatrix} 0 & 1 & 0 & 0 \\ \vdots & 0 & 1 & 0 \\ \vdots & \vdots & 0 & 1 \\ \vdots & \vdots & \vdots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ 0 & & & \end{bmatrix}$$

$$S^T = \begin{bmatrix} 0 & & & \\ 1 & 0 & & 0 \\ & 1 & \ddots & \\ & & \ddots & \ddots \\ 0 & & & 1 & 0 \end{bmatrix}$$

$S^T(x_1, x_2, \dots) \mapsto (0, x_1, x_2)$ is injective. $\ker(S^T) = \{0\}$. Hence 0 is no eigenvalue. 0 is right-sided eigenvalue of S , but not left-sided eigenvalue.

Remark 10.1. The theory of eigenvalues in infinite-dimensional spaces is more complex than the finite-dimensional case.

Definition 10.3. For $A \in \mathbb{K}^{n \times n}$.

$$\begin{aligned} \text{spec}(A) &= \{\text{right-sided eigenvalue of } A\} \\ &= \{\text{left-sided eigenvalue of } A\} \end{aligned}$$

is called spectrum of A .

Remark 10.2 (Proof exercise). $\dim V = n, f \in \text{End}(V), B$ is basis of V .

$$\implies \text{spec}(f) = \text{spec}(\Phi_B^B(f))$$

Remark 10.3. Recall that any basis exchange can be done using some T :

$$\Phi_B^B(A) = T^{-1}AT$$

Corollary 10.1. The spectrum does not depend on the choice of the basis. Hence,

$$\text{spec}(T^{-1}AT) = \text{spec}(A)$$

Direct proof. Let x be an eigenvector of A .

$$\begin{aligned} Ax = \lambda x &\iff A \cdot I \cdot x = \lambda x \iff A \cdot T \cdot T^{-1} \cdot x = \lambda x \\ &\iff T^{-1} \cdot A \cdot T \cdot T^{-1} \cdot x = T^{-1} \cdot \lambda x = \lambda \cdot T^{-1} \cdot x \\ &\iff y := T^{-1}x \text{ is eigenvector of } T^{-1}AT \\ &\iff T^{-1}ATy = \lambda y \\ &\iff \lambda \text{ is eigenvalue of } T^{-1}AT \end{aligned}$$

□

Remark 10.4. λ is eigenvalue of A .

$$\begin{aligned} &\iff \ker(\lambda \cdot I - A) \neq \{0\} \\ &\iff \text{rank}(\lambda \cdot I - A) < n \\ &\iff \det(\lambda \cdot I - A) = 0 \end{aligned}$$

10.2 Characteristic polynomial

Theorem 10.1 (Theorem and definition).

1. $\chi_A(\lambda) := \det(\lambda \cdot I - A)$ is a polynomial function and is called characteristic polynomial of A .
2. λ is eigenvector $\iff \chi_A(\lambda) = 0$

Example 10.3.

$$A = \begin{bmatrix} -1 & 1 & 2 \\ -1 & -5 & 2 \\ 2 & -2 & -4 \end{bmatrix}$$

$$\begin{aligned} \chi_A(\lambda) = \det(\lambda I - A) &= \begin{vmatrix} \lambda + 1 & -1 & -2 \\ 1 & \lambda + 5 & -2 \\ -2 & 2 & \lambda + 4 \end{vmatrix} = \begin{vmatrix} \lambda + 1 & -1 & -2 \\ 1 & \lambda + 5 & -2 \\ 0 & 2\lambda + 12 & \lambda \end{vmatrix} \\ &= \begin{vmatrix} \lambda & -\lambda - 6 & 0 \\ 1 & \lambda + 5 & -2 \\ 0 & 2\lambda + 12 & \lambda \end{vmatrix} = \lambda \cdot \begin{vmatrix} \lambda + 5 & -2 \\ 2\lambda + 12 & \lambda \end{vmatrix} - \begin{vmatrix} -\lambda - 6 & 0 \\ 2\lambda + 12 & \lambda \end{vmatrix} \\ &= \lambda \cdot [\lambda^2 + 5\lambda + 4\lambda + 24] - \lambda(-\lambda - 6) \\ &= \lambda(\lambda^2 + 5\lambda + 4\lambda + 24 + \lambda + 6) \\ &= \lambda(\lambda^2 + 10\lambda + 30) \\ x_1 = 0 \quad \lambda_{2,3} &= \frac{-10 \pm \sqrt{10^2 - 120}}{2} = \frac{-10 \pm 2\sqrt{-5}}{2} = -5 \pm i\sqrt{5} \end{aligned}$$

Thus, the existence of eigenvalues depends on the field.

Remark 10.5. A square matrix A is invertible if and only if 0 is not an eigenvalue of A .

10.3 Symmetrical minor

Theorem 10.2. Let $A \in \mathbb{K}^{n \times n}$.

$$\implies \chi_A(x) = \det(x \cdot I - A) \text{ is polynomial of degree } n$$

specifically, $\chi_A(x) = \sum_{k=0}^n (-1)^{n-k} c_k(A) \cdot x^k$ with $c_k(A) = \sum_{\substack{j \in \{1, \dots, n\} \\ |j| = n-k}} \det(A_{jj})$ with $A_{jj} = (a_{ij})_{\substack{i \in J \\ j \in J}}$ are called symmetrical minors.

What are values of c_i ?

$$\begin{aligned} c_0 &= \det(A) \\ C_n &= 1 \\ C_{n-1} &= \sum a_{ii} = \text{Tr}(A) \end{aligned}$$

Proof. The proof is given using the Leibniz formula for determinants.

$$\begin{aligned}
\det(x \cdot I - A) &= \sum_{\pi \in \sigma_n} (-1)^\pi \prod_{i=1}^n \underbrace{(x \cdot I - A)_{\pi(i),i}}_{x \cdot \delta_{\pi(i),i} - a_{\pi(i),i}} \\
&= (x - a_{11})(x - a_{22}) \dots (x - a_{nn}) + \underbrace{\sum_{\substack{\pi \in \sigma_n \\ \pi \neq \text{id}}} (-1)^\pi \prod_{i=1}^n (x \delta_{\pi(i),i} - a_{\pi(i),i})}_{\text{for at least 2 } i, \delta_{\pi(i),i} = 0} \\
&= \text{expression of degree } n + \text{expression of degree } n - 2
\end{aligned}$$

Hence x^n stays the same. Hence the degree of $\chi_A(x)$ is n .

$$\begin{aligned}
\det \prod_{i=1}^n (x \delta_{\pi(i),i} - a_{\pi(i),i}) &= \# \{i \mid \pi(i) = i\} \\
&= \# \text{fixedpoints}(\pi)
\end{aligned}$$

Let s_1, \dots, s_n be the columns of A .

$$\det(xI - A_i) = \Delta(x \cdot e_1 - s_1, x \cdot e_2 - s_2, \dots, x \cdot e_n - s_n) = \sum_{I \subseteq \{1, \dots, n\}} \Delta(y_1, \dots, y_n)$$

$$y_i = \begin{cases} x \cdot e_i & i \in I \\ -s_{i_k} & i \in I^c \end{cases}$$

Let $k \in I$.

$$\Delta(y_1, \dots, y_{k-1}, x \cdot e_k, y_{k+1}, \dots, y_n) = \begin{vmatrix} y_1 & y_2 & \dots & y_{k-1} & 0 & y_{k+1} & \dots & y_n \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & 0 & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & x & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & 0 & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & 0 & \vdots & \vdots & \vdots \end{vmatrix}$$

Permute the k -th column into the first column: $(-1)^{k-1}$.

Permute the k -th row into the first row: $(-1)^{k-1}$.

$$= \begin{vmatrix} x & \tilde{y}_1 & \tilde{y}_2 & \dots & \tilde{y}_{k-1} & \tilde{y}_{k+1} & \dots & \tilde{y}_n \\ 0 & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{vmatrix} = x \cdot \begin{vmatrix} \tilde{y}_1 & \dots & \tilde{y}_{k-1} & \tilde{y}_{k+1} & \vdots & \tilde{y}_n \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{vmatrix}$$

where \tilde{y} is the permutation of y_i such that the k -th row moved to the first.

Every time, one x is eliminated, the corresponding row and column of A is removed. In the end,

$$x^{|I|} \cdot \underbrace{\det A_{I^c I^c}}_{\text{minor of the complement } |I^c| = n - k} \cdot (-1)^{|I^c|}$$

$$\Rightarrow \chi_A(x) = \sum_{I \subseteq \{1, \dots, n\}} x^{|I|} \cdot \det[A_{I^c I^c}] (-1)^{|I^c|} = \sum_{k=0}^n x^k (-1)^{n-k} c_k(A)$$

with $c_k(A) = \sum_{|J|=n-k} \det[A_{JJ}]$. □

↓ This lecture took place on 2018/05/16.

$$Ax = \lambda x$$

$$x \in \ker(\lambda \cdot I - A)$$

$$\chi_A(\lambda) = \det(\lambda I - A) = x^n - \text{Tr}(A)x^{n-1} + \dots (-1)^n \det(A)$$

Characteristic polynomial: $= \sum_{k=0}^n (-1)^{n-k} c_k x^k$

$$c_k = \sum_{|J|=n-k} \det[A_{JJ}]$$

$$T^{-1}AT \cdot T^{-1}x = \lambda T^{-1}x$$

Lemma 10.3.

$$\chi_{T^{-1}AT}(x) = \chi_A(x)$$

Proof.

$$\begin{aligned}
\chi_{T^{-1}AT}(x) &= \det(xI - T^{-1}AT) \\
&= \det(xT^{-1}T - T^{-1}AT) \\
&= \det(T^{-1}(x \cdot I)T - T^{-1}(A)T) \\
&= \det(T^{-1}(x \cdot I - A) \cdot T) \\
&= \det(T^{-1}) \cdot \det(xI - A) \cdot \det(T) \\
&= \frac{1}{\det T} \cdot \chi_A(x) \cdot \det T = \chi_A(x)
\end{aligned}$$

□

$$A = \begin{pmatrix} a_{11} & & 0 \\ & \ddots & \\ 0 & & a_{nn} \end{pmatrix} \rightsquigarrow \text{spec}(A) = \{a_{11}, \dots, a_{nn}\}$$

Eigenvector: e_1, \dots, e_n .

Remark 10.6 (Question). Does a change of basis exist, hence $T \in \text{GL}(n)$, such that $T^{-1}AT = \text{diag}(\lambda_1, \dots, \lambda_n)$? Then the eigenvalues are necessarily on the diagonal.

10.4 Diagonalizable matrix

Definition 10.4. A is called diagonalizable if $\exists T \in \text{GL}(n)$ such that $T^{-1} \cdot AT$ is a diagonal matrix, i.e. A is similar to a diagonal matrix.

Remark 10.7 (Recall).

Equivalence $A = PBQ$ with invertible $P, Q \iff \text{rank}(A) = \text{rank}(B)$.

Congruence $A = A^*, B = B^*$.

\exists regular $C : A = C^*BC$

\rightsquigarrow index and Sylvester's law of inertia.

Similarity $A = TBT^{-1}$ with regular T . This is related to eigenvalues.

Later on $\exists T$ such that $T^* = T^{-1}$ unitary. $T^*T = I$.

Lemma 10.4. A is diagonalizable $\iff \exists$ basis of eigenvectors.

Proof. B is regular such that

$$B^{-1}AB = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \iff \begin{cases} \exists \text{ columns } b_1, \dots, b_n \text{ define a basis} \\ AB = B \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \\ A \cdot \begin{bmatrix} b_1 & b_2 & \dots & b_n \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} = \begin{bmatrix} b_1 & b_2 & \dots & b_n \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} \cdot \begin{bmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix} \\ \begin{bmatrix} Ab_1 & Ab_2 & \dots & Ab_n \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} = \begin{bmatrix} b_1\lambda_1 & b_2\lambda_2 & \dots & b_n\lambda_n \\ \vdots & \vdots & \vdots & \vdots \end{bmatrix} \end{cases}$$

$$\iff \begin{cases} \exists \text{ basis } b_1, \dots, b_n \\ A \cdot b_i = \lambda_i \cdot b_i \quad i = 1, \dots, n \end{cases}$$

□

Example 10.4.

$$A = \begin{bmatrix} 1 & 2 & 4 \\ 4 & -3 & -8 \\ -2 & 2 & 5 \end{bmatrix}$$

$$\begin{aligned} \chi_A(\lambda) &= \det(\lambda I - A) = \begin{vmatrix} \lambda + 1 & -2 & -4 \\ -4 & \lambda + 3 & 8 \\ 2 & -2 & \lambda - 5 \end{vmatrix} = \begin{vmatrix} \lambda - 1 & -2 & -4 \\ \lambda - 1 & \lambda + 3 & 8 \\ 0 & -2 & \lambda - 5 \end{vmatrix} \\ &= (\lambda - 1) \begin{vmatrix} 1 & -2 & -4 \\ 1 & \lambda + 3 & 8 \\ 0 & -2 & \lambda - 5 \end{vmatrix} = (\lambda - 1) \begin{vmatrix} 1 & -2 & -4 \\ 0 & \lambda + 5 & 12 \\ 0 & -2 & \lambda - 5 \end{vmatrix} \\ &= (\lambda - 1)(\lambda^2 - 25 + 24) = (\lambda - 1)(\lambda^2 - 1) = (\lambda - 1)^2(\lambda + 1) \end{aligned}$$

Eigenvalue $(\lambda - 1)$ has multiplicity 2.

Eigenvector: $\ker(\lambda \cdot I - A)$

Eigenvalue: $\lambda = \pm 1$

Consider $\lambda = +1$: $\ker(I - A)$

Homogeneous equation system:

$$\begin{array}{ccc|c} 2 & -2 & -4 & 0 \\ -4 & 4 & 8 & 0 \\ 2 & -2 & -4 & 0 \\ \hline 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array}$$

$$\dim \ker(I - A) = 2. \quad 2x_1 = 2x_2 + 4x_3.$$

Basis:

$$\begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} \quad \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}$$

Consider $\lambda = -1$: $\ker(-I - A)$

$$\begin{array}{ccc|c} 0 & -2 & -4 & 0 \\ -4 & 2 & 8 & 0 \\ 2 & -2 & -6 & 0 \\ \hline 0 & -2 & -4 & \\ 0 & -2 & -4 & \\ \hline 0 & 0 & 0 & \end{array}$$

$$\dim \ker(-I - A) = 1.$$

Basis:

$$\begin{aligned} x_3 &= 1 \\ x_2 &= -2x_3 = -2 \\ x_1 &= \frac{2x_2 + 6x_3}{2} = 1 \end{aligned}$$

$$b_3 = \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix}$$

$$\text{with } B = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 0 & -2 \\ 0 & 1 & 1 \end{bmatrix} \text{ it holds that } B^{-1}AB = \begin{bmatrix} 1 & & \\ & 1 & \\ & & -1 \end{bmatrix}.$$

Example 10.5 (Application).

$$A = B^{-1} \cdot \underbrace{\begin{bmatrix} \Lambda_1 & & \\ & \ddots & \\ & & \Lambda_n \end{bmatrix}}_{\Lambda} \cdot B$$

$$A^2 = B^{-1} \Lambda B \cdot B^{-1} \Lambda B = B^{-1} \Lambda^2 B$$

$$A^3 = B^{-1} \Lambda^3 B$$

\vdots

$$A^k = B^{-1} \Lambda^k B$$

$$e^A = \sum_{k=0}^{\infty} \frac{A^k}{k!} = \sum_{k=0}^{\infty} \frac{B^{-1} \Lambda^k B}{k!} = B^{-1} \sum_{k=0}^{\infty} \frac{\Lambda^k}{k!} B = B^{-1} \begin{bmatrix} e^{\Lambda_1} & & \\ & \ddots & \\ & & e^{\Lambda_n} \end{bmatrix}$$

Remark 10.8. *Leonardo Pisano (1170–1250) wrote his book “Liber Abbaci” (1202) to introduce the Arabic numbers (and zero) in Europe. He also introduced the Fibonacci sequence using the growth of a rabbit population.*

10.5 Fibonacci sequence and golden ratio

Remark 10.9 (Fibonacci sequence).

$$F_0 = F_1 = 1$$

$$F_n = F_{n-1} + F_{n-2}$$

Can we find a formula for F_n ?

Remark 10.10. *Pingala (200 BC)*

How many ways are there for the equation $x_1 + \cdots + x_k = n$ for given n and x_i in $\{1, 2\}$? The answer is the Fibonacci sequence.

His application was the number of long syllables (2) or short syllables (1) in a sentence of given length in Sanskrit.

Remark 10.11 (Growth of Fibonacci sequence).

$$F_{n+1} = F_n + F_{n-1}$$

$$F_n = F_n$$

$$\begin{aligned} \begin{pmatrix} F_{n+1} \\ F_n \end{pmatrix} &= \begin{pmatrix} F_n + F_{n-1} \\ F_n \end{pmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{pmatrix} F_n \\ F_{n-1} \end{pmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^2 \begin{pmatrix} F_{n-1} \\ F_{n-2} \end{pmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^3 \begin{pmatrix} F_{n-2} \\ F_{n-3} \end{pmatrix} \\ &= \cdots = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n \begin{pmatrix} F_1 \\ F_0 \end{pmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}^n \begin{pmatrix} 1 \\ 1 \end{pmatrix} \end{aligned}$$

and where $\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$ is diagonalizable.

$$\chi_A(\lambda) = \begin{vmatrix} \lambda - 1 & -1 \\ -1 & \lambda \end{vmatrix} = \lambda^2 - \lambda - 1$$

$$\lambda_{1,2} = \frac{1 \pm \sqrt{1+4}}{2} = \frac{1 \pm \sqrt{5}}{2}$$

Eigenvector:

$$\begin{aligned} \lambda_1 &= \frac{1 + \sqrt{5}}{2} \\ \begin{vmatrix} \frac{1+\sqrt{5}}{2} - 1 & -1 \\ -1 & \frac{1+\sqrt{5}}{2} \end{vmatrix} &\begin{vmatrix} 0 \\ 0 \end{vmatrix} \end{aligned}$$

$$x_1 = \frac{1 + \sqrt{5}}{2} x_2 \quad b_1 = \begin{bmatrix} \frac{1 + \sqrt{5}}{2} \\ 1 \end{bmatrix}$$

$$\lambda_2 = \frac{1 - \sqrt{5}}{2}$$

$$\begin{array}{cc|c} \frac{1 - \sqrt{5}}{2} & -1 & 0 \\ -1 & \frac{1 - \sqrt{5}}{2} & 0 \end{array}$$

$$x_1 = \frac{1 - \sqrt{5}}{2} x_2$$

$$b_2 = \begin{bmatrix} \frac{1 - \sqrt{5}}{2} \\ 1 \end{bmatrix}$$

$$B = \begin{bmatrix} \frac{1 + \sqrt{5}}{2} & \frac{1 - \sqrt{5}}{2} \\ 1 & 1 \end{bmatrix}$$

$$\det B = \frac{1 + \sqrt{5}}{2} - \frac{1 - \sqrt{5}}{2} = \sqrt{5}$$

$$B^{-1} = \frac{1}{\sqrt{5}} \begin{bmatrix} 1 & \frac{-1 + \sqrt{5}}{2} \\ -1 & \frac{1 + \sqrt{5}}{2} \end{bmatrix}$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

$$B^{-1}AB = \begin{bmatrix} \frac{1 + \sqrt{5}}{2} & 0 \\ 0 & \frac{1 - \sqrt{5}}{2} \end{bmatrix}$$

$$\begin{pmatrix} F_{n+1} \\ F_n \end{pmatrix} = A^n \begin{pmatrix} 1 \\ 1 \end{pmatrix} = B \begin{bmatrix} \left(\frac{1 + \sqrt{5}}{2}\right)^n & \\ & \left(\frac{1 - \sqrt{5}}{2}\right)^n \end{bmatrix} \cdot B^{-1} \cdot \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

$$F_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1 + \sqrt{5}}{2} \right)^{n+1} - \left(\frac{1 - \sqrt{5}}{2} \right)^{n+1} \right]$$

$$\frac{F_{n+1}}{F_n} = \frac{\left(\frac{1 + \sqrt{5}}{2} \right)^{n+2} - \left(\frac{1 - \sqrt{5}}{2} \right)^{n+2}}{\left(\frac{1 + \sqrt{5}}{2} \right)^{n+1} - \left(\frac{1 - \sqrt{5}}{2} \right)^{n+1}} \xrightarrow{n \rightarrow \infty} \frac{1 + \sqrt{5}}{2}$$

is the Golden ratio. This is the ratio:

$$\frac{a}{a + b} = \frac{b}{a}$$

$$\frac{F_n}{F_{n-1}} = \frac{1}{1 + \frac{1}{1 + \dots}}$$

Theorem 10.3. *Eigenvectors corresponding to different eigenvalues are linear independent.*

Proof. Let $\lambda_1, \dots, \lambda_s$ be different eigenvalues. Let v_1, \dots, v_r be the respective eigenvectors.

Induction over r .

Case $r = 1$ immediate, $v_1 \neq 0$.

Case $r - 1 \rightarrow r$ Let $\alpha_1 v_1 + \dots + \alpha_r v_r = 0$.

$$\implies A(\alpha_1 v_1 + \dots + \alpha_r v_r) = 0$$

$$\alpha_1 \cdot A v_1 + \dots + \alpha_r A v_r = 0$$

$$\alpha_1 \lambda_1 v_1 + \dots + \alpha_r \lambda_r v_r = 0$$

$$\begin{array}{rcl} (1) \alpha_1 v_1 + \alpha_2 v_2 + \dots & & \alpha_r v_r = 0 \\ (2) \lambda_1 \alpha_1 v_1 + \lambda_2 \alpha_2 v_2 + \dots & & \lambda_r \alpha_r v_r = 0 \\ \hline (2) - \lambda_r (1) (\lambda_1 - \lambda_r) \alpha_1 v_1 + (\lambda_2 - \lambda_r) \alpha_2 v_2 + \dots + (\lambda_{r-1} - \lambda_r) \alpha_{r-1} v_{r-1} + (\lambda_r - \lambda_r) \alpha_r v_r & & = \end{array}$$

By induction hypothesis: v_1, \dots, v_{r-1} are linear independent.

$$\implies (\lambda_1 - \lambda_r) \alpha_1 = 0$$

$$(\lambda_2 - \lambda_r) \alpha_2 = 0$$

$$\vdots$$

$$(\lambda_{r-1} - \lambda_r) \alpha_{r-1} = 0$$

By hypothesis: $\lambda_i - \lambda_r \neq 0 \forall i < r$

$$\implies \alpha_1 = \alpha_2 = \dots = \alpha_{r-1} = 0$$

$$(1) \implies \alpha_r \cdot v_r = 0 \implies \alpha_r = 0 \text{ because } v_r \neq 0$$

□

Corollary 10.2. *An $n \times n$ matrix with n different eigenvalues is diagonalizable.*

Hence, for every eigenvalue there exists some eigenvector. They are linear independent and n elements. Hence they define a basis.

Example 10.6.

$$A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

$$\chi_A(\lambda) = \begin{vmatrix} \lambda & -1 \\ 0 & \lambda \end{vmatrix} = \lambda^2$$

$$\text{spec}(A) = \{0\}$$

$$\dim \ker(A) = 1$$

is not a basis of eigenvectors.

$$A^2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}^2 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

A is nilpotent, hence a square matrix M such that $M^k = 0$ for any $k \in \mathbb{N}_{\geq 1}$.

10.6 Multiplicities of eigenvalues

Definition 10.5. Let λ be the eigenvalue of a matrix $A \implies \chi_A(\lambda) = 0$.

$$d(\lambda) = \dim \ker(\lambda I - A) > 0$$

is called geometric multiplicity of the eigenvalue.

$k(\lambda)$ is the multiplicity of λ as root of $\chi_A(\lambda)$ and is called algebraic multiplicity of the eigenvalue.

$$d(\lambda) \leq k(\lambda)$$

Lemma 10.5. A matrix is diagonalizable if and only if for different eigenvalues $\lambda_1, \dots, \lambda_r$ it holds that

$$d(\lambda_1) + d(\lambda_2) + \dots + d(\lambda_r) = n$$

Proof.

Direction \implies : There exists a basis of eigenvectors b_1, \dots, b_n .

$$V = \eta_{\lambda_1} + \dots + \eta_{\lambda_r} \quad \eta_{\lambda_i} = \ker(\lambda_i I - A)$$

is a direct sum (because eigenvectors for different eigenvalues are linear independent). Let $v_1 \in \eta_{\lambda_1}, \dots, v_r \in \eta_{\lambda_r}$ such that $v_1 + \dots + v_r = 0$.

$$Av_i = \lambda_i v_i \implies v_1, \dots, v_r \text{ are linear independent} \implies \text{all } v_i = 0$$

$$\implies n = \dim V = \dim(\eta_{\lambda_1}) + \dots + \dim(\eta_{\lambda_r}) = d(\lambda_1) + \dots + d(\lambda_r)$$

Direction \Leftarrow . Let B_j be the basis of η_{λ_j} , hence $|B_j| = d(\lambda_j)$. The sum $\eta_{\lambda_1} + \dots + \eta_{\lambda_r}$ is direct. $\implies B_1 \cup \dots \cup B_r$ is linear independent.

$$|B_1 \cup \dots \cup B_r| = \sum_{j=1}^r d(\lambda_j) \underbrace{=}_{\text{by induction}} n$$

$B_1 \cup \dots \cup B_n$ is basis of \mathbb{K}^n of eigenvectors.

□

Remark 10.12. *Why are there at most n eigenvalues?*

We know the characteristic polynomial has degree n .

For every eigenvalue, we have an algebraic multiplicity. Every algebraic multiplicity is at least 1. If there are more than n eigenvalues, namely $n + 1$, the sum of algebraic multiplicities of eigenvalues is at least $n + 1$. Thus the degree of the characteristic polynomial (which is the sum of algebraic multiplicities of eigenvalues), is at least $n + 1$ which contradicts our previous proof.

Theorem 10.4. *For every eigenvalue, it holds that*

$$d(\lambda) \leq k(\lambda)$$

Hence, the geometric multiplicity is smaller than the algebraic multiplicity.

Proof. Let $\lambda \in \text{spec}(A)$. Let $d = d(\lambda)$. (b_1, \dots, b_d) is basis of $\ker(\lambda I - A)$. We extend this vector to a basis of $\mathbb{K}^n : (b_1, \dots, b_d, \dots, b_n)$.

$$\begin{aligned}
 B &= \begin{pmatrix} b_1 & b_2 & \dots & b_n \\ \vdots & \vdots & \vdots & \vdots \end{pmatrix} \\
 AB &= \begin{bmatrix} Ab_1 & Ab_2 & \dots & Ab_d & Ab_{d+1} & \dots & Ab_n \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix} \\
 &= \begin{bmatrix} \lambda b_1 & \lambda b_2 & \dots & \lambda b_d & Ab_{d+1} & \dots & Ab_n \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix} \\
 &= \begin{bmatrix} b_1 & \dots & b_d & b & \dots & b_n \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \end{bmatrix} \begin{bmatrix} \lambda & & & & & \dots \\ & \lambda & & & & \dots \\ & & & \lambda & & \dots \\ 0 & 0 & 0 & 0 & \dots & \dots \\ & \dots & 0 & \dots & & \dots \\ 0 & 0 & 0 & 0 & \dots & \dots \end{bmatrix} \text{ where } \lambda \text{ occurs in } d \text{ different columns} \\
 B^{-1}AB &= \begin{bmatrix} \lambda & & & \dots \\ & \ddots & & \dots \\ & & \lambda & \dots \\ 0 & 0 & 0 & \tilde{A} \\ 0 & 0 & 0 & \tilde{A} \\ 0 & 0 & 0 & \tilde{A} \end{bmatrix} =: M
 \end{aligned}$$

$$\begin{aligned}
\chi_A(x) &= \chi_{B^{-1}AB}(x) = \det(x \cdot I - M) \\
&= \det \begin{bmatrix} x - \lambda & & & \dots \\ & \ddots & & \dots \\ & & x - \lambda & \dots \\ 0 & 0 & 0 & xI_{n-d} - \tilde{A} \end{bmatrix} \\
&= (x - \lambda)^d \det(xI - \tilde{A})
\end{aligned}$$

$\implies x - \lambda$ is d -multiple factor of $\chi_A(x) \implies k(\lambda) \geq d(\lambda)$. □

↓ This lecture took place on 2018/05/23.

Remark 10.13 (Revision). A is diagonalizable iff $\exists T$:

$$T^{-1}AT = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}$$

where T is basis of eigenvectors.

$$\iff d(\lambda) = \text{geometric multiplicity} = \dim \eta_\lambda$$

$$\stackrel{!}{=} k(\lambda) = \text{algebraic multiplicity}$$

Example 10.7.

$$A = \begin{bmatrix} \lambda & 1 & 0 \\ 0 & \lambda & 1 \\ 0 & 0 & \lambda \end{bmatrix}$$

Eigenvalues:

$$\chi_A(x) = \begin{vmatrix} x - \lambda & -1 & 0 \\ 0 & x - \lambda & -1 \\ 0 & 0 & x - \lambda \end{vmatrix} = (x - \lambda)^3$$

The only eigenvalue: λ . $k(\lambda) = 3$.

$$\ker \eta_\lambda = \ker \begin{bmatrix} 0 & -1 & 0 \\ & 0 & -1 \\ & & 0 \end{bmatrix}$$

$$d(\lambda) = \dim \ker \eta_\lambda = 1 \implies \text{not diagonalizable}$$

Remark 10.14. Camille Jordan (1838–1922): Jordan curve theorem.

11 Jordan Normal Form (JNF)

Remark 11.1. *Causs-Willhelm-Jordan (1842–1899)*

Nilpotent matrix:

$$\begin{bmatrix} 0 & * & * \\ 0 & 0 & * \\ 0 & 0 & 0 \end{bmatrix}^2 = \begin{bmatrix} 0 & 0 & * \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} 0 & 0 & * \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}^2 = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

11.1 Invariant subspaces

Definition 11.1. Let $f : V \rightarrow V$ (or matrix $A \in \mathbb{K}^{n \times n}$) be linear. A subspace $U \subseteq V$ is called *invariant under f* , if $f(U) \subseteq U$.

$$\iff Ax \in U \quad \forall x \in U$$

Example 11.1. 1. $\{0\}$. $f(0) = 0 \in \{0\}$. V is trivially invariant.

2. $\ker f$. Let $x \in \ker f \implies f(x) = 0 \implies f(f(x)) = 0 \implies f(x) \in \ker f$.
image f is invariant. $y \in \text{image } f \implies f(y) \in \text{image } f$.

3. Eigenspaces are invariant.

$$f(x) = \lambda \cdot x \implies f(f(x)) = f(\lambda \cdot x) = \lambda \cdot f(x) \implies f(x) \in \eta_\lambda$$

4. If U are invariant with $\dim U = 1$, then $U = \mathcal{L}(x)$ with x as eigenvector.
If $x \in U \setminus \{0\} (\implies U = \mathcal{L}(x))$, $f(x) \in U$. $\exists \lambda \cdot f(x) = \lambda \cdot x \implies x$ is eigenvector.

$$5. A = \begin{bmatrix} a_{11} & \dots & \dots & \dots \\ & a_{22} & \dots & \dots \\ & & \ddots & \dots \\ & & & a_{nn} \end{bmatrix}$$

$$A \cdot e_1 = a_{11} \cdot e_1 \implies e_1 \text{ is eigenvalue} \implies \mathcal{L}(e_1) \text{ is invariant}$$

$$A(\lambda_1 e_1 + \lambda_2 e_2) = \lambda_1 a_{11} e_1 + a_{12} \lambda_2 e_2 + a_{22} \lambda_2 e_2 \in \mathcal{L}(e_1, e_2) \implies \mathcal{L}(e_1, e_2) \text{ is invariant}$$

$$A \cdot e_k \in \mathcal{L}(e_1, \dots, e_k) \implies \forall k : \mathcal{L}(e_1, \dots, e_k) \text{ is invariant}$$

Numerically unstable:

$$\begin{bmatrix} \lambda & & \\ & \ddots & \\ & & \lambda \end{bmatrix} \text{ is diagonalizable}$$

$$\begin{bmatrix} \lambda & \varepsilon \\ 0 & \lambda \end{bmatrix} \text{ is not diagonalizable}$$

Theorem 11.1. Let $A \in \mathbb{K}^{n \times n}$, $V = \mathbb{K}^n$.

1. If $U \subseteq V$ is invariant and $p(x) \in \mathbb{K}[x]$, then U is invariant under $p(A)$.

$$p(x) := \sum_{k=0}^n a_k x^k \quad p(A) = \sum_{k=0}^n a_k A^k \quad \psi_A : \mathbb{K}[x] \rightarrow \mathbb{K}^{n \times n} \quad x \mapsto A$$

2. U_1, \dots, U_k are invariant subspaces.

$$\implies \begin{matrix} U_1 \cap \dots \cap U_k \\ U_1 + \dots + U_k \end{matrix} \text{ are invariant in regards of } A$$

Proof. 1. Let $x \in U$.

$$\implies Ax \in U \quad A^2 \cdot x = A \cdot \underbrace{(Ax)}_{\in U} \in U \quad \dots \quad A^k x \in U \implies \sum a_k A^k x \in U$$

because it is a linear combination of elements of U where U is the subspace with $A^k x \in U$.

2. Let $x \in \bigcap_{i=1}^k U_i$. $\implies \forall i : x \in U_i \implies \forall i : Ax \in U_i \implies Ax \in \bigcap_{i=1}^k U_i$. Let $x \in U_1 + \dots + U_k \implies x = u_1 + u_2 + \dots + u_k$ for $u_i \in U_i$.

$$\begin{aligned} \implies Ax &= \underbrace{(Au_1)}_{\in U_1} + \underbrace{(Au_2)}_{\in U_2} + \dots + \underbrace{(Au_k)}_{\in U_k} \in U_1 + \dots + U_k \\ \implies U_1 + \dots + U_k &\text{ is invariant} \end{aligned}$$

□

Lemma 11.1. Let $f : V \rightarrow V$ and $U \subseteq V$ is an invariant subspace. $\implies f|_U : U \rightarrow U$ is homomorphism. (If U is not invariant, $\varphi|_U : U \rightarrow V$ must not map $U \rightarrow U$.)

Theorem 11.2. Let $f : V \rightarrow V$. Let $U, W \subseteq V$ be invariant with $V = U + W$. Let $B = \{b_1, \dots, b_m\}$ be a basis of U , $B' = \{b'_1, \dots, b'_n\}$ is basis of W . $\implies B \cup B'$ is basis of V .

$$\Phi_{B \cup B'}^{B \cup B'}(f) = \left[\begin{array}{c|c} \Phi_B^B(f|_U) & 0 \\ \hline 0 & \Phi_{B'}^{B'}(f|_W) \end{array} \right]$$

Proof of Theorem 11.1. In the first m columns, we have the images of b_i (basis of U)

$$\begin{aligned} U \text{ invariant} &\implies f(b_i) \in U \\ \implies \text{coordinates in regards of } b'_1 \dots b'_n &\text{ are 0} \end{aligned}$$

$$\begin{bmatrix} f(b_1) & \dots & f(b_m) & f(b'_1) & \dots & f(b'_m) \\ & \ddots & & 0 & & 0 \\ & & & 0 & & 0 \\ 0 & \dots & 0 & \ddots & & \\ 0 & \dots & 0 & & \ddots & \\ 0 & \dots & 0 & & & \ddots \end{bmatrix}$$

In the last n columns, we can find the images of b'_j . W is invariant $\implies f(b'_j) \in W \implies$ coordinate in regards of b_1, \dots, b_m are 0. \square

Corollary 11.1. Let $f : V \rightarrow V$. $U_1, \dots, U_k \subseteq V$ is invariant with $V = U_1 + U_2 + \dots + U_k$. Let B_i be basis of $U_i \implies B = B_1 \cup \dots \cup B_k$ is basis of V and

$$\Phi_B^B(f) = \left[\begin{array}{c|ccc} \Phi_{B_1}^{B_1}(f|_{U_1}) & 0 & 0 & 0 \\ \hline 0 & \Phi_{B_2}^{B_2}(f|_{U_2}) & 0 & 0 \\ 0 & \vdots & \ddots & 0 \\ 0 & & & \Phi_{B_k}^{B_k}(f|_{U_k}) \end{array} \right]$$

Hence, if V can be decomposed into a direct sum of invariant subspaces, then A can be transformed into block diagonal form. (A is diagonalizable $\iff V$ can be decomposed into direct sum of one-dimensional subspaces)

Corollary 11.2. Corollary related to Corollary 11.1.

$$\chi_f(x) = \prod_{i=1}^k \chi_{f|_{U_i}}(x)$$

11.2 Fitting lemma

Remark 11.2. Hans Fitting (1906–1938)

Lemma 11.2 (Fitting lemma). Let $\dim V = n$, $f \in \text{End}(V)$.

1. $\{0\} \subseteq \ker f \subseteq \ker f^2 \subseteq \ker f^3 \subseteq \dots$
 $\text{image } f \supseteq \text{image } f^2 \supseteq \text{image } f^3 \supseteq \dots$
2. $\exists m \leq n : \ker f^m = \ker f^{m+1}$
3. The following statements are equivalent:
 - (a) $\ker f^m = \ker f^{m+1}$
 - (b) $\text{image } f^m = \text{image } f^{m+1}$
 - (c) $\ker f^m = \ker f^{m+k} \forall k \geq 1$

$$(d) \text{ image } f^m = \text{image } f^{m+k} \forall k \geq 1$$

$$(e) \ker f^m \cap \text{image } f^m = \{0\}$$

$$(f) V = \ker f^m + \text{image } f^m$$

Proof. 1. Let $x \in \ker f$. Then $f^2(x) = f(f(x)) = f(0) = 0$.
Let $y \in \text{image } f^2$. Then $\exists x : y = f(f(x))$. Thus $f(x) \in \text{image } f$.

$$2. \text{ If } \{0\} \subsetneq \ker f \subsetneq \ker f^2 \subsetneq \dots \subsetneq \ker(f^m)$$

$$\implies 0 < \dim \ker f < \dim \ker f^2 < \dots < \dim \ker f^m$$

$$\implies m \leq n$$

3. We prove a set of equivalences.

- We prove (a) \iff (b). Because of (1.), we know

$$\ker(f^m) \subseteq \ker(f^{m+1})$$

$$\implies \ker(f^m) = \ker(f^{m+1}) \iff \dim \ker f^m = \dim \ker f^{m+1}$$

$$\iff n - \dim \text{image}(f^m) = n - \dim \text{image}(f^{m+1})$$

$$\iff \dim \text{image}(f^m) = \dim \text{image}(f^{m+1})$$

Because of (1.), $\text{image } f^m \supseteq f^{m+1}$

$$\iff \text{image}(f^m) = \text{image}(f^{m+1})$$

- The proof of (c) \iff (d) follows analogously. The proofs (a) \iff (c) and (d) \iff (b) are trivial.
- We prove (a) \iff (c):

$$0 \subseteq \ker f \subseteq \ker f^2 \subseteq \ker f^3 \subseteq \dots$$

$$m_0 = \min \{ m \mid \ker(f^m) = \ker(f^{m+1}) \}$$

Claim:

$$\ker f^{m_0+k} = \ker f^{m_0+k+1} \forall k \geq 0$$

Direction \subseteq is immediate. Direction \supseteq : Let $x \in \ker f^{m_0+k+1} \implies f^{m_0+k+1}(x) = f^{m_0+1}(f^k(x)) = 0$.

$$\implies f^k(x) \in \ker f^{m_0+1} = \ker f^{m_0} \implies f^{m_0+k}(x) = 0 \implies x \in \ker f^{m_0+k}$$

with $f^k(x) \in \ker f^{m_0+1} = \ker f^{m_0}$ following from the definition of m_0 .

- We prove (b) \iff (d). Let $m_0 = \min \{m \mid \text{image } f^m = \text{image } f^{m+1}\}$.
 Claim: $\text{image } f^{m_0+k} = \text{image } f^{m_0+k+1} \forall k \geq 0$. Direction \supseteq is trivial.
 Direction \subseteq : Let $y \in \text{image } f^{m_0+k}$.

$$\implies \exists x : y = f^{m_0+k}(x) = f^k(\underbrace{f^{m_0}(x)}_{\in \text{image } f^{m_0} = \text{image } f^{m_0+1}})$$

hence $\exists z : f^{m_0}(x) = f^{m_0+1}(z)$.

$$\implies y = f^k(f^{m_0+1}(z)) = f^{m_0+k+1}(z) \in \text{image } f^{m_0+k+1}$$

- We prove $\{(a), (b), (c), (d)\} \iff$ (e). Let $w = \text{image } f^m$ is invariant under f^m .

$$g := f^m|_w \in \text{Hom}(W, W)$$

$$\ker g = \ker(f^m) \cap W = \ker f^m \cap \text{image } f^m$$

$$\ker f^m \cap \text{image } f^m = \{0\} \iff \ker g = \{0\}$$

$$\iff g \text{ linear} \wedge g \text{ injective} \wedge g \text{ surjective} \iff \text{image } g = W$$

$$\iff f^m(f^m(V)) = f^m(V) \iff f^{2m}(w) = f^m(w)$$

$$f^{m+m}(v) = f^m(v)$$

$$\text{image } f^{m+m} = \text{image } f^m$$

This is equivalent to statements 1 and 3, thus rendering our assumption $\ker f^m \cap \text{image } f^m = \{0\}$ valid.

- We prove (d) \iff (b).

$$\text{image } f^{m+m} = \text{image } f^m \iff \text{image } f^{m+1} = \text{image } f^m$$

- (f) \implies (e) is trivial.
- We prove (e) \iff (f).

$$\left. \begin{array}{l} \dim \text{image } f^m + \dim \ker f^m = n \\ \text{image } f^m \cap \ker f^m = \{0\} \end{array} \right\} \implies \text{image } f^m + \ker f^m = V$$

because of dimensionality reasons.

□

Remark 11.3. The intersection of image and kernel of a linear function is only $\{0\}$.

↓ This lecture took place on 2018/05/28.

Fitting Lemma:

$$\begin{aligned}\ker A &\subseteq \ker A^2 \subseteq \cdots \subseteq \ker A^r \\ \text{image } A &\supseteq \text{image } A^2 \supseteq \cdots \supseteq \text{image } A^r = \text{image } A^{r+1} \\ \ker A^r \oplus \text{image } A^r &= V \\ \ker A^r \cap \text{image } A^r &= \{0\}\end{aligned}$$

Example 11.2.

$$\begin{aligned}A &= \begin{bmatrix} 0 & 1 & 1 \\ & 0 & 1 \\ & & 0 \end{bmatrix} & \ker A &= \mathcal{L}\left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}\right) & \text{image } A &= \mathcal{L}\left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}\right) \\ A^2 &= \begin{bmatrix} 0 & 0 & 1 \\ & 0 & 0 \\ & & 0 \end{bmatrix} & \ker A^2 &= \mathcal{L}\left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}\right) & \text{image } A^2 &= \mathcal{L}\left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}\right) \\ A^3 &= \begin{bmatrix} 0 & 0 & 0 \\ & 0 & 0 \\ & & 0 \end{bmatrix} & \ker A^3 &= \mathbb{K}^3 & \text{image } A^3 &= \{0\}\end{aligned}$$

$$x \in \ker A^k \implies A \cdot x \in \ker A^{k-1}$$

Example 11.3.

$$\begin{aligned}A &= \begin{bmatrix} \lambda & 1 & \\ & \lambda & 1 \\ & & \lambda \end{bmatrix} & \text{eigenvalue: } \lambda & \lambda I - A &= \begin{bmatrix} 0 & -1 & \\ & 0 & -1 \\ & & 0 \end{bmatrix} \\ \ker(\lambda I - A) &= \mathcal{L}\left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}\right) & \ker((\lambda I - A)^2) &= \mathcal{L}\left(\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}\right) \\ x \in \ker((\lambda I - A)^2) &\implies (\lambda I - A)x \in \ker(\lambda I - A) \\ &\implies Ax = \lambda x + b & b &\in \ker(\lambda I - A)\end{aligned}$$

11.3 Main space

Definition 11.2. Let $A \in \mathbb{K}^{n \times n}$, $\lambda \in \text{spec}(A)$. Then $\ker(\lambda I - A)^n$ is called main space (dt. Hauptraum) of A for eigenvalue λ . The elements are called generalized eigenvectors.

Actually, $\ker(\lambda I - A)^n = \ker(\lambda I - A)^r$ and a generalized eigenvector satisfies

$$A \cdot x = \lambda \cdot x + y \quad y \in \ker(\lambda I - A)^{r-1}$$

with r as first index such that $\ker(\lambda I - A)^r = \ker(\lambda I - A)^{r+1}$.

Fitting:

$$\mathbb{K}^n = \ker(\lambda I - A)^r \oplus \text{image}(\lambda I - A)^r$$

Next step: decomposition for different eigenvalues.

Lemma 11.3. Let $\lambda_1, \dots, \lambda_k$ be different eigenvalues of A and $\ker(\lambda_i I - A)^{r_i}$ the corresponding main spaces where

$$\begin{aligned} \ker(\lambda_i I - A)^{r_{i-1}} &\subsetneq \ker(\lambda_i I - A)^{r_i} = \ker(\lambda_i \cdot I - A)^{r_i+1} \\ \Rightarrow \bigcap_{i=1}^k \text{image}(\lambda_i I - A)^{r_i} \cap \ker((\lambda_1 I - A)^{r_1} (\lambda_2 I - A)^{r_2} \dots (\lambda_k I - A)^{r_k}) &= \{0\} \end{aligned}$$

Remark 11.4.

$$\begin{aligned} \ker(\lambda_1 I - A)^{r_1} \dots (\lambda_k I - A)^{r_k} &\supseteq \ker(\lambda_i I - A)^{r_i} \forall i \\ &\supseteq \ker(\lambda_1 I - A)^{r_1} + \dots + \ker(\lambda_k I - A)^{r_k} \end{aligned}$$

if $(\lambda_i I - A)^{r_i} \cdot x = 0$,

$$\begin{aligned} &(\lambda_1 I - A)^{r_1} \dots (\lambda_k I - A)^{r_k} \cdot x \\ &= (\lambda_1 I - A)^{r_1} \dots (\lambda_{i-1} I - A)^{r_{i-1}} (\lambda_{i+1} I - A)^{r_{i+1}} \dots (\lambda_k I - A)^{r_k} \cdot \underbrace{(\lambda_i \cdot I - A)^{r_i} x}_{=0} = 0 \end{aligned}$$

If one of the factors is zero, the product is zero.

$$p(A) \cdot q(A) = q(A) \cdot p(A) \text{ for arbitrary polynomials } p(x) \text{ and } q(x)$$

especially: $p_i(x) = (\lambda_i - x)^{r_i}$.

Example 11.4.

$$A = \begin{bmatrix} 1 & 1 & & & & & \\ & 1 & & & & & \\ & & 1 & & & & \\ & & & 2 & 1 & & \\ & & & 2 & 1 & & \\ & & & & 2 & & \\ & & & & & 0 & -1 \\ & & & & & 1 & 0 \end{bmatrix} \text{ over } \mathbb{R}$$

$$\text{spec}(A) = \{1, 2\} \cup \underbrace{(\{\pm i\})}_{\notin \mathbb{R}}$$

Consider $\lambda = 1$.

$$\ker(I - A) = \ker \begin{bmatrix} 0 & -1 & & & & & & \\ & 0 & & & & & & \\ & & 0 & & & & & \\ & & & -1 & -1 & & & \\ & & & & -1 & -1 & & \\ & & & & & -1 & & \\ & & & & & & 1 & 1 \\ & & & & & & -1 & 1 \end{bmatrix}$$

$$\ker(I - A) = \mathcal{L}(e_1, e_2) \quad \text{the two zero columns}$$

$$\ker(I - A)^2 = \ker \begin{bmatrix} 0 & 0 & & & & & & \\ 0 & 0 & & & & & & \\ & & 0 & & & & & \\ & & & -1 & -1 & & & \\ & & & & -1 & -1 & & \\ & & & & & -1 & & \\ & & & & & & 1 & 1 \\ & & & & & & -1 & 1 \end{bmatrix}$$

$$\ker(I - A)^2 = \mathcal{L}(e_1, e_2, e_3)$$

$$\text{image}(I - A)^2 = \mathcal{L}(e_4, e_5, e_6, e_7, e_8)$$

Consider $\lambda = 2$.

$$\ker(2I - A) = \ker \begin{bmatrix} 1 & -1 & & & & & & \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 0 & -1 & & & \\ & & & & 0 & -1 & & \\ & & & & & 0 & & \\ & & & & & & 2 & 1 \\ & & & & & & -1 & 2 \end{bmatrix}$$

$$\ker(2I - A) = \mathcal{L}(e_4)$$

$$\ker(2I - A)^2 = \ker \begin{bmatrix} 1 & 1 & & & & & & \\ & 1 & & & & & & \\ & & 1 & & & & & \\ & & & 0 & 0 & 1 & & \\ & & & & 0 & 0 & & \\ & & & & & 0 & & \\ & & & & & & & \text{[regular]} \end{bmatrix}$$

$$\begin{aligned}
\ker(2I - A)^2 &= \mathcal{L}(e_4, e_5) \\
\ker(2I - A)^3 &= \mathcal{L}(e_4, e_5, e_6) \\
\ker(2I - A)^3 &= \mathcal{L}(e_1, e_2, e_3, e_7, e_8) \\
\bigcap_{i=1}^2 \text{image}(\lambda_i I - A)^{r_i} &= \mathcal{L}(e_1, e_2, e_3, e_7, e_8) \cap \mathcal{L}(e_4, e_5, e_6, e_7, e_8) = \mathcal{L}(e_7, e_8)
\end{aligned}$$

$$\begin{aligned}
(I - A)^2 \cdot (2I - A)^3 &= \begin{bmatrix} 0 & 0 & & & & & & \\ 0 & 0 & & & & & & \\ & & 0 & & & & & \\ & & & [regular\ 3 \times 3] & & & & \\ & & & & [regular\ 2 \times 2] & & & \end{bmatrix} \\
&\cdot \begin{bmatrix} [regular\ 2 \times 2] & & & & & & & \\ & 1 & & & & & & \\ & & 0 & 0 & 0 & & & \\ & & 0 & 0 & 0 & & & \\ & & 0 & 0 & 0 & & & \\ & & & & & [regular\ 2 \times 2] & & \end{bmatrix} \\
&= \begin{bmatrix} 0 & 0 & & & & & & \\ 0 & 0 & & & & & & \\ & & 0 & & & & & \\ & & & 0 & 0 & 0 & & \\ & & & 0 & 0 & 0 & & \\ & & & 0 & 0 & 0 & & \\ & & & & & & [regular\ 2 \times 2] \end{bmatrix}
\end{aligned}$$

$$\ker(I - A)^2 \cdot (2I - A)^3 = \mathcal{L}(e_1, \dots, e_6) = \left(\bigcap (\lambda_i I - A)^{r_i} \right)^c$$

Proof of Lemma 11.3. Show: If $x \in \bigcap_{i=1}^k \text{image}(\lambda_i I - A)^{r_i}$ and $(\lambda_1 I - A)^{r_1} \dots (\lambda_k I - A)^{r_k} \cdot x = 0$, then $x = 0$.

Proof by induction over k :

Case $k = 1$

$$x \in \text{image}(\lambda_1 I - A)^{r_1} \wedge (\lambda_1 I - A)^{r_1} x = 0$$

$$\xrightarrow{\text{Fitting}} x = 0$$

Case $k \rightarrow k + 1$ Let $x \in \bigcap_{i=1}^k \text{image}(\lambda_i I - A)^{r_i}$ and $(\lambda_1 I - A)^{r_1} \dots (\lambda_{k+1} I - A)^{r_{k+1}} x = 0$.

$$\text{Let } y = (\lambda_{k+1} I - A)^{r_{k+1}} x \implies y \in \ker(\lambda_1 I - A)^{r_1} \dots (\lambda_k I - A)^{r_k}.$$

$$\forall i \in \{1, \dots, k+1\} \exists u_i : x = (\lambda_i I - A)^{r_i} \cdot u_i$$

$$\begin{aligned}
y &= (\lambda_{k+1} I - A)^{r_{k+1}} x = (\lambda_{k+1} I - A)^{r_{k+1}} (\lambda_i I - A)^{r_i} \cdot u_i \\
&= (\lambda_i I - A)^{r_i} (\lambda_{k+1} I - A)^{r_{k+1}} u_i \\
&\in \text{image}(\lambda_i I - A)^{r_i}
\end{aligned}$$

$$\begin{aligned}
p(A)q(A) &= q(A)p(A) \\
p(x) &= (\lambda_{k+1} - x)^{r_{k+1}} \quad q(x) = (\lambda_i - x)^{r_i} \\
\implies y &\in \bigcap_{i=1}^k \text{image}(\lambda_i I - A)^{r_i}
\end{aligned}$$

By the induction hypothesis, $y = 0$.

$$\begin{aligned}
\implies x &\in \ker(\lambda_{k+1} I - A)^{r_{k+1}} \wedge x \in \text{image}(\lambda_{k+1} I - A)^{r_{k+1}} \\
&\xRightarrow{\text{Fitting}} x = 0
\end{aligned}$$

□

- Lemma 11.4.** 1. $\forall \lambda \neq \mu \forall k, l \geq 1 : \ker(\lambda I - A)^k \cap \ker(\mu I - A)^l = \{0\}$
2. The sum $\ker(\lambda_1 I - A)^{r_1} + \dots + \ker(\lambda_k I - A)^{r_k}$ is direct for arbitrary pairwise different $\lambda_1, \dots, \lambda_k$.

Proof. Proof of the first statement. Induction over $m = k + l$.

Induction base Consider $m = 2, k = l = 1$.

$$\ker(\lambda I - A) \cap \ker(\mu I - A) = \{0\}$$

The eigenvectors for different eigenvalues are linear independent.

Induction step $m - 1 \rightarrow m$: Consider $m \geq 3$. Without loss of generality: $k \geq 2$. Let $x \in \ker(\lambda I - A)^k \cap \ker(\mu I - A)^l$. Let $y = (\lambda I - A)x \in \ker(\lambda I - A)^{k-1} \cap \ker(\mu I - A)^l$. Then,

$$(\mu I - A)^l \cdot y = (\mu I - A)^l (\lambda I - A) \cdot x = (\lambda I - A) \underbrace{(\mu I - A)^l}_{=0} \cdot x = 0$$

Let $k - 1 + l = m - 1$. By induction hypothesis, $y = 0$.

$$\implies x \in \ker(\lambda I - A)$$

$$\begin{aligned}
\implies x &\in \ker(\lambda I - A) \cap \ker(\mu I - A)^l \xRightarrow{\text{induction hypothesis}} x = 0 \\
1 + l &\leq m - 1
\end{aligned}$$

Proof of the second statement. Induction over k .

Induction base $k = 1$: trivial

Induction step $k \rightarrow k+1$

Show: if $v_i \in \ker(\lambda_i I - A)^{r_i}$ $i = 1, \dots, k+1$ and $v_1 + \dots + v_{k+1} = 0 \implies$
all $v_i = 0$.

Let $w_i = (\lambda_{k+1} I - A)^{r_{k+1}} v_i \implies w_{k+1} = 0$.

$$\sum_{i=1}^k w_i = \sum_{i=1}^{k+1} w_i = (\lambda_{k+1} I - A)^{r_{k+1}} \underbrace{\sum_{i=1}^{k+1} v_i}_{=0} = 0$$

$$(\lambda_i - A)^{r_i} w_i = (\lambda_i I - A)^{r_i} (\lambda_{k+1} I - A)^{r_{k+1}} v_i = (\lambda_{k+1} I - A)^{r_{k+1}} \underbrace{(\lambda_i I - A)^{r_i} \cdot v_i}_{=0} = 0$$

$$p(x) = (\lambda_i - x)^{r_i} \quad q(x) = (\lambda_{k+1} - x)^{r_{k+1}}$$

$$\implies w_i \in \ker(\lambda_i I - A)^{r_i}$$

$$\xRightarrow{\text{induction hypothesis}} w_i = 0 \forall i$$

$$\implies v_i \in \ker(\lambda_{k+1} I - A)^{r_{k+1}}$$

$$v_i \in \ker(\lambda_i - A)^{r_i}$$

$$\implies v_i \in \ker(\lambda_{k+1} I - A)^{r_{k+1}} \cap \ker(\lambda_i I - A)^{r_i} = \{0\}$$

$$\implies v_i = 0 \quad \forall i = 1, \dots, k$$

$$\implies 0 + \dots + 0 + v_{k+1} = 0 \implies v_{k+1} = 0$$

□

Theorem 11.3. Let $\lambda_1, \dots, \lambda_k$ be pairwise different eigenvalues of $A \in \mathbb{K}^{n \times n}$.

1.

$$V = \ker(\lambda_1 I - A)^n \oplus \dots \oplus \ker(\lambda_k I - A)^n \oplus \underbrace{\bigcap_{i=1}^k \text{image}(\lambda_i I - A)^n}_{=: W}$$

Compare with example $(I - A)^2$:

$$\begin{bmatrix} 0 & & & & \\ & 0 & & & \\ & & 0 & & \\ & & & \dots & \\ & & & & \dots \end{bmatrix}$$

$(2I - A)^2$:

$$\begin{bmatrix} [\dots] & & & \\ & 1 & & \\ & & 0 & \\ & & & \underbrace{[\dots]}_W \end{bmatrix}$$

2. W is invariant under A and $\lambda_i \notin \text{spec}(A|_W) \forall i = 1, \dots, k$

Proof. 1. Induction over k

Induction base $k = 1$:

$$V = \ker(\lambda_i I - A)^n \oplus \text{image}(\lambda_i I - A)^n \quad (\text{Fitting})$$

Induction step $k \rightarrow k + 1$: We assume:

$$V = \ker(\lambda_1 I - A)^n \oplus \dots \oplus \ker(\lambda_k I - A)^n \oplus W_k$$

$$W_k = \bigcap_{i=1}^k \text{image}(\lambda_i I - A)^n$$

W_k is invariant: $y \in W_k \xRightarrow{!} Ay \in W_k$. Let $y \in W_k \implies \forall i = 1, \dots, k : \exists x_i : y = (\lambda_i I - A)^n x_i$.

$$\implies Ay = A \cdot (\lambda_i I - A)^n x_i = (\lambda_i I - A)^n \cdot Ax_i \in \text{image}(\lambda_i I - A)^n$$

For all $i = 1, \dots, k$ it holds that

$$\implies Ay \in \bigcap_{i=1}^k \text{image}(\lambda_i I - A)^n$$

$$p(x) = x \quad q(x) = (\lambda_i - x)^n$$

Consider $g : W_k \rightarrow W_k$ with $x \mapsto Ax$ with $\dim(W_k) \leq n$.

$$\text{Fitting} \implies \ker(\lambda_{k+1} I - g)^n \oplus \text{image}(\lambda_{k+1} I - g)^n = W_k$$

where $\text{image}(\lambda_{k+1} I - g)^n \subseteq \text{image}(\lambda_{k+1} I - A)^n$.

$$\subseteq \ker(\lambda_{k+1} I - A)^n + (\text{image}(\lambda_{k+1} I - A)^n) \cap W_k$$

$$= \ker(\lambda_{k+1} I - A)^n + \bigcap_{i=1}^{k+1} \text{image}(\lambda_i I - A)^n$$

$$\implies V = \ker(\lambda_1 I - A)^n + \dots + \ker(\lambda_k I - A)^n + \ker(\lambda_{k+1} I - A)^n + W_{k+1}$$

Claim: This sum is direct.

Let $x_i \in \ker(\lambda_i I - A)^n$ and $i = 1, \dots, k+1$. Let $y \in W_{k+1} = \bigcap_{i=1}^{k+1} \text{image}(\lambda_i I - A)^{r_i}$. Show that all $x_i = 0$ and $y = 0$. Thus $\sum_{i=1}^{k+1} x_i + y = 0$.

$$\begin{aligned} 0 &= \prod_{i=1}^{k+1} (\lambda_i I - A)^n \left(\sum_{i=1}^{k+1} x_i + y \right) = \sum_{i=1}^{k+1} 0 + \prod_{i=1}^{k+1} (\lambda_i I - A)^n \cdot y \\ \implies j &\in \ker \prod_{i=1}^{k+1} (\lambda_i I - A)^n \cap \bigcap_{i=1}^{k+1} \text{image}(\lambda_i I - A)^n \xrightarrow{\|\cdot\|} y = 0 \\ &\implies \sum_{i=1}^{k+1} x_i = 0 \xrightarrow{\text{Lemma 11.4}} \text{TODO} \end{aligned}$$

2. W_k is invariant, see proof of part (1)

$$\ker(\lambda_i - A) \cap W_k \subseteq \ker(\lambda_i - A)^n \cap \{0\}$$

\implies no eigenvector for λ_1 in W_k .

□

↓ This lecture took place on 2018/05/30.

The sum of main spaces $\ker(\lambda_1 I - A)^n + \dots + \ker(\lambda_k I - A)^n + W$ is direct. The main spaces are invariant and also $W = \bigcap_{i=1}^k \text{image}(\lambda_i I - A)^n$ and the restriction $A|_W$ has no λ_i as eigenvalue.

Let B_0 be a basis of W , B_i is a basis of $\ker(\lambda_i I - A)^n$. Then $B := B_1 \cup B_2 \cup \dots \cup B_k \cup B_0$ is a basis and in this basis,

$$\Phi_B^B(A) = \begin{bmatrix} [B_1] & & & & \\ & [B_2] & & & \\ & & \ddots & & \\ & & & [B_k] & \\ & & & & [B_0] \end{bmatrix}$$

If $x \in \mathcal{L}(B_i) \implies Ax \in \mathcal{L}(B_i)$. By invariance, $\Phi_B^B(A)$ has block diagonal form.

$$\ker(\lambda_i I - A)^n = \mathcal{L}(B_i)$$

$$\implies (\lambda_i I - A)^n|_{\mathcal{L}(B_i)} = 0 \implies \text{nilpotent}$$

Theorem 11.4. Let \mathbb{K} be algebraically closed (hence, every other matrix has eigenvalue) and let $\lambda_1, \dots, \lambda_k$ all eigenvalues of a matrix $A \in \mathbb{K}^{n \times n}$.

$$\implies \mathbb{K}^n = \ker(\lambda_1 I - A)^n \oplus \dots \oplus \ker(\lambda_k I - A)^n$$

Proof.

$$\mathbb{K}^n = \ker(\lambda_1 I - A)^n \oplus \cdots \oplus \ker(\lambda_n I - A)^n \oplus W$$

$$W = \bigcap_i \text{image}(\lambda_i I - A)^n$$

$A|_W$ has no eigenvalue (because eigenvalue of $A|_W$ are also eigenvalues of A , but none of λ_i is an eigenvalue of $A|_W$), otherwise the sum is not direct. $\implies W$ is trivial ($W = \{0\}$). \square

11.4 Nilpotent matrix

Theorem 11.5. *A matrix/linear map $f : V \rightarrow V$ is called nilpotent, if $\exists k \in \mathbb{N} : f^k = 0$. The smallest k is called index of f .*

$(\lambda_i I - A)|_{i\text{-th main space}}$ is nilpotent

Goal: Structure of nilpotent matrices:

$$\begin{bmatrix} 0 & * & \ddots & 0 \\ & \ddots & \ddots & \\ & & \ddots & * \\ 0 & & & 0 \end{bmatrix}$$

Lemma 11.5. *Let $\ker(f^m) \subseteq \ker(f^{m+1}) \subseteq \ker(f^{m+2})$*

$u_1 \dots u_p \dots$ basis of $\ker f^n$

$u_1 \dots u_p v_1 \dots v_k \dots$ basis of $\ker f^{m+1}$

$u_1 \dots u_p v_1 \dots v_k w_1 \dots w_r \dots$ basis of $\ker f^{m+2}$

Then $(u_1 \dots u_p, f(w_1), \dots, f(w_r))$ is linear independent.

Proof. Immediate: $f(w_i) \in \ker f^{m+1}$, thus $f(\ker f^{m+2}) \subseteq \ker f^{m+1}$.

Show that: $\sum_{i=1}^p \lambda_i u_i + \sum_{j=1}^r \mu_j f(w_j) = 0 \stackrel{!}{\implies} \text{all } \lambda_i = 0, \mu_j = 0$.

$$\begin{aligned} \implies & \underbrace{f^m(\dots)}_{= \sum_{j=1}^p \lambda_i f^m(u_i) + \sum_{j=1}^r \mu_j f^{m+1}(w_j) = 0} = 0 \\ \implies & \sum_{i=1}^r \mu_j w_j \in \ker f^{m+1} \end{aligned}$$

but $\ker f^{m+2} = \ker f^{m+1} \oplus \underbrace{\mathcal{L}(w_1, \dots, w_r)}_{w_j \in \mathcal{L}(w_1, \dots, w_r)}$. Hence, $\ker f^{m+1} \cap \mathcal{L}(w_1, \dots, w_r) = \{0\}$.

$$\begin{aligned} \Rightarrow \sum_{i=1}^r \mu_j w_j = 0 &\xrightarrow{w_j \text{ linear indep.}} \text{all } \mu_j = 0 \\ \Rightarrow \sum_{i=1}^p \mu_i u_i = 0 &\xrightarrow{u_i \text{ linear indep.}} \text{all } \lambda_i = 0 \end{aligned}$$

□

11.5 Jordan's normal form

Theorem 11.6. Jordan's normal form is a nilpotent matrix. Let $\dim V = n$. $f : V \rightarrow V$ is nilpotent of index p ($f^p = 0$). $d = \dim \ker f$. Then there exists a basis B of V such that

$$\Phi_B^B(f) = \begin{bmatrix} [N_1] & & & \\ & [N_2] & & \\ & & \ddots & \\ & & & [N_d] \end{bmatrix}$$

where

$$N_i = \begin{bmatrix} 0 & 1 & \ddots & 0 \\ & 0 & 1 & \\ & & \ddots & 1 \\ 0 & & & 0 \end{bmatrix}_{n_i \times n_i}$$

$$p = n_1 \geq n_2 \geq \dots \geq n_d \geq 1$$

$$n_1 + \dots + n_d = n$$

Proof. Let $U_k = \ker f^k$, $\dim U_k = m_k$. $U_1 \subseteq U_2 \subseteq \dots \subseteq U_p = V$. $d = m_1 \leq m_2 \leq m_3 \leq \dots \leq m_p = n$.

$$f(U_i) \subseteq U_{i-1}$$

$$\underbrace{\underbrace{[[u_1 \dots u_{m_1}] u_{m_1+1} \dots u_{m_2}] \dots u_{m_{p-1}+1} \dots u_{m_p}}_{U_1}}_{U_2} \dots \underbrace{\phantom{u_1 \dots u_{m_p}}}_{U_p}$$

$u_1 \dots u_{m_k}$ is basis of U_k .

We start from behind:

$$\ker f^{p-2} \leq \ker f^{p-1} \leq \ker f^p$$

We apply Lemma 11.5.

$$u_1 \dots u_{m_{p-2}} | u_{m_{p-2}+1} \dots u_{m_{p-1}} | u_{m_{p-1}+1} \dots u_{m_p}$$

$$v_1^{(p)} := u_{m_{p-1}+1} \quad v_2^{(p)} = u_{m_{p-1}+2} \dots v_{m_p-m_{p-1}}^{(p)} := u_{m_p}$$

is basis of $U_p \ominus U_{p-1}$.

$$v_1^{(p+1)} = f(v_1^{(p)}) \quad v_2^{(p+1)} = f(v_2^{(p)}) \dots v_{m_p-m_{p-1}}^{(p+1)} \in U_{p-1} \underbrace{\ominus}_{(*)} U_{p-2}$$

(*) by Lemma 11.5 $f(v_j^{(p)})$ linear independent of $u_1 \dots u_{m_{p-2}}$.

And these $v_i^{(p-1)}$ are linear independent of $u_1 \dots u_{m_{p-2}}$. Extend $u_1 \dots u_{m_{p-2}} v_1^{(p-1)} \dots v_{m_p-m_{p-1}}^{(p-1)}$ to basis of U_{p-1} : $v_{m_p-m_{p-1}+1}^{(p+1)} \dots v_{m_{p-1}-m_{p-2}}^{(p+1)}$ chosen from $u_{m_{p-2}+1} \dots u_{m_{p-1}}$.

$$m_{p-2} + \dots + m_{p-1} - m_{p-2} = m_{p-1}$$

$$u_1 \dots u_{m_{p-2}} | u_{m_{p-2}+1} \dots u_{m_{p-1}} | u_{m_{p-1}+1} \dots u_{m_p}$$

$$\underbrace{u_1 \dots u_{m_{p-2}}}_{U_{p-2}} \underbrace{v_1^{(p-1)} \dots v_{m_{p-1}-m_{p-2}}^{(p-1)} u_{m_{p-1}+1} \dots u_{m_p}}_{f(n_{m_{p-1}+1}) \dots f(u_{m_p}) U_{p-1}}$$

$$\underbrace{\hspace{10em}}_{U_p}$$

where $u_{m_{p-1}+1} = v_1^{(p)} \dots u_{m_p} = v_{m_p-m_{p-1}}^{(p)}$.

Iteration:

$$v_1^{(p-2)} = f(v_1^{(p-1)}) \in U_{p-2} \ominus U_{p-3}$$

$$v_2^{(p-2)} = f(v_2^{(p-1)})$$

$$\vdots$$

$$v_{m_{p-1}-m_{p-2}}^{(p-2)} = f(v_{m_{p-1}-m_{p-2}}^{(p-1)})$$

$$\rightsquigarrow u_1 \dots u_{m_{p-3}} v_1^{(p-2)} \dots v_{m_{p-1}-m_{p-2}}^{(p-2)} \subseteq U_{p-2}$$

are linear independent. \rightarrow extend to basis of U_{p-2} :

$$u_1 \dots u_{m_{p-3}} v_1^{(p-2)} \dots v_{m_{p-2}-m_{p-3}}^{(p-2)}$$

and so on and so forth.

In the end, we get a basis:

$$\begin{array}{ccccccc}
 v_1^{(p)} & v_2^{(p)} & \dots & v_{m_p-m_{p-1}}^{(p)} & & & \\
 v_1^{(p-1)} & v_2^{(p-1)} & \dots & v_{m_p-m_{p-1}}^{(p-1)} & \dots & v_{m_{p-1}-m_{p-2}}^{(p-1)} & \\
 v_1^{(p-2)} & v_2^{(p-2)} & \dots & v_{m_p-m_{p-1}}^{(p-2)} & \dots & v_{m_{p-1}-m_{p-2}}^{(p-2)} & \dots v_{m_{p-2}-m_{p-3}}^{(p-2)} \\
 \vdots & & & & & & \\
 v_1^{(1)} & v_2^{(1)} & & & & & v_{m_1}^{(1)}
 \end{array}$$

where each successive row can be reached by applying f . The last row represents the basis of U_1 , all rows give the basis of U_p .

1. The last row is basis of U_1
2. f maps k -th row to $k-1$ -th column.

$$B = \begin{bmatrix} \vdots & & \\ \vdots & \vdots & \\ \vdots & \vdots & \vdots \end{bmatrix}$$

$$B = v_1^{(1)} v_n^{(2)} \dots v_1^{(p)} v_2^{(2)} v_2^{(2)} \dots v_2^{(p)} \dots v_k^{(1)} v_k^{(2)} \dots v_k^{(p-1)} \dots v_{M_i}$$

$$B = v_1^{(i)} \dots v_1^{(p)} v_2^{(i)} \dots v_2^{(n_2)} v_3^{(1)} \dots v_3^{(n_3)} \dots v_d^{(1)} \dots v_d^{(n_d)}$$

$$n_3 \leq n_2 \leq n_1$$

$$f(v_i^{(i)}) = 0 \forall i = 1, \dots, d$$

$$f(v_i^{(2)}) = v_i^{(1)} \quad f(v_i^{(3)}) = v_1^{(2)}$$

$$\Phi_B^B(f) = \begin{bmatrix} 0 & 1 & & & \vdots \\ & 0 & 1 & & \vdots \\ & & \ddots & \ddots & \vdots \\ & & & 0 & \vdots \\ \vdots & & & 0 & 1 \\ \vdots & & & & 0 & 1 \\ \vdots & & & & & \ddots \\ \vdots & & & & & & 1 \\ \vdots & & & & & & 0 \\ 0 & 0 & \dots & 0 & & & \end{bmatrix}$$

where this matrix goes on with these block matrices in the diagonal from n_1 to n_d . \square

Example 11.5.

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 3 & 0 & -2 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 & 1 & 3 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -4 & 0 & 0 & 0 & -4 & 0 & 0 \end{bmatrix} \rightsquigarrow \begin{bmatrix} 0 & 0 & 3 & 0 & -2 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 3 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$x_2 = -x_6 \quad x_5 = 0 \quad x_7 = 2x_3 = 0 \quad x_3 = 0$$

Bases of $\ker(A) = e_1, e_4, e_8, -e_2 + e_6 =: \{u_1, u_2, u_3, u_4\}$.

$$\ker A = \ker N_1, N_1 = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & -2 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \text{ are pivot rows}$$

$$\ker A^2 = \ker N_1 \cdot A$$

$$\text{because : } A^2x = 0 \iff Ax \in \ker A \iff Ax \in \ker N_1 \iff N_1 \cdot Ax = 0$$

$$\ker(N_1 \cdot AU) = \ker \begin{bmatrix} 0 & 0 & 1 & 0 & -2 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

$$\ker A^2 : x_3 = 2x_5$$

$$\text{Basis of } \ker A^2 : e_1, e_2, e_4, e_6, e_7, e_8, 2e_3 + e_5$$

$$\begin{array}{cccc} u_1u_2 & u_3u_4 & u_5u_6 & u_7 \\ e_1e_4 & e_8 - e_2 + e_6 & e_2e_7 & 2e_3 + e_5 \end{array}$$

$$\text{Basis of } U_2 \quad m_2 = 7$$

$$A^3 = 0$$

$$U_3 = \ker A^3 = \mathbb{R}^8$$

Basis of U_3 .

$$\begin{array}{cccc} u_1u_2 & u_3u_4 & u_5u_6 & u_7u_8 \\ e_1e_4 & e_8 - e_2 + e_6 & e_2e_7 & 2e_3 + e_5e_3 \end{array}$$

$$p = 3 \quad d = 4$$

→ 4 blocks, $n_i \leq 3$.

$$A \cdot v_1^{(3)} = A \cdot e_3 = 3e_2 - 2e_6$$

$$\begin{array}{cccc} u_1 u_2 & u_3 u_4 & u_1^{(2)} u_2^{(2)} & v_3^{(3)} v_1^{(3)} \\ e_1 e_4 & e_8 - e_2 + e_6 & 3e_2 - 2e_6 e_7 & 2e_3 + e_5 e_3 \end{array}$$

$$v_1^{(3)} = v_8 = e_3$$

$$v_1^{(2)} = A \cdot v_1^{(3)} = 3e_2 - 2e_6 \quad v_2^{(2)} = e_7 \quad v_3^{(2)} = 2e_3 + e_5$$

$$v_1^{(1)} = A v_1^{(2)} = e_1 + 3e_4 - 4e_8 \quad v_2^{(1)} = -e_2 + e_6 \quad v_3^{(1)} = 4e_2 + e_4 - 4e_6 \quad v_4^{(1)} = e_1$$

$$v_1^{(1)} = A \cdot v_1^{(2)} = A \cdot \begin{bmatrix} 0 \\ 3 \\ 0 \\ 0 \\ 0 \\ -2 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 3 \\ 0 \\ 0 \\ 0 \\ -4 \end{bmatrix}$$

$$v_2^{(1)} = A \cdot v_2^{(2)} = A e_7 = -e_2 + e_6$$

$$v_3^{(2)} = A \cdot \begin{bmatrix} 0 \\ 0 \\ 2 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 4 \\ 0 \\ 1 \\ 0 \\ 0 \\ -4 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{array}{cccccc} u_1 & u_2 u_3 & u_4 u_5 & u_6 u_7 & u_8 \\ \text{step } 2e_1 & e_4 e_8 & -e_2 + e_6 3e_2 - 2e_6 & e_7 2e_3 + e_5 & e_3 \\ \text{step } 3e_1 + 3e_4 - 4e_8 & -e_2 + e_6 4e_2 + e_4 - 4e_6 & e_1 3e_2 - 2e_6 & e_7 2e_3 + e_5 & e_3 \end{array}$$

$$B = \underbrace{e_1 + 3e_4 - 4e_8, 3e_2 - 2e_6, e_3}_{n_1=3} \mid \underbrace{-e_2 + e_6, e_7}_{n_2=2} \mid \underbrace{4e_2 + e_4 - 4e_6, 2e_3 + e_5}_{n_3=2} \mid \underbrace{e_1}_{n_4=1}$$

$$\Phi_B^B(A) = \begin{bmatrix} 0 & 1 & & & \\ & 0 & 1 & & \\ & & 0 & 1 & \\ & & 0 & 0 & \\ & & & 0 & 1 \\ & & & 0 & 0 \\ & & & & & 0 \end{bmatrix} \text{ is the Jordan norm form of } A$$

↓ This lecture took place on 2018/06/04.

$$A \text{ nilpotent} \implies A^k = 0$$

∃ basis B such that

$$B^{-1}AB = \begin{bmatrix} 0 & 1 & & & \\ & 0 & 1 & & \\ & & \ddots & \ddots & \\ & & & 0 & 1 \end{bmatrix}$$

In general: Decomposition in main spaces:

$$U_i = \ker(\lambda_i I - A)^n \quad V = \bigoplus_{i=1}^l U_i$$

$$(\lambda_i I - A)|_{U_i} \text{ is nilpotent}$$

↪ basis B_i such that

$$(\lambda_i I - A)|_{U_i} = \begin{bmatrix} 0 & 1 & & & \\ & 0 & \ddots & & \\ & & 0 & 0 & \\ & & & 0 & 1 \\ & & & & \ddots & \ddots \end{bmatrix}$$

$$A|_{U_i} \sim \begin{bmatrix} \lambda_i & 0 & & & \\ & \lambda_i & 1 & & \\ & & \ddots & \ddots & \\ & & & \ddots & 1 \\ & & & & \lambda_i & 1 \end{bmatrix}$$

11.6 Jordan block

Definition 11.3. A matrix of form $J_k(\lambda) = \begin{bmatrix} \lambda & 1 & & & \\ & \lambda & 0 & & \\ & & \lambda & 1 & \\ & & & \lambda & 1 \\ & & & & \ddots & \ddots \end{bmatrix} \in \mathbb{K}^{n+m}$ is called

Jordan block of length k to eigenvalue λ .

Remark 11.5. 1. $\chi_{J_k(\lambda)}(x) = (x - \lambda)^k$
2. $J_k(\lambda) - \lambda \cdot I$ is nilpotent with index k .

Theorem 11.7. Let \mathbb{K} be an algebraically closed field (hence, every polynomial has roots). Then every matrix $A \in \mathbb{K}^{n+m}$ is similar to a matrix of Jordan normal form.

$$\implies \exists B \in GL(\mathbb{K}, n) : B^{-1}AB = \begin{bmatrix} J_1 & & \\ & J_2 & \\ & & \ddots \\ & & & J_n \end{bmatrix}$$

where every J_i is a Jordan block to eigenvalue of A .

Proof. Let $\lambda_1, \dots, \lambda_m$ be the different eigenvalues of A . Let U_i be the main spaces.

$$U_i = \ker(\lambda_i I - A)^n$$

$$V = U_1 \oplus \dots \oplus U_m$$

By Theorem 11.4 and the field is algebraically closed,

$$\implies U_i \text{ are invariant } \wedge (\lambda_i I - A)|_{U_i} \text{ is nilpotent}$$

By Theorem 11.6, \exists basis B_i of U_i ,

$$\Phi_{B_i}^{B_i}((\lambda_i I - A)|_{U_i}) = \begin{bmatrix} N_{n_{i_1}} & & \\ & N_{n_{i_2}} & \\ & & \ddots \\ & & & N_{n_{i_d}} \end{bmatrix}$$

with

$$N_k = \begin{bmatrix} 0 & 1 & & \\ & \ddots & \ddots & \\ & & \ddots & 1 \\ & & & 0 \end{bmatrix} = J_k(0)$$

where d_i is the geometric multiplicity of eigenvalue $\lambda_i = \dim \ker(\lambda_i I - A)$.

$$B = B_1 \cup B_2 \cup \dots \cup B_n$$

$$\Rightarrow \Phi_B^B(A) = B^{-1}AB = \begin{bmatrix} J_{n_1 1}(\lambda_1) & & & & \\ & J_{n_1 2}(\lambda_1) & & & \\ & & \ddots & & \\ & & & J_{n_1 d_1}(\lambda_1) & \\ & & & & \ddots \\ & & & & & J_{n_n 1}(\lambda_n) \\ & & & & & & \ddots \\ & & & & & & & J_{n_n d_n}(\lambda_n) \end{bmatrix}$$

where columns of n_1 give U_1 and columns of n_n give U_n . \square

Theorem 11.8. Let $B^{-1}AB = \begin{bmatrix} J_1 & & \\ & \ddots & \\ & & J_q \end{bmatrix} \in \mathbb{K}^{n+m}$ be a Jordan normal form with

$$J_i = J_{k_i}(\lambda_i).$$

1. $\sum_{i=1}^q k_i = n$
2. $\text{spec}(A) = \{\lambda_i\}$, potentially with repetitions.

$$\forall \lambda \in \text{spec}(A) : d(\lambda) = \#\{i : \lambda_i = \lambda\}$$

$$k(\lambda) = \sum_{\lambda_i = \lambda} k_i$$

Geometric multiplicity of λ equals the number of corresponding Jordan blocks.
Diagonal multiplicity of $\lambda = \sum$ of size of corresponding Jordan blocks.

3. The smallest exponent r such that

$$\ker((\lambda I - A)^r) = \ker((\lambda I - A)^{r+1})$$

is the largest length of a corresponding Jordan block.

$$\min \{r : \ker((\lambda I - A)^r) = \ker((\lambda I - A)^{r+1})\} = \max \{k_i : \lambda_i = \lambda\}$$

The reason is given in an example:

Example 11.6.

$$\begin{aligned} A &= \begin{bmatrix} J_{k_1}(\lambda) & \\ & J_{k_2}(\lambda) \end{bmatrix} & A - \lambda I &= \begin{bmatrix} J_{k_1}(0) & \\ & J_{k_2}(0) \end{bmatrix} \\ (A - \lambda I)^r &= \begin{bmatrix} J_{k_1}(0)^r & \\ & J_{k_2}(0)^r \end{bmatrix} \stackrel{!}{=} 0 \\ \Rightarrow J_{k_1}(0)^r &= 0 \wedge J_{k_2}(0)^r = 0 \\ \Rightarrow r \geq k_1 \wedge r \geq k_2 &\Rightarrow r = \max \{k_1, k_2\} \end{aligned}$$

$$4. \# \{i : \lambda_i = \lambda \wedge k_i \geq k + 1\} = \text{rank}(\lambda I - A)^k - \text{rank}(\lambda I - A)^{k+1}$$

5. The Jordan blocks are uniquely determined (except for the order)

Proof. 1. Immediate.

2. For every Jordan block, there exists exactly one eigenvector.

$$(\text{rank}(J_k(\lambda) - \lambda I_k)) = k = 1$$

$$\chi_{J_{k_i}(\lambda_i)}(x) = (x - \lambda)^k$$

$$\Rightarrow \chi_A(x) = \prod_{i=1}^q \chi_{J_{k_i}(\lambda_i)}(x) = \prod_{i=1}^q (x - \lambda_i)^{k_i}$$

$$3. \text{ Let } A = \begin{bmatrix} J_1 & & \\ & \ddots & \\ & & J_q \end{bmatrix}.$$

$$(\lambda I - A)^k = \begin{bmatrix} (\lambda - J_1)^k & & \\ & \ddots & \\ & & (\lambda - J_q)^k \end{bmatrix}$$

If $\lambda \neq \lambda_i$, then $\lambda - J_k(\lambda_i)$ is regular (triangular matrix with all entries on the main diagonal $\neq 0$). If $\lambda = \lambda_i$, then $\lambda - J_{k_i}(\lambda_i)$ is nilpotent.

$$\text{rank}((\lambda - J_{k_i}(\lambda_i))^k) = \begin{cases} k_i - k & k_i > k \\ 0 & \text{else} \end{cases}$$

4.

$$\text{rank}(\lambda_i - J_{k_i}(\lambda_i))^k - \text{rank}(\lambda_i - J_{k_i}(\lambda_i))^{k+1} = \begin{cases} 1 & k_i \geq k + 1 \\ 0 & k_i \leq k \end{cases}$$

$$\text{rank}(\lambda - J_{k_i}(\lambda_i))^k = \begin{cases} k_i & \text{if } \lambda \neq \lambda_i \\ k_i - k & \text{if } \lambda = \lambda_i \text{ and } k_i \geq k \\ 0 & \text{if } \lambda = \lambda_i \wedge k_i < k \end{cases}$$

$$\text{rank}(\lambda - A)^k = \sum_{\lambda \neq \lambda_j} k_j + \sum_{\lambda = \lambda_i} \begin{cases} k_i - k & \text{if } k_i \geq k \\ 0 & \text{if } k_i < k \end{cases}$$

$$\text{rank}(\lambda - A)^{k+1} = \sum_{\lambda \neq \lambda_j} k_j + \sum_{\lambda = \lambda_i} \begin{cases} k_i - (k + 1) & \text{if } k_i \geq k \\ 0 & \text{else} \end{cases}$$

$$\begin{aligned}
\text{rank}(\lambda - A)^k - \text{rank}(\lambda - A)^{k+1} &= \overbrace{0}^{\text{first sum}} + \sum_{\lambda=\lambda_i} (k_i - k) - (k_i - (k+1)) \\
&= \sum_{\lambda=\lambda_i} 1 \text{ if } k_i > k+1 \\
&= \{i : k_i > k+1\}
\end{aligned}$$

5. Left as an exercise to the reader.

□

Lemma 11.6. Let $A \in \mathbb{K}^{n \times n}$ matrix.

$$\Psi_A : \mathbb{K}[x] \rightarrow \mathbb{K}^{n \times n}$$

$$p(x) \mapsto p(A)$$

$$a_0 + a_1x + \cdots + a_kx^k \mapsto a_0 \cdot I + a_1A + \cdots + a_kA^k$$

$$1. \ p \left(\begin{bmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix} \right) = \begin{bmatrix} p(\lambda_1) & & \\ & \ddots & \\ & & p(\lambda_n) \end{bmatrix}$$

$$2. \ p \left(\begin{bmatrix} A_1 & & \\ & \ddots & \\ & & A_n \end{bmatrix} \right) = \begin{bmatrix} p(A_1) & & \\ & \ddots & \\ & & p(A_n) \end{bmatrix}$$

$$3. \ A = T^{-1}BT \implies p(A) = T^{-1}p(B)T$$

Proof. Because of linearity, it suffices to show that this holds for basis polynomials x^k .

1. Immediate.

$$2. \ \begin{bmatrix} A_1 & & \\ & \ddots & \\ & & A_m \end{bmatrix}^k = \begin{bmatrix} A_1^k & & \\ & \ddots & \\ & & A_m^k \end{bmatrix}$$

3.

$$\begin{aligned}
A^k &= (T^{-1}BT)^k \\
&= (T^{-1}BT)(T^{-1}BT) \cdots (T^{-1}BT) \\
&= T^{-1}B^kT
\end{aligned}$$

$$\implies p(A) \text{ can be reduced to } p(\text{JNF}).$$

□

Lemma 11.7. For some Jordan block $J_k(\lambda)$ it holds that

$$p(J_k(\lambda))_{i,j} = \begin{cases} \frac{p^{(j-i)}(\lambda)}{(j-i)!} & j \geq i \\ 0 & j < i \text{ (below the diagonal)} \end{cases}$$

Proof.

$$(A + B)^M = \sum_{k=0}^M \binom{M}{k} A^k B^{M-k}$$

In general, $(A + B)^2 = AA + AB + BA + BB$, but here $A = I$ and therefore $AB = BA$.

$$\begin{aligned} (\lambda I + N)^M &= \sum_{k=0}^M \binom{M}{k} \lambda^{M-k} \cdot I \cdot N^k \\ &= \begin{bmatrix} \lambda^M & \binom{M}{1} \lambda^{M-1} & & \\ & \ddots & & \\ & & \lambda^M & \end{bmatrix} & k=0 \implies \text{diag}(\lambda^M), N^1 = I \\ &= \sum_{k=0}^M \frac{M(M-1)(M-2) \dots (M-k+1)}{k!} \cdot \lambda^{M-k} \cdot N^k \\ &= \sum_{k=0}^M \frac{(\lambda^M)^{(k)}}{k!} \end{aligned}$$

□

Example 11.7 (Application). 1. *Fibonacci \rightarrow Tribonacci:* $T_n = T_{n-1} + T_{n-2} + T_{n-3}$
(see practicals)

2. *Discrete dynamic systems: Predator-prey equations:*

$$F_n := \text{number of foxes} \quad H_n := \text{number of rabbits}$$

When is F_n and H_n stationary?

$$\begin{aligned} F_{n+1} &= p \cdot F_n + q \cdot H_n \\ H_{n+1} &= -t \cdot F_n + g \cdot H_n \\ \rightarrow \begin{pmatrix} F_{n+1} \\ H_{n+1} \end{pmatrix} &= \begin{bmatrix} p & q \\ -t & g \end{bmatrix} \cdot \begin{pmatrix} F_n \\ H_n \end{pmatrix} \end{aligned}$$

When it is balanced? This depends on the matrix ...

$$\begin{bmatrix} p & q \\ -t & g \end{bmatrix} \sim \begin{bmatrix} \lambda_1 & 1 \\ & \lambda_2 \end{bmatrix} \xrightarrow{n \rightarrow \infty} \begin{cases} 0 & |\lambda_1|, |\lambda_2| < 1 \\ \infty & |\lambda_1|, |\lambda_2| > 1 \\ \text{balance} & \text{if } |\lambda_1| = 1 \\ \text{dynamic balance} & \text{if } |\lambda_1| = |\lambda_2| = 1 \end{cases}$$

$$\vec{x}' = A \cdot \vec{x}$$

$$\frac{d}{dx} = A \cdot x \rightarrow \text{solution } x = e^{A \cdot t} \cdot x_0$$

11.7 Matrix exponentiation

Theorem 11.9 (Matrix exponential function). *In general: A is diagonalizable.*

$$\begin{aligned} \Rightarrow A &= T^{-1} \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} T \\ \rightarrow f(A) &= T^{-1} \begin{bmatrix} f(\lambda_1) & & \\ & \ddots & \\ & & f(\lambda_n) \end{bmatrix} T \end{aligned}$$

If not diagonalizable: Jordan norm form.

$$\leadsto f(J_k(\lambda)) = ?$$

- For polynomials, immediate.
- Otherwise, only works for analytical functions. Hence,

$$f(x) = \text{Taylor series} = \sum_{k=0}^{\infty} a_k (x - \lambda)^k$$

$$f(J_k(\lambda)) = \sum_{k=0}^{\infty} a_k \underbrace{(J_k(\lambda) - \lambda)^k}_{\text{nilpotent}}$$

nilpotent \Rightarrow series escapes \Rightarrow convergent.

$$f(x) = e^x \Rightarrow e^A = \sum_{k=0}^{\infty} \frac{A^k}{k!}$$

$$JNF \Rightarrow A = B \begin{bmatrix} J_1 & & \\ & \ddots & \\ & & J_n \end{bmatrix} B^{-1}$$

We have to determine e^{J_i} , because $e^A = B \cdot \begin{bmatrix} e^{J_1} & & \\ & \ddots & \\ & & e^{J_k} \end{bmatrix} B^{-1}$.

$$\begin{aligned}
 J &= \lambda \cdot I + N \\
 \rightarrow e^J &= e^{\lambda I + N} \\
 &= e^{\lambda I} \cdot e^N && \text{because } \lambda I \text{ and } N \text{ commute} \\
 &= e^\lambda \cdot \sum_{k=0}^{\infty} \frac{N^k}{k!} \\
 &= e^\lambda \cdot \sum_{k=0}^{M-1} \frac{N^k}{k!} && \text{if } N \text{ is nilpotent} \\
 &= e^\lambda \cdot \begin{bmatrix} 1 & \frac{1}{2} & \dots \\ & \ddots & \ddots \\ & & \frac{1}{2} & 1 \end{bmatrix}
 \end{aligned}$$

↓ This lecture took place on 2018/06/06.

Example 11.8.

$$e^A \begin{cases} \frac{dx}{dt} = Ax \leadsto x(t) = e^{A \cdot t} x_0 \\ x(0) = x_0 \end{cases}$$

for $t \rightarrow \infty : e^{At} x_0$ Asymptotically, depends on the eigenvalue.

$$A = B^{-1} \underbrace{J}_{JNF} B \quad e^{At} = B^{-1} e^{Jt} B = B^{-1} \begin{bmatrix} e^{J_1 t} & & \\ & e^{J_2 t} & \\ & & \ddots \\ & & & e^{J_k t} \end{bmatrix} \cdot B$$

$$J = \begin{bmatrix} J_1 & & \\ & J_2 & \\ & & \ddots \\ & & & J_k \end{bmatrix} \text{ Jordan blocks}$$

$$J_k = \begin{bmatrix} \lambda & 1 & & \\ & \ddots & \ddots & \\ & & \ddots & \ddots \\ & & & \ddots & 1 \\ & & & & \lambda \end{bmatrix}$$

$$e^{J_k t} = e^{\lambda t} \cdot e^{\left[\begin{array}{c} \overbrace{\begin{bmatrix} 0 & 1 & & \\ & \ddots & \ddots & \\ & & \ddots & 1 \\ & & & 0 \end{bmatrix}}^{\text{polynomial}} \\ t \end{array} \right]}$$

$$e^{\lambda t} = e^{(\xi + i\eta)t} \xrightarrow{t \rightarrow \infty} \begin{cases} \infty & \text{if } \xi > 0 \\ 0 & \text{if } \xi < 0 \end{cases}$$

$$e^{(\xi + i\eta)t} = e^{\xi t} \cdot e^{i\eta t}$$

with $|e^{i\eta t}| = 1$ and $\xi = \Re \lambda$. \rightarrow if $\Re \lambda < 0 \forall$ eigenvalue λ_i .

$$e^{A t} x_0 \xrightarrow{t \rightarrow \infty} 0$$

for arbitrary x_0 . If $\Re \lambda_i > 0 \forall$ eigenvalue $\implies e^{A t} x_0 \xrightarrow{t \rightarrow \infty} \infty$.

If $\Re \lambda_i < 0$ for some specific λ_i and $\Re \lambda_i > 0$ for other λ_i . Asymptotically depends on the initial value x_0 .

Example 11.9 (Pendulum).

$$m \cdot l \cdot \ddot{\varphi} = -m \cdot g \cdot \sin \varphi \approx -g \cdot \varphi$$

$$l \cdot \ddot{\varphi} = -g \cdot \varphi = -\omega^2 \cdot \varphi$$

$$\psi = \frac{\dot{\varphi}}{\omega} \quad \dot{\varphi} = \omega \cdot \psi$$

$$\dot{\psi} = \frac{\ddot{\varphi}}{\omega} = -\frac{\omega^2 \cdot \varphi}{\omega} = -\omega \cdot \varphi$$

$$\frac{d}{dt} \begin{pmatrix} \varphi \\ \psi \end{pmatrix} = \begin{pmatrix} \omega \psi \\ -\omega \varphi \end{pmatrix} = \begin{bmatrix} 0 & \omega \\ -\omega & 0 \end{bmatrix} \begin{pmatrix} \varphi \\ \psi \end{pmatrix}$$

$$\begin{pmatrix} \varphi(t) \\ \psi(t) \end{pmatrix} = e^{\begin{bmatrix} 0 & \omega \\ -\omega & 0 \end{bmatrix} t} \begin{bmatrix} \varphi_0 \\ \psi_0 \end{bmatrix}$$

eigenvalue: $\lambda^2 + \omega^2 = 0$, $\omega = \pm i\omega$.

$$\varphi(t) \sim \Re(c \cdot e^{i\omega t} \sim a \cos \omega t + b \cdot \sin \omega t)$$

$$\omega = \sqrt{\frac{g}{l}}$$

Example 11.10.

$$\begin{aligned}\frac{\partial T(x, t)}{\partial t} &= \Delta T(x, t) \\ \Delta T(x, t) &= \frac{\partial^2}{\partial x^2} T(x, t) \\ \Delta &= \sum \frac{\partial^2}{\partial x_i^2} \quad \text{Laplace operator} \\ T(x, t) &= e^{t\Delta} T(x, 0)\end{aligned}$$

Example 11.11 (Schrödinger's equation).

$$-\frac{\hbar}{i} \frac{\partial \psi}{\partial t} = H\psi \rightsquigarrow \psi(t) = e^{-\frac{i}{\hbar} \Delta t} \psi_0$$

Hamilton:

$$H = \Delta + V(x)$$

11.8 Minimal polynomial and annihilator

Definition 11.4 (Theorem and definition). Let $A \in \mathbb{K}^{N \times N}$.

1. $\exists p(x) \in \mathbb{K}[x] : p(A) = 0$
2. \exists a unique polynomial $m_A(x) \in \mathbb{K}[x]$ with minimal degree and leading coefficient 1. $m_A(x)$ is called minimal polynomial of A .
3. $\text{Ann}(A) = \{p(x) \in \mathbb{K}[x] \mid p(A) = 0\}$ is called annihilator of A and it holds that $p(x) \in \text{Ann}(A) \iff m_A(x) \mid p(x)$
4. $m_A(\lambda) = 0 \forall \lambda \in \text{spec}(A)$

Proof. 1. $A^0, A^1, A^2, A^3, \dots \in \mathbb{K}^{N \times N}$. Infinitely many elements of a finite dimensional vector space are linear dependent.

$$\implies \exists n \exists a_0, a_1, \dots, a_n : a_0 A^0 + \underbrace{a_1 A^1}_{=p(A)} + \dots + a_n A^n = 0$$

$$p(x) = a_0 + a_1 x + \dots + a_n x^n$$

2. + 3. Let n be minimal such that A^0, \dots, A^n are linear dependent ($\implies n \leq N^2$) and $a_0 I + a_1 A + \dots + a_n A^n = 0$ with $a_n \neq 0$ (this will be shown in the practicals).

$$\implies m_A(x) = \frac{a_0}{a_n} + \frac{a_1}{a_n} x + \dots + \frac{a_{n-1}}{a_n} x^{n-1} + x^n$$

is the unique minimal polynomial.

Assume $p(x) \in \mathbb{K}[x]$ with $p(A) = 0 \implies \deg(p(x)) \geq n$.

By the division algorithm: $\exists q(x) \in \mathbb{K}[x] \exists r(x) \in \mathbb{K}[x]$ with $\deg(r(x)) < \deg(m_A(x))$ and $p(x) = q(x)m_A(x) + r(x)$. Insert A :

$$\begin{aligned} p(A) &= q(A) \cdot m_A(A) + r(A) \\ 0 &= 0 + r(A) \\ \implies r(A) &= 0 \\ \deg(r(x)) < n &\stackrel{\text{minimality } n}{\implies} r(x) \equiv 0 \\ \implies p(x) &= q(x) \cdot m_A(x) \\ \implies m_A(x) | p(x) &\implies (c) \end{aligned}$$

Especially, if $\deg(p(x)) = n = \deg(m_A(x)) \implies \deg(q(x)) = 0 \implies p(x) = c \cdot m_A(x)$.

4. Will be shown in the practicals.

□

11.9 Cayley-Hamilton Theorem

Theorem 11.10 (Cayley-Hamilton Theorem).

$$\chi_A(A) = 0 \quad (\iff \chi_A(x) \in \text{Ann}(A))$$

Corollary 11.3.

$$m_A(x) | \chi_A(x)$$

and therefore the roots of $m_A(x)$ are the eigenvalues of A .

Proof. Three different proofs will be given:

1. $\chi_A(x) = \det(xI - A)$ and $\chi_A(A) = \det(AI - A) = 0$ (incorrect, used among physicists)
2. Using Jordan's normal form (if \mathbb{K} is algebraically closed¹¹):

$$\begin{aligned} A &= B \cdot J \cdot B^{-1} \\ \chi_A(A) &= B \cdot \chi_A(J) \cdot B^{-1} \\ J &= \begin{bmatrix} J_1 & & \\ & \ddots & \\ & & J_q \end{bmatrix} \rightsquigarrow \chi_A(J) = \begin{bmatrix} \chi_A(J_1) & & \\ & \ddots & \\ & & \chi_A(J_q) \end{bmatrix} \end{aligned}$$

¹¹And therefore the third proof is required.

$$J_i = \begin{bmatrix} \lambda_k & 1 & & \\ & \ddots & \ddots & \\ & & \ddots & 1 \\ & & & \lambda_{n_i} \end{bmatrix}$$

We know that $\sum_{i, \lambda_i = \lambda} n_i = k(\lambda)$ algebraic multiplicity of eigenvalue λ . $\chi_A(J_i)$.

$$\begin{aligned} \chi_A(x) &= \prod_j (\lambda - \lambda_j)^{k_j} \\ \chi_A(J_i) &= \prod_{j \neq i} (J_i - \lambda_j I)^{k_j} \cdot \underbrace{(J_i - \lambda_i I)^{k_i}}_{\begin{bmatrix} 0 & 1 & & \\ & \ddots & \ddots & \\ & & \ddots & 1 \\ & & & 0 \end{bmatrix}^{k_i} = 0} \end{aligned}$$

$n_i \leq k_i$

$\chi_A(j) = 0$ for all Jordan blocks of A .

$$\chi_A(A) = B \begin{bmatrix} \chi_A(J_1) & & \\ & \ddots & \\ & & \chi_A(J_q) \end{bmatrix} B^{-1} = 0$$

3. Complementary matrix:

$$A \cdot \widehat{A} = \det(A) \cdot I$$

$$\widehat{A}_{ij} = -(-1)^{i+j} \det(A_{ji})$$

where A_{ji} denotes removing the j -th row and i -th column.

$$xI - A \in \mathbb{K}[x]^{n \times n} = [b_{ij}(x)]_{i,j=1,\dots,n}$$

$$b_{ij}(x) = (-1)^{i+j} \det(xI - A)_{ji} \in \mathbb{K}[x] \text{ with degree } \leq n-1$$

$$b_{ij}(x) = \sum_{k=0}^{n-1} b_{ijk} x^k$$

$$\begin{aligned} xI - A &= \left[\sum_{k=0}^{n-1} b_{ijk} x^k \right]_{i,j=1,\dots,n} \\ &= \sum_{k=0}^{n-1} [b_{ijk}]_{i,j=1,\dots,n} \cdot x^k \\ &= \sum_{k=0}^{n-1} B_k \cdot x^k \end{aligned}$$

Example 11.12.

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$\begin{aligned} \widehat{xI - A} &= \begin{bmatrix} \widehat{x - a} & -b \\ -c & \widehat{x - d} \end{bmatrix} \\ &= \begin{bmatrix} x - d & b \\ c & x - a \end{bmatrix} \in \mathbb{K}[x]^{2 \times 2} \quad \text{matrix with polynomial entries} \\ &= \underbrace{\begin{bmatrix} -d & b \\ c & -a \end{bmatrix}}_{B_0} + \underbrace{\begin{bmatrix} 1 & \\ & 1 \end{bmatrix}}_{B_1 x} x \in \mathbb{K}^{2 \times 2}[x] \end{aligned}$$

$$\begin{aligned} (xI - A)(\widehat{xI - A}) &= \det(xI - A) \cdot I \\ &= \chi_A(x) \cdot I \end{aligned}$$

$$\text{Let } \chi_A(x) = c_0 + c_1x + \cdots + c_{n-1}x^{n-1} + x^n.$$

$$\implies (xI - A)(B_0 + B_1x + B_2x^2 + \cdots + B_{n-1}x^{n-1}) = (c_0 + c_1x + \cdots + c_{n-1}x^{n-1} + x^n)I$$

$$-AB_0 + (B_0 - AB_1)x + (B_1 - AB_2)x^2 + \cdots + (B_{n-2} - AB_{n-1})x^{n-1} + B_{n-1}x^n = c_0I + c_1Ix + \cdots + I \cdot x^n$$

We apply coefficient comparison (multiply with A^k from left):

$$\begin{aligned} c_0 \cdot I &= -AB_0 & c_0I &= -AB_0 \\ c_1 \cdot I &= B_0 - AB_1 & +c_1A &= AB_0 - A^2B_1 \\ c_2 \cdot I &= B_1 - AB_2 & +c_2A^2 &= A^2B_1 - A^3B_2 \\ &\vdots & \\ c_{n-1} \cdot I &= B_{n-2} - AB_{n-1} & +c_{n-1}A^{n-1} &= A^{n-1}B_{n-2} - A^nB_{n-1} \\ I &= B_{n-1} & +A^n &= A^nB_{n-1} \end{aligned}$$

$$\chi_A(A) = \sum_{k=0}^{n-1} c_k A^k + A^n = -AB_0 + \sum_{k=1}^{n-1} (A^k B_{k-1} - A^{k+1} B_k) + A^n B_{n-1} = 0$$

Thus this proof has proven it for every zero-divisor-free field.

□

Corollary 11.4 (Corollary for second proof).

1. The minimal polynomial has the structure $m_A(x) = \prod (\lambda - \lambda_i)^{m_i}$ where m_i is the smallest exponent for $\ker(\lambda_i - A)^m = \ker(\lambda_i - A)^{m+1}$, hence this equals the largest length of a Jordan block for eigenvalue λ_i .

2. A is diagonalizable \iff all $m_i = 1$ $\iff m_A(x) = \prod_{i=1}^k (\lambda - \lambda_i)$ $\iff m_A(x)$ has only simple roots.

Example 11.13 (Application). Let $A \in \mathbb{K}^{2 \times 2}$. We consider $A \in \mathbb{C}^{2 \times 2}$.

$$e^{\alpha I + A} = e^\alpha \cdot e^A$$

Without loss of generality: $\text{Tr}(A) = 0$. Otherwise consider $\mathring{A} = A - \frac{\text{Tr}(A)}{2} \cdot I$.

$$\implies \text{Tr}(\mathring{A}) = \text{Tr}(A) - \frac{\text{Tr}(A)}{2} \cdot \text{Tr}(I) = 0$$

$$e^A = e^{\frac{\text{Tr}(A)}{2}} \cdot e^{\mathring{A}}$$

Without loss of generality: $A = \begin{bmatrix} a & b \\ c & -a \end{bmatrix}$.

$$\begin{aligned} \chi_A(A) &= (X - a)(X + a) - bc \\ &= x^2 - a^2 - bc \\ &= x^2 - \delta \end{aligned}$$

$$\delta = a^2 + bc = -\det(A)$$

$$e^A = \sum_{n=0}^{\infty} \frac{A^n}{n!}$$

Cayley-Hamilton Theorem:

$$\chi_A(A) = 0$$

$$A^2 - \delta I = 0 \implies A^2 = \delta I, A^3 = \delta A, A^4 = (A^2)^2 = \delta^2 \cdot I$$

$$A^{2n} = \delta^n \cdot I$$

$$A^{2n+1} = \delta^n \cdot A$$

$$\begin{aligned} e^A &= \sum_{n=0}^{\infty} \frac{A^n}{n!} = \sum_{k=0}^{\infty} \frac{A^{2k}}{2k!} + \sum_{k=0}^{\infty} \frac{A^{2k+1}}{(2k+1)!} \\ &= \sum_{k=0}^{\infty} \frac{\delta^k}{2k!} I + \sum_{k=0}^{\infty} \frac{\delta^k}{(2k+1)!} \cdot A = \dots \end{aligned}$$

$$Ax = \lambda x$$

$$A^2 x = \lambda^2 x$$

$$A^k x = \lambda^k x$$

$$p(A) \cdot x = p(\lambda) \cdot x$$

if $\lambda \in \text{spec}(A)$, then $p(\lambda) \in \text{spec}(p(A))$.

11.10 Spectrum mapping theorem

Theorem 11.11 (Spectrum mapping theorem). For $A \in \mathbb{K}^{n \times n}$ and \mathbb{K} be algebraically closed and $p(x) \in \mathbb{K}[x]$ is $\text{spec}(p(A)) = p(\text{spec}(A)) = \{p(\lambda) \mid \lambda \in \text{spec}(A)\}$.

Proof. \supseteq

$$\forall \lambda \in \text{spec}(A) : p(\lambda) \in \text{spec}(p(A))$$

\subseteq

$$\forall \mu \in \text{spec}(p(A)) : \exists \lambda \in \text{spec}(A) : p(\lambda) = \mu$$

□

Example 11.14. If \mathbb{K} is not algebraically closed, then \subseteq does not hold! For example, consider $A = \begin{bmatrix} & 1 \\ -1 & \end{bmatrix} \in \mathbb{R}^{2 \times 2}$.

$$\text{spec}(A) = \{\pm i\} \implies \text{spec}_{\mathbb{R}}(A) = \emptyset$$

$$p(x) = x^2 \implies A^2 = -I$$

has eigenvalue $\{-1\}$ but there exists no $\lambda \in \text{spec}_{\mathbb{R}}(A)$ such that $\lambda^2 = -1$.

Let $\mu \in \text{spec}(p(A))$.

$$q(x) = p(x) - \mu = (x - \mu_1) \dots (x - \mu_m)$$

where μ_i are the roots of $q(x)$.

$$\implies q(A) = p(A) - \mu I \text{ is not invertible}$$

$$q(A) = (A - \mu_1 I)(A - \mu_2 I) \dots (A - \mu_m I) \text{ not invertible}$$

$$\implies \exists i : (A - \mu_i I) \text{ not invertible}$$

$$\implies \mu_i \in \text{spec}(A)$$

$$\implies q(\mu_i) = 0$$

$$q(\mu_i) = p(\mu_i) - \mu$$

$$\implies \mu = p(\mu_i) \text{ and } \mu_i \in \text{spec}(A)$$

↓ This lecture took place on 2018/06/11.

12 Normal matrices

Definition 12.1 (Revision: Inner product).

$$\langle x, y \rangle \text{ sesquilinear}$$

where the first argument is linear and the second argument is antilinear.

$$\begin{aligned}\langle \lambda x + \mu y, z \rangle &= \lambda \langle x, z \rangle + \mu \langle y, z \rangle \\ \langle x, y \rangle &= \overline{\langle y, x \rangle} \\ \langle x, x \rangle &\geq 0 \text{ or equivalently, } \langle x, x \rangle = 0 \iff x = 0\end{aligned}$$

Adjoint map: $f : V \rightarrow W$ where B is the basis of V and C is the basis of W .

$$\langle f(x), y \rangle = \langle x, f^*(y) \rangle$$

f^* is linear: $W \rightarrow V$. $f \mapsto f^*$ is antilinear.

$\Phi_C^B(f)$ is its matrix representation

$$\begin{aligned}\Phi_B^C(f^*) &= \Phi_C^B(f)^* \\ (A^*)_{ij} &= \overline{a_{ji}}\end{aligned}$$

Definition 12.2. A linear map $f : V \rightarrow V$ (and accordingly a matrix A) is called normal if

$$f \circ f^* = f^* \circ f \quad AA^* = A^*A$$

Special case: $A = A^*$ is self-adjoint.

A real-valued self-adjoint matrix is called symmetrical: $A = A^T$.

Example 12.1. Unitary matrices are normal: $U^*U = I \implies UU^* = I$. A, B normal $\implies A \cdot B$.

$$\begin{aligned}(AB)^* \cdot AB &\stackrel{?}{=} AB(AB)^* = ABB^*A^* \\ &= (AB)^* \cdot AB = B^*A^* \cdot AB \text{ only if } AB = BA\end{aligned}$$

Example 12.2. $A + B$ is normal if and only if $AB = -BA$.

$$A, B \text{ self-adjoint} \implies A + B \text{ self-adjoint}$$

$$(A + B)^* = A^* + B^* = A + B$$

Lemma 12.1. $A \in \mathbb{C}^{n \times n}$ is normal.

1. $\ker A = \ker A^*$
2. $\ker A = \ker A^2$

Corollary 12.1. $A \in \mathbb{C}^{n \times n}$ is normal. (So $\lambda I + A$ is normal)

1. $\ker(\lambda I - A) = \ker(\bar{\lambda}I - A^*)$
2. $\ker(\lambda I - A)^2 = \ker(\lambda I - A)$

\implies main spaces = eigenspaces
 \implies diagonalizable

Proof. 1.

$$\begin{aligned}
 x \in \ker A &\iff Ax = 0 \\
 &\iff \|Ax\|^2 = 0 \\
 &\iff \langle Ax, Ax \rangle = 0 \\
 &\iff \langle x, A^*Ax \rangle = 0 \\
 &\iff \langle x, AA^*x \rangle = 0 \\
 &\iff \langle A^*x, A^*x \rangle = 0 \\
 &\iff \|A^*x\|^2 = 0 \\
 &\iff \|A^*x\| = 0 \\
 &\iff A^*x = 0 \\
 &\iff x \in \ker A^*
 \end{aligned}$$

2. $\ker A \subseteq \ker A^2$ immediate (Fitting)

$$\begin{aligned}
 \text{Let } x \in \ker A^2 &\implies A^2x = 0 \\
 &\implies Ax \in \ker A \\
 &\implies Ax \in \ker A^* \\
 &\implies A^*Ax = 0 \\
 &\implies \langle A^*Ax, x \rangle = 0 \\
 &\implies \langle Ax, Ax \rangle = 0 \\
 &\implies \|Ax\|^2 = 0 \\
 &\implies Ax = 0 \\
 &\implies x \in \ker A
 \end{aligned}$$

□

Lemma 12.2. Let A be normal.

$$\lambda \neq \mu \in \text{spec } A \implies \ker(\lambda I - A) \perp \ker(\mu I - A)$$

Proof. Let $Ax = \lambda x$ and $Ay = \mu y$

$$\xRightarrow{\text{Corollary 12.1}} A^*y = \bar{\mu}y$$

$$\begin{aligned} \langle Ax, y \rangle &= \langle \lambda x, y \rangle = \lambda \langle x, y \rangle \\ \langle Ax, y \rangle &= \langle x, A^*y \rangle = \langle x, \bar{\mu}y \rangle = \bar{\mu} \langle x, y \rangle \\ \implies \lambda \langle x, y \rangle &= \bar{\mu} \langle x, y \rangle \\ \implies \underbrace{(\lambda - \mu)}_{\neq 0} \langle x, y \rangle &= 0 \implies \langle x, y \rangle = 0 \end{aligned}$$

□

Remark 12.1 (Summary). *Let A be normal. Then main spaces are eigenspaces. So they are diagonalizable. Then there exists some basis of eigenvectors. Eigenspaces are always orthogonal, so there exists an orthogonal basis of eigenvectors.*

Theorem 12.1. *For $A \in \mathbb{C}^{n \times n}$, the following statements are equivalent:*

1. A is normal.
2. \exists orthonormal basis of eigenvectors
3. \exists unitary matrix U such that $U^*AU = \text{diag}(\lambda_1, \dots, \lambda_n)$, hence A is unitarily diagonalizable.

Proof. 1. \rightarrow 2. A is diagonalizable, so there exists a basis of eigenvectors and accordingly, $V = \mathbb{C}^n$ is the orthogonal sum of eigenspaces. Construct an orthonormal basis in every eigenspace (Gram-Schmidt process). Thus, the union of these bases gives an orthonormal basis of V .

2. \rightarrow 3. Let B be an orthonormal basis of eigenvectors.

$$\implies U = \begin{pmatrix} b_1 & \dots & b_n \\ \vdots & & \vdots \end{pmatrix} \implies AU = U \cdot \Lambda$$

$$U^{-1} = U^* \implies AU = U\Lambda$$

$$\text{where } \Lambda = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}.$$

3. \rightarrow 1.

$$U^*AU = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}$$

$$\Rightarrow A = U \begin{pmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{pmatrix} U^*$$

$$A^* = U \begin{bmatrix} \bar{\lambda}_1 & & \\ & \ddots & \\ & & \bar{\lambda}_n \end{bmatrix} U^*$$

$$\begin{aligned} AA^* &= U \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \cdot \underbrace{U^* \cdot U}_I \begin{bmatrix} \bar{\lambda}_1 & & \\ & \ddots & \\ & & \bar{\lambda}_n \end{bmatrix} U^* \\ &= U \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \begin{bmatrix} \bar{\lambda}_1 & & \\ & \ddots & \\ & & \bar{\lambda}_n \end{bmatrix} U^* \\ &= U \begin{bmatrix} \lambda_1 \bar{\lambda}_1 & & \\ & \ddots & \\ & & \lambda_n \bar{\lambda}_n \end{bmatrix} \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} U^* \\ &= U \begin{bmatrix} \bar{\lambda}_1 & & \\ & \ddots & \\ & & \bar{\lambda}_n \end{bmatrix} U^* \cdot U \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} U^* \\ &= A^* A \end{aligned}$$

□

Remark 12.2. Issai Schur (1875–1941)

12.1 Schur's decomposition, QR decomposition

Theorem 12.2 (Schur's decomposition). *Let $A \in \mathbb{C}^{n \times n}$.*

1. $\Rightarrow \exists U \in \mathcal{U}(n) : U^* A U = R$ is an upper triangular matrix.

$$U^* A U = \begin{bmatrix} \lambda_1 & & * \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix} \quad \lambda_i = \text{eigenvalue of } A$$

is called Schur decomposition, Schur normal form or QR decomposition.

2. *If $A \in \mathbb{R}^{n \times n}$ and $\chi_A(\lambda)$ decomposes into linear factors.*

$\Rightarrow U$ has real-valued entries, thus $U \in O(n)$

Proof. Proof by induction over n .

Induction base $n = 1$ immediate.

Induction step Let $\lambda \in \text{spec}(A)$. Let u be an eigenvector with $\|u\| = 1$.

$$Au = \lambda u$$

Extend u to an orthonormal basis of \mathbb{C}^n

$$(u, w_1, \dots, w_{n-1})$$

$$U_1 = \begin{bmatrix} u & & \\ \vdots & W & \\ \vdots & & \end{bmatrix}$$

$$W = \begin{bmatrix} w_1 & \dots & w_{n-1} \\ \vdots & & \vdots \end{bmatrix} \in \mathbb{C}^{n \times (n-1)}$$

$$U_1^* = \begin{bmatrix} u^* & \dots & \dots \\ & W^* & \end{bmatrix} \text{ where } W^* \in \mathbb{C}^{(n-1) \times n}$$

$$U_1^* A U_1 = \begin{bmatrix} u^* & \dots & \dots \\ & W^* & \end{bmatrix} A \begin{bmatrix} u \\ \vdots \\ \vdots & W \end{bmatrix}$$

$$= \begin{bmatrix} u^* & \dots & \dots & \dots \\ & & W^* & \end{bmatrix} \begin{bmatrix} \lambda u \\ \vdots \\ \vdots & AW \end{bmatrix} \text{ where } AW \in \mathbb{C}^{n \times (n-1)} = \begin{bmatrix} \lambda & u^* A W \\ 0 & \\ \vdots & [W^* A W] \\ 0 & \end{bmatrix} \text{ where } W^* A W \in \mathbb{C}^{(n-1) \times (n-1)}$$

$$I = U^* U = \begin{bmatrix} u^* & \dots & \dots & \dots \\ & & W^* & \end{bmatrix} \begin{bmatrix} u \\ \vdots \\ \vdots & W \end{bmatrix} = \begin{bmatrix} 1 & u^* W \\ 0 & \\ \vdots & W^* w \\ 0 & \end{bmatrix}$$

By induction hypothesis, there exists $U_2 \in \mathcal{U}(n-1)$ TODO

□

Corollary 12.2. *A matrix is normal \iff Schur normal form = diagonal matrix.*

$$U^*AU = R \implies A = URU^*$$

$$A^* = UR^*U^*$$

$$A^*A = AA^* \iff R^*R = RR^* \iff R \text{ is diagonal matrix}$$

The proof is left to the reader as an exercise.

Theorem 12.3. *Let $A \in \mathbb{C}^{n \times n}$, $A = A^* \implies \text{spec}(A) \subseteq \mathbb{R}$.*

Compare with $z \in \mathbb{C} : z \in \mathbb{R} \iff z = \bar{z}$.

Proof. A is normal.

$$\begin{aligned} \ker(\lambda I - A) &= \ker(\lambda I - A)^* \\ &= \ker(\bar{\lambda} I - A^*) \\ &= \ker(\bar{\lambda} I - A) \end{aligned}$$

hence if $x \neq 0$ and $Ax = \lambda x \implies Ax = \bar{\lambda}x \implies \lambda = \bar{\lambda}$. □

Remark 12.3. *Matrix is real-valued \implies eigenvalues are real-valued.*

Corollary 12.3. *If $A \in \mathbb{R}^{n \times n}$ is symmetrical.*

$$\implies \exists Q \in O(n) : Q^T A Q = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \text{ with } \lambda_i \in \mathbb{R}$$

Remark 12.4 (Revision). *A matrix is called positive definite, if $A = A^*$ and $\langle Ax, x \rangle > 0 \forall x \neq 0$. Let x be an eigenvector. $Ax = \lambda x$ and $\langle Ax, x \rangle = \lambda \langle x, x \rangle = \lambda \cdot \|x\|^2 > 0$*

Theorem 12.4. *Let $A \in \mathbb{C}^{n \times n}$ self-adjoint.*

1. $A > 0 \iff \text{spec}(A) \subseteq]0, \infty[$, hence eigenvalues are > 0 .
2. $A \geq 0 \iff \text{spec}(A) \subseteq]0, \infty[$, hence eigenvalues ≥ 0 .
3. Analogously for negatively (semi)definite.
4. A is indefinite $\iff \exists$ at least one positive and one negative eigenvalue.

$$\text{spec}(A) \cap (-\infty, 0) \neq \emptyset \text{ and } \text{spec}(A) \cap (0, \infty) \neq \emptyset$$

Proof. 1. Direction \implies .

$$A > 0 \implies A \text{ is self-adjoint} \implies \text{spec}(A) \subseteq \mathbb{R}$$

$$\text{and } Ax = \lambda x \implies \lambda > 0 \text{ (self-adjoint)}$$

Direction \Leftarrow . A is self-adjoint, $\text{spec}(A) \subseteq (0, \infty)$. Show that $\langle Ax, x \rangle > 0 \forall x \neq 0$.

A is self-adjoint \Rightarrow there exists an orthonormal basis of eigenvectors (u_1, \dots, u_n) . Let $x \in \mathbb{C}^n \setminus \{0\} \Rightarrow x = \sum_{i=1}^n \alpha_i u_i$ and $A \cdot u_i = \lambda_i \cdot u_i$.

$$\begin{aligned} \Rightarrow \langle Ax, x \rangle &= \left\langle A \sum_{i=1}^n \alpha_i u_i, \sum_{j=1}^n \alpha_j u_j \right\rangle \\ &= \left\langle \sum_{i=1}^n \alpha_i \lambda_i u_i, \sum_{j=1}^n \alpha_j u_j \right\rangle \\ &= \sum_{i=1}^n \sum_{j=1}^n \alpha_i \lambda_i \underbrace{\overline{\alpha_j} \langle u_i, u_j \rangle}_{\delta_{ij}} \\ &= \sum_{i=1}^n |\alpha_i|^2 \lambda_i > 0 \text{ (if all } \lambda_i > 0) \\ &= \sum_{i=1}^n |\alpha_i|^2 \lambda_i \geq 0 \text{ (if all } \lambda_i \geq 0) \end{aligned}$$

This proves (1.) and (2.)

□

Remark 12.5 (Application: Taylor expansion).

$$\begin{aligned} f(x) &= f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2!}(x - x_0)^2 - \dots \\ f(x, y) &= f(x_0, y_0) + \frac{\partial f}{\partial x}(x_0, y_0)(x - x_0) + \frac{\partial f}{\partial y}(x_0, y_0)(y - y_0) + \frac{1}{2} \frac{\partial^2 f}{\partial x^2}(x_0, y_0)(x - x_0)^2 + \frac{1}{2} \frac{\partial^2 f}{\partial y^2}(x_0, y_0)(y - y_0)^2 \\ &\quad + \left(\frac{\partial^2 f}{\partial x \partial y}(x_0, y_0) + \frac{\partial^2 f}{\partial y \partial x}(x_0, y_0) \right) (x - x_0)(y - y_0) \end{aligned}$$

Gradient:

$$\begin{aligned} \nabla f &= \begin{pmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \end{pmatrix} \\ H_f(x_0, y_0) &= \begin{bmatrix} \frac{\partial^2 f}{\partial x^2} & \frac{\partial^2 f}{\partial x \partial y} \\ \frac{\partial^2 f}{\partial y \partial x} & \frac{\partial^2 f}{\partial y^2} \end{bmatrix} \quad \text{“Hesse matrix”} \\ f(x, y) &= f(x_0, y_0) + (\nabla f)^T \cdot \underbrace{\begin{pmatrix} x - x_0 \\ y - y_0 \end{pmatrix}}_{\Delta \vec{x}} + \frac{1}{2} (\Delta \vec{x})^T H_f \Delta \vec{x} + O((\Delta x)^2) \end{aligned}$$

$H_f^T = H_f$ has real eigenvalues. Extrema are found if $\nabla f = 0$. Minimum at $(x_0, y_0) \iff H_f(x_0, y_0) > 0$. Maximum at (x_0, y_0) if $H_f(x_0, y_0) < 0$.

↓ This lecture took place on 2018/06/13.

$$A = A^*$$

$$\mathbb{K} = \mathbb{R} : A = A^T \quad \text{Taylor series}$$

Example 12.3 (Bicycling). Consider the wheel when bicycling. It has radius r , ω is the angle. \vec{v} is the tangent.

Angular momentum:

$$\vec{L} = m \cdot \vec{r} \times \vec{v}$$

is maintained.

$$|\vec{L}| = m \cdot r \cdot v = m \cdot r^2 \cdot \omega$$

$\omega = \dot{\theta}$ is angular speed. $v = r \cdot \omega$.

$$\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{c}) \cdot \vec{b} - (\vec{a} \cdot \vec{b}) \cdot \vec{c}$$

$$\vec{v} = \vec{\omega} \times \vec{r}$$

$$\vec{L} = m \cdot \vec{r} \times \vec{v}$$

$$= m \cdot \vec{r} \times (\vec{\omega} \times \vec{r})$$

$$= m \cdot \|\vec{r}\|^2 \times \vec{\omega} - (\vec{r} \cdot \vec{\omega}) \cdot \vec{r}$$

$$= m \cdot \left(\|\vec{r}\|^2 I - \vec{r} \vec{r}^T \right) \cdot \vec{\omega}$$

$$= \underbrace{H}_{\text{inertia tensor}} \cdot \vec{\omega}$$

$$H = r^2 I - \vec{r} \cdot \vec{r}^T$$

\Rightarrow at least 3 main axes.

12.2 Application: Conic section

Example 12.4 (Conic section). Radius: $r = z$

Cone: $\{(x, y, z) | x^2 + y^2 = z^2\}$.

Plane: $E = \{u + \xi v + \mu w | \xi, \mu \in \mathbb{R}\}$. Without loss of generality $v \perp w$. $\|v\| = \|w\| = 1$.

$$u = \begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix} \quad v = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} \quad w = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}$$

Intersection:

$$x^2 = (u_1 + \xi v_1 + \eta w_1)^2 + y^2 = (u_2 + \xi v_2 + \eta w_2)^2 = z^2 = (u_3 + \xi v_3 + \eta w_3)^2$$

\Rightarrow quadratic equation.

$$a\xi^2 + 2b\xi\eta + c\eta^2 + d\xi + e\eta = f$$

$$\begin{pmatrix} \xi & \eta \end{pmatrix} \begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix} + \begin{pmatrix} d & e \end{pmatrix} \begin{pmatrix} \xi \\ \eta \end{pmatrix} = f$$

Which curve?

Step 1: Move the center to the origin Hence, apply translation such that $d = e =$

0. Let $\begin{pmatrix} x_0 \\ y_0 \end{pmatrix}$ be the center. By translation, let $\xi = x_0 + x$ and $\eta = y_0 + y$. Choose x_0, y_0 such that d and e are different.

$$a(x + x_0)^2 + 2b(x + x_0)(y + y_0) + c(y + y_0)^2 + d(x + x_0) + e(y + y_0) + f = 0$$

$$ax^2 + 2ax_0x + ax_0^2 + 2bxy + 2by_0x + 2bx_0y + 2bx_0y_0 + xy^2 + 2cy_0y + cy_0^2 + dx + dx_0 + ey + ey_0 - f = 0$$

$$ax^2 + 2bxy + cy^2 + (2ax_0 + 2by_0 + d)x + (2bx_0 + 2cy_0 + e)y + ax_0^2 + 2bx_0y_0 + cy_0^2 + dx_0 + ey_0 - f = 0$$

x_0, y_0 such that

$$2ax_0 + 2by_0 = -d \quad 2bx_0 + 2cy_0 = -e$$

$$\begin{bmatrix} a & b \\ b & c \end{bmatrix} \begin{pmatrix} x_0 \\ y_0 \end{pmatrix} = \begin{pmatrix} -\frac{d}{2} \\ -\frac{e}{2} \end{pmatrix}$$

is solvable?

Case 1: determinant is zero

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = 0$$

Existence of such a solution is not guaranteed. So, 0 is an eigenvalue. There exists an orthogonal matrix Q (rotation/reflection).

$$Q^t \begin{pmatrix} a & b \\ b & c \end{pmatrix} Q = \begin{bmatrix} \lambda_1 & \\ & 0 \end{bmatrix}$$

Rotation of planes:

$$\begin{pmatrix} x \\ y \end{pmatrix} = Q^t \begin{pmatrix} \xi \\ \eta \end{pmatrix} \quad \Rightarrow \quad \begin{pmatrix} \xi \\ \eta \end{pmatrix} = Q \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\begin{pmatrix} \tilde{d} \\ \tilde{e} \end{pmatrix} = Q^t \cdot \begin{pmatrix} d \\ e \end{pmatrix}$$

$$\begin{pmatrix} x & y \end{pmatrix} Q^t A Q \begin{pmatrix} x \\ y \end{pmatrix} + \underbrace{\begin{pmatrix} d & e \end{pmatrix} Q}_{\begin{pmatrix} \tilde{d} \\ \tilde{e} \end{pmatrix}} \begin{pmatrix} x \\ y \end{pmatrix} = f$$

$$\lambda_1 x^2 + \tilde{d}x + \tilde{e}y = f$$

If $\tilde{e} = 0$:

$$\lambda_1 x^2 + \tilde{d}x = f \text{ and } y \text{ arbitrary}$$

$$x = \frac{-\tilde{d} \pm \sqrt{\tilde{d}^2 + 4\lambda_1 f}}{2\lambda_1}$$

if $\tilde{d}^2 + 4\lambda_1 f < 0$, no solution. 1 or 2 lines.

If $\tilde{e} \neq 0$:

$$y = \frac{f - \tilde{d}x - \lambda_1 x^2}{\tilde{e}}$$

is a parabola.

Case 2: determinant is non-zero

$$\begin{vmatrix} a & b \\ b & c \end{vmatrix} \neq 0$$

\leadsto equation by translation.

$$ax^2 + 2bxy + cy^2 = g$$

$$\begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} a & b \\ b & c \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = g$$

$\leadsto Q$ orthogonal such that

$$Q^T \begin{pmatrix} a & b \\ b & c \end{pmatrix} Q = \begin{bmatrix} \lambda_1 & \\ & \lambda_2 \end{bmatrix} \text{ with } \lambda_1, \lambda_2 \neq 0$$

$$\begin{pmatrix} \tilde{x} \\ \tilde{y} \end{pmatrix} = Q^t \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = Q \begin{pmatrix} \tilde{x} \\ \tilde{y} \end{pmatrix}$$

$$\begin{pmatrix} \tilde{x} & \tilde{y} \end{pmatrix} Q^t \begin{pmatrix} a & b \\ b & c \end{pmatrix} Q \begin{pmatrix} \tilde{x} \\ \tilde{y} \end{pmatrix} = g$$

$$\lambda_1 \tilde{x}^2 + \lambda_2 \tilde{y}^2 = g$$

Case 2a: $g = 0$

$$\tilde{y}^2 = -\frac{\lambda_1}{\lambda_2}\tilde{x}^2$$

if $\text{sign } \lambda_1 = \text{sign } \lambda_2 \implies \tilde{x} = \tilde{y} = 0$. If $\text{sign } \lambda_1 \neq \text{sign } \lambda_2 \leadsto \tilde{y} = \pm \frac{\lambda_1}{\lambda_2}\tilde{x}$.

$$\frac{\lambda_1}{g}\tilde{x}^2 + \frac{\lambda_2}{g}\tilde{y}^2 = 1$$

Classification: $A > 0$ or $A < 0$ gives an ellipsis, point or the empty set. If $A \geq 0$ or $A \leq 0$, a parabola or line or the empty set is given. If A is indefinite, a hyperbola is given.

Example 12.5. Analogously: Quadric in \mathbb{R}^3 . A three-dimensional conic section.

Theorem 12.5. Let $A \geq 0$. Then $\exists! B \geq 0 : B^2 = A$.

$$B := A^{\frac{1}{2}}$$

Proof. We prove the existence of B .

A is self-adjoint, so there exists a unitary matrix U such that $U^*AU = \text{diag}(\lambda_1, \dots, \lambda_n)$ with all $\lambda_i \geq 0$.

$$A = U \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} U^*$$

$$B = U \begin{bmatrix} +\sqrt{\lambda_1} & & \\ & \ddots & \\ & & +\sqrt{\lambda_n} \end{bmatrix} U^*$$

$M \geq 0 \implies C^*MC' \geq 0$. B satisfies the condition $B^2 = A$ with $B \geq 0$.

Conditionless $B \geq 0$: $\leadsto 2^{\text{rank}(A)}$ different solutions.

We prove uniqueness of B .

Let $B \geq 0$ and $B^2 = a$. Let u_1, \dots, u_n be orthonormal basis of eigenvectors for eigenvalues μ_1, \dots, μ_n of B with $\lambda_i \geq 0$.

$$\implies Bu_i = \mu_i u_i \quad \mu_i \geq 0$$

$$Au_i = B^2u_i = \mu_i^2u_i \implies \mu_i^2 = \lambda_i \implies \mu_i = +\sqrt{\lambda_i}$$

$$\implies U^*BU = \begin{bmatrix} \mu_1 & & \\ & \ddots & \\ & & \mu_n \end{bmatrix} \wedge U^*AU = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}$$

□

Remark 12.6 (Another solution). Find B such that $B^*B = A$. There are infinitely many solutions.

12.3 Cholesky decomposition

Remark 12.7. *André-Louis Cholesky (1875–1918) descending from the Cholewsky (Polish) family*

Theorem 12.6 (Cholesky decomposition). *Let $A > 0 \iff \exists C \in \mathbb{C}^{n \times n}$ (lower triangular matrix) such that $A = C \cdot C^*$. Compare with LU decomposition.*

Remark 12.8 (Algorithm). *Determine the matrix C or abort if A is not positive definite.*

Remark 12.9 (Schur complement).

$$M = \begin{bmatrix} A & B \\ C & D \end{bmatrix}$$

$$\implies M/D := A - BD^{-1}C$$

is the Schur complement of block D in M where M is a $(p+q) \times (p+q)$ matrix and M/D is a $p \times p$ matrix.

Lemma 12.3. *Let $A \in \mathbb{C}^{n \times n}$. $A > 0$, $b \in \mathbb{C}^n$, $\gamma > 0$.*

$$\det \left[\begin{array}{c|c} A & B \\ \hline b^* & \gamma \end{array} \right] = \det A \cdot (\gamma - b^* A^{-1} b)$$

$$\begin{vmatrix} A & b \\ b^* & \gamma \end{vmatrix} = \begin{vmatrix} I & 0 \\ -b^* A^{-1} & 1 \end{vmatrix} \begin{vmatrix} A & b \\ b^* & \gamma \end{vmatrix} = \begin{vmatrix} A & b \\ \underbrace{-b^* A^{-1} A + b^*}_{=0} & -b^* A^{-1} b + \gamma \end{vmatrix} = \det A \cdot (-b^* A^{-1} b + \gamma)$$

Proof by Cholesky. By complete induction:

Case $n = 1$

$$[a_{11}] > 0$$

$$\implies a_{11} > 0$$

$$C = [e^{i\theta} \sqrt{a_{11}}] \text{ is unique}$$

$$C^* = [e^{-i\theta} \sqrt{a_{11}}]$$

Case $k \rightarrow k+1$

$$A_{k+1} = \begin{bmatrix} A_k & b \\ b^* & \gamma \end{bmatrix} \quad A_k \in \mathbb{C}^{n \times n}, A_k > 0$$

By induction hypothesis: $\exists C_k : A_k = C_k C_k^*$.

$$\text{Find } C_{k+1} = \begin{pmatrix} & 0 \\ C_k & \vdots \\ & 0 \\ C^* & \alpha \end{pmatrix} \text{ such that } C_{k+1} C_{k+1}^* = A_{k+1}, \alpha > 0$$

$$C_{k+1} C_{k+1}^* = \begin{bmatrix} & 0 \\ C_k & \vdots \\ & 0 \\ C^* & \alpha \end{bmatrix} \begin{bmatrix} & c \\ C_k^* & \\ 0 & \dots & 0 & \alpha \end{bmatrix} = \left[\begin{array}{c|c} C_k C_k^* & C_k \cdot c \\ \hline C^* C_k^* & C^* C + \alpha^2 \end{array} \right] \stackrel{!}{=} \begin{bmatrix} A_k & b \\ b^* & \gamma \end{bmatrix}$$

Requirement: $C_k C \dot{C} = b$. $c^* c + \alpha^2 = \gamma$. Choose $c = C_k^{-1} b$.

$$\begin{aligned} \alpha^2 &= \gamma - c^* c &= \gamma - b^* C_k^{*-1} C_k^{-1} b \\ &= \gamma - b^* A^{-1} b \\ &= \frac{\det A_{k+1}}{\det A_k} \\ &> 0 \end{aligned}$$

$$\begin{aligned} A_k &= C_k C_k^* & A_k^{-1} &= C_k^{*-1} C_k^{-1} \\ \implies \alpha &= \sqrt{\gamma - b^* A^{-1} b} = \sqrt{\frac{\det A_{k+1}}{\det A_k}} \end{aligned}$$

□

Remark 12.10 (Application in practice). Find C such that $C \cdot C^* = A$. $c_{ij} = 0$ if $j > i$. $c_{ii} > 0$.

$$\begin{aligned} a_{ij} &= \sum_{k=1}^n c_{ik} (c^*)_{kj} \\ &= \sum_{k=1}^n c_{ik} \overline{c_{jk}} \\ &= \sum_{k=1}^{\min(i,j)} c_{ik} \overline{c_{jk}} \end{aligned}$$

The algorithm fills up columns from top to bottom and the columns are fill from left to right.

1st column

$$a_{11} = c_{11} \cdot \overline{c_{11}} = c_{11}^2 \implies c_{11} = \sqrt{a_{11}}$$

$$a_{21} = c_{21} \cdot \overline{c_{11}} = c_{21} \cdot c_{11} \implies c_{21} = \frac{a_{21}}{c_{11}}$$

$$a_{31} = \sum_{k=1}^{\min(3,1)} c_{3k} \overline{c_{jk}} = c_{31} c_{11} \implies c_{31} = \frac{a_{31}}{c_{11}}$$

$$c_{n1} = \frac{a_{n1}}{c_{11}} = \frac{a_{n1}}{\sqrt{a_{11}}}$$

$$c_{11} = \sqrt{a_{11}} \quad c_{21} = \frac{a_{21}}{\sqrt{a_{11}}} \quad \dots \quad c_{n1} = \frac{a_{n1}}{\sqrt{a_{11}}}$$

Induction Assume columns $1, \dots, j-1$ are determined, i.e. c_{ik} is known for $k = 1, \dots, j-1$ and $i = 1, \dots, n$.

$$a_{jj} = \sum_{k=1}^j c_{jk} \overline{c_{jk}} = \sum_{k=1}^{j-1} |c_{jk}|^2 + c_{jj}^2$$

$$\implies c_{jj} = \sqrt{a_{jj} - \sum_{k=1}^{j-1} |c_{jk}|^2}$$

for $i > j$:

$$a_{ij} = \sum_{k=1}^j c_{ik} \overline{c_{jk}} = \sum_{k=1}^{j-1} c_{ik} \overline{c_{jk}} + c_{ij} c_{jj}$$

$$c_{ij} = \frac{a_{ij} - \sum_{k=1}^{j-1} c_{ik} \overline{c_{jk}}}{c_{jj}}$$

where c_{jj} was already determined and the values of the enumerator are already known.

↓ This lecture took place on 2018/06/18.

Cholesky decomposition:

1. Given $A > 0$
2. Find C such that $A = CC^*$. Recursively:

$$c_{ij} = \sqrt{a_{ii} - \sum_{k=1}^{i-1} |c_{ik}|^2}$$

$$c_{ij} = \frac{1}{c_{jj}} (a_{ij} - \sum_{k=1}^{j-1} c_{ik} \overline{c_{jk}})$$

Example 12.6.

$$A = \begin{bmatrix} 4 & 12 & -16 \\ 12 & 37 & -43 \\ -16 & -43 & 98 \end{bmatrix} = \begin{bmatrix} c_{11} & c_{21} & c_{31} \\ c_{21} & c_{22} & c_{32} \\ c_{31} & c_{32} & c_{33} \end{bmatrix} \cdot \begin{bmatrix} c_{11} & \overline{c_{21}} & \overline{c_{31}} \\ 0 & c_{22} & \overline{c_{32}} \\ 0 & 0 & c_{33} \end{bmatrix}$$

$$= \begin{bmatrix} c_{11}^2 & c_{11}\overline{c_{21}} & c_{11}\overline{c_{31}} \\ c_{21}c_{11} & |c_{21}|^2 + c_{22}^2 & c_{31}\overline{c_{21}} + c_{32}c_{22} \\ c_{31}c_{11} & c_{31}\overline{c_{21}} + c_{32}c_{22} & |c_{31}|^2 + |c_{32}|^2 + c_{33}^2 \end{bmatrix}$$

with

$$\begin{aligned} c_{11} &= \sqrt{4} = 2 \\ 12 &= c_{11} \cdot \overline{c_{21}} \implies c_{21} = \frac{12}{c_{11}} = 6 \\ -16 &= c_{11}\overline{c_{31}} \implies c_{31} = -\frac{16}{c_{11}} = -8 \\ |c_{21}|^2 &= 37 \\ c_{22} &= \sqrt{37 - 6^2} = 1 \\ c_{31}\overline{c_{21}} + c_{32}c_{22} &= -43 \\ c_{32} &= \frac{-43 - c_{31}\overline{c_{21}}}{c_{22}} = \frac{-43 + 8 \cdot 6}{1} = 5 \\ |c_{31}|^2 + |c_{32}|^2 + c_{33}^2 &= 98 \implies c_{33} = \sqrt{98 - (-8)^2 - 5^2} = 3 \end{aligned}$$

If some root of a negative number needs to be taken, the matrix was not positive definite.

Remark 12.11. If $A \in \mathbb{R}^{n \times n}$, then

1. also $C \in \mathbb{R}^{n \times n}$.
2. C is uniquely determined.
3. Cholesky decomposition is an LU decomposition

Corollary 12.4. If $A > 0$, then $\det A \leq a_{11}a_{22} \dots a_{nn}$.

Proof. $A = CC^*$ is a Cholesky decomposition.

$$c_{ii} = \sqrt{a_{ii} - \sum_{k=1}^{i-1} |c_{ik}|^2} \leq \sqrt{a_{ii}}$$

□

Remark 12.12. We apply the product law.

$$\begin{aligned}
 \det(A) &= \det(CC^*) \\
 &= \det(C) \cdot \det(C^*) \\
 &= |\det(C)|^2 \\
 &= \left| \prod_{i=1}^n c_{ii} \right|^2 = \prod c_{ii}^2 \leq \prod a_{ii}
 \end{aligned}$$

Remark 12.13. J. S. Hadamard (1865–1963)

Corollary 12.5 (Hadamard's inequality). Let $A \in \mathbb{C}^{n \times n}$ with column vectors a_1, \dots, a_n .

$$|\det A| \leq \prod_{i=1}^n \|a_i\|$$

$$\|a_i\| = \sqrt{\langle a_i, a_i \rangle} = \sqrt{a_i^* \cdot a_i} = \text{Euclidean norm}$$

Proof. **Case 1: A is singular** trivial ($\det(A) = 0$)

Case 2: A is regular

$$\implies B = A^*A \text{ is regular} \implies \text{positive definite}$$

$$\det B \leq \prod_{i=1}^n b_{ii} \qquad \qquad \qquad = \det(A^*A) = |\det A|^2$$

$$b_{ii} = \text{i-th row of } A^* \times \text{i-th column of } A$$

$$= a_i^* \cdot a_i = \|a_i\|^2 \implies |\det A|^2 \leq \prod_{i=1}^n \|a_i\|^2$$

□

In this chapter: $AA^* = A^*A$.

There exists an orthonormal basis of eigenvectors:

$$\begin{aligned}
Ax_i &= \lambda_i x_i \\
\langle x_i, x_j \rangle &= \delta_{ij} \\
x \in \mathbb{C}^n \rightsquigarrow x &= \sum \alpha_i x_i \quad \alpha_i = \langle x, x_i \rangle \\
Ax &= \sum \alpha_i Ax_i = \sum \alpha_i \lambda_i x_i \\
&= \sum \lambda_i \langle x, x_i \rangle x_i \\
\implies A &= \sum_{i=1}^n \lambda_i \langle \cdot, x_i \rangle x_i
\end{aligned}$$

where \cdot is a placeholder. It represents the map $\langle \cdot, x_i \rangle : \mathbb{C}^n \rightarrow \mathbb{C}$ with $x \mapsto \langle x, x_i \rangle$.

Remark 12.14. Let $z \in \mathbb{C}$.

$$\begin{aligned}
z &= r \cdot e^{i\theta} \\
r &= |z| = \sqrt{z\bar{z}} \\
e^{i\theta} &= \frac{z}{|z|}
\end{aligned}$$

Theorem 12.7 (Polar decomposition). Let $A \in \mathbb{C}^{n \times n}$.

$$|A| := (A^*A)^{\frac{1}{2}} \quad (\text{unique, positive semidefinite root})$$

Then $\exists U \in \mathcal{U}(n)$ such that $A = U \cdot |A|$.

Proof. **Case 1: A is regular** AA^* is positive definite.

$$\begin{aligned}
\rightsquigarrow A^*A &= V \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} V^* \quad \lambda_i > 0 \\
|A| &:= (A^*A)^{\frac{1}{2}} = V \begin{bmatrix} \sqrt{\lambda_1} & & \\ & \ddots & \\ & & \sqrt{\lambda_n} \end{bmatrix} V^* \\
U &= A \cdot |A|^{-1} \text{ is unitary} \\
U &= A \cdot (A^*A)^{-\frac{1}{2}} \\
U^*U &= (A(A^*A)^{-\frac{1}{2}})^*(A(A^*A)^{-\frac{1}{2}}) = ((A^*A)^{-\frac{1}{2}} \cdot A^*)(A(A^*A)^{-\frac{1}{2}}) = (A^*A)^{-\frac{1}{2}}(A^*A)(A^*A)^{-\frac{1}{2}} \\
&= V \begin{bmatrix} \lambda_1^{-\frac{1}{2}} & & \\ & \ddots & \\ & & \lambda_n^{-\frac{1}{2}} \end{bmatrix} V^* V \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} V^* V \begin{bmatrix} \lambda_1^{-\frac{1}{2}} & & \\ & \ddots & \\ & & \lambda_n^{-\frac{1}{2}} \end{bmatrix} V^* = V \begin{bmatrix} 1 & & \\ & \ddots & \\ & & 1 \end{bmatrix} V^* = I \\
U \cdot |A| &= A \cdot |A|^{-1} \cdot |A| = A
\end{aligned}$$

Case 2: A is singular Let $A^*A \geq 0$ and some $\lambda_i = 0$. By change of basis, $A = V \text{diag}(\lambda_1, \dots, \lambda_k, 0, \dots, 0)V^*$.

$$|A| = V \text{diag}(\lambda_1^{\frac{1}{2}}, \dots, \lambda_k^{\frac{1}{2}}, 0, \dots, 0)V^*$$

$$|A| = \begin{bmatrix} P & \\ & 0 \end{bmatrix} \quad U = V \begin{bmatrix} P^{-1} & \\ & 0 \end{bmatrix} \simeq \begin{bmatrix} \tilde{U} & \\ & I \end{bmatrix}$$

$$\Rightarrow A = U|A|$$

□

12.4 Singular value decomposition

Remark 12.15 (Singular value decomposition).

$$A \in \mathbb{C}^{n \times n} \quad A = U(A^*A)^{\frac{1}{2}} \quad U \text{ unitary}$$

$(A^*A)^{\frac{1}{2}} \geq 0$ with eigenvalue $s_i \geq 0$ called singular values of A

$$(A^*A)^{\frac{1}{2}} = \sum s_i \langle \cdot, x_i \rangle x_i \text{ where } x_i \text{ is an orthonormal basis}$$

$$A = U \cdot (A^*A)^{\frac{1}{2}}$$

$$= \sum_{i=1}^n s_i \langle \cdot, x_i \rangle \overbrace{U x_i}^{=: y_i}$$

$$= \sum_{i=1}^n s_i \langle \cdot, x_i \rangle y_i$$

$y_i = U x_i$ is also an orthonormal basis

$A \cdot y_i = s_i \cdot x_i \rightarrow$ numerically stable

$$A^{-1} = \sum s_i^{-1} \langle \cdot, y_i \rangle x_i$$

Remark 12.16. Singular value decomposition is numerically stable and therefore very desirable.

It furthermore has a very important application in medicine: CT scans. Inside the CT tube, X-rays are sent from all directions to all directions. You can only determine how much the

$$\int_{\gamma} f(x(t), y(t)) dt = Rf(\psi, \Theta)$$

Is linear: $R(f_1 + f_2) = Rf_1 + Rf_2$. Radon transformation. $\lambda_i \rightarrow 0$.

You need to invert the integral. The SVD is the only numerically stable method to achieve it. Other methods will trigger numerical errors that will amplify and therefore give wrong images.

13 Eigenvalue estimates

Definition 13.1. Let $A \in \mathbb{C}^{n \times n}$.

$$W(A) = \{ \langle Ax, x \rangle \mid \|x\| = 1 \} \subseteq \mathbb{C}$$

is called numerical range of A .

$$w(A) = \sup \{ |z| \mid z \in W(A) \}$$

is called numerical radius of A .

Lemma 13.1.

$$\text{spec}(A) \subseteq W(A)$$

Proof. $\lambda \in \text{spec}(A)$, eigenvector x such that $Ax = \lambda x$, $\|x\| = 1$.

$$\implies \langle Ax, x \rangle = \langle \lambda x, x \rangle = \lambda \in W(A)$$

□

Theorem 13.1 (Theorem by Toeplitz-Hausdorff). $W(A)$ is convex.

Remark 13.1. The following implications are left as an exercise to the reader: A is

$$\text{normal} \implies W(A) = \underbrace{\text{convex spec}(A)}_{\text{convex hull}} = \left\{ \sum_{i=1}^n \alpha_i \underbrace{\lambda_i}_{\text{eigenvalue}} \mid 0 \leq \alpha_i \leq 1, \sum \alpha_i = 1 \right\} = \text{convex set that contains spec}(A).$$

Remark 13.2. J. W. Strutt (1842–1919) aka 3. Lord Rayleigh

Remark 13.3. W. Ritz (1878–1909), discovered the element Argon (\rightarrow Nobel prize)

Theorem 13.2 (Rayleigh–Ritz Theorem). Let $A \in \mathbb{C}^{n \times n}$ be self-adjoint $\implies W(A)$. $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$. Then

$$\lambda_1 = \min_{x \neq 0} \underbrace{\frac{\langle Ax, x \rangle}{\langle x, x \rangle}}_{\text{Rayleigh quotient}} = \min \{ \langle Ax, x \rangle \mid \|x\| = 1 \} = \min W(A)$$

\vdots

$$\lambda_n = \max_{x \neq 0} \frac{\langle Ax, x \rangle}{\langle x, x \rangle} = \max \{ \langle Ax, x \rangle \mid \|x\| = 1 \} = \max W(A)$$

where

$$A = A^* \implies \langle Ax, x \rangle = \langle x, Ax \rangle = \overline{Ax, x}$$

Proof. $\lambda_1, \lambda_n \in W(A)$ because $\text{spec}(A) \subseteq W(A)$.

$$\implies \min W(A) \leq \lambda_1 \quad \max W(A) \geq \lambda_n$$

Show that $\forall x : \lambda_1 \leq \frac{\langle Ax, x \rangle}{\langle x, x \rangle} \leq \lambda_n$.

Let u_1, \dots, u_n be an orthonormal basis of eigenvectors. Let $x \in \mathbb{C}^n$, $\|x\| = 1$.
 $x = \sum_{i=1}^n \alpha_i x_i \implies \|x\|^2 = \sum_{i=1}^n |\alpha_i|^2 = 1$.

$$\begin{aligned} \langle Ax, x \rangle &= \left\langle A \sum \alpha_i x_i, \sum \alpha_j x_j \right\rangle \\ &= \left\langle \sum \alpha_i \lambda_i x_i, \sum \alpha_j x_j \right\rangle \\ &= \sum_i \sum_j \lambda_i \alpha_i \bar{\alpha}_j \langle x_i, x_j \rangle \\ &= \sum_i \lambda_i |\alpha_i|^2 \end{aligned} \quad \begin{aligned} &\leq \max(\lambda_i) \sum |\alpha_i|^2 = \max \lambda_i \\ &\geq \min(\lambda_i) \sum |\alpha_i|^2 = \min \lambda_i \end{aligned}$$

□

Remark 13.4. How can we obtain the other eigenvalues?

For $x \in \mathcal{L}(u_2, \dots, u_n)$, $\langle Ax, x \rangle \geq \lambda_2 \|x\|^2$.

For $x \in \mathcal{L}(u_1, \dots, u_{n-1}) = \{x \mid \langle x, u_n \rangle = 0\} = \{u_n\}^\perp$, $\langle Ax, x \rangle \leq \lambda_{n-1}$.

Remark 13.5. Richard Courant (1888–1972)

E. Fischer (1875–1954)

H. Weyl (1885–1955)

All of them worked in Göttingen, Germany.

Theorem 13.3 (Courant–Fischer–Weyl min–max principle). Let $A \in \mathbb{C}^{n \times n}$ be self-adjoint.

$$\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$$

Then it holds that

1. $\lambda_k = \max_{\substack{W \subseteq V \\ \dim W = k-1}} \min_{x \in W^\perp \setminus \{0\}} \frac{\langle Ax, x \rangle}{\langle x, x \rangle}$. Special case $k = 1$: $W = \{0\} \rightarrow \lambda_1 = \min_x \frac{\langle Ax, x \rangle}{\langle x, x \rangle}$.
2. $\lambda_{n+1-k} = \min_{\substack{W \subseteq V \\ \dim W = k-1}} \max_{x \in W^\perp \setminus \{0\}} \frac{\langle Ax, x \rangle}{\langle x, x \rangle}$. Special case $k = 1$: $\lambda_n = \max_x \frac{\langle Ax, x \rangle}{\langle x, x \rangle}$.

This theorem is more generic than the Rayleigh–Ritz Theorem.

Proof. For $W \subseteq V$. Let $m_A(W) = \min \left\{ \frac{\langle Ax, x \rangle}{\langle x, x \rangle} \mid x \in W^\perp \setminus \{0\} \right\}$. For vectors $m_A(w_1, \dots, w_k) = m_A(\mathcal{L}(w_1, \dots, w_k))$. For some orthonormal basis u_1, \dots, u_n of eigenvectors,

$$\lambda_k = m_A(u_1, \dots, u_{k-1}) \stackrel{\text{see proof of Theorem 13.2}}{=} \min \left\{ \frac{\langle Ax, x \rangle}{\langle x, x \rangle} \mid x \in \underbrace{\mathcal{L}(u_k, \dots, u_n)}_{\{u_1, \dots, u_{k-1}\}^\perp} \right\}$$

$$\implies \lambda_k \leq \max_{\substack{W \subseteq V \\ \dim W = k-1}} m_A(W)$$

Show that: $\forall W \subseteq V$ with $\dim(W) = k-1$, $m_A(w) \leq \lambda_k$.

$$\dim W^\perp = n - k + 1 \implies W^\perp \cap \mathcal{L}(u_1, \dots, u_k) \neq \{0\}$$

$$v = \sum_{i=1}^k \alpha_i u_i \in W^\perp \cap \mathcal{L}(u_1, \dots, u_k) \text{ with } \|v\| = 1$$

$$\langle Av, v \rangle = \sum_{i=1}^k |\alpha_i|^2 \cdot \lambda_i \leq \lambda_k \underbrace{\sum_{i=1}^k |\alpha_i|^2}_{=1} = \lambda_k$$

$$m_A(w) = \min_{x \in W^\perp} \langle Ax, x \rangle \leq \langle Av, v \rangle = \lambda_k$$

□

↓ This lecture took place on 2018/06/20.

The Rayleigh-Ritz coefficient gives us $\lambda_1 = \min_{\langle x, x \rangle = 1} \langle Ax, x \rangle = \min W(A)$ and $\lambda_n = \max_{\langle x, x \rangle = 1} \langle Ax, x \rangle$ with $A = A^*$ and $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$. $x = \sum \alpha_i u_i$ with $\sum |\alpha_i|^2 = 1$ is an orthonormal basis u_1, \dots, u_n of eigenvectors. $\langle Ax, x \rangle = \sum_1^n \lambda_i |\alpha_i|^2$. For $x \in \mathcal{L}(u_2, \dots, u_n) \rightarrow \geq \lambda_2$.

This is used by the Courant–Fischer–Weyl min–max principle.

1. First statement:

$$\lambda_k = \max_{\substack{W \subseteq V \\ \dim W = k-1}} \min_{\substack{x \in W^\perp \\ \langle x, x \rangle = 1}} \langle Ax, x \rangle$$

2. Second statement:

$$\lambda_{n+1-k} = \min_{\substack{W \subseteq V \\ \dim W = k-1}} \max_{\substack{x \in W^\perp \\ \langle x, x \rangle = 1}} \langle Ax, x \rangle$$

Proof of Theorem 13.3 continued. The second statement follows from the first: $-A$ has eigenvalues: $-\lambda_n \leq -\lambda_{n-1} \leq \dots \leq -\lambda_2 \leq -\lambda_1$.

We apply the first statement on $-A$.

$$\begin{aligned} \underbrace{\lambda_k(-A)}_{-\lambda_{n+1-k}} &= \max_{\substack{W \subseteq V \\ \dim W = k-1}} \min_{\substack{x \in W^\perp \\ \langle x, x \rangle = 1}} \langle -Ax, x \rangle \\ &= \max_{\substack{W \subseteq V \\ \dim W = k-1}} \left(- \max_{\substack{x \in W^\perp \\ \langle x, x \rangle = 1}} \langle Ax, x \rangle \right) \\ &= - \min_{\substack{W \subseteq V \\ \dim W = k-1}} \max_{\substack{x \in W^\perp \\ \langle x, x \rangle = 1}} \langle Ax, x \rangle \end{aligned}$$

□

Corollary 13.1 (Nesting Theorem (dt. Schachtelungssatz von Cauchy)). Let $A \in \mathbb{C}^{n \times n}$, $A = A^*$. $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$. Let $B = [a_{ij}]_{i,j=1,\dots,n-1}$. Thus, the last row and column was removed. The dimension is reduced by 1. $B = B^*$. Eigenvalues: $\mu_1 \leq \mu_2 \leq \dots \leq \mu_{n-1}$. Then it holds that $\lambda_1 \leq \mu_1 \leq \lambda_2 \leq \mu_2 \leq \lambda_3 \leq \dots \leq \lambda_{n-1} \leq \mu_{n-1} \leq \lambda_n$. In general: If P is an orthogonal projection on a subspace of dimension $n - 1$.

For example,

$$P = \begin{bmatrix} 1 & & \\ & \ddots & \\ & & 1 & \\ & & & 0 \end{bmatrix}$$

then the eigenvalues (except for eigenvalue 0) of PAP are nested like above.

Proof.

$$A = \begin{bmatrix} [B] & b \\ b^* & \gamma \end{bmatrix}$$

Let $w_1, \dots, w_{n-1} \in \mathbb{C}^{n-1}$ be an orthonormal basis of eigenvectors of B .

$$Bw_i = \mu_i w_i \quad u_i = \begin{pmatrix} w_i \\ 0 \end{pmatrix} \in \mathbb{C}^n \quad u_n = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

$\leadsto u_1, \dots, u_{n-1}, u_n$ is an orthonormal basis of \mathbb{C}^n . Attention! There is no eigenvector of A .

$$W_k = \mathcal{L}(u_1, \dots, u_{k-1}, u_n)$$

$$\text{By Theorem 13.3} \implies \lambda_{k+1} = \max_{\substack{W \subseteq \mathbb{C}^n \\ \dim W = k}} \min_{x \in W^\perp \setminus \{0\}} \frac{\langle Ax, x \rangle}{\langle x, x \rangle} \geq \min_{x \in W_k^\perp \setminus \{0\}} \frac{\langle Ax, x \rangle}{\langle x, x \rangle}$$

$$x \in W_k^\perp \iff \langle x, u_i \rangle = 0 \quad i = 1, \dots, k-1 \wedge \langle x, u_n \rangle = x_n = 0$$

$$\iff x = \begin{bmatrix} y \\ 0 \end{bmatrix}, y \in \{w_1, \dots, w_{k-1}\}^\perp \quad \underbrace{\quad}_{w_1, \dots, w_{n-1} \text{ is ONB of } \mathbb{C}^{n-1}} = \mathcal{L}(w_k, \dots, w_{n-1}) \subseteq \mathbb{C}^{n-1}$$

$$y = \sum_{i=k}^{n-1} \alpha_i w_i$$

$$\frac{\langle Ax, x \rangle}{\langle x, x \rangle} = \frac{\left\langle \begin{bmatrix} B & b \\ b^* & \gamma \end{bmatrix} \begin{bmatrix} y \\ 0 \end{bmatrix}, \begin{bmatrix} y \\ 0 \end{bmatrix} \right\rangle}{\left\langle \begin{bmatrix} y \\ 0 \end{bmatrix}, \begin{bmatrix} y \\ 0 \end{bmatrix} \right\rangle} = \frac{\left\langle \begin{bmatrix} B & y \\ b^* & -y \end{bmatrix} \begin{bmatrix} y \\ 0 \end{bmatrix} \right\rangle}{\langle y, y \rangle} = \frac{\langle By, y \rangle}{\langle y, y \rangle}$$

$$= \frac{\langle B \sum_{i=k}^{n-1} \alpha_i w_i, \sum_{j=k}^{n-1} \alpha_j w_j \rangle}{\sum_{j=1}^{n-1} \text{TODO}} = \frac{\sum_{i=k}^{n-1} \mu_i |\alpha_i|^2}{\sum \text{TODO}} \geq \mu_k$$

Inversion: $-A$ has eigenvalues $-\lambda_n \leq -\lambda_{n-1} \leq \dots \leq -\lambda_2 \leq -\lambda_1$. $-B$ has eigenvalues $-\mu_{n-1} \leq -\mu_{n-2} \leq \dots \leq -\mu_2 \leq -\mu_1$.

We apply step 1 on $-A$ and $-B$:

$$\begin{aligned} \lambda_{k+1}(-A) &\geq \lambda_k(-B) \\ -\lambda_{n-k} &\geq -\mu_{n-k} \\ \implies \lambda_{n-k} &\leq \mu_{n-k} \quad \forall k = 1, \dots, n-1 \\ \implies \lambda_k &\leq \mu_k \quad \forall k = 1, \dots, n-1 \end{aligned}$$

□

Corollary 13.2. A, B are self-adjoint $\in \mathbb{C}^{n \times n}$.

$$\lambda_k(A) + \lambda_1(B) \leq \lambda_k(A + B) \leq \lambda_k(A) + \lambda_n(B)$$

Proof.

$$\lambda_1(B) \leq \frac{\langle Bx, x \rangle}{\langle x, x \rangle} \leq \lambda_n(B)$$

because of Theorem 13.2.

$$\begin{aligned} \lambda_k(A + B) &= \max_{\dim W = k-1} \min_{x \in W^\perp \setminus \{0\}} \frac{\langle (A + B)x, x \rangle}{\langle x, x \rangle} \\ \frac{\langle Ax, x \rangle}{\langle x, x \rangle} + \lambda_n(B) &\leq \frac{\langle (A + B)x, x \rangle}{\langle x, x \rangle} \leq \frac{\langle Ax, x \rangle}{\langle x, x \rangle} + \lambda_n(B) \end{aligned}$$

This goes on and on $\leadsto \lambda_k(A) + \lambda_l(B)$.

$$\leq \lambda_l(A) + \lambda_n(B)$$

□

Corollary 13.3. If $B \geq 0$, $A = A^*$, then $\lambda_k(A) \leq \lambda_k(A + B) \forall k$.

Remark 13.6. Semën Aranovič Geršgorin (1901–1933)

13.1 Geršgorin theorem

Theorem 13.4 (Geršgorin Theorem). Let $A \in \mathbb{C}^{n \times n}$.

$$r_i = \sum_{\substack{j=1 \\ j \neq i}}^n |a_{ij}|$$

$$\begin{bmatrix} |a_{11}| & |a_{12}| & \dots & |a_{1n}| \\ |a_{21}| & |a_{22}| & \dots & |a_{2n}| \\ \vdots & & \ddots & \vdots \\ |a_{n1}| & |a_{n2}| & \dots & |a_{nn}| \end{bmatrix}$$

we remove the diagonal elements. The rows become:

$$\begin{bmatrix} \sum_{j \neq 1} |a_{1j}| \\ \sum_{j \neq 2} |a_{2j}| \\ \vdots \\ \sum_{j \neq n} |a_{nj}| \end{bmatrix}$$

$$\text{spec}(A) \subseteq \bigcup_{k=1}^n \{z \in \mathbb{C} \mid |z - a_{ii}| \leq r_i\}$$

This yields the so-called Gershgorin discs.

Proof. Show that $\forall \lambda \in \text{spec}(A) \exists i$ such that $|\lambda - a_{ii}| \leq r_i$. Let x be an eigenvector: $Ax = \lambda x$. Without loss of generality: $\max_i |x_i| = 1$. Hence $\forall j : |x_j| \leq 1$ and $\exists i : |x_i| = 1$.

$$\underbrace{(Ax)_i}_{\sum_j a_{ij}x_j} = \lambda \cdot x_i \text{ because it is an eigenvector}$$

$$\implies \sum_{\substack{j=1 \\ j \neq i}}^n a_{ij}x_j = \lambda x_i - a_{ii}x_i$$

$$\implies \underbrace{|\lambda x_i - a_{ii}x_i|}_{=|\lambda - a_{ii}||x_i|} = \left| \sum_{j \neq i} a_{ij}x_j \right|$$

$$|\lambda - a_{ii}| \cdot \underbrace{|x_i|}_{=1} \leq \sum_{j \neq i} |a_{ij}| \underbrace{|x_j|}_{\leq 1}$$

$$|\lambda - a_{ii}| \leq \sum_{j \neq i} |a_{ij}| = r_i$$

□

Remark 13.7. For dimension greater 5, it becomes infeasible to determine the characteristic polynomial to retrieve the eigenvalues.

In theory:

1. $\chi_A(x) = \det(x \cdot I - A)$ (difficult if precision is necessary)

2. find roots (cumbersome or infeasible)
3. find eigenvectors (numerically unstable)

In practice, we apply an iterative approach:

Given A

Find x such that $Ax = \lambda x$, $A^2x = \lambda Ax$, $A^3x = \lambda A^2x \leadsto A^\infty x = \lambda A^\infty x$ then $A^\infty x$ is eigenvector. As mathematicians we need to ask ourselves, whether $A^\infty x$ converges? The answer is no, not always. We need to fix x to converge to infinity, not 0.

Choose initial vector x_0 with $\|x_0\| = 1$.

$$w_{k+1} = Ax_k \quad x_{k+1} = \frac{w_{k+1}}{\|w_{k+1}\|} \implies \forall n : \|x_n\| = 1$$

Thus we bounded all the points lie inside the unit sphere. By the Heine-Borel Theorem, a sequence in a bounded and closed (thus compact) space has a convergent subsequence.

Claim. x_n converges assuming $|\lambda_1| > |\lambda_2| \geq \dots \geq |\lambda_n|$ and A diagonalizable.

Let v_1, \dots, v_n be a basis of eigenvectors.

$$x_0 = \alpha_0 v_1 + \dots + \alpha_n v_n$$

$$x_k = \frac{A^k x_0}{\|A^k x_0\|}$$

converges towards v_1 .

$$A^k v_i = \lambda_i^k v_i$$

$$\begin{aligned} A^k x_0 &= \alpha_1 A^k v_1 + \dots + \alpha_n A^k v_n \\ &= \alpha_1 \lambda_1^k v_1 + \alpha_2 \lambda_2^k v_2 + \dots + \alpha_n \lambda_n^k v_n \\ x_k &= \frac{A^k x_0}{\|A^k x_0\|} = \frac{\alpha_1 \lambda_1^k v_1 + \alpha_2 \lambda_2^k v_2 + \dots + \alpha_n \lambda_n^k v_n}{\|\alpha_1 \lambda_1^k v_1 + \alpha_2 \lambda_2^k v_2 + \dots + \alpha_n \lambda_n^k v_n\|} \\ &= \frac{\lambda_1 \cdot \lambda_1^k}{|\alpha_1 \cdot \lambda_1^k|} \cdot \frac{v_1 + \frac{\alpha_2}{\alpha_1} \left(\frac{\lambda_2}{\lambda_1}\right)^k v_2 + \dots + \frac{\alpha_n}{\alpha_1} \left(\frac{\lambda_n}{\lambda_1}\right)^k v_n}{\left\|v_1 + \frac{\alpha_2}{\alpha_1} \left(\frac{\lambda_2}{\lambda_1}\right)^k v_2 + \dots + \frac{\alpha_n}{\alpha_1} \left(\frac{\lambda_n}{\lambda_1}\right)^k v_n\right\|} \\ &\stackrel{k \rightarrow \infty}{\approx} \frac{\alpha_1 \lambda_1^k}{|\lambda_1^k|} \cdot v_1 \end{aligned}$$

$$|\lambda_i| > |\lambda_1| \forall i \geq 2$$

$$\rightarrow \frac{|\lambda_i|}{|\lambda_1|} < 1$$

The smaller $\frac{|\lambda_2|}{|\lambda_1|}$ is, the faster it converges.

$$\left(\frac{|\lambda_i|}{|\lambda_1|}\right)^k \xrightarrow{k \rightarrow \infty} 0$$

$|\lambda_1| - |\lambda_2|$ is called spectral gap.

14 Matrix norms

$$\|x\| = \sqrt{\langle x, x \rangle}$$

Norm on a vector space is a map $\|\cdot\| : V \rightarrow [0, \infty[$:

1. $\|v\| = 0 \iff v = 0$
2. $\|\lambda v\| = |\lambda| \cdot \|v\|$
3. $\|v + w\| \leq \|v\| + \|w\|$

Definition 14.1. Let V, W be vector spaces with norms $\|\cdot\|_V$ and $\|\cdot\|_W$. A norm on $\text{Hom}(V, W)$ is compatible with $\|\cdot\|_V$ and $\|\cdot\|_W$ if

$$\forall v \in V \forall f \in \text{Hom}(V, W) : \|f(v)\|_W \leq \|f\| \cdot \|v\|_V$$

$$\implies \|f(x) - f(y)\|_W = \|f(x - y)\|_W \leq \|f\| \cdot \|x - y\|_V$$

Hence f is Lipschitz continuous with constant $\leq \|f\|$. Specifically, we define the Lipschitz constant uniquely with

$$\inf \left\{ C \mid \|f(v)\|_W \leq C \cdot \|v\|_V \forall v \in V \right\}$$

Example 14.1. In $\mathbb{C}^{n \times n}$. Scalar product $\langle A, B \rangle = \text{Tr}(B^* A)$. Frobenius Hilbert-Schmidt:

$$\|A\|_F = \sqrt{\text{Tr}(A^* A)} = \left(\sum_{ij} |a_{ij}|^2 \right)^{\frac{1}{2}}$$

is compatible with Euclidean norm on \mathbb{C}^n .

Let $x \in \mathbb{C}^n$.

$$\begin{aligned}
\|Ax\|_2^2 &= \sum_{i=1}^n |(Ax)_i|^2 \\
&= \sum_{i=1}^n \left| \sum_j a_{ij} x_j \right|^2 \\
&\stackrel{(\text{CBS inequality, Theorem 8.2})}{\leq} \sum_{i=1}^n \sum_j |a_{ij}|^2 - \sum_k |x_k|^2 \\
&= \|A\|_F^2 - \|x\|_2^2
\end{aligned}$$

Lemma 14.1. Let V, W be normed vectorspaces. $f \in \text{Hom}(V, W)$. Then $\|f\|_{V,W}$ defines a compatible norm and is called induced norm.

$$\|f\|_{V,W} = \inf \left\{ C \mid \|f(v)\|_W \leq C \cdot \|v\|_V \ \forall v \in V \right\}$$

And specifically,

$$\begin{aligned}
\|f\|_{V,W} &= \sup \left\{ \frac{\|f(v)\|_W}{\|v\|} \mid v \in V \setminus \{0\} \right\} \\
&= \sup \left\{ \|f(v)\|_W \mid v \in V, \|v\| = 1 \right\}
\end{aligned}$$

Proof. 1.

$$\sup_{v \neq 0} \frac{\|f(v)\|_W}{\|v\|_V} = \sup_{\|v\|_V=1} \|f(v)\|_W \text{ is immediate}$$

2. Let $M_f = \inf \left\{ C \mid \|f(v)\|_W \leq \|v\|_V \ \forall v \in V \right\}$. Show that $M_f = \sup_{v \neq 0} \frac{\|f(v)\|_W}{\|v\|_V}$.

Let $C \geq M_f$.

$$\begin{aligned}
&\implies \|f(v)\|_W \leq C \cdot \|v\|_V \ \forall v \in V \\
&\implies \frac{\|f(v)\|_W}{\|v\|_V} \leq C \ \forall v \neq 0 \\
&\implies \sup_{v \neq 0} \frac{\|f(v)\|_W}{\|v\|_V} \leq C
\end{aligned}$$

holds for $\forall C \geq M_f$

$$\implies \sup_{v \neq 0} \frac{\|f(v)\|_W}{\|v\|_V} = M_f$$

□

↓ This lecture took place on 2018/06/25.

Matrix norms:

A norm on $\mathbb{K}^{m \times n}$ (with $\mathbb{K} = \mathbb{C}$ or \mathbb{R}) is *compatible* with norms on \mathbb{K}^n and \mathbb{K}^m if

$$\underbrace{\|Ax\|_m}_{\text{vector norm on } \mathbb{K}^m} \leq \underbrace{\|A\|_{m \times n}}_{\text{matrix norm}} \cdot \underbrace{\|x\|_n}_{\text{vector space on } \mathbb{K}^n}$$

$$\|A\|_F = \text{Tr}(A^*A)^{\frac{1}{2}} = \left(\sum |a_{ij}|^2 \right)^{\frac{1}{2}}$$

is compatible with the Euclidean norm.

Optimal norm on $\mathbb{K}^{m \times n}$.

$$\|A\| = \inf \{ C > 0 \mid \forall x \in \mathbb{K}^n : \|A \cdot x\|_m \leq C \cdot \|x\|_n \} := \sup_{\|x\|_n \leq 1} \|Ax\|_m$$

$$f : V \rightarrow W \quad \|f\| = \sup_{\substack{x \in V \\ \|x\|_V \leq 1}} \|f(x)\|_W$$

Exercise for the practicals:

$$\underbrace{V}_{\|\cdot\|_V} \xrightarrow{f} \underbrace{W}_{\|\cdot\|_W} \xrightarrow{g} \underbrace{Z}_{\|\cdot\|}$$

Example 14.2. 1. The Frobenius norm $\|A\|_F = \text{Tr}(A^*A)^{\frac{1}{2}}$ is not optimal.

$$\text{id} : \mathbb{C}^n \rightarrow \mathbb{C}^n$$

$$\|\text{id}\|_{2 \rightarrow 2} = \sup_{\|x\|_2 \leq 1} \|x\|_2 = 1$$

$$\text{id}_F = \|I\|_F = \text{Tr}(I^2)^{\frac{1}{2}} = \sqrt{n}$$

2. The norm induced by the Euclidean norm

$$\begin{aligned} \|A\|_{2 \rightarrow 2} &= \sup \{ \|Ax\|_2 \mid \|x\|_2 \leq 1 \} \\ &= \sup_{x \neq 0} \frac{\|Ax\|_2}{\|x\|_2} = \sup_{x \neq 0} \frac{\langle Ax, Ax \rangle^{\frac{1}{2}}}{\langle x, x \rangle^{\frac{1}{2}}} \\ &= \sup_{x \neq 0} \frac{\langle A^*Ax, x \rangle^{\frac{1}{2}}}{\langle x, x \rangle^{\frac{1}{2}}} = \sqrt{\sup_{x \neq 0} \frac{\langle A^*Ax, x \rangle}{\langle x, x \rangle}} \\ &= \sqrt{\text{largest eigenvalue of } A^*A} = \sqrt{\text{largest singular value of } A} \end{aligned}$$

3. $\|A\|_{\infty \rightarrow \infty}$ on \mathbb{K}^n : $\|x\|_{\infty} = \max |x_i|$ and $\|x\|_1 = \sum |x_i|$.

$$\|x\|_p = \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} \xrightarrow{p \rightarrow \infty} \max |x_i| \quad (1 \leq p \leq \infty)$$

This convergence is left to the reader as an exercise.

$$\|A\|_{\infty \rightarrow \infty} = \sup \{ \|Ax\|_{\infty} \mid \|x\|_{\infty} \leq 1 \}$$

$$\begin{aligned} \|Ax\|_{\infty} &= \max_i |(Ax)_i| = \max_i \left| \sum_{j=1}^n a_{ij} x_j \right| \\ &\leq \max_i \sum_{j=1}^n |a_{ij}| \underbrace{|x_j|}_{\leq \max_j |x_j|} \leq \max_j |x_j| \underbrace{\max_i \sum_{j=1}^n |a_{ij}|}_{=\|x\|_{\infty}} \\ &\implies \forall x \in \mathbb{K}^n : \|Ax\|_{\infty} \leq \max_i \sum_j |a_{ij}| \cdot \|x\|_{\infty} \\ &\implies \|A\|_{\infty \rightarrow \infty} \leq \max_i \sum_j |a_{ij}| \end{aligned}$$

Claim. $\|A\|_{\infty \rightarrow \infty} = \max_i \sum_j |a_{ij}|$

Proof. Find vector \tilde{x} such that $\|A\tilde{x}\|_{\infty} = \max_i \sum_j |a_{ij}| \cdot \|\tilde{x}\|_{\infty}$. Choose i_0 such that $\sum_i |a_{ij}| = \max!$.

$$\tilde{x}_j = \begin{cases} \frac{|a_{i_0 j}|}{a_{i_0 j}} & a_{i_0 j} \neq 0 \\ 0 & \text{else} \end{cases}$$

\tilde{x}_j are not all zero, $|\tilde{x}_j| \in \{0, 1\} \forall j$.

$$\begin{aligned} (A \cdot \tilde{x})_{i_0} &= \sum_j a_{i_0 j} \tilde{x}_j = \sum_j a_{i_0 j} a_{i_0 j} \frac{|a_{i_0 j}|}{a_{i_0 j}} = \sum_j |a_{i_0 j}| = \max_i \sum_j |a_{ij}| \\ &\implies \|A\tilde{x}\|_{\infty} \geq |(A\tilde{x})_{i_0}| = \max_i \sum_j |a_{ij}| \cdot \underbrace{\|\tilde{x}\|_{\infty}}_{=1} \\ &\implies \|A\|_{\infty \rightarrow \infty} \geq \max_i \sum_j |a_{ij}| \\ &\implies \|A\|_{\infty \rightarrow \infty} = \max_i \sum_j |a_{ij}| = \max \{ \|z_i\|_1 \mid z_i \text{ row of } A \} \end{aligned}$$

□

4. $\|A\|_{1 \rightarrow 1} = \max_j \sum_i |a_{ij}|$. The proof is left as an exercise to the reader.

$$\begin{aligned} Ax = \lambda x &\implies \|Ax\| = |\lambda| \cdot \|x\| \\ &\implies \sup_{x \neq 0} \frac{\|Ax\|}{\|x\|} \geq |\lambda| \end{aligned}$$

Definition 14.2.

$$\rho(A) := \max \{ |\lambda| \mid \lambda \in \text{spec}(A) \}$$

is called spectral radius of A .

Remark 14.1. Why is it called radius? Consider the complex plane. This value is the radius of the smallest circle with center 0 that contains all eigenvalues.

Lemma 14.2. 1. For every compatible matrix norm: $\|A\| \geq \rho(A)$

2. $\rho(A)$ is not a matrix norm (for some nilpotent matrix, $\rho(A) = 0$ but $A \neq 0$, hence it cannot be a norm)
3. $\forall A \in \mathbb{C}^{n \times n} \forall \varepsilon > 0 \exists$ norm on \mathbb{C}^n : the induced matrix norm satisfies $\|A\| \leq \rho(A) + \varepsilon$

Proof. Proof of point 3.

$$\begin{aligned} \exists \text{ regular } T \in \mathbb{C}^{n \times n} : T^{-1}AT = J = \begin{bmatrix} \lambda_1 & \eta_k & & 0 \\ & \ddots & \eta_{23} & \\ & & \ddots & \ddots \\ 0 & & & \ddots & \eta_{n-1,n} \\ & & & & \lambda_n \end{bmatrix} \quad \eta_{ij} \in \{0, 1\} \\ D_\varepsilon = \text{diag}(1, \varepsilon, \varepsilon^2, \dots, \varepsilon^{n-1}) \\ D_\varepsilon^{-1} J D_\varepsilon = \text{diag}\left(1, \frac{1}{\varepsilon}, \frac{1}{\varepsilon^2}, \dots, \frac{1}{\varepsilon^{n-1}}\right) \begin{bmatrix} \lambda_1 & \eta_{12} & & \\ & \lambda_2 & \eta_{23} & \\ & & \ddots & \\ & & & \lambda_{n-1} & \eta_{n-1,n} \\ & & & & \eta_n \end{bmatrix} \text{diag}(1, \varepsilon, \varepsilon^2, \dots, \varepsilon^{n-1}) \\ = \begin{bmatrix} \lambda_1 & \eta_{12}\varepsilon & & & \\ \frac{\lambda_2}{\varepsilon} & \frac{\eta_{23}}{\varepsilon} & \varepsilon^2 & & \\ & \frac{\varepsilon_3}{\varepsilon^2} & \varepsilon^2 \frac{\eta_{34}}{\varepsilon^2} & \varepsilon^3 & \\ & & & \ddots & \\ & & & & \ddots & \eta_{n-1}\varepsilon \\ & & & & & \lambda_n \end{bmatrix} \end{aligned}$$

Define a norm on \mathbb{C}^n .

$$\|x\|_\varepsilon = \|D_\varepsilon^{-1}T^{-1}x\|_\infty$$

1. $\|x\|_\varepsilon \geq 0$ is immediate.
 $\|x\|_\varepsilon = 0 \implies D_\varepsilon^{-1}T^{-1}x = 0 \implies x = 0$
2. $\|\lambda x\|_\varepsilon = \|D_\varepsilon^{-1}T^{-1}\lambda x\| = |\lambda| \cdot \|x\|_\varepsilon$
3. $\|x + y\|_\varepsilon = \|D_\varepsilon^{-1}T^{-1}(x + y)\|_\infty \leq \|D_\varepsilon^{-1}T^{-1}x\|_\infty + \|D_\varepsilon^{-1}T^{-1}y\|_\infty = \|x\|_\varepsilon + \|y\|_\varepsilon$

$$\|A\|_{\varepsilon \rightarrow \varepsilon} = \sup_{x \neq 0} \frac{\|Ax\|_\varepsilon}{\|x\|_\varepsilon}$$

$$\begin{aligned} x = \underbrace{TD_\varepsilon}_{\text{regular}} y &= \sup_{y \neq 0} \frac{\|ATD_\varepsilon y\|_\varepsilon}{\|TD_\varepsilon y\|_\varepsilon} = \sup_{y \neq 0} \frac{\|D_\varepsilon^{-1}T^{-1}(ATD_\varepsilon y)\|_\infty}{\|D_\varepsilon^{-1}T^{-1}(TD_\varepsilon y)\|_\infty} = \sup_{y \neq 0} \frac{\|J_\varepsilon \cdot y\|_\infty}{\|y\|_\infty} = \sup_{y \neq 0} \frac{\|J_\varepsilon y\|_\infty}{\|y\|_\infty} \\ &\leq \|J_\varepsilon\|_{\infty \rightarrow \infty} = \max_i \sum_j |(J_\varepsilon)_{ij}| \leq \underbrace{\max |\lambda_i| + \varepsilon}_{=\rho(A)+\varepsilon} \end{aligned}$$

□

Remark 14.2. *Israel Gelfand (1913–2009)*

Remark 14.3.

$$\begin{aligned} \rho(A) &\leq \|A\| \\ \rho(A^2) &= \rho(A)^2 \quad \rho(A^3) = \rho(A)^3 \\ \text{spec}(A^2) &= \{\lambda^2 \mid \lambda \in \text{spec}(A)\} \\ \max(|\lambda_i|^2) &= (\max |\lambda_i|)^2 \end{aligned}$$

$$\begin{aligned} \rho(A^2) &\leq \|A\|^2 \quad (\leq \|A\|)^2 \\ \rho(A^3) &\leq \|A\|^3 \quad \underbrace{\rho(A^k)}_{\rho(A)^k} \leq \|A^k\| \\ &\implies \rho(A) \leq \|A^k\|^{\frac{1}{k}} \quad \forall k \end{aligned}$$

Gelfand showed:

$$\lim_{k \rightarrow \infty} \|A^k\|^{\frac{1}{k}} = \rho(A)$$

Lemma 14.3. 1. $\|A\|_{2 \rightarrow 2} = \rho(A^*A)^{\frac{1}{2}}$ is the spectral norm

2. If A is normal, then $\|A\|_{2 \rightarrow 2} = \rho(A)$

Proof. 1. Already shown

$$\begin{aligned} 2. \operatorname{spec}(A^*A) &= \{|\lambda|^2 \mid \lambda \in \operatorname{spec}(A)\} \text{ is left to be proved by the reader. Then} \\ &= \{\bar{\lambda} \cdot \lambda \mid \lambda \in \operatorname{spec}(A)\} \implies \rho(A^*A) = \rho(A)^2 \end{aligned}$$

□

Remark 14.4. Let V be a vector space with dimension ∞ with a norm. For example, $l^2 = \{(x_n) \in \mathbb{R}^\infty \mid \sum_1^\infty |x_n|^2 < \infty\}$ with norm $\|(x_n)\| = \left(\sum_1^\infty |x_n|^2\right)^{\frac{1}{2}}$ is a Hilbert space.

$$l^\infty = \{(x_n) \mid \sup |x_n| < \infty\} \quad \|(x_n)\|_\infty = \sup |x_n|$$

Example 14.3.

$$V = C^\infty[0, 1]$$

$$\frac{d}{dx} : f \mapsto f' \text{ is linear}$$

$\left\|\frac{d}{dx}\right\| = \infty$ it does not matter which norm on $C^\infty[0, 1]$ is defined.

$$\rho\left(\frac{d}{dx}\right) = \infty \quad f' = \lambda f \quad f(x) = e^{\lambda x} \in C^\infty \implies \operatorname{spec}\left(\frac{d}{dx}\right) = \mathbb{C}$$

Remark 14.5. Carl Neumann (1832–1925)

Corollary 14.1. For compatible norms, if $\|A\| < 1$ then $I - A$ is invertible and $(I - A)^{-1} = \frac{1}{I - A} = \sum_{n=0}^\infty A^n$ and $\|(I - A)^{-1}\| \leq \frac{1}{1 - \|A\|}$

$$1. \rho(A) \leq \|A\| < 1 \implies 1 \notin \operatorname{spec}(A) \implies I - A \text{ invertible}$$

$$2. \text{ Neumann series: } \frac{1}{1-x} = \sum_{n=0}^\infty x^n \text{ with } |x| < 1$$

Remark 14.6.

$$y' = \lambda y \quad \int y' = \lambda \cdot \int y \implies \int y = \frac{1}{\lambda} \cdot y$$

Every differentiation can be converted into integration. An integration is bounded (λ becomes $\frac{1}{\lambda}$).

Proof. $\rho(A) < 1 \implies I - A$ invertible.

Claim.

$$\sum_{n=0}^\infty A^n \text{ converges}$$

$$\begin{aligned} \sum_{n=1}^{N+m} A^n - \sum_{n=1}^N A^n &= \left\| \sum_{n=N+1}^{N+m} A^n \right\| \leq \sum_{n=N+1}^{N+m} \|A^n\| \leq \sum_{n=N+1}^{N+m} \|A\|^n \leq \sum_{n=N+1}^{\infty} \|A\|^n \\ &\leq \frac{\|A\|^{N+1}}{1 - \|A\|} \xrightarrow{N \rightarrow \infty} 0 \end{aligned}$$

Thus, the sequence of partial sums is Cauchy and therefore convergent.

$$(I - A) \sum_{n=0}^{\infty} A^n = \sum_{n=0}^{\infty} A^n - \underbrace{\sum_{n=0}^{\infty} A^{n+1}}_{=\sum_{n=1}^{\infty} A^n} = A^0 = I$$

□

$$Ax = \tilde{b} = b + \text{error}$$

$$\tilde{x} - x = A^{-1}(\tilde{b} - b) \implies \|\tilde{x} - x\| \leq \|A^{-1}\| \cdot \|\tilde{b} - b\|$$

$$\|\tilde{x} - x\| \stackrel{?}{\leq} C \cdot \|\tilde{b} - b\|$$

Corollary 14.2. Let A be invertible and B be arbitrary with $\|B\| < \|A^{-1}\|^{-1}$. $\implies A+B$ is invertible and

$$\|(A+B)^{-1}\| \leq \frac{\|A^{-1}\|}{1 - \|A^{-1} \cdot \|B\|\|}$$

$$\|A\| \sim \max |\lambda_i|$$

$$\|A^{-1}\| \sim \max \frac{1}{|\lambda_i|} = \frac{1}{\min |\lambda_i|}$$

$$\|A^{-1}\|^{-1} = \min |\lambda_i|$$

Compare with $A = I$ as special case (this was covered in Corollary 14.1).

$$\|B\| < 1 \quad \|(I+B)^{-1}\| \leq \frac{1}{1 - \|B\|}$$

$$A+B = A \cdot (I + A^{-1}B)$$

$$(A+B)^{-1} = (I + A^{-1}B)^{-1} \cdot A^{-1}$$

$I + A^{-1}B$ is invertible because $\|A^{-1}B\| \leq \|A^{-1}\| \cdot \|B\| < 1$.

$$\implies \|(A+B)^{-1}\| \leq \|(I + A^{-1}B)^{-1}\| \cdot \|A^{-1}\| \leq \frac{1}{1 - \|A^{-1}B\|} \|A^{-1}\| \leq \frac{1}{1 - \|A^{-1}\| \|B\|} \|A^{-1}\|$$

$$x < y \quad 1 - x > 1 - y$$

$$\frac{1}{1 - x} < \frac{1}{1 - y}$$

↓ This lecture took place on 2018/06/30.

$$\|A\| < 1 \implies I - A \text{ invertible}$$

$$(I - A)^{-1} = \sum_{n=0}^{\infty} A^n$$

$$\limsup \|A^n\|^{\frac{1}{n}} = g(A) < 1$$

$\sum a_n$ converges absolutely if $\limsup |a_n|^{\frac{1}{n}} < 1$. So it holds.

A_0 is invertible, $\|A\| < \|A_0^{-1}\|^{-1}$.

$$A_0 + A \text{ invertible}$$

$$\|(A_0 + A)^{-1}\| \leq \frac{\|A_0^{-1}\|}{1 - \|A_0^{-1}\| \cdot \|A\|} < 1$$

Remark 14.7 (Sensitivity of linear equation systems). Instead of $Ax = b$, we consider $\tilde{A}\tilde{x} = \tilde{b}$ with $\|A - \tilde{A}\|$ and $\|b - \tilde{b}\|$ as “small” values. Does this imply $\|x - \tilde{x}\|$ is small?

Let A be a regular matrix and $\|\cdot\|$ is a compatible matrix norm. Let x be the unique solution of $Ax = b$. Let \tilde{x} be the unique solution of $\tilde{A}\tilde{x} = \tilde{b}$.

$$\implies \tilde{b} = b + \underbrace{\Delta b}_{\text{error}}$$

$$\Delta x = \tilde{x} - x = \text{error in the solution}$$

The relative error is interesting. The error relative to the solution is given by:

$$\frac{\|\Delta x\|}{\|x\|} \leq \underbrace{\|\Delta\|}_{\substack{\text{largest} \\ \text{singular value}}} \cdot \underbrace{\|A^{-1}\|}_{\substack{\text{reciprocal of} \\ \text{smallest} \\ \text{singular value}}} \cdot \frac{\|\Delta b\|}{\|b\|}$$

Definition 14.3.

$$\kappa(A) := \|A\| \cdot \|A^{-1}\| \geq 1$$

is called condition number of A . If $\kappa(A)$ is large, then the problem is badly conditioned.

Example 14.4. For diagonal matrices:

$$\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$$

$$\|\Lambda\|_{2 \rightarrow 2} = \max |\lambda_i|$$

$$\|\Lambda^{-1}\|_{2 \rightarrow 2} = \frac{1}{\min |\lambda_i|}$$

$$\implies \kappa(A) = \frac{\max |\lambda_i|}{\min |\lambda_i|}$$

Proof of Remark 14.7.

$$Ax = b \quad A\tilde{x} = \tilde{b}$$

$$A(x + \Delta b) = b + \Delta b$$

$$\iff Ax + A\Delta x = b + \Delta b$$

$$\iff A\Delta x = \Delta b$$

$$\implies \Delta x = A^{-1}\Delta b$$

$$\implies \|\Delta x\| \leq \|A^{-1}\| \cdot \|\Delta b\|$$

$$\|b\| = \|Ax\| \leq \|A\| \cdot \|x\|$$

$$\implies \frac{1}{\|x\|} \leq \frac{\|A\|}{\|b\|}$$

$$\implies \frac{\|\Delta x\|}{\|x\|} = \|A^{-1}\| \cdot \|\Delta b\| \cdot \frac{\|A\|}{\|b\|}$$

□

Remark 14.8 (General case).

$$\tilde{A} = A + \Delta A \quad Ax = b$$

$$\tilde{b} = b + \Delta b \quad \tilde{A}\tilde{x} = \tilde{b}$$

$$\Delta\tilde{x} = \tilde{x} - x$$

Requirement:

$$\|\Delta A\| < \|A^{-1}\|^{-1} \text{ such that } \tilde{A} \text{ invertible}$$

$$\implies \frac{\|\Delta A\|}{\|A\|} \leq \frac{1}{\|A\| \cdot \|A^{-1}\|} = \frac{1}{\kappa(A)}$$

Then this modified system becomes solvable.

All these times, we use the inequality:

$$\|A \cdot B\| \leq \|A\| \cdot \|B\|$$

$$\|A \cdot x\| \leq \|A\| \cdot \|x\|$$

$$(A + \Delta A)(x + \Delta x) = b + \Delta b$$

$$\begin{aligned} &\iff (I + A^{-1}\Delta A)(x + \Delta x) = x + A^{-1}\Delta b \\ \iff x + \Delta x + A^{-1}\Delta Ax + A^{-1}\Delta A\Delta x &= x + A^{-1}\Delta b \\ \iff \Delta x &= A^{-1}\Delta b - A^{-1}\Delta A(x + \Delta x) \end{aligned} \tag{4}$$

$$\|\Delta x\| \leq \underbrace{\|A^{-1}\| \cdot \|\Delta b\|}_{\text{Eq. (5)}} + \underbrace{\|A^{-1}\| \cdot \|\Delta A\| \cdot \|x + \Delta x\|}_{\text{Eq. (6)}}$$

$$\begin{aligned} \|A^{-1}\| \cdot \|\Delta b\| &= \frac{\|A^{-1}\| \cdot \|b\| \cdot \|\Delta b\|}{\|b\|} \\ &\leq \frac{\|A^{-1}\| \cdot \|Ax\| \cdot \|\Delta b\|}{\|b\|} \\ &\leq \|A^{-1}\| \cdot \|A\| \cdot \|x\| \cdot \frac{\|\Delta b\|}{\|b\|} \\ &\leq \kappa(A) \cdot \|x\| \cdot \frac{\|\Delta b\|}{\|b\|} \end{aligned} \tag{5}$$

$$\begin{aligned} \|x + \Delta x\| &\stackrel{(4)}{=} \|(1 + A^{-1}\Delta A)^{-1} \cdot (x + A^{-1}\Delta b)\| \\ &\leq \|(1 + A^{-1}\Delta A)^{-1}\| \cdot (\|x\| + \|A^{-1}\| \cdot \|\Delta b\|) \\ &\leq \frac{1}{1 - \|A^{-1}\Delta A\|} \cdot \|x\| \cdot \left(1 + \|x\| \cdot \|A^{-1}\| \cdot \|b\| \cdot \frac{\|\Delta b\|}{\|b\|}\right) \\ &\leq \frac{1}{1 - \|A^{-1}\| \cdot \|A\| \cdot \frac{\|\Delta A\|}{\|A\|}} \cdot \|x\| \cdot \left(1 + \kappa(A) \cdot \frac{\|\Delta b\|}{\|b\|}\right) \\ &\leq \frac{1}{1 - \kappa(A) \cdot \frac{\|\Delta A\|}{\|A\|}} \cdot \|x\| \cdot \left(1 + \kappa(A) \cdot \frac{\|\Delta b\|}{\|b\|}\right) \\ &\leq \kappa(A) \cdot \frac{\|\Delta b\|}{\|b\|} \cdot \|x\| + \frac{1}{1 - \kappa(A) \cdot \frac{\|\Delta A\|}{\|A\|}} \left(1 + \kappa(A) \cdot \frac{\|\Delta b\|}{\|b\|}\right) \cdot \|x\| \tag{7} \\ \iff \frac{\|\Delta x\|}{\|x\|} &\leq \kappa(A) \cdot \frac{\|\Delta b\|}{\|b\|} + \frac{1 + \kappa(A) \cdot \frac{\|\Delta b\|}{\|b\|}}{1 - \kappa(A) \cdot \frac{\|\Delta A\|}{\|A\|}} \end{aligned}$$

Remark 14.9 (Sensitivity of eigenvalues). *Let $A \in \mathbb{C}^{n \times n}$, diagonalizable.*

$$\rightarrow B^{-1}AB = \text{diag}(\lambda_1, \dots, \lambda_n)$$

$$\tilde{A} = A + \Delta A$$

$$\lambda \in \text{spec}(\tilde{A}) \implies \exists j : |\lambda - \lambda_j| \leq \|B\| \cdot \|B^{-1}\| \cdot \|\Delta A\|$$

($\|\cdot\| = \|\cdot\|_{2 \rightarrow 2}$ or $\|\cdot\|_{\infty \rightarrow \infty}$)

Lemma 14.4 (Special case).

$$A \text{ normal} \implies B \text{ unitary} \implies \text{isometric} \quad \|u_x\| = \|x\| \quad \forall x$$

$$\implies \|B\| = 1, \|B^{-1}\| = 1$$

Proof. 1. If $\lambda \in \text{spec}(A)$, then nothing to show.

2. If $\lambda \notin \text{spec}(A)$, then $A - \lambda I$ is regular, $\tilde{A} - \lambda I$ is non-regular.

$$\begin{aligned} \underbrace{\tilde{A} - \lambda I}_{\text{not invertible}} &= (A - \Delta A) - \lambda I \\ &= (A - \lambda I)(A - \lambda I)^{-1} \cdot (A - \lambda I + \Delta A) \\ &= (A - \lambda I)(I + (A - \lambda I)^{-1} \Delta A) \\ &\implies I + (A - \lambda I)^{-1} \cdot \Delta A \text{ is not invertible} \\ &\xRightarrow{\text{Neumann negated}} \|(A - \lambda I)^{-1} \cdot \Delta A\| \geq 1 \end{aligned}$$

$$\begin{aligned} 1 &\leq \|(A - \lambda I)^{-1} \Delta A\| \\ &= \|(B \Lambda B^{-1} \cdot \lambda B B^{-1})^{-1} \cdot \Delta A\| \\ &= \|(B(\Lambda - \lambda I)B^{-1})^{-1} \cdot \Delta A\| \\ &= \|(B(\Lambda - \lambda I^{-1})B^{-1}) \Delta A\| \\ &\leq \|B\| \cdot \|(\Lambda - \lambda I)^{-1}\| \cdot \|B^{-1}\| \cdot \|\Delta A\| \\ &= \|B\| \cdot \|B^{-1}\| \cdot \frac{1}{\min |\lambda_i - \lambda|} \cdot \|\Delta A\| \\ &\implies \min |\lambda_i - \lambda| \leq \kappa(B) \cdot \|\Delta A\| \end{aligned}$$

Recall that,

$$\begin{aligned}
 (\Lambda - \lambda I)^{-1} &= \begin{bmatrix} \lambda_1 - \lambda & & \\ & \ddots & \\ & & \lambda_n - \lambda \end{bmatrix}^{-1} \\
 &= \begin{bmatrix} \frac{1}{\lambda_1 - \lambda} & & \\ & \ddots & \\ & & \frac{1}{\lambda_n - \lambda} \end{bmatrix} \\
 \|(\Lambda - \lambda I)^{-1}\| &= \max \left| \frac{1}{\lambda_i - \lambda} \right| = \frac{1}{\min |\lambda_i - \lambda|}
 \end{aligned}$$

□

15 Non-negative matrices

Definition 15.1. $A \in \mathbb{K}^{n \times n}$ is called non-negative if $a_{ij} \geq 0 \forall i, j$. We denote $A \geq 0$. Do not mix this up with positive definiteness!

Example 15.1 (Markov chains).

$$a_{ij} = W_s k$$

Manhattan: $a_{ij} = W_s k$ that you can reach node j from node i .

$$a_{ij} \geq 0 \forall i, j$$

For fixed i : $\sum_j a_{ij} = 1$.

Matrix A is called row-stochastic.

A has eigenvector:

$$A \cdot \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{bmatrix}$$

$$\Rightarrow 1 \text{ is eigenvalue, } \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix} =: \Gamma \text{ is eigenvector of } 1$$

A directed graph is called strongly connected if you can reach every node from each other.

Theorem 15.1 (Perron-Frobenius theorem). If the graph is strongly connected, then eigenvalue 1 has geometric and algebraic multiplicity 1, all the other eigenvalues satisfy $|\lambda| < 1$.

$$A_v^n \rightarrow_{n \rightarrow \infty} \mathbf{1} \forall v$$

Index

- Adjoint map, 91
- Adjugate matrix, 34
- Algebra, 104
- Algebra homomorphism, 107
- Alternating group, 15
- Annihilator, 160
- Associative algebra, 104

- Basis of a polynomial, 106
- Bilinearity, 5

- Cauchy product, 105
- Characteristic polynomial, 120
- Chebyshev polynomials, 88
- Cholesky decomposition, 177
- Cofactor, 34
- Commutative algebra, 104
- Commutator, 104
- Compatible norms, 191
- Complementary matrix, 34, 162
- Complex numbers, 103
- Condition number of a matrix, 199
- Congruence of matrices, 53
- Conjugate transpose, 51
- Cross product, 42

- Degree of a polynomial, 106
- Determinant, 16
- Determinant form, 9
- Determinant of a matrix, 4
- Diagonalizable matrix, 123

- Eigen function, 117
- Eigenspace, 116
- Eigenvalue, 115
- Eigenvector, 115
- Euclidean norm, 48
- Euclidean space, 65

- Factorization of a polynomial, 114
- Formal power series algebra, 105

- Generalized eigenvectors, 138

- Geometric multiplicity of an eigenvalue, 129
- Geršgorin disc, 188
- Geršgorin Theorem, 188
- Golden ratio, 127
- Gram matrix of tuple v_1, v_2, \dots, v_m , 79
- Gram-Schmidt process, 85

- Hadamard's inequality, 181
- Hermitian form, 46
- Hermitian matrix, 51
- Hilbert space, 197

- Indefinite inner product, 46
- Index of a nilpotent matrix, 145
- Induced norm, 192
- Inner product, 166
- Invariant subspace, 132
- Involution, 93
- Irreducible polynomial, 114
- Isometry, 96, 98

- Jacobian identity, 104
- Jordan algebra, 104
- Jordan block, 152
- Jordan's normal form, 146

- Left-sided eigenvalue, 118
- Lie algebra, 104
- Lie groups, 104
- Linear dependence, 7
- Linear isometry, 95

- Main space, 138
- Malposition, 12
- Matrix norm, 191
- Method by Cardano/del Ferro, 110
- Minimal polynomial, 160
- Minor of a square matrix, 63
- Multilinearity, 9

- Negative definite inner product, 46
- Nilpotent matrix, 145

- Non-negative matrix, 203
- Norm, 48
- Normal linear map, 166
- Normal matrix, 166
- Numerical range, 184

- Orthogonal complement, 69
- Orthogonal matrix, 96
- Orthonormal, 67
- Orthonormal basis, 67
- Outer product, 42

- Permanent of a square matrix, 26
- Polar decomposition, 182
- Polynomial algebra, 105
- Polynomial division, 111
- Polynomial function, 106
- Positive inner product, 46
- Positive semidefinite inner product, 46

- Quaternions, 103

- Reducible polynomial, 114
- Right-sided eigenvalue, 118
- Root of a polynomial, 109
- Row-stochastic matrix, 203

- Schur complement, 177
- Self-adjoint map, 95
- Self-adjoint matrix, 51
- Signature of π , 12
- Singular value decomposition, 183
- Special linear group, 36
- Spectral gap, 191
- Spectral norm, 196
- Spectral radius, 195
- Spectrum, 115
- Strongly connected graph, 203
- Symmetric group, 10
- Symmetrical minors, 120

- Unitary matrix, 96
- Unitary product, 46
- Unitary space, 65
- Unitary transformation, 95