Supplementary numerical results

Revision: 8 November 2026. Saved evidence through 31 October 2026. This appendix includes all 21 central paired comparisons. It is not a count of every historical fit. The full-precision CSV and evidence map preserve source links. Differences are graph minus the paired comparator; intervals condition on fixed fitted predictions. AP uses positive-outcome participants and macro-Brier uses all participants. No numerical result is regenerated by training in this documentation build.

Earlier graph versus late-fusion sequence; one epoch, seed 20261003

Fold	Graph AP	Comparator AP	AP difference (95% interval)	Macro-Brier difference (95% interval)
outer-0	0.475384	0.394764	+0.080620 [+0.071498, +0.089245]	-0.001953 [-0.002211, -0.001708]
outer-1	0.460125	0.378379	+0.081746 [+0.073130, +0.090150]	-0.002153 [-0.002476, -0.001836]
outer-2	0.453164	0.379705	+0.073460 [+0.064308, +0.082407]	-0.001930 [-0.002230, -0.001638]
outer-3	0.471442	0.360977	+0.110465 [+0.103420, +0.117004]	-0.002562 [-0.002933, -0.002200]
outer-4	0.474226	0.387856	+0.086370 [+0.077628, +0.094814]	-0.001964 [-0.002228, -0.001717]

Fold	Graph macro-Brier	Comparator macro-Brier	AP-defined / all participants
outer-0	0.021945	0.023899	130 / 130
outer-1	0.020059	0.022213	140 / 141
outer-2	0.019544	0.021474	124 / 124
outer-3	0.019757	0.022319	129 / 130
outer-4	0.022488	0.024452	127 / 127

Paired audit sources: outer-0, outer-1, outer-2, outer-3, outer-4.

Earlier graph versus late-fusion sequence; one epoch, seed 20261004

Fold	Graph AP	Comparator AP	AP difference (95% interval)	Macro-Brier difference (95% interval)
outer-0	0.473152	0.394613	+0.078540 [+0.069529, +0.087102]	-0.002178 [-0.002464, -0.001901]
outer-1	0.459892	0.369684	+0.090208 [+0.081627, +0.098271]	-0.002430 [-0.002807, -0.002070]
outer-2	0.443559	0.373283	+0.070275 [+0.061924, +0.078368]	-0.001721 [-0.001970, -0.001469]
outer-3	0.468932	0.362545	+0.106387 [+0.099619, +0.112682]	-0.002504 [-0.002872, -0.002147]

Fold	Graph AP	Comparator AP	AP difference (95% interval)	Macro-Brier difference (95% interval)
outer-4	0.474380	0.386917	+0.087463 [+0.079215, +0.095616]	-0.002267 [-0.002577, -0.001964]

Fold	Graph macro-Brier	Comparator macro-Brier	AP-defined / all participants
outer-0	0.022039	0.024217	130 / 130
outer-1	0.019980	0.022410	140 / 141
outer-2	0.019955	0.021676	124 / 124
outer-3	0.019784	0.022288	129 / 130
outer-4	0.022523	0.024790	127 / 127

Paired audit sources: outer-0, outer-1, outer-2, outer-3, outer-4.

Graph versus time-aligned pooling; two epochs, seed 20261029

Fold	Graph AP	Comparator AP	AP difference (95% interval)	Macro-Brier difference (95% interval)
outer-0	0.498579	0.507520	-0.008941 [-0.009766, -0.008136]	-0.000118 [-0.000177, -0.000060]
outer-1	0.487658	0.493560	-0.005902 [-0.006855, -0.004940]	+0.000174 [+0.000146, +0.000204]
outer-2	0.473438	0.484578	-0.011140 [-0.012849, -0.009501]	+0.000152 [+0.000118, +0.000188]
outer-3	0.498401	0.505002	-0.006601 [-0.007319, -0.005948]	+0.000078 [+0.000039, +0.000113]
outer-4	0.505888	0.511579	-0.005691 [-0.006656, -0.004758]	+0.000111 [+0.000090, +0.000131]

Fold	Graph macro-Brier	Comparator macro-Brier	AP-defined / all participants
outer-0	0.020534	0.020652	130 / 130
outer-1	0.019487	0.019313	140 / 141
outer-2	0.019139	0.018987	124 / 124
outer-3	0.019143	0.019065	129 / 130
outer-4	0.020478	0.020367	127 / 127

Paired audit sources: outer-0, outer-1, outer-2, outer-3, outer-4.

Graph versus time-aligned pooling; two epochs, seed 20261030

Fold	Graph AP	Comparator AP	AP difference (95% interval)	Macro-Brier difference (95% interval)
outer-0	0.498684	0.507538	-0.008854 [-0.010005, -0.007736]	+0.000039 [+0.000010, +0.000068]
outer-1	0.486196	0.497349	-0.011153 [-0.012896, -0.009550]	+0.000186 [+0.000161, +0.000212]
outer-2	0.471794	0.493181	-0.021387 [-0.023610, -0.019330]	+0.000340 [+0.000306, +0.000377]
outer-3	0.494227	0.505545	-0.011318 [-0.012458, -0.010314]	+0.000146 [+0.000093, +0.000195]
outer-4	0.507687	0.516353	-0.008666 [-0.009782, -0.007650]	+0.000158 [+0.000137, +0.000183]

Fold	Graph macro-Brier	Comparator macro-Brier	AP-defined / all participants
outer-0	0.020588	0.020549	130 / 130
outer-1	0.019484	0.019298	140 / 141
outer-2	0.019203	0.018862	124 / 124
outer-3	0.019222	0.019076	129 / 130
outer-4	0.020452	0.020293	127 / 127

Paired audit sources: outer-0, outer-1, outer-2, outer-3, outer-4.

Graph versus pooling; eight-epoch CPU follow-up, seed 20261031

Fold	Graph AP	Comparator AP	AP difference (95% interval)	Macro-Brier difference (95% interval)
outer-0	0.533263	0.533979	-0.000716 [-0.001317, -0.000106]	+0.000014 [+0.000001, +0.000028]

Fold	Graph macro-Brier	Comparator macro-Brier	AP-defined / all participants
outer-0	0.019720	0.019706	130 / 130

Paired audit sources: outer-0.

Two-seed participant summary

Each participant's paired effects were averaged across the two fixed seeds before participant resampling. This differs from averaging fold APs and does not make the seeds independent cohorts. The mean AP difference was -0.0098996662, with interval [-0.01047669424739838, -0.009345412841435134]; macro-Brier difference was 0.0001264322, with interval [0.0001096006568634377, 0.0001428661822493643]. Source summary.

Auxiliary evidence

Historical baseline, event-summary/residual, recovered GRU, routing, endpoint, event-age, event-input, explanation and external-transfer evidence is itemized with source links and scope in the master research record. The sequence comparison plot is a separate supplementary figure. Those diagnostics are not pooled into this 21-pair numerical table.