



Jan 21, 2021

nPCR measurement method

PLOS One

lyzhaojh¹¹, Zhejiang People's Hospital, Yi-Wen Li

1

Works for me

dx.doi.org/10.17504/protocols.io.bp94mr8w



lyzhaojh

EXTERNAL LINK

<https://doi.org/10.1371/journal.pone.0245370>

DOI

[dx.doi.org/10.17504/protocols.io.bp94mr8w](https://doi.org/10.17504/protocols.io.bp94mr8w)

EXTERNAL LINK

<https://doi.org/10.1371/journal.pone.0245370>

DOCUMENT CITATION

lyzhaojh 2021. nPCR measurement method. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bp94mr8w>



EXTERNAL LINK

<https://doi.org/10.1371/journal.pone.0245370>

LICENSE

This is an open access document distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Dec 01, 2020

LAST MODIFIED

Jan 21, 2021

DOCUMENT INTEGER ID

45084

1. Clinically, we will encounter only limited data. For example, we only have data on urea nitrogen before hemodialysis and KT/V for single dialysis. How to calculate the normalized protein catabolic rate (nPCR)?
2. We introduce a simple formula, as followed.

$$nPCR(g/kg/d) = C0/[a + b * KT/V + c/(KT/V)] + 0.168, \text{ where } C0$$

indicates pre-dialysis blood nitrogen in md/dl; a,b,c has different coefficients depending on the time of the dialysis schedule.

3. In our blood purification center, patients are dialyzed three times a week, either on Mondays, Wednesdays, Fridays or either Tuesdays, Thursdays, Saturdays, so that the time between the first dialysis sessions at the beginning of the week is longer.
4. Therefore, the following formula is used: beginning-of-week:

$$nPCR(g/kg/d) = C0/[36.3 + 5.48 * KT/V + 53.5/(KT/V)] + 0.168$$

5. As 1 mmol/l urea nitrogen equal 2.802mg/dl,

$$nPCR(g/kg/d) = 2.802 * BUN/[36.3 + 5.48 * KT/V + 53.5/(KT/V)] + 0.168$$

(urea nitrogen is in mmol/l).

