

Oct 01, 2024

SA-B-Gal Activity

DOI

dx.doi.org/10.17504/protocols.io.3byl49n8ogo5/v1

Karyna Tarasova¹, Sinan Gültekin², iris.gerner Gerner³, Florian Jenner⁴

¹Veterinary Medicine University; ²Vetmeduni Vienna; ³Vetmeduni; ⁴University of Veterinary Medicine Vienna

VETERM



Karyna Tarasova

Veterinary Medicine University

OPEN  ACCESS



DOI: **dx.doi.org/10.17504/protocols.io.3byl49n8ogo5/v1**

Protocol Citation: Karyna Tarasova, Sinan Gültekin, iris.gerner Gerner, Florian Jenner 2024. SA-B-Gal Activity. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.3byl49n8ogo5/v1>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: October 01, 2024

Last Modified: October 01, 2024

Protocol Integer ID: 108717

Abstract

To measure cellular senescence



Senescence associated SA- β -Galactosidase Activity

1

1.1 The cells were washed twice with ice cold PBS and then incubated with 60 μ L of cell lysis buffer provided with the cellular senescence assay kit (Fluorometric format) (Cell Biolabs, USA) for 10 minutes at 4 °C.



1.2 The whole cell lysates were centrifuged 15 minutes at 14000 g at 4 °C. Total protein concentrations were determined by Qubit Protein Assay Kit (ThermoFisher, **Q33211**) using the Qubit 4 Fluorometer (Thermo Fisher Scientific, Singapore, shipped from Germany) according to the manufacturer's instructions.



1.3 Following protein quantification, the assay buffer (Cell Biolabs, Inc., CBA-231, San Diego, USA) was added 1:1 to the cell lysates and incubated for 3h in the dark at 37 °C.



1.4 The reaction was stopped by adding 120 μ L of the Stop solution (provided with the cellular senescence assay kit) per 30 μ L of the reaction mixture.

1.5 Fluorescence was measured using a plate reader at 360 nm (Excitation) / 465 nm (Emission). SA-b-Gal activity was normalized to total protein concentrations.

