



VERSION 1

JUN 05, 2023

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.dm6gpj4n5gzp/v1

External link:
https://en.tiagen.com/upload/file/20220509/20220509164548_88950.pdf

Protocol Citation: TIANGEN Biotech 2023. TIANprep Mini Plasmid Kit Protocol.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.dm6gpj4n5gzp/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working


Created: Jun 04, 2023

Last Modified: Jun 05, 2023

TIANprep Mini Plasmid Kit Protocol V.1

TIANGEN
 Biotech¹

¹TIANGEN Biotech (Beijing) Co., Ltd.



Chottiwatt Jittprasong

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

TIANprep Mini Plasmid Kit is based on alkaline lysis technology followed by adsorption of DNA onto silica membrane in the presence of high salt. Plasmid DNA purified with TIANprep Mini Plasmid Kit is immediately ready for use. Phenol extraction and ethanol precipitation are not required. High quality plasmid DNA is eluted in a small volume of Tris Buffer or deionized water. This protocol is designed for purification of plasmid DNA from 1-5ml overnight cultures of E. coli in LB (Luria-Bertani) medium. Plasmid DNA prepared by TIANprep Mini Plasmid Kit is suitable for a variety of routine applications including restriction enzyme digestion, sequencing, library screening, ligation and trans-formation, in vitro translation, and transfection of robust cells.

GUIDELINES

Applicable to TIANprep Mini Plasmid Kit manufactured by TIANGEN.

PROTOCOL integer ID:
82861

Keywords: plasmid DNA

MATERIALS

RNase A, Buffer BL, Buffer P1, Buffer P2, Buffer P3, Buffer PD, Buffer PW, Buffer EB, Spin Columns CP3, Collection Tubes 2ml

SAFETY WARNINGS



Avoid direct contact with Buffer P2 and P3.


BEFORE START INSTRUCTIONS

Buffer P1 must be activated with RNase A solution before use (one vial per one bottle) and store at 2-8 degrees celsius after activation.

Preparation of Bacterial Cells


16m

1

Centrifugation of  1-5 mL of bacterial cells in a microcentrifuge tube at

1m



 12000 rpm, Room temperature, 00:01:00 , table-top microcentrifuge

2

Direct drainage of supernatant by opening and inverting the tube.

Expected result

Bacteria pellet should be present in the microcentrifuge tube.

3


Complete resuspension of the bacteria pellet in  250 μ L of **Buffer P1**.



Note

- Buffer P1 must be pre-activated with **RNase A**.
- No cell clumps should be visible after resuspension. Vortex and pipette can be used to help homogenize the mixture.

4

Add  250 μ L of **Buffer P2** (lysis buffer) and mix by inverting the tube 6-8 times.

5m






Safety information

Avoid direct contact with Buffer P2

Note

- **DO NOT** vortex or violently mix the reaction.
- **DO NOT** allow the lysis reaction to continue for more than

5

Add  350 μL of **Buffer P3** and mix immediately by inverting the tube 6-8 times to neutralize Buffer P2.




Safety information

Avoid direct contact with Buffer P3

Expected result

There should be no white precipitation left in the supernatant. If there is, the mixture should be centrifuged again.

6

Centrifuge at  12000 rpm, Room temperature, 00:10:00 , table-top microcentrifuge



10m

Preparation of Spin Column CP3

1m

7 Place Spin Column CP3 in a clean collection tube.


8

Add  500 μL of **Buffer BL** to CP3.



9



Centrifuge at  12000 rpm, Room temperature, 00:01:00 , table-top microcentrifuge

1m

10 Discard the flow-through and put the Spin Column CP3 back into the collection tube.

Plasmid Extraction


9m

11 Transfer of supernatant from **Step 6** to Spin Column CP3 with collection tube attached.




12



Centrifuge at  12000 rpm, Room temperature, 00:01:00 , table-top microcentrifuge


1m

13 Discard the flow-through, retain Spin Column CP3 with collection tube attached.

14 **OPTIONAL:** Wash the Spin Column CP3 by adding  500 μ L of **Buffer PD** and centrifuge at

1m



 12000 rpm, Room temperature, 00:01:00 , table-top microcentrifuge and discard the flow-through.

Note

Recommended to remove trace nuclease activity when using **endA+** strains such as the JM series, HB101 and its derivatives, or any wild-type strain, which have high levels of nuclease activity or high carbohydrate content.

15 Wash the column with Buffer PW

15.1 Add 600 µL of Buffer PW



Note

Buffer PW must be pre-treated with 96-100% ethanol.

15.2 Centrifuge at 12000 rpm, Room temperature, 00:01:00 , table-top microcentrifuge



1m

15.3 Discard the flow-through, retain Spin Column CP3 with collection tube attached.

16 Repeat **Step 15**.

17 Centrifuge at 12000 rpm, Room temperature, 00:02:00 , table-top microcentrifuge to remove residual Buffer PW.



2m

18 Air dry the column at room temperature for a while to allow residual ethanol from Buffer PW to evaporate.


Note

Residual ethanol from Buffer PW may inhibit subsequent enzymatic reactions.

19 Place Spin Column CP3 in a clean 1.5 ml microcentrifuge tube. Discard the collection tube.


20 Add  50-100 μ L of **Buffer EB** to the center of the Spin Column CP3.



21 Incubate for  00:02:00



2m

22 Centrifuge at  12000 rpm, Room temperature, 00:02:00 , table-top microcentrifuge



2m