



Aug 24, 2020

nCoV-2019 McGill Artic PCR Protocol, 5 ul RT and V3 only + LA1 at 63C

Forked from nCoV-2019 McGill Artic PCR Protocol, 5 ul RT and V3 only + LA1

Sarah J Reiling¹, Josh Quick²

¹McGill University; ²University of Birmingham

1 Works for me dx.doi.org/10.17504/protocols.io.bjkrkkv6

McGill Genome Centre

Sarah Reiling
McGill University

ABSTRACT

V3 only primers can be found here:

https://github.com/sarahreiling/artic-ncov2019/blob/master/primer_schemes/nCoV-2019/V3/nCoV-2019_V3only.scheme.bed

DOI

dx.doi.org/10.17504/protocols.io.bjkrkkv6

PROTOCOL CITATION

Sarah J Reiling, Josh Quick 2020. nCoV-2019 McGill Artic PCR Protocol, 5 ul RT and V3 only + LA1 at 63C.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bjkrkkv6>

FORK FROM

Forked from nCoV-2019 McGill Artic PCR Protocol, 5 ul RT and V3 only + LA1, Sarah Reiling

LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Aug 11, 2020

LAST MODIFIED

Aug 24, 2020

PROTOCOL INTEGER ID

40305

MATERIALS

NAME	CATALOG #	VENDOR
Q5 High-Fidelity 2X Master Mix - 500 rxns	M0492L	New England Biolabs
nuclease-free water		
Fresh 80% Ethanol		
Quant-iT™ PicoGreen™ dsDNA Assay Kit	P11496	Invitrogen - Thermo Fisher
AmpureXP beads	A63880	Beckman Coulter

1 PRIMER POOL PREPARATION

If required resuspend lyophilised primers at a concentration of 100 μM each



V3 only primers for this protocol were designed using [Primal Scheme](#) and generate overlapping 400 nt amplicons. Primer names and dilutions are listed in the table below.
https://github.com/sarahreiling/artic-ncov2019/blob/master/primer_schemes/nCoV-2019/V3/nCoV-2019_V3only.scheme.bed

2

Generate primer pool stocks by adding **5 μL** of each primer pair to a **1.5 mL** Eppendorf labelled either "Pool 1 (100 μM)" or "Pool 2 (100 μM)". Total volume should be **490 μL** for Pool 1 (100 μM) and **490 μL** for Pool 2 (100 μM). These are your 100 μM stocks of each primer pool.

Make another primer pool named "Pool LA1 (100 μM)" that contains 5 μL of primer pairs 5, 17, 23, 26, 66, 70, 74, 91, 97, and 10 μL of primer pair 64.



Primers should be diluted and pooled in the **mastermix cabinet** which should be cleaned with decontamination wipes and UV sterilised before and after use.

3

Dilute this primer pool 1:10 in molecular grade water, to generate 10 μM primer stocks. It is recommend that multiple aliquots of each primer pool are made to in case of degradation or contamination.

LA1 primer pool will be diluted to 1 μM primer stock.



Primers need to be used at a final concentration of 0.015 μM per primer. In this case both pools have 98 primers in so the requirement is 3.65 μL primer pools (10 μM) per 25 μL reaction. For other schemes, adjust the volume added appropriately.

Multiplex PCR

4

MULTIPLEX PCR













In the **extraction and sample addition cabinet** add **5 μL** RT product to each tube and mix well by pipetting.



The **extraction and sample addition cabinet** should be cleaned with decontamination wipes and UV sterilised before and after use.

5

In the mastermix hood set up the multiplex PCR reactions as follows in 0.2mL 8-strip PCR tubes:

Component uM]	Pool 1 [10 uM primer]	Pool 2 [10 uM]	Pool LA1 [1
Q5 Hot Start High-Fidelity 2X Master Mix	 12.5 µl	 12.5 µl	 12.5 µl
Primer Pool 1 or 2 (10µM pool 1+2; 1µM LA1)	 3.7 µl	 3.7 µl	 3.7 µl
Nuclease-free water	 3.8 µl	 3.8 µl	 3.8 µl
Total	 20 µl	 20 µl	 20 µl








Add 5 ul RT product as mentioned in step 10.



A PCR mastermix for each pool should be made up in the **mastermix cabinet** and aliquoted into PCR strip tubes. Tubes should be wiped down when entering and leaving the mastermix cabinet.

6 Pulse centrifuge the tubes to collect the contents at the bottom of the tube.

7 Set-up the following program on the thermal cycler:

Step	Temperature	Time	Cycles
Heat Activation	 98 °C	 00:00:30	1
Denaturation	 98 °C	 00:00:15	36
Annealing	 63 °C	 00:05:00	36
Hold	 4 °C	Indefinite	1




Cycle number should be 25 for Ct 18-21 up to a maximum of 36 cycles for Ct 35

PCR clean-up

8

PCR CLEANUP

Combine the entire contents of "Pool 1" and "Pool 2" PCR reactions for each biological sample into to a single 1.5 mL Eppendorf tube. **Keep Pool LA1 separate from the combined Pool 1+2 until after the clean-up!!**

9 Clean-up the amplicons using the following protocol:

Add an equal volume (1:1) of AmpureXP beads to the sample tube and mix by pipetting.
Incubate for 5 min at room temperature.
Pellet on magnet for 5 min. Remove supernatant.

Add 200 µl of 80% ethanol to the pellet and wash twice.
Let the beads dry for 3 min.
Add 30 µl elution buffer and resuspend the beads. Incubate for 3 minutes.
Pellet on magnet for 5 min. Remove and keep eluate (30 µl).



Amplicon clean-up should be performed in the **post-PCR cabinet** which should be cleaned with decontamination wipes and UV sterilised before and after use.

Amplicon Quantification and normalisation

10



AMPLICON QUANTIFICATION AND NORMALIZATION

Quantify the amplicon pools using a fluorimetric dsDNA assay. (e.g: PicoGreen with a standard curve 0-200ng)

We expect following concentrations:

Pool 1+2 combined:

100-150 ng/µl for Ct 14-24

30-80 ng/µl for Ct 25-29

10-30 ng/µl for Ct 30-36

Pool LA1:

1-10 ng/µl for all Ct

11



After quantification of Pool 1+2 and Pool LA1, mix them together in following ratio: 89.8% Pool 1+2 and 10.2% Pool LA1. For this, take a new plate and add 135 ng of Pool 1+2 and 15.3 ng of Pool LA1, and add up with nuclease-free water to a total volume of **30 µl (= 150 ng or 5 ng/µl)**.