

Jul 10, 2024

Protein extraction and Western blotting

DOI

dx.doi.org/10.17504/protocols.io.rm7vzj784lx1/v1

Shiyi Wang¹

¹Duke University

ASAP Collaborative Rese...



Shiyi Wang

Duke University

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.rm7vzj784lx1/v1

Protocol Citation: Shiyi Wang 2024. Protein extraction and Western blotting. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.rm7vzj784lx1/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: July 10, 2024

Last Modified: July 10, 2024

Protocol Integer ID: 103174

Keywords: ASAPCRN

Funders Acknowledgement:
Aligning Science Across
Parkinson's (ASAP) initiative
Grant ID: ASAP-020607



Disclaimer

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to **protocols.io** is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with **protocols.io**, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

Protein extraction and Western blotting

- 1 ****Prepare Membrane Solubilization Buffer**** - 25 mM Tris (pH 7.4) - 150 mM NaCl - 1 mM CaCl₂ - 1 mM MgCl₂ - 0.5% NP-40 - Protease inhibitors
- 2 ****Wash Cultured Astrocytes**** - Wash cultured rat astrocytes twice with ice-cold TBS containing 1 mM CaCl₂ and 1 mM MgCl₂.
- 3 ****Extract Protein**** - Incubate astrocytes on ice in membrane solubilization buffer for 20 minutes with occasional agitation.
- 4 ****Collect Cell Lysates**** - Vortex briefly. - Centrifuge at 4°C at high speed for 10 minutes to pellet non-solubilized material.
- 5 ****Store Supernatant**** - Collect the supernatant and store at -80°C.
- 6 ****Determine Protein Concentration**** - Use the Pierce BSA Protein Assay Kit (Thermo Fisher).
- 7 ****Prepare Protein Samples**** - Mix lysates with 4x Pierce™ Reducing Sample Buffer (Thermo Scientific). - Incubate at 45°C for 45 minutes to denature proteins.
- 8 ****Load Protein into Gels**** - Load 7-10 µg of protein (cultured astrocyte lysates) into Bolt™ 4–12% Bis-Tris Plus gels (Thermo Scientific). - Run at 150 V for 1 hour.
- 9 ****Transfer Proteins to Membrane**** - Transfer proteins at 100 V to PVDF membrane (Millipore) for 1.5 hours.
- 10 ****Block Membrane**** - Block in 5% BSA in TBST (137 mM NaCl, 2.68 mM KCl, 24.7 mM Tris-Base, 0.1% (w/v) Tween 20) for 1 hour.
- 11 ****Incubate with Primary Antibodies**** - Incubate overnight at 4°C with the following primary antibodies: - Anti-LRRK2 (Rabbit, 1:500; ab133474, Abcam) - GAPDH (Mouse, 1:5000; ab8245, Abcam) - β-actin (Mouse, 1:5000; A5441, Millipore Sigma)
- 12 ****Wash Membranes**** - Wash membranes with TBST.
- 13 ****Incubate with Secondary Antibodies**** - Incubate in HRP secondary antibodies (Thermo Fisher Scientific) for 2 hours.



- 14 ****Wash and Image Membranes**** - Wash membranes in TBST. - Image on a Biorad Gel Doc imaging system.
- 15 ****Quantify Protein Levels**** - Quantify protein levels using FIJI software.