

Jul 19, 2024

Wastewater grab sample processing with PEG-8000 precipitation

DOI

dx.doi.org/10.17504/protocols.io.4r3l2q3ojl1y/v1

Dilip Abraham¹, Nirmal kumar¹, Vinoth kumar¹, Ganesh Rajamanickam¹, Venkata Raghava Mohan²

¹Wellcome Trust Research Laboratory, Christian Medical College Vellore, India;

²Department of Community Health and Development, Christian Medical College Vellore, India

ES_multipathogen



Dilip Abraham

CMC Vellore

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.4r3l2q3ojl1y/v1

Protocol Citation: Dilip Abraham, Nirmal kumar, Vinoth kumar, Ganesh Rajamanickam, Venkata Raghava Mohan 2024. Wastewater grab sample processing with PEG-8000 precipitation . **protocols.io** <https://dx.doi.org/10.17504/protocols.io.4r3l2q3ojl1y/v1>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: July 11, 2024

Last Modified: July 19, 2024

Protocol Integer ID: 103219

Keywords: Wastewater, PEG-8000, Environmental sample concentration

Funders Acknowledgement:

**Bill & Melinda Gates
Foundation**




Grant ID: INV-049093



Abstract

PEG (Polyethylene Glycol) is a chemically inert, nontoxic, water-soluble synthetic polymer and has been used in aqueous polymer two-phase systems that help concentrating and isolating viruses from a variety of environmental samples. PEG is known as a good inductor of attractive interactions that crystallize viruses in the interpolymer spaces between PEG molecules. So, the combination of PEG concentration and RNA extraction steps enable 900– 1500X concentration of wastewater samples and sufficiently eliminate most of the organic matter, which could inhibit the subsequent qPCR assay.

Materials

1.  PEG 8000 **Merck MilliporeSigma (Sigma-Aldrich) Catalog #2139-1KG**
2.  Sodium Chloride Fisher BioReagents™ **Fisher Scientific Catalog #BP358-1**
3.  1X PBS (Phosphate-buffered saline)
4. Ethanol Absolute Honeywell Catalog #02875
5. Whirl-Pak® Sample Bag Merck Catalog #WPB01027WA-500EA
6. Nalgene® centrifuge bottles, Style 3122 Merck Catalog #B0408-4EA
7. SPINIX™ – Vortex Shaker Tarsons Catalog #3020
8. Eppendorf® Centrifuge 5910 R Merck Catalog #EP5943000246
9. Nalgene Centrifuge Bottle Sealing Cap Assembly Catalog number: DS3131-0038.





Sample


1 Wastewater grab samples






Procedural steps - PEG 8000 concentration method:



2h 15m

2 Collect  350 mL of sewage samples in a sterile Whirl-Pak bag and disinfect the sampling bag with ethanol to avoid contamination.


3 Transfer the bag to the laboratory in a cold chain  4 °C .

4 The sewage sample is transferred to a sterile  250 mL Nalgene centrifuge bottle.




5 Add  25 g of PEG 8000 and  5.6 g of NaCl in  250 mL of the sample to get the final concentrations of  10 % (w/v) and  0.3 Molarity (M) respectively.


6 Vortex the mixture at  500 rpm for  00:15:00 until the reagents are completely dissolved in the suspension.

15m

7 Thereafter, seal the Nalgene bottles using the cap assembly, and centrifuge the mixture at  12000 x g, 02:00:00 until the pellet is visibly seen.

2h

8 Resuspend the pellet in  3 mL of PBS ( 7.4). Then, aliquot  1 mL of the suspension into each of 3 screw cap tubes.

9 The resulting pellet is stored at  -20 °C until further process.

Freeze-thawing of primary concentrates is limited to once.



Protocol references

1. Wu F, Zhang J, Xiao A, Gu X, Lee WL, Armas F, et al. SARS-CoV-2 Titers in Wastewater Are Higher than Expected from Clinically Confirmed Cases. *mSystems* 2020 Aug 25 [cited 2020 Nov 30];5(4)
2. Kumar M, Patel AK, Shah AV, Raval J, Rajpara N, Joshi M, et al. First proof of the capability of wastewater surveillance for COVID-19 in India through detection of genetic material of SARS-CoV-2. *Science of The Total Environment*. 2020 Dec 1;746:141326