

Apr 22, 2024

DNA extraction and detection of Mansonella perstans from Dry Blood Spots (DBS)

DOI

dx.doi.org/10.17504/protocols.io.x54v928b4l3e/v1

Millicent Opoku¹, Dziedzom De Souza²

¹La Trobe University, Australia;

²Department of Parasitology, Noguchi Memorial Institute for Medical Research, University of Ghana, Ghana



Millicent Opoku

La Trobe University

OPEN ACCESS



DOI: dx.doi.org/10.17504/protocols.io.x54v928b4l3e/v1

Document Citation: Millicent Opoku, Dziedzom De Souza 2024. DNA extraction and detection of Mansonella perstans from Dry Blood Spots (DBS). **protocols.io** https://dx.doi.org/10.17504/protocols.io.x54v928b4l3e/v1

License: This is an open access document distributed under the terms of the <u>Creative Commons Attribution License</u>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Created: April 04, 2024

Last Modified: April 22, 2024

Document Integer ID: 97770

Keywords: DNA, Dried Blood Spot, Mansonella, Nested PCR, Excised Gels



Disclaimer

DISCLAIMER - FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to protocols.io is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with protocols.io, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

Mansonella perstans is a filarial parasite that causes mansonellosis in humans. Unlike other filarial parasitic diseases, mansonellosis is rarely symptomatic and hence poorly studied. This protocol describes the detection of M. perstans in blood. Firstly, DNA was extracted from dried blood blots (DBS). The DNA were screened by a nested PCR and subsequently sanger sequencing was done on excised gels. The sequences were then compared to reference sequences deposited in NCBI.



Detection of Mansonella perstans fro... 23KB

Protocol references

- 1. Quick-DNATM MiniPrep (Zymo Research, USA)
- 2. Tang T-HT, López-Vélez R, Lanza M, Shelley AJ, Rubio JM, Luz SLB. Nested PCR to detect and distinguish the sympatric filarial species Onchocerca volvulus, Mansonella ozzardi and Mansonella perstans in the Amazon Region. Memórias do Instituto Oswaldo Cruz. 2010;105(6):823-8.