

Jul 23, 2024

Immunohistochemistry of free floating slices

DOI

dx.doi.org/10.17504/protocols.io.rm7vzj6rrlx1/v1

[louis-eric.trudeau](#)¹

¹Department of Pharmacology and Physiology and Department of Neurosciences, Faculty of Medicine, Université de Montréal, Montreal, QC, Canada. SNC and CIRCA Research Groups, Université de Montréal, Montréal, QC, Canada. Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD, 20815, USA.

ASAP Collaborative Rese...



Nicolas Giguère

Trudeau Lab

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.rm7vzj6rrlx1/v1

Protocol Citation: louis-eric.trudeau 2024. Immunohistochemistry of free floating slices. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.rm7vzj6rrlx1/v1>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: June 20, 2024

Last Modified: July 23, 2024

Protocol Integer ID: 102402

Keywords: ASAPCRN

Abstract

This protocol details the immunohistochemistry of free floating slices.



Steps

2w 0d 5h 40m

1

2w

Note



Step 1 is for fresh tissue only, otherwise, start directly to step 2

Fix with paraformaldehyde (PFA) 4% Overnight at Room temperature .

- 4 g of PFA
- Dissolve in 100 mL de PBS (pH 7.3) previously heated to around 60 °C .
- Mix the solution until it completely clears up.
- Filter and store at 4 °C for a duration of 336:00:00 .

2 Rinse with PBS 1X (3X 10 min) under agitation.

2.1 Rinse with PBS 1X for 00:10:00 under agitation (1/3).

10m



2.2 Rinse with PBS 1X for 00:10:00 under agitation (2/3).

10m



2.3 Rinse with PBS 1X for 00:10:00 under agitation (3/3).

10m



3 Permeabilize and block the non-specific primary antibodies bindings in a 24-well multiwell plate by incubating under agitation for 01:00:00 with 0.5 mL to 1 mL of a BSA enriched ab solution (10 mL of ab solution for 1 g of BSA).


1h






4 All the subsequent steps are done using the following ab solution:

- 95 mL of PBS 7.3
- 20 mg of NaN3 (final 0.02 %)
- 300 µL of Triton X-100 (final 0.3 %)
- 500 mg of BSA (final 0.5 %)



-  5 mL of goat serum (final 5 %)


5 Dilute the primary antibodies in the ab solution and apply  0.5 mL per well (up to 10 slices)  Overnight at  Room temperature under agitation.

1h



6 Rinse the slices with PBS (3X 10 min) under agitation.



6.1 Rinse the slices with PBS for  00:10:00 under agitation (1/3).

10m



6.2 Rinse the slices with PBS for  00:10:00 under agitation (2/3).




10m



6.3 Rinse the slices with PBS for  00:10:00 under agitation (3/3).

10m



7 Dilute the secondary antibodies in the ab solution and apply  0.5 mL per well (up to 10 slices) for  02:00:00 at  Room temperature under agitation.

2h



8 Rinse the slices with PBS (3X 10 min) under agitation.



8.1 Rinse the slices with PBS for  00:10:00 under agitation (1/3).


10m



8.2 Rinse the slices with PBS for  00:10:00 under agitation (2/3).

10m




8.3 Rinse the slices with PBS for  00:10:00 under agitation (3/3).

10m





9 Mount the slices on charged (+) glass slides submerged in PBS.

10 Let the mounted slices dry (about  00:10:00).

10m

**Note**

*Don't wait too long otherwise salt deposits will start to appear and affect your staining.

- 11 Seal the slides with  100 μL -  120 μL /slide of Fluoromount G and a rectangular glass coverslip and seal the coverslip to the slide with nail polish.