



JAN 05, 2024

OPEN ACCESS



**DOI:**  
[dx.doi.org/10.17504/protocols.io.e6nvwdnzzlmk/v1](https://dx.doi.org/10.17504/protocols.io.e6nvwdnzzlmk/v1)

**Protocol Citation:** Federico N. Soria, Mario Fernandez-Ballester 2024. Rat organotypic cultures for AAV-mediated vital labeling of the extracellular matrix.

**protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.e6nvwdnzzlmk/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working  
 We use this protocol and it's working

**Created:** Jan 05, 2024

## 🌐 Rat organotypic cultures for AAV-mediated vital labeling of the extracellular matrix

Federico N. Soria<sup>1</sup>, Mario Fernandez-Ballester<sup>1</sup>

<sup>1</sup>Achucarro Basque Center for Neuroscience

ASAP Collaborative Research Network



Federico N. Soria

Achucarro Basque Center for Neuroscience

### ABSTRACT

In this protocol we describe the preparation and maintenance of rat organotypic cultures (**cortico-striatal**) for AAV-infection and imaging of the extracellular matrix. The AAV, in this case, induces the expression of a exportable hyaluronan-binding protein (Neurocan) fused with GFP, that allows vital labeling (and hence live imaging) of the extracellular matrix. The organotypic cultures can be maintained for several weeks, allowing for repeated imaging and longitudinal studies.

Last Modified: Jan 05, 2024









PROTOCOL integer ID:  
92986

Keywords: ASAPCRN


Funders  
Acknowledgement:  
Aligning Science Across  
Parkinson  
Grant ID: ASAP-020505

MATERIALS

Organotypic culture medium

- 50%  MEM, HEPES, no glutamine Gibco,  
ThermoFisher Catalog #32360026
- 25%  EBSS, calcium, magnesium, phenol red Thermo  
Fisher Catalog #24010043
- 25%  Horse Serum, heat inactivated, New Zealand origin Thermo  
Fisher Catalog #26050088
- 1x  B-27 Supplement Gibco - Thermo  
Fischer Catalog #17504044
- 1x  GlutaMAX™ Supplement Thermo Fisher Scientific Catalog #35050061
- 36 mM  Glucose Solution Thermo Fisher Catalog #A2494001
- 50 U/mL  Penicillin-Streptomycin (5,000 U/mL) Thermo  
Fisher Catalog #15070063
- 1.25 ug/mL  Fungizone (Amphotericin B) Thermo Fisher  
Scientific Catalog #15290018

Membrane Inserts

-  Millipore Cell Culture Inserts (0.4 um pore size, 30 mm diameter) Merck  
MilliporeSigma (Sigma-Aldrich) Catalog #PICM0RG50

Tissue Chopper

Equipment

McIlwain Tissue Chopper

NAME


Cavey Laboratory

BRAND


Engineering

51350


SKU




## PROTOCOL MATERIALS

 Fungizone (Amphotericin B) Thermo Fisher Scientific Catalog #15290018

Materials

 HBSS, no calcium, no magnesium, no phenol red Thermo Fisher Scientific Catalog #14175095


Step 2

 B-27 Supplement Gibco - Thermo Fischer Catalog #17504044


Materials

 GlutaMAX™ Supplement Thermo Fisher Scientific Catalog #35050061

Materials

 Millipore Cell Culture Inserts (0.4 µm pore size, 30 mm diameter) Merck MilliporeSigma (Sigma-Aldrich) Catalog #PICM0RG50


Materials

 EBSS, calcium, magnesium, phenol red Thermo Fisher Catalog #24010043


Materials

 Penicillin-Streptomycin (5,000 U/mL) Thermo Fisher Catalog #15070063

Materials

 Horse Serum, heat inactivated, New Zealand origin Thermo Fisher Catalog #26050088


Materials

 MEM, HEPES, no glutamine Gibco, ThermoFisher Catalog #32360026

Materials


 Glucose Solution Thermo Fisher Catalog #A2494001

Materials

 Cytosine β-D-arabinofuranoside (AraC) Merck MilliporeSigma (Sigma-Aldrich) Catalog #C1768




Step 11

## Culture preparation

- 1 Prepare a 6-well culture plate with Millipore 0.4 µm culture inserts placed on 1 mL of organotypic medium per well, and incubate it at  and 5% CO<sub>2</sub> to warm up.

### Note

It is important to minimize air bubbles between the medium and the membrane insert, as this will prevent access of nutrients to the slice.

- 2 Prepare a p100 and p60 petri dishes with 25 mL and 5 mL, respectively, of  
 HBSS, no calcium, no magnesium, no phenol red Thermo Fisher Scientific Catalog #14175095  
and keep them  On ice .
- 3 Prepare the **McIlwain Tissue Chopper** (see Materials) with a fresh blade and clean holder plates.
- 4 Sacrifice P5-P7 rat pups by rapid decapitation and extract the brain without damaging the cortex.
- 5 Put the brain in the on a filter paper and cut through the midline to separate the cerebral hemispheres. Remove the midbrain, separating it from the cortex with a scalpel.
- 6 Place both hemispheres in the same holder plate, with the medial face downward. Hold the plate with both hemispheres perpendicular to the blade (coronal sections) and section the brain at 350  $\mu$ m width.
- 7 Pour the sliced brain carefully into the p100 petri dish with HBSS  On ice . You may use a Pasteur pipette to moist the brain and displace it more easily.
- 8 Separate slices under the magnification microscope with a couple of spatulas (preferentially plastic-ones, which can be custom crafted cutting the back of a Pasteur pipette). Use a blade and clean cuts to remove unwanted regions (such as hippocampus or SVZ). Transfer the slices into the p60 petri dish (you can use an inverted glass pipette with a suction rubber bulb).


#### Note

It is important to remove the subventricular zone (SVZ) and the meninges with the scalpel to avoid cell proliferation.

- 9 **Under the laminar flow hood**, transfer the selected slices (undamaged slices from the preferred rostro-caudal levels) to preheated culture inserts with medium (Step 1). Each insert can only hold two cortico-striatal slices. Remove excess dissection buffer from the top of the insert with a sterile tip.

#### Note

Leaving too much HBSS solution on top of the insert will eventually dilute the feeding medium.

- 10 Incubate the culture at  37 °C and 5% CO<sub>2</sub>.

#### Expected result

Typically, 4-6 cortico-striatal slices can be obtained from each hemisphere

## Maintenance and AAV-infection

- 11 At DIV1 (24h after culture preparation), replace the feeding medium with 1 mL of the same medium but adding antimetabolic



Cytosine β-D-arabino-furanoside (AraC) Merck MilliporeSigma (Sigma-Aldrich) Catalog #C1768

4,4 μm per well.

#### Expected result

AraC will prevent the overproliferation of astrocytes, minimizing the formation of a glial scar on top of the slice.

- 12 At DIV3, replace the AraC-containing medium with fresh medium. Then add 1 μL of the virus of choice (in our case, AAV-Ncan-GFP 10<sup>12</sup> vg/mL) over the slice.

### Safety information

AAVs have to be manipulated in BSL1 facilities.

### Note

To improve spatial precision (e.g. when infecting a selected region from the slice), AAV infection can be done with a micromanipulator and a glass pipette.

**13** To maintain the culture, change medium three times a week (1 mL per well).

### Expected result

Slices will flatten overtime, and become more transparent. Cortical regions will have more cell density than striatal regions.

AAV expression become visible aprox. at DIV10, with maximum expression from DIV14 onwards (it depends on AAV serotype and promoter).

## Imaging

**14** Imaging can be performed in two ways:

- **IN-PLATE:** The insert **is not removed from the plate**, and the imaging is performed through the plastic. This is usually done in *widefield* setups, with low magnification. It is not optimal in terms of resolution, but it is very useful for repeated imaging (longitudinal studies) without compromising sterility.
- **OFF-PLATE:** The insert is **removed from the plate** and placed in a smaller petri-dish or similar holding chamber. Imaging is usually done from above, in upright microscopes (objective on-top), with water-based objectives. Since CO<sub>2</sub> is usually not available, a buffer medium with HEPES is recommended. Phenol-red should be avoided as well, since it interferes with fluorescence. This type of configuration is typically used for *confocal* or *2-photon* imaging of the slices.