




DEC 14, 2023

## DAB ANTIBODY (IHC) STAINING PROTOCOL

Michael Lee<sup>1</sup>

<sup>1</sup>University of Minnesota

Team Lee



jbalster

### DISCLAIMER

OPEN ACCESS



**DOI:**  
[dx.doi.org/10.17504/protocols.io.kxygzqyzv8j/v1](https://dx.doi.org/10.17504/protocols.io.kxygzqyzv8j/v1)

**Protocol Citation:** Michael Lee 2023. DAB ANTIBODY (IHC) STAINING PROTOCOL. **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.kxygzqyzv8j/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working  
 We use this protocol and it's working

**Created:** Jun 07, 2022

**Last Modified:** Dec 14, 2023

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

### ABSTRACT

This is the basic protocol for antibody staining of formalin fixed paraffin embedded (FFPE) tissue.

### ATTACHMENTS

[452-959.docx](#)

**PROTOCOL integer ID:**  
64108

**Keywords:** DAB ANTIBODY  
(IHC) STAINING PROTOCOL,  
Formalin fixed paraffin  
embedded (FFPE) tissue,  
FFPE, ASAPCRN

**Funders**

**Acknowledgement:**

ASAP

## GUIDELINES

### Principle:

For antibody staining to be successful, most FFPE tissue requires antigen retrieval of some kind. Formalin fixation cross-links proteins during the course of fixation. Antigen retrieval unlinks the proteins and opens up the antigen sites so that the antibody will be able to bind to them.

### References:

Carson, Freida, *Histotechnology – A Self-Instructional Text*, 2<sup>nd</sup> Edition, ASCP Press, 1997

## MATERIALS

### **Specimen Preparation:**

10% Neutral buffered formalin fixed tissue, or 4% paraformaldehyde fixed tissue, paraffin embedded sections cut at ~ 5-6 microns and mounted on charged slides.

### **Materials and Equipment:**

- Charged slides
- Coverslips
- Drying oven, 60 degrees
- Fume Hood
- Gloves
- Microtome
- Staining racks
- Timer

### **Reagents:**

- Xylene
- 100% Alcohol
- 95% Alcohol
- 80% Alcohol
- Harris Hematoxylin
- Background Sniper (Biocare Medical)
- TRIS with tween 20X (Biolegend,[hydroxymethyl aminomethane])
- Secondary antibody; i.e. Donkey anti-Rabbit
- Tertiary link; i.e., Vector Laboratories Vectastain ABC-HRP kit
- 3% Hydrogen peroxide
- DAB (Diaminobenzidine) from Biolegend
- DAB substrate Buffer (Biolegend)
- Permout or other xylene compatible mounting media

### **Pretreatment reagents:**

- Reveal Decloaker (Biocare Medical) preferred
- Citrate buffer solution if you don't use the Reveal
- Specific case for Beta-amyloid antibody
- 70% Formic acid treatment

## SAFETY WARNINGS



### Precautions:

#### Personal Protection:

Gloves, lab coat, goggles, fume hood, and use of universal precaution practices.

#### Chemical Wastes:

Dispose of alcohols, dyes, and xylene in appropriate labeled waste containers as directed by the University of Minnesota Hazardous Chemical Waste Management Manual 5<sup>th</sup> Edition.

#### Hazards:





- Xylene = Flammable, Carcinogen, Skin irritant
- Eosin & Alcohols = Flammable, Skin irritant
- Hematoxylin = Skin irritant
- Avoid strong oxidizers with all listed chemicals

## Deparaffinize tissue: Day 1

45m

### 1 2 options:

45m

1) Either place slides on a slide warmer with temperature set to approx.  57 °C . Leave the slides on the warming plate until the paraffin looks melted on all of the slides (about 10 -  00:15:00 ) or 2) Put slides in a  60 °C oven for about  00:30:00 .

2 Place slides vertically in the gray plastic slide holders and run the slides through the following solutions in the staining set up located in the fume hood.



A	B
Xylene	5 minutes
Xylene	5 minutes
Xylene	10 minutes
100% Alcohol	3 - 5 minutes
100% Alcohol	3 - 5 minutes
95% Alcohol x 2	3 - 5 minutes


A	B
80% Alcohol	3 - 5 minutes
Filtered water	3 - 5 minutes

## STEP CASE

### Preferred Antigen retrieval steps:

34 steps

- 3 Fill the vegetable steamer with deionized water to the second line in the transparent corner section of the steamer.


- 3.1 Turn steamer on and push the up arrow button until the time is about  00:40:00 .

40m

#### Note

This will allow the steamer to heat up and be ready for the prepared slides.

- 4 Add capillary gap slides (same number as the number of slides you are staining) in the slide holder to wet the slides.

- 4.1 Fill the plastic "boat" containers with approx.  20 mL of one of the antigen retrieval solutions (Reveal or citrate buffer).





- 4.2 Pick up a set of slides (rough side of the capillary gap slide facing the tissue section slide) and put the slides into one of the troughs in the boat container.


- 4.3 Push slides toward each other and allow the fluid to come up slowly between the two slides.

#### Note


Try not to get bubbles between the slides or this will obstruct the antigen retrieval process and give you uneven staining.

4.4 Steam for about  00:30:00 . 30m

5 Remove clear basket from the steamer and allow slides to cool for about  00:20:00 to  Room temperature . 20m

6 Transfer the “boats” of slides to the inner steamer container and rinse with running water for about  00:20:00 . 20m





6.1 Unpeel the slides from each other and transfer the slides to the gray slide holder for a final rinse ( 00:05:00 ). 5m







## Staining steps:

2h 38m



7 Dilute or use prediluted TRIS (1x concentration) for  00:10:00 , approx 250-  300  $\mu\text{L}$  per slide. 10m



8 100% Background Sniper for nearly  00:13:00 . Do not go longer than  00:15:00 . (alternat 1h 28m  
10% normal goat serum made in 1X TRIS or PBS can be substituted for  01:00:00 ).

9 Make up antibody solution at desired concentration in a diluent of 5% Sniper in 1X TRIS solution. Using approx.  100  $\mu\text{L}$  per slide, make up enough antibody to cover all slides.



10 Cover the slides with either a glass coverslip (24 x 60 mm) or parafilm and put in a  4  $^{\circ}\text{C}$  . refrigerator 1h  
 Overnight .



## DAY 2:

1h 20m

- 11 Take slides out of the refrigerator and warm up to Room temperature about 00:15:00 - 00:20:00 . Remove the cover slips and rinse the slides with 1X TRIS solution, then incubate slides in 1X TRIS solution for about 00:05:00 x 2. 40m



- 12 **Secondary reagent/antibody for** 00:30:00 30m
- Apply the secondary reagent appropriate to the antibody, i.e., if your antibody was raised in a rabbit, you will apply an anti- Rabbit link.



### Note

You will need to make up enough reagent for all the slides with approx. 250-300 microliter per slide or enough of the reagent to totally cover the tissue. If your antibody was raised in some other animal, you will have to find a secondary to that animal, i.e., donkey anti-mouse, goat anti-chicken, etc. Usually a dilution of 1:500 in a diluent of 5% Sniper made in 1X TRIS is a good starting concentration to use. Jackson ImmunoResearch Labs supply a host of secondary antibodies (**make sure the secondary is biotinylated**).

- 13 Rinse slides with 1X TRIS, 00:10:00 . 10m





- 14 3% Hydrogen Peroxide made in 1X TRIS for 00:10:00 x 2. Rinse between with 1X TRIS. 10m




- 15 Rinse slides with 1X TRIS, 5 minutes x 2.



15.1 Rinse slides with 1X TRIS,  00:05:00 x 2 (1/2). 5m




15.2 Rinse slides with 1X TRIS,  00:05:00 x 2 (2/2). 5m

16 TERTIARY Reagent for  00:30:00 . 30m

16.1 The Tertiary Reagent is made up about  00:25:00 before you apply it. 25m





#### Note

We typically use the Vectastain Elite ABC kit, Peroxidase (standard) from VECTOR Laboratories. The staining concentration is  100  $\mu$ L reagent A to  5 mL diluent of 5% Sniper made in 1X TRIS, then add  100  $\mu$ L B to the solution, mix and let sit until applied to the tissue.


17 Rinse slides with 1X TRIS, 5 minutes x 2.



17.1 Rinse slides with 1X TRIS,  00:05:00 x 2 (1/2). 5m

17.2 Rinse slides with 1X TRIS,  00:05:00 x 2 (2/2). 5m





18 DAB (DIAMINOBENZINE) chromogen for approximately  00:03:00 to develop.

3m





### Safety information


DAB is carcinogenic – wear gloves and make sure to dispose of waste solutions, pipets, and syringes in appropriate hazardous waste containers.

18.1 Make up enough DAB solution to cover all the slides, using  250  $\mu\text{L}$  -  300  $\mu\text{L}$  per slide as a guide.



### Note

A dilution of  40  $\mu\text{L}$  of concentrated DAB (Biolegend) per  1 mL DAB substrate Buffer (Biolegend) is usually good.

18.2 Filter the DAB solution with a  0.22  $\mu\text{L}$  syringe filter into a new vial before use.

### Note

Microscopic checks to check and stop the development of the chromogen is advised. If the chromogen develops very quickly, then you will want to dilute the antibody concentration.

18.3 After development put slides in slide holder in water container.

18.4 Fill clear basket with water then submerge slides in running water in sink.

18.5 Now counterstain.

## Counterstaining with hematoxylin and coverslipping:


1h 20m

19

Filtered Harris Hematoxylin – several dips, return to water and check after a few minutes to see if the hematoxylin is strong enough, if not, return for more dips and repeat.

20



Tap water rinse for about  00:05:00 to both “blue” the hematoxylin and rinse the slides until the water runs clear. Put slides in container with filtered water for dehydration and clearing steps as follows: 5m

A	B
80% Alcohol	2 minutes
95% Alcohol	2 minutes
100% Alcohol	2 minutes
100% Alcohol	2 minutes
100% Alcohol	2 minutes
100% Alcohol	2 minutes
Xylene	2 minutes
Xylene	2 minutes
Xylene	2 minutes

### Note


Leave slides in last container of xylene until they are mounted with Permount. Usually 2 drops of Permount is enough. Wipe off excess mounting media and allow to dry flat. Leave slides/slide holder to evaporate Xylene fumes in hood for a while.

### Result examples:


Nuclei = Blue

Positive staining antigen = Brown

Background & cytoplasm= pale blue to colorless



Tau Stain



IBA1 stain