



Version 1

Jul 15, 2020

MARVICS: A Robust and Safe Magnetic Nanoparticle based RNA Extraction Method Compatible with Phenol-chloroform Inactivated Infectious Samples V.1

Mo Li¹, Gerardo Ramos-Mandujano²¹King Abdullah University of Science and Technology;²Laboratory of Stem Cell and Regeneration, Biological and Environmental Science and Engineering Division, King Abdullah University of Science and Technology (KAUST)

1

Works for me

dx.doi.org/10.17504/protocols.io.bh82j9ye

Mo Li

King Abdullah University of Science and Technology

ABSTRACT

Diagnosis and surveillance of emerging pathogens such as SARS-CoV-2 depend on nucleic acid isolation from clinical and environmental samples. Under normal circumstances, samples would be processed using commercial proprietary reagents in Biosafety 2 (BSL-2) or higher facilities. A pandemic at the scale of COVID-19 has caused a global shortage of proprietary reagents and BSL-2 laboratories to safely perform testing. Therefore, alternative solutions are urgently needed to address these challenges. We developed an open-source method called Magneticnanoparticle-Aided Viral RNA Isolation of Contagious Samples (MAVRICS) that is built upon reagents that are either readily available or can be synthesized in any molecular biology laboratory with basic equipment. Unlike conventional methods, MAVRICS works directly in samples inactivated in acid guanidinium thiocyanate-phenol-chloroform (e.g., TRIzol), thus allowing infectious samples to be handled safely without biocontainment facilities.

EXTERNAL LINK

<https://www.medrxiv.org/content/10.1101/2020.06.28.20141945v1>

DOI

dx.doi.org/10.17504/protocols.io.bh82j9ye

PROTOCOL CITATION

Mo Li, Gerardo Ramos-Mandujano 2020. MARVICS: A Robust and Safe Magnetic Nanoparticle based RNA Extraction Method Compatible with Phenol-chloroform Inactivated Infectious Samples. **protocols.io**
dx.doi.org/10.17504/protocols.io.bh82j9ye



EXTERNAL LINK

<https://www.medrxiv.org/content/10.1101/2020.06.28.20141945v1>

KEYWORDS

SARS-CoV-2, MAVRICS, COVID-19, RNA extraction, Magnetic nanoparticle synthesis, Silica magnetic nanoparticles (SiMNP)

LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Jul 05, 2020

LAST MODIFIED

Jul 15, 2020

MATERIALS

NAME	CATALOG #	VENDOR
Bis-Tris	B-020	Gold Biotechnology
TRIzol [®] Reagent	15596018	Thermo Fisher
SuperScript [®] IV Reverse Transcriptase	18090010	Thermo Fisher
TaqMan [®] Fast Advanced Master Mix	4444556	Thermo Fisher
Hydrochloric Acid	A144S	Fisher Scientific
Sodium hydroxide	306576	Sigma-aldrich
Ethanol absolute ≥99.8% AnalaR NORMAPUR [®] ACS Reag. Ph. Eur. analytical reagent	20821.330DP	
Tetraethyl orthosilicate ≥99.0% (GC)	86578	Sigma-aldrich
Iron (III) chloride anhydrous Extra Pure	10224390	Fisher Scientific
Guanidine Hydrochloride	BP178-500	Fisher Scientific
RNaseOUT [™] Recombinant Ribonuclease Inhibitor	10777019	Invitrogen - Thermo Fisher
RNase H	M0297	New England Biolabs
2019-nCoV RUO Kit	10006605	
Iron(II) chloride tetrahydrate ≥98%		


1 Silica magnetic nanoparticles (SiMNP) synthesis.

SiMNP synthesis was done following the published protocols in [BOMB.bio: BOMB magnetic core nanoparticles synthesis](#) and [BOMB coating ferrite MNPs with silica oxide](#).

 **Overnight**

2

COVID-19 patient samples.





Oropharyngeal or nasopharyngeal swabs were steeped in  **1 mL** acid guanidinium thiocyanate-phenol-chloroform (AGPC, e.g., TRIzol Reagent or TRI reagent).



Follow CDC or institutional safety guidelines when handling potential infectious samples. AGPC, TRIzol and TRI reagent contain phenol. Follow local safety guidelines when handling and disposing these reagents.

 **00:01:00**

3 Making Bis-Tris Buffer **50 mL**

3.1 Dissolve  **14.33 g** guanidine hydrochloride and  **104.6 mg** Bis-Tris in  **45 mL** of 100% ethanol.  **00:10:00**



If Bis-Tris is not available, it may be substituted by Tris Base (10 mM final concentration)



Add 40 ml of 100% ethanol to the other chemicals, and wait for guanidine hydrochloride to completely dissolve and add the remaining volume of 100% Ethanol.

3.2 Adjust pH (<6.5) with HCl, and adjust the volume with H₂O to 50 mL .

00:05:00

4 Magnetic-nanoparticle-Aided Viral RNA Isolation of Contagious Samples.

4.1 In an Eppendorf tube add 200 μ l clinical sample, 200 μ l Bis-Tris buffer, mix well by vortexing.

00:01:00



We recommend Lo-Bind Eppendorf tubes or similar low binding tubes.



Samples contain phenol. Cap the tubes slowly. Make sure the tubes can be closed securely during vortexing.

4.2 Add 40 μ l SiMNP, mix at 1300 rpm 00:05:00



The stock SiMNP is composed of 1 volume of SiMNP in 1 volume of RNase-free water. The stock is further diluted to be used in RNA extraction. Typical dilution ranges from 1:4 to 1:10. The optimal ratio should be determined empirically.



4.3 Spin the tube for 2-3 seconds, settle the SiMNPs on a magnetic stand and remove the supernatant.

00:02:00



The supernatant contains phenol. Follow local safety guidelines when handling and disposing these reagents.

4.4

Mix  **200 µl** of AGPC (TRIzol or TRI reagent) and  **200 µl** Bis-Tris buffer, add to the SiMNPs, mix well by vortexing. ⌚ **00:01:00**


4.5

Settle the SiMNPs on a magnetic stand and remove the supernatant. ⌚ **00:02:00**



The supernatant contains phenol. Follow local safety guidelines when handling and disposing these reagents.

4.6

Add  **400 µl** 90% ethanol, spin for 2-3 seconds, settle the siMNPs on a magnetic stand and remove the supernatant. ⌚ **00:02:00**

4.7

Repeat Setp 4.6 three more times for a total of 4 ethanol washes ⌚ **00:06:00**

4.8

After removing the supernatant from the last ethanol wash, dried the SiMNPs on a heat block at 50°C. Keep the lid open, no shaking. Do not elute before the SiMNPs are dried. ⌚ **00:20:00**




Drying may take less than 20 min. Monitor the appearance of the SiMNPs during drying. The SiMNPs appear in a rusty brownish color when dried.

4.9 To elute the RNA, add **40 µl** nuclease-free water, and mix at **1300 rpm** ⌚ **00:05:00**

4.10 Settle the SiMNPs on a magnetic stand and transfer the eluted RNA to a new RNase-free tube. ⌚ **00:02:00**

4.11






Analyze RNA concentration and purity using a Qubit fluorometer or Nanodrop. ⌚ **00:10:00**

- 5 **Reverse transcription (RT):** use  **4 µl** of eluted RNA and follow the instructions for [SuperScript™ IV Reverse Transcriptase](#) adding the RNase H incubation step. ⌚ **01:00:00**



User should optimize the input RNA amount.

- 6 

Real-time PCR: For each  **10 µl** qPCR reaction mix  **1.5 µl** cDNA,  **0.5 µl** [SARS-CoV-2 \(2019-nCoV\) CDC qPCR Probe Assay](#),  **5 µl** TaqMan Fast Advanced Master Mix, and  **1.5 µl** nuclease-free water. ⌚ **01:20:00**