



Aug 04, 2021

## SEGA protocol collection

Carolyn N Bayer<sup>1</sup>, Maja Rennig<sup>1</sup>, Anja Ehrmann<sup>1</sup>, Morten Norholm<sup>1</sup><sup>1</sup>Technical University of Denmark

1 Works for me

Share

[dx.doi.org/10.17504/protocols.io.bvxhn7j6](https://dx.doi.org/10.17504/protocols.io.bvxhn7j6)

SEGA

Carolyn Bayer  
Technical University of Denmark

### DISCLAIMER

#### DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](https://protocols.io) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](https://protocols.io), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

### ABSTRACT

SEGA, the Standardized Genome Engineering Architecture, is a comprehensive strain collection that enables genome engineering by combining only two reagents: a DNA fragment that can be ordered from a commercial vendor and a stock solution of bacterial cells followed by incubation on agar plates. Recombinant genomes are identified by visual inspection using green-white colony screening akin to classical blue-white screening for recombinant plasmids. The modular nature of SEGA allows precise multi-level control of transcriptional, translational, and post-translational regulation. The SEGA architecture simultaneously supports increased standardization of genetic designs and a broad application range by utilizing well-characterized parts optimized for robust performance in the context of the bacterial genome.

This protocol is a collection of all important SEGA protocols.

### DOI

[dx.doi.org/10.17504/protocols.io.bvxhn7j6](https://dx.doi.org/10.17504/protocols.io.bvxhn7j6)

### COLLECTION CITATION

Carolyn N Bayer, Maja Rennig, Anja Ehrmann, Morten Norholm 2021. SEGA protocol collection.  
**protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.bvxhn7j6>

### LICENSE

This is an open access collection distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

### CREATED

Jun 18, 2021

LAST MODIFIED

Aug 04, 2021

COLLECTION INTEGER ID

50889

DISCLAIMER:

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK




The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](https://protocols.io) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](https://protocols.io) can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

SEGA, the Standardized Genome Engineering Architecture, is a comprehensive strain collection that enables genome engineering by combining only two reagents: a DNA fragment that can be ordered from a commercial vendor and a stock solution of bacterial cells followed by incubation on agar plates. Recombinant genomes are identified by visual inspection using green-white colony screening akin to classical blue-white screening for recombinant plasmids. The modular nature of SEGA allows precise multi-level control of transcriptional, translational, and post-translational regulation. The SEGA architecture simultaneously supports increased standardization of genetic designs and a broad application range by utilizing well-characterized parts optimized for robust performance in the context of the bacterial genome.

This protocol is a collection of all important SEGA protocols.

FILES

-  **Integration of a landing pad brick**  
**Version 1**  
by Carolyn Bayer, Technical University of Denmark
-  **Integration of a cargo brick**  
**Version 1**  
by Carolyn Bayer, Technical University of Denmark
-  **Integration of a control brick**  
**Version 1**  
by Carolyn Bayer, Technical University of Denmark