

Jul 08, 2024

Phosphoglycerate kinase is a central leverage point in Parkinson's Disease driven neuronal metabolic deficits.

DOI

dx.doi.org/10.17504/protocols.io.3byl495nzgo5/v1

Alexandros C Kokotos^{1,2}, ama^{2,3}, Santiago Unda^{2,3}, Myung Soo Ko¹, Daehun Park^{2,4}, David Eliezer¹, Michael G. Kaplitt^{2,3}, Pietro De Camilli^{2,5}, Tim Ryan^{1,2}

¹Department of Biochemistry, Weill Cornell Medicine, New York, NY 10065;

²Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, Maryland 20815, USA;

³Department of Neurosurgery, Weill Cornell Medicine, New York, NY 10065;

⁴Department of Medical and Biological Sciences, The Catholic University of Korea, Bucheon, 14662, Korea;

⁵Departments of Neuroscience and Cell Biology, Howard Hughes Medical Institute, Program in Cellular Neuroscience, Neurodegeneration and Repair, Yale University School of Medicine, New Haven, Connecticut 06520, USA

ASAP Collaborative Rese...



Alexandros C Kokotos

Weill Cornell Medicine

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.3byl495nzgo5/v1

Collection Citation: Alexandros C Kokotos, ama, Santiago Unda, Myung Soo Ko, Daehun Park, David Eliezer, Michael G. Kaplitt, Pietro De Camilli, Tim Ryan 2024. Phosphoglycerate kinase is a central leverage point in Parkinson's Disease driven neuronal metabolic deficits.. protocols.io <https://dx.doi.org/10.17504/protocols.io.3byl495nzgo5/v1>

License: This is an open access collection distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this collection and it's working

Created: July 08, 2024



Last Modified: July 08, 2024

Collection Integer ID: 103017

Keywords: ASAPCRN

Funders Acknowledgement:

ASAP

Grant ID: 000580

ASAP

Grant ID: 020608

Abstract

The current collection contains all experimental protocols associated with the study "Phosphoglycerate kinase is a central leverage point in Parkinson's Disease driven neuronal metabolic deficits".

Files

 SEARCH

Protocol



NAME

Immunofluorescence for adherent cells

VERSION 1

CREATED BY



Alexandros C Kokotos
Weill Cornell Medicine

OPEN →

Protocol



NAME

Fast rodent genotyping

VERSION 1

CREATED BY



Alexandros C Kokotos
Weill Cornell Medicine

OPEN →

Protocol



NAME

Primary hippocampal and cortical neuronal culture and transfection

VERSION 1

CREATED BY



Alexandros C Kokotos
Weill Cornell Medicine

OPEN →

Protocol



NAME

Protocol for Neuronal Live-imaging of primary cultures

VERSION 1

CREATED BY

 **Camila Pulido**
Weill Cornell Medicine


[OPEN](#) →

Protocol

 NAME
AAV Craniotomy

VERSION 1

CREATED BY

 **Eileen Ruth Torres**
Weill Cornell Medicine


[OPEN](#) →

Protocol

 NAME
6-OHDA lesion in medial forebrain bundle

VERSION 1

CREATED BY

 **Eileen Ruth Torres**
Weill Cornell Medicine


[OPEN](#) →

Protocol

 NAME
Apomorphine-induced rotations


VERSION 1

CREATED BY

 **Eileen Ruth Torres**
Weill Cornell Medicine

[OPEN](#) →

Protocol

 NAME
Immunohistochemistry for brain sections

VERSION 1

CREATED BY



Eileen Ruth Torres
Weill Cornell Medicine

OPEN →

Protocol



NAME

Genetic expression suppressor screen

VERSION 1

CREATED BY



Alexandros C Kokotos
Weill Cornell Medicine

OPEN →

Protocol



NAME

vGlut1-pH imaging experiment analysis

VERSION 1

CREATED BY



Alexandros C Kokotos
Weill Cornell Medicine

OPEN →

Protocol



NAME

Synapto-iATPSnFR2-miRFP670nano3 analysis

VERSION 1

CREATED BY



Alexandros C Kokotos
Weill Cornell Medicine

OPEN →