



MAR 28, 2023

OPEN  ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.8epv5jp94l1b/v1

Protocol Citation: Geórgia Hackradt Rêgo, Alice Leite, Maryana Thalyta Ferreira Câmara de Oliveira, Beatriz Helena Dantas Rodrigues de Albuquerque, Larissa Alves Honorato Ferreira, Yago Tomaz Vieira da Silva, Eduardo Pereira de Azevedo, Janaína Ferreira Aderaldo, Daniel Carlos Ferreira Lanza, Ricardo Ney Cobucci 2023. Protocol for Papillomavirus DNA extraction from cervical brushes using the Wizard® Genomic DNA Purification Kit. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.8epv5jp94l1b/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Mar 28, 2023

Protocol for Papillomavirus DNA extraction from cervical brushes using the Wizard® Genomic DNA Purification Kit

Geórgia Hackradt Rêgo¹, Alice Leite²,
 Maryana Thalyta Ferreira Câmara de Oliveira³,
 Beatriz Helena Dantas Rodrigues de Albuquerque⁴,
 Larissa Alves Honorato Ferreira⁵, Yago Tomaz Vieira da Silva⁵,
 Eduardo Pereira de Azevedo⁶, Janaína Ferreira Aderaldo⁷,
 Daniel Carlos Ferreira Lanza⁸, Ricardo Ney Cobucci⁹

¹Graduate program in sciences applied to women's health (PPGCASM), Federal University of Rio Grande do Norte (UFRN), Natal, RN, Brazil;

²Medical residency in gynecology and obstetrics, Maternity School Januário Cicco (MEJC), Federal University of Rio Grande do Norte (UFRN), Natal, RN, Brazil;

³Laboratory of Applied Molecular Biology - LAPLIC, Federal University of Rio Grande do Norte, Natal, RN, Brazil;

⁴Graduate Program in Biochemistry and Molecular Biology - PPGBqBM, Federal University of Rio Grande do Norte (UFRN), Natal, RN, Brazil.

⁵Graduate Program in Biotechnology (PPGB), Potiguar University (UnP), Anima, Natal, Brazil;

⁵Laboratory of Applied Molecular Biology - LAPLIC, Federal University of Rio Grande do Norte, Natal, RN, Brazil.;

⁶Graduate Program in Biotechnology (PPGB), Potiguar University (UnP), Anima, Natal, Brazil;

⁷Brazilian Company of Hospital Services (EBSERH), Maternity School Januário Cicco (MEJC), Federal University of Rio Grande do Norte (UFRN), Natal, RN, Brazil.;

⁸Graduate Program in Biochemistry and Molecular Biology - PPGBqBM, Federal University of Rio Grande do Norte (UFRN), Natal, RN, Brazil.;

⁹Graduate program in sciences applied to women's health (PPGCASM), Federal University of Rio Grande do Norte (UFRN), Natal, RN, Brazil.



RICARDO COBUCCI

PROTOCOL integer ID:
79605

DNA analysis for early diagnosis of cervical cancer is a promising alternative for greater treatment efficiency. Biological samples obtained from the endocervix are commonly used for diagnosis, even though such samples are still little explored in the molecular context. This work aims to describe a modified protocol for extracting HPV DNA from infected endocervical cells after collection with endocervical brushes. The Wizard® Genomic DNA Purification Kit (Promega, CAT.# A1125) was used to extract DNA from endocervical samples and the extraction protocol was adapted from the manufacturer's instructions. Forty-one samples were collected from women with a mean age of 35 ± 7.8 years. It was observed that samples containing higher blood content resulted in inhibiting the amplification signal, however, all samples had enough material for DNA extraction, despite the blood contamination. The quality of the extraction was confirmed in 100% of the samples with the amplification of the actin gene, indicating that enough DNA was extracted. Considering the absorbance ratio of 260/280, the average purification of the samples was reasonably good. The protocol was able to extract and detect HPV DNA with the potential to be used in the detection of high-risk HPV from cervical brushes with a high cost-effectiveness ratio.

COLLECTION OF SAMPLES

- 1 For the collection of endocervical material, the speculum is inserted into the vaginal canal
- 2 After viewing the cervix, the brush is introduced into the cervical canal
- 3 Then the brush is rotated 360 degrees by eight turns to the right and eight to the left in order
- 4 After collection, the brush is transferred to a clean collection tube and stored at -80°C

DNA EXTRACTION

- 5** Add 600µl of Nuclei Lysis Solution to a 1.5 ml microcentrifuge tube free of DNases, RNases and pyrogens
- 6** Add the endocervical brush to the tube containing the Nuclei Lysis Solution and homogenize the sample by rotating the brush in the solution
- 7** Remove the brush and add 3µl of RNase Solution and mix the sample by inverting the tube 5 times
- 8** Incubate the mixture for 30 minutes at 37°C. Allow the sample to cool to room temperature for 5 minutes before proceeding
- 9** To the room temperature sample, add 200µl of Protein Precipitation Solution and vortex vigorously at high speed for 20 seconds. Chill sample on ice for 5 minutes
- 10** Centrifuge at room temperature at 14,000 x g for 4 minutes
- 11** Transfer supernatant to a fresh tube containing 600µl of room temperature 99.5% isopropanol
- 12** Mix gently by inversion and centrifuge for 1 minute at 14,000 × g at room temperature
- 13** Decant the supernatant and add 600µl of room temperature 70% ethanol, rinsing the tube walls and homogenizing by inversion

- 14** Centrifuge for 1 minute at $14,000 \times g$ at room temperature and aspirate the ethanol
- 15** Invert the tube on clean absorbent paper, and air-dry the pellet for 15 minutes
- 16** Rehydrate the DNA in 100 μ l of DNA Rehydration Solution for 1 hour at 65°C. Periodically mix the solution by gently tapping the tube
- 17** Store the DNA at - 20°C