



Jun 04, 2022


# Bacterial Delivery System

contact.microbialtec<sup>1</sup><sup>1</sup>Creative Biogene

1

[dx.doi.org/10.17504/protocols.io.kqdg3pyjpl25/v1](https://dx.doi.org/10.17504/protocols.io.kqdg3pyjpl25/v1)

Microbialtec

 contact.microbialtec

Protein synthesis can be controlled with the bacterial translation machinery. Control of invasion is necessary to carry the produced proteins into cells. Invasion requires both flagella and the type III secretion system-1 (T3SS-1). Flagella are essential for cell invasion because they sense the cell surface and determine the optimal location for invasion<sup>32</sup>. T3SS-1 is a needle apparatus that initiates invasion by injecting effector proteins into cells. These proteins rearrange the actin cytoskeleton and induce endocytosis of the bacteria. Production of these two bacterial structures is controlled by the factors *fliZ* and *hilD*, which are, in turn, controlled by the master regulator *flhDC*. Protein release requires the activation of bacterial genes specifically inside cells. In *Salmonella*, this can be controlled with the promoters of SPI2 genes.

DOI

[dx.doi.org/10.17504/protocols.io.kqdg3pyjpl25/v1](https://dx.doi.org/10.17504/protocols.io.kqdg3pyjpl25/v1)<https://www.microbialtec.com/bacterial-delivery-system.html>

contact.microbialtec 2022. Bacterial Delivery System. **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.kqdg3pyjpl25/v1>



Bacterial Delivery System

 protocol ,

Jun 04, 2022

Jun 04, 2022

63868