



JAN 16, 2024

🌐 Anatomical Verification (Mendonça et al 2024)

Marcelo D
Mendonça^{1,2,3}


¹Champalimaud Research, Champalimaud Foundation, Lisbon 1400-038, Portugal;

²Champalimaud Clinical Centre, Champalimaud Foundation, Lisbon 1400-038, Portugal;

³NOVA Medical School | Faculdade de Ciências Médicas, Universidade Nova de Lisboa, Lisbon 1169-056, Portugal

OPEN  ACCESS

ASAP Collaborative Research Network



c.weber-schmidt
Allen Institute, ASAPCRN

ABSTRACT

Protocol for anatomical verification (sample prep, immunostaining and imaging) in Mendonça et al 2024.

MATERIALS

Standard perfusion setup
Vibrotome

BEFORE START INSTRUCTIONS

Animals were euthanized after completion of the behavioural tests.

DOI:
dx.doi.org/10.17504/protocols.io.14egn3o3ql5d/v1

Protocol Citation: Marcelo D Mendonça 2024. Anatomical Verification (Mendonça et al 2024). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.14egn3o3ql5d/v1>

MANUSCRIPT CITATION:
bioRx:
<https://doi.org/10.1101/2021.04.20.440527>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jan 12, 2024

Last Modified: Jan 16, 2024

PROTOCOL integer ID:
93475

Keywords: ASAPCRN

Sample Preparation

- 1 Anaesthetize mice with isoflurane, followed by intraperitoneal injection of ketamine-xylazine (5 mg/kg xylazine; 100 mg/kg ketamine).
- 2 Perfuse transcardially in 1% phosphate buffered saline (PBS) and then 4% paraformaldehyde (PFA) in PBS.
- 3 Dissect brains after perfusion is completed.
- 4 Keep brains in 4% PFA solution overnight and then transferred to 1x PBS solution.
- 5 Section brains coronally in 30 to 50-um slices (using a Leica vibratome VT1000S) and keep in PBS before mounting or immunostaining.

Immunostaining for TH (Day 1)

- 6 Rinse 3 times with TBS (10 min each)

- 7 Peroxides treatment (inhibition peroxidase and permeabilization) **5 min**
 - 1.5 mL methanol
 - 450 μ L H₂O₂
 - 13.05 mL TBS(this step is for DAB staining but we kept it because it favors the penetration of the Ab)
- 8 Rinse 3 times with TBS (10 min each)
- 9 Membrane permeabilization treatment **15min** (or 20min for Ab that don't penetrate well)
 - 300 μ L Triton 10%
 - 14.7 mL TBS
- 10 Rinse 3 times with TBS (10 min each)
- 11 Blockade with BSA 3% or other blocking solutions **1h** (if needed)
- 12 Primary Antibody (diluted in TBS): **overnight at 4°C** (or longer period if needed)
TH (1:1000) Pel-Freez, Rogers, AR

Immunostaining for TH & Mounting (Day 2)

- 13 Rinse 3 times with TBS (10 min each)
- 14 Secondary Antibody (diluted in TBS without) **40-45 min Room Temperature** (1:400)

15 Rinse 2 times with TBS (10 min each) AND Rinse 2 times with TB (10 min each)

16 Mounting with **Vectashield** or **PVA DABCO**

Imaging

17 Images were taken using a wide-field fluorescence microscope (Zeiss AxioImager).