



NOV 23, 2023

OPEN  ACCESS

**DOI:**  
[dx.doi.org/10.17504/protocols.io.dm6gp35jdvzp/v1](https://dx.doi.org/10.17504/protocols.io.dm6gp35jdvzp/v1)

**Protocol Citation:** Laura Mahoney-Sanchez 2023. iPSC-derived neurons sample preparation for ELISA assays. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.dm6gp35jdvzp/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working  
We use this protocol and it's working


**Created:** Nov 07, 2023

## iPSC-derived neurons sample preparation for ELISA assays

Laura Mahoney-Sanchez<sup>1,2</sup>

<sup>1</sup>The Francis Crick Institute, 1 Midland Road, London NW1 1AT;

<sup>2</sup>Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology, Queen Square, London, WC1N 3BG






Laura Mahoney-Sanchez

The Francis Crick Institute, 1 Midland Road, London NW1 1AT,...

### ABSTRACT

Cell lysis sample preparation for ELISA assays

## Cell lysis

- 1 Cells were washed with cold PBS then lysed with  300  $\mu$ L cold RIPA buffer (Thermo Fisher Scientific) supplemented with 1x of a cocktail of phosphatase and protease inhibitors (Life Technologies).
- 2 Cell lysates were transferred into 1.5ml eppendorf tubes and centrifuged for  00:05:00 at 3500rp  to remove cell debris.
- 3 Following centrifugation, the supernatant was collected and protein concentration were measured using a BCA or Bradford protein assay (Thermo Fisher Scientific).

## ELISA assay

- 4 Equal amounts of protein lysate,  20  $\mu$ g, were used to perform ELISA assays according to manufacturer's instructions.