


Jul 24, 2024

Optimizing the Effect of Additives on qPCR Detection

DOI

dx.doi.org/10.17504/protocols.io.j8nlk8nkxl5r/v1



Jonathan Phillips¹, Gregor Blaha¹

¹University of California, Riverside



Gregor Blaha

University of California, Riverside

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.j8nlk8nkxl5r/v1

Protocol Citation: Jonathan Phillips, Gregor Blaha 2024. Optimizing the Effect of Additives on qPCR Detection. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.j8nlk8nkxl5r/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: July 21, 2024

Last Modified: July 24, 2024

Protocol Integer ID: 103797

Keywords: qPCR, optimization, additive

Funders Acknowledgement:
California Department of Food
and Agriculture
Grant ID: 21-0001-056-SF



Abstract

Optimizing qPCR reactions has two key objectives: (1) to handle a wider range of samples, and (2) to minimize variability between replicates. [Anonymous 2009; Minas K., et al. 2011; Nolan T., et al. 2006]. Anecdotal evidence suggests that certain additives can enhance the PCR reactions and other DNA- and RNA-polymerase-based reactions, resulting in the detection of even smaller quantities of nucleic acids with higher confidence [Villalava C., et al. 2001; Rapley R., et al. 1994]. Although several mechanisms of action for these additives have been evoked, their effect on a particular assay cannot be predicted and must be determined experimentally. At the end of this protocol, we provide a table with some additives that may improve PCR-based reaction.

Attachments



04_optimizing the ef...

387KB

Image Attribution

For permission to use the image or to contact the creator of the image, irod, please contact the lead author at gregor.blaha@ucr.edu



Protocol references

Gallup J.M. (2011) Difficult Templates and Inhibitors of PCR. in “PCR Troubleshooting and Optimization: The Essential Guide” edited by Kennedy S. and Oswald N. published in Caister Academic Press, U.K.

Minas K., et al. (2011) Optimization of a high-throughput CTAB-based protocol for the extraction of qPCR-grade DNA from rumen fluid, plant and bacterial pure cultures. FEMS Microbiology Letters, 325, 162–169, PMID: 22029887. doi: 10.1111/j.1574-6968.2011.02424.x

Nolan T, et al. (2006) Quantification of mRNA using real-time RT-PCR. Nat Protoc., 1, 1559-82. PMID: 17406449. doi: 10.1038/nprot.2006.236.

Rapley, R. et al. (1994) Enhancing PCR amplification and sequencing using DNA-binding proteins. Mol Biotechnol 2, 295–298. doi: 10.1007/BF02745882

Anonymous (2009). Thermo Fisher qPCR Optimization guide
<https://tools.thermofisher.com/content/sfs/manuals/D13002~.pdf>

Villalva C., et al. (2001) Increased yield of PCR products by addition of T4 gene 32 protein to the SMART PCR cDNA synthesis system. Biotechniques, 31, 81-83. PMID 11464525. doi: 10.2144//01311st04.