


Aug 19, 2024

Crystallisation protocol for SARS-CoV-2 nsp3 macrodomain in P43

DOI

dx.doi.org/10.17504/protocols.io.e6nvw1qb2lmk/v1



Jasmin Aschenbrenner^{1,2}, Peter Marples^{1,2}, Charlie Tomlinson^{1,2}, Lizbé Koekemoer³, Daren Fearon^{1,2}

¹Diamond Light Source; ²Research Complex at Harwell; ³Centre of Medicines Discovery, University of Oxford Jasmin Aschenbrenner: The principle crystallographer on the SARS-CoV-2 nsp3 macrodomain project;

ASAP Discovery



Lizbé Koekemoer
University of Oxford

OPEN ACCESS



DOI: dx.doi.org/10.17504/protocols.io.e6nvw1qb2lmk/v1

External link: https://asapdiscovery.org/outputs/target-enabling-packages/#ASAP-SARS-COV-2-NSP3-MAC1

Protocol Citation: Jasmin Aschenbrenner, Peter Marples, Charlie Tomlinson, Lizbé Koekemoer, Daren Fearon 2024. Crystallisation protocol for SARS-CoV-2 nsp3 macrodomain in P43. **protocols.io** https://dx.doi.org/10.17504/protocols.io.e6nvw1qb2lmk/v1

License: This is an open access protocol distributed under the terms of the <u>Creative Commons Attribution License</u>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's

working

Created: August 15, 2024

Last Modified: August 19, 2024

Protocol Integer ID: 105614

Keywords: crystallisation, XChem, ASAP, AViDD, CMD, Diamond Light Source, i04-1, SARS-CoV-2, nsp3, macrodomain



Funders Acknowledgement: National Institutes of Health/National Institute Of Allergy and Infectious Diseases (NIH/NIAID) Grant ID: Grant ID: U19AI171399

Disclaimer

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Acknowledgements:

Diamond Light Source Ltd, Harwell Science and Innovation Campus, Didcot OX11 0QX, UK Research Complex at Harwell, Harwell Science and Innovation Campus, Didcot OX11 0FA, UK Oxford Lab Technologies crystal shifter https://doi.org/10.1107/S2059798320014114

Abstract

The COVID-19 pandemic has demonstrated the need for novel therapeutic interventions and improved pandemic preparedness strategies against severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). This protocol details an optimized crystallization method for the SARS-CoV-2 nsp3 macrodomain, a potential drug target. Using sitting drop vapor diffusion, we describe specific buffer conditions and procedures to consistently produce high-quality crystals suitable for XChem fragment screening. The method yields crystals that diffract to an average resolution of 1.2 Å, enabling high-resolution structural studies.

All structures solved during the development of tool compounds for the SARS-CoV-2 nsp3 macrodomain are deposited on the PDB (Group deposition: G_1002283).

Materials

SwissCl 3 lens crystallization plates https://swissci.com/product/3-lens-crystallisation-plate/ Codes: Midi: UVXPO-3LENS 3W96T-PS 3W96T-UVP

[M] 1 Molarity (M) CHES, рн 9.5, Molecular Dimensions, Catalog # MD2-008-PH 9.5 50% w/v PEG 3000, Molecular Dimensions, Catalog # MD2-100-8

Purified SARS-CoV-2 nsp3 macrodomain protein ([M] 47 mg/mL) in [M] 10 millimolar (mM) HEPES, PH 7.5 , [M] 0.5 Molarity (M) NaCl, 5% glycerol, [M] 0.5 millimolar (mM) TCEP



Safety warnings




• Follow all handling warning for the chemicals used in the crystalllisation screen composition.



SARS-CoV-2 nsp3 macrodomain expression and purification

The protein usedfor crystallisation was expressed and purified using the following protocol.



Equipment needed

2 <u>Formulatrix Rock Imager</u> (or incubator of choice)

SPT mosquito

P100 multi-channel pipette

SwissCI 3 lens plate

Crystallization experiment



3 Protein and buffer requirements:



4 **Crystallisation screen composition:**

```
[м] 100 millimolar (mM) CHES, Ср. 9.5
30% w/v PEG 3000
```

Stock solutions used:

```
[м] 1 Molarity (M) CHES adjusted to Срн 9.5 with NaOH
50% w/v PEG 3000
```



Note

The crystallisation screen can be stored in a Duran bottle or aliquoted into 96 deep well block for easy dispensing into SwissCl 3 lens plates.

For long-term storage keep the Crystallisation screen in the fridge at 4°C.

5 Dispense 🚨 20 µL Crystallisation screen into SwissCl 3 lens plate reservoir wells using a 100 µl multi-channel pipette.

Dispense 4 150 nL [M] 47 mg/mL Sample to each lens using the SPT mosquito.

Dispense 4 150 nL Crystallisation screen to each lens using the SPT mosquito.

Drop ratio: 1:1 ratio (150 nL Sample : 150 nL reservoir solution)

Final drop volume: 300 nl

6 Incubate at \$\mathbb{g}\$ 20 °C for \(\bar{\chi} \) 48:00:00 h in Formulatrix Rock Imager.

Imaging Schedule: The first images are taken after 12 h and the imaging schedule follows a Fibonacci sequence of days for further collections.

7 Crystal form after ~24 h. 2d



Expected result

The crystals reach their maximum size after 48 h.

Crystals typically form as single crystals.

Morphology: typically large cubes or rectangles.

Size: \sim 300 μm in length and \sim 150 μm in width, depth of the crystals is \sim 100 μm


Appearance: large crystal chunks.

Average resolution: 1.2 Å

Space group: P4₃

Unit cell: 89 Å, 89 Å, 39 Å

90.00°, 90.00°, 90.00°



An example of a drop containing SARS-CoV-2 nsp3 macrodomain crystals



Data Collection at Synchrotron

8 Diamond Light Source

> **Unattended Data Collection (UDC) Data Collection Temperature: 100K Detector: DECTRIS EIGER2 X 9M**

Beamline: 104-1

Wavelength: 0.9212 Å Resolution (Å): 1.62 Beam Size (µm): 60 X 50 Number of images: 3600

Oscillation: 0.10° **Exposure (s):** 0.0020 Transmission (%): 100 **Flux (ph/s):** 3.80e+12

Protocol references

Protocol based on

Marion Schuller et al., Fragment binding to the Nsp3 macrodomain of SARS-CoV-2 identified through crystallographic screening and computational docking. Sci. Adv. 7, eabf8711 (2021).

DOI: 10.1126/sciadv.abf8711