

Aug 04, 2021

© Change in treatment burden among people with multimorbidity: protocol of a follow up survey and development of efficient measurement tools for primary care.

H.O.Hounkpatin ¹, *¹, Paul Roderick¹, James E Morris¹, James E Morris¹, Scott Harris¹, Forbes Watson¹, Hajira Dambha-Miller¹, Helen Roberts¹

¹University of Southampton

H.O.Hounkpatin



Treatment burden is the effort required of patients to look after their health and the impact this has on their functioning and wellbeing. It is likely treatment burden changes over time as circumstances change for patients and health services. However, there are a lack of population-level studies of treatment burden change and factors associated with this change over time. Furthermore, there are currently no practical screening tools for treatment burden in time-pressured clinical settings or at population level.

Methods and analysis

This is a three-year follow-up of a cross-sectional survey of 700+ people with multimorbidity (defined as three or more long-term conditions; LTCs) registered at GP practices in in Dorset, England. The survey will repeat collection of information on treatment burden (using the 10-item Multimorbidity Treatment Burden Questionnaire (MTBQ) and a novel single-item screening tool), sociodemographics, medications, LTCs, health literacy and financial resource, as at baseline. Descriptive statistics will be used to compare change in treatment burden since the baseline survey in 2019 and associations of treatment burden change will be assessed using regression methods. Diagnostic test accuracy metrics will be used to evaluate the single-item treatment burden screening tool using the MTBQ as the gold-standard. Routine primary care data (including demographics, medications, LTCs, and healthcare usage data) will be extracted from medical records for consenting participants. A forward-stepwise, likelihood-ratio logistic regression model building approach will be employed in order to assess the utility of routine data metrics in quantifying treatment burden in comparison to self-reported treatment burden using the MTBQ.

Impact

To the authors' knowledge, this will be the first study investigating longitudinal aspects of treatment burden. Findings will improve understanding of the extent to which treatment burden changes over time for people with multimorbidity and factors contributing to this change, as well as allowing better identification of people at risk of high treatment burden.

DO!

dx.doi.org/10.17504/protocols.io.bw5fpg3n

PROTOCOL CITATION

H.O.Hounkpatin , *, Paul Roderick, James E Morris, James E Morris, Scott Harris, Forbes Watson, Hajira Dambha-Miller, Helen Roberts 2021. Change in treatment burden among people with multimorbidity: protocol of a follow up survey and development of efficient measurement tools for primary care..

protocols.io

https://dx.doi.org/10.17504/protocols.io.bw5fpg3n

LICENSE

•

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Aug 04, 2021

LAST MODIFIED

Aug 04, 2021

PROTOCOL INTEGER ID

52103

1